

Technical University of Denmark



60-year record of iodine-129 in Philippines terrestrial sediment revealing human nuclear activities history at low latitude area

Hou, Xiaolin; Zhang, Luyuan; Li, Hongchun

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Hou, X., Zhang, L., & Li, H. (2017). 60-year record of iodine-129 in Philippines terrestrial sediment revealing human nuclear activities history at low latitude area. Abstract from 14th International Conference on Accelerator Mass Spectrometry, Ottawa, Canada.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

60-year record of iodine-129 in Philippines terrestrial sediment revealing human nuclear activities history at low latitude area

Global distribution of iodine-129 (^{129}I) is important to apply it as an environmental tracer and for its environmental safety assessment and management [1]. Despite a number of ^{129}I data reported throughout many regions [2,3], less is known at low latitude area. A sediment core collected from the Taal Lake in the north of Philippines was analyzed for ^{129}I and ^{127}I aiming to investigate the influence of human nuclear activities on terrestrial systems of low latitude regions. Concentrations of ^{129}I in the core ranged from $(0.5\pm 0.07)\times 10^5$ atoms/g in the pre-nuclear era to $(215.92\pm 20.16)\times 10^5$ atoms/g. Compared to other marine and terrestrial cores, levels of ^{129}I in the Taal Lake sediment were as similar as that in Mississippi river (USA), but 1-3 orders of magnitude lower than the European lakes and sea sediments [2,3]. Taking into account the factors of distance from sites of human nuclear activities, climate and geological settings of Taal Lake, the increased $^{129}\text{I}/^{127}\text{I}$ ratios indicated that the origin of ^{129}I is predominated by testing of the Pacific Proving Ground, global fallout of other nuclear testing and European nuclear reprocessing plants, which were primarily transported by air. Easterly equatorial trade wind and East Asian Winter Monsoon are important significant transportation pathways of ^{129}I to the southeast Asia. The lowest $^{129}\text{I}/^{127}\text{I}$ ratio of $(2.04\pm 0.42)\times 10^{-12}$ at the bottom of the sediment suggested less migration of ^{129}I from upper to deep layers and negligible influence of nuclear tests until 1956, and also imply that incursion of seawater might occur in Taal Lake. Volcanic eruption and direct geothermal fluid activity at the Taal Lake might change the sedimentation pattern and influence the retention of iodine isotopes.

Dr. ZHANG, Luyuan (State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences); HOU, Xiaolin (Technical University of Denmark); Prof. LI, Hongchun
