

Technical University of Denmark



## Re-thinking stressor interactions: The role of groundwater contamination impacting stream ecosystems

McKnight, Ursula S.; Sonne, Anne Thobo; Rønde, Vinni Kampman; Rasmussen, Jes J.; Traunspurger, Walter; Höss, Sebastian; Bjerg, Poul Løgstrup

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
McKnight, U. S., Sonne, A. T., Rønde, V. K., Rasmussen, J. J., Traunspurger, W., Höss, S., & Bjerg, P. L. (2016). Re-thinking stressor interactions: The role of groundwater contamination impacting stream ecosystems. Abstract from Groundwater Quality Conference (GQ16), Shenzhen Kylin Villa , China.

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Re-thinking stressor interactions: The role of groundwater contamination impacting stream ecosystems

Ursula S. McKnight<sup>1</sup>, Anne T. Sonne<sup>1</sup>, Vinni Rønde<sup>1</sup>, Jes J. Rasmussen<sup>2</sup>, Walter Traunspurger<sup>3</sup>, Sebastian Höss<sup>4</sup>, Poul L. Bjerg<sup>1</sup>

<sup>1</sup> Department of Environmental Engineering, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

<sup>2</sup> Department of Bioscience, Aarhus University, Vejlssøvej 25, 8600 Silkeborg, Denmark

<sup>3</sup> University of Bielefeld, Department of Animal Ecology, Konsequenz 45, 33615 Bielefeld, Germany

<sup>4</sup> Ecosa, Giselastr. 6, Starnberg, Germany

Keywords: groundwater contamination; stream ecosystems; stressors

Streams are significantly impacted by a large number of micropollutants and their transformation products (e.g. pharmaceuticals, chlorinated solvents, pesticides/biocides, nutrients, heavy metals), leading to a severe degradation of essential ecological functions and services. The large diversity of compounds and their sources makes the quantification of their occurrence and toxicity a challenging task. Conventionally, chemical impacts to ecosystems are assessed one compound/group of compounds or one source type at a time. We presume this will give us a good indication of the impact of a particular stressor. Our hypothesis, however, is that this will underestimate the combined impact caused by chemical cocktail effects, and interactions between stressors (e.g. contaminant-effected change in redox conditions releasing heavy metals). Moreover, a stream system impacted by multiple stressors has a high chronic stress level, so even small perturbations on top of changes in water flow or additional chemical stressors may be detrimental to the stream health.

To address this issue, we identified contaminant sources and chemical stressors along a 16-km groundwater-fed stream stretch (Grindsted, Denmark) to quantify the contaminant discharges, and potentially link the chemical impact and stream water quality. Potential pollution sources include two contaminated sites (Grindsted factory/landfill), aquaculture, wastewater discharges, and diffuse sources from agriculture and urban areas. Data for xenobiotic organic groundwater contaminants, pesticides, heavy metals, general water chemistry, physical conditions and stream flow from three campaigns in 2012 and 2014 were assessed. The measured chemical concentrations were converted to toxic units (TU) based on 48-h acute toxicity tests with *Daphnia magna*. Ecological status was determined by monitoring meiobenthic (i.e. nematodes) and macrobenthic invertebrate communities.

The results show a substantial impact on Grindsted stream from multiple sources of many origins. The groundwater plume from the Grindsted factory site caused elevated concentrations of chlorinated ethenes, benzene and site-specific pharmaceuticals in both the hyporheic zone and the stream water for several km downstream of the discharge area. Heavy metal concentrations were found around or above the threshold values for barium, copper, lead, nickel and zinc in the stream water, hyporheic zone and streambed sediment. The calculated TU was generally similar along the stream, but for arsenic and nickel higher values were observed where the groundwater plume discharges into the stream. Also, log sum TU<sub>*D.magna*</sub> values for organic contaminants were elevated in both the hyporheic zone and stream. Thus, the overall chemical stress in the main discharge area is much higher than upstream, while it gradually decreases downstream.

The ecological results indicate a change in community composition for both meio- and macrobenthic fauna, pointing towards the presence of a local impact resulting from the discharging contaminated groundwater. Ecological impacts could be related to xenobiotic compounds coming from groundwater, as well as the presence of trace metals of diffuse and/or biogenic origin in the same area, but linkage to one specific factor was not possible. This work highlights the importance of a holistic assessment of stream water quality to identify and quantify the main contaminant sources and resulting chemical stream stressors leading to potential ecological impacts.