

Technical University of Denmark



## Novel method reveals a narrow phylogenetic distribution of bacterial dispersers in environmental communities exposed to low hydration conditions

Krüger, U. S.; Bak, F.; Aamand, J.; Nybroe, O.; Badawi, N.; Smets, Barth F.; Dechesne, Arnaud

*Published in:*  
Applied and Environmental Microbiology

*Link to article, DOI:*  
[10.1128/AEM.02857-17](https://doi.org/10.1128/AEM.02857-17)

*Publication date:*  
2018

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Krüger, U. S., Bak, F., Aamand, J., Nybroe, O., Badawi, N., Smets, B. F., & Dechesne, A. (2018). Novel method reveals a narrow phylogenetic distribution of bacterial dispersers in environmental communities exposed to low hydration conditions. *Applied and Environmental Microbiology*. DOI: 10.1128/AEM.02857-17

**DTU Library**  
Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Novel method reveals a narrow phylogenetic distribution of bacterial**  
2 **dispersers in environmental communities exposed to low hydration**  
3 **conditions**

4 U. S. Krüger<sup>1,2</sup>, F. Bak<sup>1,2</sup>, J. Aamand<sup>1</sup>, O. Nybroe<sup>2</sup>, N. Badawi<sup>1</sup>, B. F. Smets<sup>3</sup>, A. Dechesne<sup>3</sup>

5 <sup>1</sup>Geological Survey of Denmark and Greenland, Copenhagen, Denmark

6 <sup>2</sup>University of Copenhagen, Department of Plant and Environmental Sciences, Copenhagen,  
7 Denmark

8 <sup>3</sup>Technical University of Denmark, Department of Environmental Engineering, Lyngby, Denmark

9 Corresponding author: Arnaud Dechesne: arde@env.dtu.dk, Urse Scheel Krüger: usk@geus.dk

10

11

12

13

14

15

16

17

18 **ABSTRACT**

19 In this study, we developed a method that provides community-level surface dispersal profiles  
20 under controlled hydration conditions from environmental samples and enables us to isolate and  
21 uncover the diversity of the fastest bacterial dispersers. The method expands on the Porous  
22 Surface Model (PSM), previously used to monitor dispersal of individual bacterial strains in liquid  
23 films at the surface of a porous ceramic disc. The novel procedure targets complex communities  
24 and captures the dispersed bacteria on a solid medium for growth and detection. The method was  
25 first validated by distinguishing motile *Pseudomonas putida* and *Flavobacterium johnsoniae* strains  
26 from their non-motile mutants. Applying the method to soil and lake water bacterial communities  
27 showed that community-scale dispersal declined as conditions became drier. However, for both  
28 communities, dispersal was detected even under low hydration conditions (matric potential: -3.1  
29 kPa), previously proven too dry for *P. putida* KT2440 motility. We were then able to specifically  
30 recover and characterize the fastest dispersers from the inoculated communities. For both soil and  
31 lake samples, 16S rRNA gene amplicon sequencing revealed that the fastest dispersers were  
32 substantially less diverse than the total communities. The dispersing fraction of the soil microbial  
33 community was dominated by *Pseudomonas* which increased in abundance at low hydration  
34 conditions, while the dispersing fraction of the lake community was dominated by *Aeromonas*  
35 and, under wet conditions (-0.5 kPa), also by *Exiguobacterium*. The results gained in this study  
36 bring us a step closer to assessing the dispersal ability within complex communities under  
37 environmentally relevant conditions.

38

39

40 **IMPORTANCE**

41 Dispersal is a key process of bacterial community assembly. Yet, very few attempts have been  
42 made at assessing bacterial dispersal at the community level as focus has previously been on pure  
43 culture studies. A crucial factor for dispersal in habitats where hydration conditions vary, such as  
44 soils, is the thickness of the liquid films surrounding solid surfaces, but little is known on how the  
45 ability to disperse in such films varies within bacterial communities. Therefore, we developed a  
46 method to profile community dispersal and identify fast dispersers on a rough surface resembling  
47 soil surfaces. Our results suggest that within the motile fraction of a bacterial community only a  
48 minority of the bacterial types are able to disperse in the thinnest liquid films. During dry periods,  
49 these efficient dispersers can gain a significant fitness advantage through their ability to colonize  
50 new habitats ahead of the rest of the community.

51

52 **KEYWORDS**

53 Community motility, *Pseudomonas putida* KT2440, liquid film, soil, lake water, succession, porous  
54 surface model.

55

56 **Introduction:**

57 Dispersal is essential in order to ensure fitness in a world of limited and heterogeneously  
58 distributed resources and is recognized as a key contributor to community dynamics (1, 2). While  
59 dispersal has long been studied as an integral part of the ecology of animals and plants, its  
60 contribution to microbial ecology has received less attention (3–5).

61

62 Dispersal is traditionally divided into passive (caused e.g. by weather or human activities) and  
63 active, also termed motility, which requires metabolic energy (6). Motility is not limited to  
64 environments saturated with water, but is also commonly found on or near surfaces in  
65 unsaturated environments, such as the thin liquid films between soil particles or on the surface of  
66 leaves (7). Bacteria have evolved diverse mechanisms of active dispersal on surfaces including  
67 swimming, swarming, twitching, sliding, and gliding, all of which have been mainly described and  
68 studied in pure culture settings (8–11) using agar plates or glass slides for capture of motile cells  
69 (8, 9, 11, 12). Hence, it remains unclear how well these methods capture dispersal potential in  
70 more natural settings, such as in soils, and how the ability to disperse is distributed within the tree  
71 of life and within individual communities. There have been a few efforts to uncover the  
72 phylogenetic distribution of flagellar motility (13, 14) and gliding motility (12, 15), but a  
73 comprehensive view of how the dispersal abilities vary across and within bacterial phyla is still  
74 lacking.

75  
76 This gap in our knowledge partly results from the lack of methods for mass assessment of  
77 dispersal potential of bacteria in environmental samples. In a community, not all bacteria have  
78 equal potential for dispersal; though this is rarely assessed. To assess the dispersal potential of a  
79 community one could, in theory, isolate and test all its members but considering that there can be  
80 up to  $10^9$  bacterial cells in a gram of productive soil (16), this would be practically unfeasible. In  
81 addition, by studying strains in isolation, the effects of interaction between strains would be  
82 missed. Indeed, most past studies of motility have focused on the motion of single strains (8, 17,  
83 18) largely neglecting the vast possibilities of interactions which have only recently been  
84 uncovered (17). Bacterial co-cultures have been observed swarming together, combining their

85 skills to conquer barriers such as antibiotics (19) or to engage in metabolic cross-feeding (20).  
86 Motile bacteria have been demonstrated to carry non-motile bacteria as cargo (21) and inter  
87 kingdom cooperation has been described such as bacterial dispersal with the help of fungi and  
88 amoeba (22–26). It would seem logical that these complex interactions occur in natural  
89 communities but only very few attempts have been made to tackle motile bacteria in  
90 environmental samples at the community level (27–31).

91 A few studies did address community-level motility in aquatic environments. Grossart *et al.* (29),  
92 Mitchell *et al.* (31) and Fenchel *et al.* (30), assessed swimming motility in ocean samples using  
93 microscopy and revealed a large percentage of motile bacteria, but did not identify these. Dennis  
94 *et al.* (27) used a syringe based-assay and 16S rRNA gene amplicon sequencing to uncover the  
95 identity of motile lake water bacteria showing a chemotactic response towards inorganic  
96 substrates. However, to our knowledge, only one study assessed dispersal and identity of  
97 dispersing bacteria in a complex natural community under conditions relevant for partially-water-  
98 saturated habitats (e.g. surface or vadose zone soils, phyllosphere) (28). Using sand microcosms,  
99 Wolf *et al.* revealed that a subset of a soil community consisting mainly of the family  
100 *Enterobacteriaceae* and the genera *Undibacterium*, *Pseudomonas* and *Massilia* were able to  
101 expand to a distance of more than 2 cm from the inoculation point within 48 h (28). While this  
102 study provided important insights into the identity of dispersers and their expansion rate, they  
103 only considered one hydration condition (7.5% moisture w/w, i.e., matric potential in the -20 to -  
104 50 kPa range based on the particle size (32)). Yet, previous studies have stated that water is one of  
105 the primary factors controlling bacterial motility (33).

106 In a non-permanently water saturated habitat such as soil, the ability to disperse is primarily  
107 dependent on the thickness of the water film surrounding solid surfaces. The hydration status of

108 soil in the vadose zone is highly variable and can increase or decrease rapidly, e.g. following  
109 rainfalls or droughts (7, 18, 33). At low matric potential, the thinning of the aqueous films between  
110 soil particles will lead to habitat fragmentation into separate micro habitats (34), with strong  
111 effect on the bacterial dispersal ability.

112 The Porous Surface Model (PSM) is a 2D model system used for studying bacterial motility on the  
113 surface of a porous ceramic disc under controlled hydration conditions that mimics unsaturated  
114 soil surfaces. Studies of fluorescently tagged pure cultures using the PSM have demonstrated  
115 that flagellar motility is restricted to a relatively narrow range of water potential (0 to -2kPa) (18,  
116 34). However, it remains unclear how this knowledge on specific flagellated bacterial isolates can  
117 be translated at the scale of complex environmental communities.

118

119 Hence, the aims of the current study were to (i) further develop the PSM for its use to assess  
120 bacterial dispersal of natural (untagged) bacterial communities and (ii) apply the method to a soil  
121 and a lake community to obtain community-level surface motility profiles under controlled  
122 hydration conditions and uncover the diversity of the fastest dispersers.

123

124

## 125 **Results**

### 126 **Developing and validating the novel method with pure cultures.**

127 The Porous Surface Model (PSM) has previously been used to monitor bacterial dispersal under  
128 controlled hydration conditions (35). The challenge of the current study was to expand the  
129 method from being solely usable with fluorescently tagged cultures to evaluate dispersal of a  
130 broader range of complex natural communities, tracking the movement from the inoculation point  
131 at the center towards the edge of the ceramic disc. To achieve this goal, we devised a procedure

132 to get an imprint of dispersal on the ceramic disc by pressing agar plates onto the PSM surface  
133 (see method overview in the supporting information, Fig. S1). This step resembles the agar lift  
134 method used for visualization of the bacterial distribution on soil surfaces described in a previous  
135 study (36).

136

137 Initial tests with *P. putida* KT2440 GFP and full agar plates pressed onto the ceramic disc of the  
138 PSM showed a clear discrepancy between the bacterial spatial distribution observed on the  
139 ceramic disc with epifluorescence microscopy before pressing and that captured on the agar plate.  
140 It was clear that pressing the agar plate onto the surface of the ceramic disc disturbed the  
141 bacterial distribution, so that the high density of cells at the inoculation point were often spread  
142 over a much larger area. To avoid this error, we developed a series of concentric annular  
143 hollowed-out agar plates, which were pressed sequentially, starting from the edge of the ceramic  
144 disc and leaving the center undisturbed until a full plate was used to capture the total community  
145 that had developed on the PSM disc (Fig. 1 and S1).

146

147 To test the method, we inoculated a mixture of the motile strain *P. putida* KT2440 GFP and its non-  
148 flagellated mutant *P. putida* K2440 dsRed *fliM*<sup>-</sup> which had previously been used for motility studies  
149 on the PSM (18, 35). This pure culture experiment demonstrated the ability of the method to  
150 clearly contrast the dispersal potential of these strains (Fig. 1 and Fig. 2). The non-motile strain  
151 generally stayed near the inoculation point. For the motile strain, the fastest dispersal was seen at  
152 -0.5 kPa with 4 out of 6 replicates reaching the edge of the pressed plate furthest from the  
153 inoculation point, i.e. the 25 - 41.3 mm section, after 40 hours incubation (Fig. 2). Dispersal at -1.2  
154 kPa was not significantly different from that at -0.5 kPa, after 24 h ( $p=0.229$ ) or 40h ( $p=0.857$ ),



155 and no dispersal was registered at -3.1 kPa. This was consistent with the threshold for flagellar  
156 motility of -2.0 kPa previously reported for *P. putida* KT2440 (18).

157

158 To test the ability of the method for capturing other types of motility than flagellum powered  
159 swimming we inoculated the gliding bacteria *Flavobacterium johnsoniae* strain CJ1827 (37) and  
160 the non-motile mutant *F. johnsoniae* strain 2122  $\Delta$ *gldK* (38) on separate PSMs incubated at -0.5  
161 kPa. After 48h incubation the non-motile mutant (n=2) stayed near the inoculation point while the  
162 gliding bacteria (n=3) were recovered in the 11.5-15 mm section and, in one case, at the edge of  
163 the pressed plate, 25-41.3 mm section.

164

#### 165 **Dispersal potential of environmental communities.**

166 Applying the novel method on extracted soil and lake bacterial communities confirmed that  
167 dispersal rate declined as conditions became drier (Fig. 3). However, surprisingly, for both  
168 community types, dispersal was detected even under the lowest hydration condition tested (-3.1  
169 kPa and one sample at -4.2 kPa, not shown), previously proven too dry for *P. putida* KT2440  
170 dispersal (Fig. 2). For the soil community, the slowest dispersal was detected at -3.1 kPa with 2  
171 out of 3 replicates dispersed to the 11.5-15mm section at 24 h, though all reached the most  
172 distant section of the plate (25 - 41.3 mm section) after 48 h (Fig. 3). To record the magnitude of  
173 the colonization of the ceramic plate sections a coverage score was introduced for the  
174 environmental samples, where the extent of colony coverage of the agar plates was roughly  
175 assigned into four categories. This scoring indicated that even though soil bacteria reached  
176 the most distant sections of the plates after 48 h, the colonization was less at -3.1 kPa (1-25% to  
177 26-50% coverage) than at -1.2 kPa and -0.5 kPa (51-76% to 76-100% coverage).

178 The lake bacterial community study was mainly included for comparison purposes, as we assumed  
179 that this community would have experienced a weaker selection for dispersal ability on dry  
180 surfaces, and to demonstrate the versatility of the method. It was based on fewer replicates which  
181 limits interpretation, but revealed a similar picture with bacteria reaching the edge of the plate  
182 after 48 h for both dry (-3.1 kPa) and wet (-0.5 kPa) conditions though the extent of colonization  
183 was less for the dry samples (Fig. 3). Direct comparisons of the soil and lake data should be done  
184 with caution because the CFU counts suggested that more cultivable cells were inoculated for the  
185 lake than for the soil samples ( $34 \times 10^3$  vs  $2\text{-}8 \times 10^3$  CFUs per inoculum, respectively). However, for  
186 both soil and lake communities it remains clear that we registered much faster dispersal at both -  
187 0.5 kPa and -3.1 kPa (Fig. 3) compared to *P. putida* KT2440 (Fig. 1).

188

#### 189 **Diversity of dispersers.**

190 DNA was extracted from the Nycodenz soil extracts and the lake filtrate used for inoculations,  
191 from agar plates reflecting the total community present on the ceramic disc (Full Plate), and from  
192 the community that developed upon inoculation of the environmental cell extracts onto a  
193 'standard' 25% R2A solid medium plate with 20 g agar l<sup>-1</sup>, which provides conditions that are not  
194 conducive to motility (39) (the No Motility Reference Plate, shortened as 'Reference Plate'). All  
195 samples were sequenced using Illumina sequencing targeting the V3-V4 regions of the 16S rRNA  
196 gene. A total of 3.8 million sequences were kept after filtering for further analysis.

197 Comparisons between the communities in the inoculum (Nycodenz extractions and the Lake  
198 filtrate) and the cultivable communities dispersed or not, confirm the expected cultivation bias  
199 (Fig S7-8). However, the cultivable community represented on the Full plates and Reference plates  
200 retained a high diversity with representatives of 261 unique genera for soil, and 143 for lake. In

201 addition, 4172 and 665 amplicon sequence variants (ASVs (40)) in soil and lake samples were not  
202 identifiable at the genus level. Moreover, the dominating genera in the cultivated soil  
203 communities, (*Pseudomonas*, *Flavobacterium* and *Paenibacillus*) were also among the abundant  
204 taxa of the Nycodenz extractions (Fig S7) and lake water filtrate (*Aeromonas Flavobacterium* and  
205 *Exiguobacterium*) (Fig S8), respectively.

206

207 For each PSM we collected the DNA of the fastest dispersers i.e. that of the colonies of the pressed  
208 agar plate the furthest from the point of inoculation that presented growth. As the method did not  
209 allow for selective recovery of the cells unable to disperse we compared these ‘dispersed’  
210 communities to the total community present on the Full Plate and the Reference Plate.

211 Sequencing results from the soil community showed a dominance of *Pseudomonas* in the  
212 dispersed communities benefitting from increasingly dry conditions, and achieving almost total  
213 dominance at the driest conditions (-3.1 kPa and -4.2 kPa: 99.4 - 98.3% after 48 h) (Fig. 4). Under  
214 wet conditions (0.0 kPa and -0.5 kPa) the dispersed bacterial community consisted, besides  
215 *Pseudomonas*, mainly of *Paenibacillus*, *Rahnella*, *Lysinibacillus* and, after 48 h (-0.5kPa), also of  
216 *Flavobacterium* and *Janthinobacterium*. At moderate dryness (-1.2 kPa), *Bacillus* was almost equal  
217 in abundance to *Pseudomonas* (47.2% and 50.4% respectively) at 24 h but were reduced over time  
218 to 3.7% at 48 h in favor of *Pseudomonas* (50.8%), *Paenibacillus* (19.1%) and to some extent  
219 *Janthinobacterium* (11.7%).

220

221 For the lake community, *Aeromonas* was the most abundant genus in all samples and almost  
222 completely dominated the dispersed community under dry conditions (91.9% after 24 h and 95.6%  
223 after 48 h, -3.1 kPa) (Fig. 5). Under wet conditions *Aeromonas* dispersed and colonized fast (79.1%,

224 after 24 h) but appeared to experience increased competition from *Exiguobacterium* over time  
225 (43.3% and 37.5% respectively after 48 h).

226

227 The bacterial diversity, calculated using the Shannon Diversity index (Fig. 6 A, B and S3 A, B),  
228 revealed that diversity was significantly affected by the matric potential in the 24 h soil samples  
229 ( $p < 0.001$ , ANOVA based on comparison of the differences between Full Plates and dispersed  
230 communities). Both the dry (-3.1 kPa) and moderately dry (-1.2 kPa) conditions were significantly  
231 different from the wet (-0.5 kPa) and very wet (0.0 kPa) conditions ( $p < 0.05$  for all pairwise  
232 comparisons). The two dry (-3.1 and -1.2 kPa) and two wet (0.0 and -0.5 kPa) soil communities did  
233 not significantly differ from each other (Fig S3 B). At 48 h, the differences between the Shannon  
234 Diversity indices at the four matric conditions were not significant ( $p = 0.121$ ) in spite of a clear  
235 trend for decreasing diversity with drier conditions (Fig. 6 B). The difference in diversity between  
236 the dispersed community and that recovered on the Full plate could only be rigorously tested at  
237 48 h, due to the significant effect of matric potential at 24 h, but showed that the dispersed soil  
238 community had a significantly lower diversity ( $p = 0.001$ ) (Fig. 6 B).

239 The lake data also indicated a trend for lower diversity in the dispersed community compared  
240 to the Full Plate at both 24 h and 48 h (Fig. 6 A and S3 A). As expected, the Shannon diversity  
241 values of the Nycodenz extract and the Lake filtrate were much higher than for the samples  
242 collected after cultivation on the agar plates ( $5.57 \pm 1.45$  S.D (n=4) and 3.82 (n=1), respectively)  
243 clearly indicating cultivation bias.

244 The phylogenetic diversity calculated using Faith's Phylogenetic Diversity index (Fig. 6 C, D and  
245 S3 C, D) revealed a consistently narrow phylogenetic diversity of the fastest dispersers at all tested

246 matric potentials, compared to the total community of the Full Plate and the motility restricted  
247 Reference Plate for both the soil and lake community.

248

249 A closer look at the phylogenetic distribution of the two dominant genera *Pseudomonas* and  
250 *Aeromonas* in soil and lake water, respectively, showed that the dispersed communities at 48 h  
251 consisted of multiple and diverse amplicon sequence variants (ASVs) (Fig. S4, Table S1 and Fig. S5,  
252 Table S2). Notably, a search of the literature uncovered that all the type strains with the closest  
253 sequence similarity to our ASVs possess the ability for active motility mainly by using flagella,  
254 except for one for which motility is unknown (table S1 and S2). Neither for the soil community nor  
255 for the lake water community was there a clear separation of ASVs between matric conditions  
256 visible in the phylogenetic trees (Fig. S4 and S5).

257

258 A comparison of the *Pseudomonas* present in the total community of the Full Plate and the  
259 dispersed soil community at 48 h shows that only 11 out of 44 ASVs were solely present in the  
260 total community, and thus did not disperse from the center of the ceramic disc (Fig S6). This  
261 supports the general notion of *Pseudomonads* as efficient dispersers. Furthermore, 9 out of 44  
262 ASVs were solely detected in the dispersed community. This is most likely because they were  
263 below detection limit in the total community, as strains present in the dispersed community must  
264 also be present in the total community. Other evidence of large enrichments in the dispersed  
265 community can be found in the heatmap (Fig. 4), where, in addition to *Pseudomonas*,  
266 *Paenibacillus* (at matric potential below -3.1) and *Bacillus* (at -1.2 kPa) also notably increased their  
267 abundance in the dispersed community compared to the total community of the Full Plate and the  
268 motility restricted Reference Plate. These results illustrate that there can be a large fitness gain

269 associated with dispersal for a motile strain i.e. going from being below detection limit to  
270 potentially very high relative abundance far from the inoculation point. By moving ahead of the  
271 pack, such strains benefit from decreased competition for nutrients and maximize their growth.

272

## 273 **Discussion**

### 274 *Performance and limitations of the method*

275 In this study, we developed a method for assessing dispersal of natural bacterial communities  
276 under controlled hydration conditions. We achieved this by expanding on the Porous Surface  
277 Model already well established for single strain motility studies (18, 35, 39), and using agar plates  
278 to get an imprint of the colonization on the surface of the ceramic disc. The method proved  
279 effective in separating the dispersal of a motile flagellated *P. putida* strain from a non-motile  
280 mutant, which stayed near the inoculation point in the center of the ceramic disc on the agar plate  
281 imprints (Fig. 1 and 2). It was also able to capture the effect of lowered matric potential, which  
282 resulted in a reduced dispersal rate of the motile strain and a cessation of all movement at -3.1  
283 kPa, in agreement with previous studies (18, 34). In addition, the method was able to detect  
284 dispersal of the gliding bacterium *F. johnsoniae*, indicating the potential for detection of other  
285 types of motility than swimming.

286

287 The possibilities for precise control of hydration conditions are one of the key points that separate  
288 this method from the few previous studies on community motility (27–29). Calculations coupled  
289 with recent measurements of the liquid film thickness on the surface of the ceramic disc in the  
290 PSM model (18, 34, 35) provides us with a unique platform to study the behavior of microbial  
291 communities on surfaces as they are affected in their microhabitats by water film thickness. While

292 we only tested the effect of fixed hydration conditions in this study, exploring dynamic conditions  
293 such as dry-wet cycles would be straightforward. Indeed, recent studies with a synthetic soil  
294 community on the PSM demonstrated a clear effect of such cycles on competition and co-  
295 existence (41).

296

297 We recognize that the results from using this method are biased by cultivation and are only valid  
298 for the fraction of bacteria able to grow under the selected growth conditions. However, we did  
299 find a high diversity of genera among the cultured community and that the most dominant genera  
300 e.g. *Pseudomonas* and *Aeromonas* were prominent parts of the original inocula. This indicates  
301 that, in spite of the existence of some cultivation bias, our method do provide information of  
302 relevance to the original communities.

303

304 In addition, nutrient supplementation is often necessary to detect dispersal (28) and one of the  
305 strengths of this setup is that it does allow for easy isolation of strains of interest as we essentially  
306 already have them on agar plates. This has led to a culture collection of soil isolates able to  
307 disperse at -0.5 and -3.1 kPa for use in future studies (data not shown). A possible venue to  
308 decrease cultivation bias is to optimize the medium. We currently use a medium with a relatively  
309 low substrate concentration (25% R2A and R2B) to avoid selection of only fast growing bacteria,  
310 but this could be further improved by e.g. using a soil extract medium (42). Results are also likely  
311 affected by the extraction methods used to obtain microbial inocula from the environment,  
312 because extraction, and especially Nycodenz extraction (43, 44), affects the composition of the  
313 inoculum. However, this is not a limit to the method itself. The method would also be applicable  
314 when using intact environmental samples (for example, soil aggregates) placed in the center of the

315 ceramic disc. Finally, while there are benefits of using the agar plate sampling method (low  
316 detection limit for cultivable bacteria), a possible improvement would be to recover the dispersed  
317 community for DNA extraction directly from the surface of the PSM. The recovery rate and  
318 detection limit would need to be evaluated carefully.

319

#### 320 *Dispersal of environmental communities under low hydration conditions*

321 When we applied the method to soil and lake water communities, the results extended previous  
322 pure culture studies in confirming that dispersal rates decline as conditions become drier.  
323 However, surprisingly, for both communities, relatively rapid dispersal was detected even under  
324 the lowest hydration conditions (-3.1 kPa). After 48 h, members of both communities had reached  
325 the maximum possible distance of 25-41.3 mm (Fig. 3). It is unlikely that the detection of cells  
326 several cm away from the inoculation point could have been caused by simple colonial growth (i.e.  
327 cell division and shoving) because colony expansion by growth only is very slow (e.g. diameter  
328 expansion rate of  $17 \mu\text{m h}^{-1}$  for a *P. putida* KT2440 at -3.6 kPa) (35). Therefore, this dispersal is  
329 likely facilitated by motility. This strong dispersal potential under low hydration conditions was  
330 particularly surprising for the lake water community because the selective value of such traits in  
331 the original habitat is not obvious.

332

333 A possible explanation for the rapid dispersal at conditions previously thought to be too dry could  
334 be a difference in cell size between the model strains and the bacteria in the environmental  
335 samples. As discussed in Dechesne *et al.* (18) the effective thickness of the liquid-film on the  
336 surface of the ceramic disc is the limiting factor for flagellar motility. Pure culture studies on the  
337 PSM using the motile strains *Pseudomonas protegens* CHAO and *Pseudomonas putida* KT2440



338 report a threshold for swimming and dispersal at -2.0 kPa (18, 34, 35). At -2.0 kPa the predicted  
339 effective liquid film thickness on the surface of the ceramic is less than 1.5  $\mu\text{m}$  and decreases to  
340 approximately 0.4  $\mu\text{m}$  at -3.6 kPa, close to the shorter dimension of *P. putida* KT2440 rods  
341 (measured by others as 0.74  $\mu\text{m}$  (rod shaped) (45) and as 0.6  $\mu\text{m}$  by us under nutrient rich  
342 conditions). Hence, motility becomes strongly limited in liquid films thinner than the cell diameter,  
343 due to exposing the cell surface to liquid-air interfaces, capillary pressure and pinning forces (18,  
344 35). As many bacteria from soil and aquatic environments are small, with diameters less than 0.4  
345  $\mu\text{m}$  and some even passing through 0.2  $\mu\text{m}$  filters (44, 46–48), it is possible for some of them to be  
346 able to actively disperse in the thinnest liquid films tested in this study. It should be noted,  
347 however, that as bacterial cell size can vary with the conditions, e.g. Pseudomonads have been  
348 known to change both size and shape as a response to starvation or other chemical stressors (45,  
349 49), the size of the bacteria used in this study should be measured under the actual imposed  
350 conditions to confirm this theory.

351

### 352 *Diversity of efficient dispersers*

353 Our results show that diversity decreased in the dispersed communities compared to the total  
354 community in the soil and lake samples. This indicates that, within natural communities, there is a  
355 less diverse sub community of bacteria with the potential for dispersal, which will most likely have  
356 important consequences for community composition, competition and microbial succession. The  
357 study by Wolf *et al.* (28), which is most comparable to ours, identified the most abundant  
358 dispersers in their soil community as members of the genera *Enterobacteriaceae*, *Pseudomonas*,  
359 *Massilia* and *Undibacterium*, with *Enterobacteriaceae* as the most dominant. Here, we also find  
360 both *Pseudomonas* and *Enterobacteriaceae* within the 20 most abundant ASVs in the dispersed

361 soil community and *Pseudomonas* is the most dominant disperser. We also find *Paenibacillus* and  
362 *Cupriavidus*, that Wolf *et al.* (28) detected in low abundance, but along with *Enterobacteriaceae*,  
363 they are only present in our study at relatively wet conditions, with matric potentials of -1.2 kPa or  
364 lower. *Undibacterium* was not present in either the initial community or in the dispersed, and  
365 *Massilia* was only detected in low numbers on one Reference Plate and in one of the Nycodenz  
366 extractions. The differences in the abundance and composition of communities between the two  
367 studies are most likely caused by a combination of various factors such as different initial  
368 communities in the inoculum, medium selection, and variation in hydration conditions.  
369 Nonetheless, it remains clear that *Pseudomonads* play a key role in the two soil communities as  
370 early colonizers of unoccupied habitats and possibly gaining a further advantage at relatively low  
371 hydration conditions where they dominate the community.

372

### 373 *Potential modes of dispersal*

374 Under dry conditions *Pseudomonas* and *Aeromonas* dominated the dispersed soil and lake  
375 communities (Fig. 4 and 5). Many members of these two genera produce biosurfactants which  
376 have been shown to facilitate dispersal on surfaces (50–53) such as leaves, an ability which has  
377 been hypothesized to increase fitness for *Pseudomonas* (54, 55). We speculate that biosurfactants  
378 also play a role in increasing the connectedness in the liquid film on surfaces. An important factor  
379 in our model system is the residual roughness of the ceramic surface, although polished, it can  
380 result in the fragmentation of the aqueous habitat as matric potential decreases along with the  
381 liquid film thickness, and the topography of the surface, as a result, becomes more apparent.  
382 Tecon *et al.* reports that a rapid decrease in connectedness of the aqueous habitat was found at -  
383 2.0 to -5.0 kPa which influenced the motility of their tested flagellated bacteria (34). Hence,

384 biosurfactant production could be a strategy to overcome dispersal limitation under dry conditions  
385 for the two genera observed in our study. In addition members of the orders *Exiguobacterales*  
386 (*Exiguobacterium*) and *Bacillales* (*Bacillus* and *Paenibacillus*), which are frequent in the dispersed  
387 lake and soil community at wet conditions (Fig. 4 and 5) have also been found to produce  
388 biosurfactants giving rise to speculation that the benefit of surfactant production for increased  
389 dispersal ability might not only be limited to dry conditions (53). While we did not look for  
390 biosurfactant production in this study, it would be straightforward to screen the obtained isolates  
391 for biosurfactant production in the future (56).

392

393 Alternative modes of surface motility apart from flagella powered swimming might play an  
394 increased role as conditions become dryer (7). Therefore, one of the strengths of the PSM for  
395 complex community studies is that it is not limited to investigate bacteria with swimming ability,  
396 as in the previous work by Grossart *et al.* (29) and Dennis *et al.* (27), but also enables studies of  
397 other modes such as sliding, gliding, biosurfactant aided movement, fungal highways (23), or even  
398 expansion by filamentous growth (32). The PSM could thus be instrumental to establish which of  
399 these modes of motility are relevant on rough unsaturated surfaces.

400 The pure culture experiment with *F. johnsoniae* CJ1827 confirmed that gliding is possible, and can  
401 provide a detectable dispersal advantage, on the rough surface of the PSM. In the soil community  
402 experiments, *Flavobacterium* was detected in low abundance in the dispersed communities at -0.5  
403 to -4.2 kPa (Fig. 4). Many members of this genus have been found to possess gliding motility (12,  
404 57, 58), while flagellar motility in the family *Flavobacteriaceae* is almost unheard of (59), and  
405 recent isolates of the order *Flavobacteriales* from leaf surfaces have also been reported as  
406 biosurfactant producers (53). While the role of chemotaxis was not directly measured, it is

407 possible that chemotactic organisms are enriched at the rim of the ceramic plate where the  
408 substrate concentration is highest thanks to the low cell density. As many pseudomonads are  
409 known to possess chemosensory systems (60), this might contribute to their prevalence in the  
410 dispersed communities.

411

412 In theory, not all the strains we observe in our dispersed community have to possess the ability for  
413 active motility themselves, they might be non- motile strains hitching a ride with their flagellated  
414 or gliding companions (21). The co-dispersal of multiple species unveils a much more complex  
415 picture of interactions that could be addressed by future studies employing the current PSM  
416 model system. A possible next step could be to test the isolates obtained in this study to establish  
417 which are able to autonomously disperse, versus those that rely on others.

418

#### 419 **Conclusion:**

420 A novel method to study motility at the community level was developed and tested on a soil and a  
421 lake microbial community. The results obtained suggest that within the motile fraction of a  
422 bacterial community only a minority of the bacteria is able to disperse under relatively low  
423 hydration conditions, previously thought too dry for flagellar motility. During dry periods, these  
424 highly efficient dispersers will gain a significant advantage with their ability to colonize new  
425 habitats ahead of the rest of the community. This highlights the need for increased focus on  
426 complex communities, rather than pure culture studies for the prediction of actual dispersal ability  
427 on solid surfaces such as soil.

428

#### 429 **Materials and Methods**

430 **Bacterial Strains.**

431 The bacterial strain *Pseudomonas putida* KT2440 GFP, a tagged-derivative of a motile bacterium  
432 initially isolated from rhizosphere soil (61) was used as a motile model strain for flagellar motility  
433 and a nonflagellated mutant *P. putida* K2440 dsRed *fliM* previously created (18) was used as a  
434 non-motile model strain.

435 The bacterial strain *Flavobacterium johnsoniae* CJ1827 (37), was used as a model strain for gliding  
436 motility, and a non-motile mutant *F. johnsoniae* 2122  $\Delta$ *gldK* (38) was used as a non-motile model  
437 strain. All strains were routinely maintained on agar plates. *P. putida* strains on R2 agar (R2A,  
438 Fluka; Sigma-Aldrich, St. Louis, USA) and *F. johnsoniae* strains on CYE agar (62) medium at 25°C.

439

440 **Visualizing dispersal of non-fluorescent bacteria from environmental samples on the PSM.**

441 The porous surface model (PSM) has previously been described and used for observing motility  
442 and growth of fluorescent strains after their inoculation at the center of a ceramic disc (diameter =  
443 41.3 mm, thickness = 7.1 mm, maximum pore size <1.5  $\mu$ m, 1 bar bubbling pressure; Soilmoisture,  
444 Santa Barbara, USA) simulating a soil surface, under controlled hydration conditions (35).

445 Imposing suction on the disc controls the thickness of the liquid film on the ceramic surface.

446

447 In this study, we have expanded the use of the PSM for environmental communities. As non-  
448 fluorescent cells are not detectable on the surface of the ceramic disc by standard microscopy, we  
449 trapped the bacteria from the PSM by pressing small agar plates on top of the ceramic disc. This  
450 allows visualizing the colonization on the ceramic disc by observing the growth on the  
451 corresponding agar plates (Fig. S1).

452

453 The agar plates were obtained by pouring 6.3 ml 25% R2A with 20 g agar l<sup>-1</sup> into the lid of a small  
454 plastic petri dish (Star<sup>TM</sup>Dish diameter, 40 mm; height, 12.5 mm; Phoenix Biomedical Products,  
455 Mississauga, Canada) filling it to the brim. To further flatten the surface of the agar, the sterile lid  
456 of a standard petri dish (diameter 90 mm; height, 14.2 mm; VWR International, Søborg, Denmark)  
457 was pressed on top of the small agar plate before it had completely solidified. After drying the  
458 small agar plate was transferred into a big petri dish for storage. The PSM reservoirs were filled  
459 with 200-250 ml 25% R2B (Alpha Biosciences, Maryland, USA) and autoclaved before use.

460

461 Preliminary tests with fluorescent strains revealed that pressing of agar plates on the ceramic discs  
462 provided a distorted image of the bacterial spatial pattern because cells are inevitably displaced  
463 along the contact plane. Therefore, we detected bacterial colonization in concentric annular  
464 sections of the PSM surface. By preparing agar plates with holes of diameters ranging from 11.5  
465 mm, 15 mm, 20 mm and 25 mm (Fig. S1) we could estimate dispersal by sequentially pressing  
466 these plates on the PSM starting with that with the biggest hole and finishing with a full plate (Full  
467 Plate).

468 The holes were punched in the agar plates with a custom-made tool consisting of a teflon handle,  
469 for safe handling during flame sterilization, fitted to brass tubes of varying diameters (length: 12.5  
470 cm; diameters: 11.5, 15, 20, 25 mm) (Fig. S2). A printed template was placed under the agar plate  
471 to help center the holes. All plates were kept for a minimum of 48 h at room temperature before  
472 use on the PSM to test for contamination.

473

474 **Proof of concept with motile and non-motile pure cultures.**

475 We tested the ability of the method for distinguishing the dispersal patterns of *P. Putida* KT2440  
476 GFP and *P. putida* K2440 dsRed *fliM*. The bacteria, cultivated on R2A plates, were suspended in  
477 0.9% NaCl solution and adjusted by optical density measurements at 600 nm to obtain a cell  
478 density of ca. 2000 cells  $\mu\text{l}^{-1}$ , as confirmed by plate counts. Before inoculation the PSMs were  
479 elevated to -4.2 kPa (the length of the hanging water column is 40 cm and equals a suction of -4.2  
480 kPa) for 20 minutes to drain excess fluid from the ceramic surface. The two bacterial suspensions  
481 were mixed in equal ratio and 0.5  $\mu\text{l}$  was inoculated in the center of the ceramic disc, where it was  
482 rapidly absorbed. The discs were then brought to matric potentials (suction) of -0.5, -1.2 or -3.1  
483 kPa (-5, -12 and -30 cm of water suction) and incubated at room temperature for 14, 24 or 40  
484 hours before sampling by pressing the suite of agar plates onto the surface. Plates were incubated  
485 at 25°C for a 48 h growth period before being stored in the fridge at 4°C until observation by  
486 microscopy.

487  
488 To test the applicability of the method for other types of motility, we tested the gliding bacterium  
489 *Flavobacterium johnsoniae* strain CJ1827 (37), and a non-motile mutant *F. johnsoniae* 2122  $\Delta\text{gldK}$   
490 (38) on separate PSMs. Bacteria were streaked from CYE agar and grown in overnight cultures at  
491 25°C in motility medium (MM) (63) and adjusted by optical density measurements at 600 nm to  
492 obtain a cell density of ca. 63000 cells  $\mu\text{l}^{-1}$ , as confirmed by plate counts, before inoculation of 1  $\mu\text{l}$   
493 in the center of the ceramic disc. The PSMs were kept at -0.5 kPa for 48h of incubation at room  
494 temperature, using 25% R2B medium in the PSM reservoirs. The 25% R2A pressed plates were  
495 kept at 25°C for a 48 h period before growth was recorded.

496

497 **Microscopy and imaging.**

498 *P. putida* KT2440 GFP and *P. putida* KT2440 dsRed *fliM* spatial patterns on the PSM and on agar  
499 plates were determined with a Leica MZ16 FA epifluorescence stereomicroscope equipped for GFP  
500 and DsRed detection and fitted with a charge-coupled device (CCD) camera. Each plate was scored  
501 for the presence or absence of each strain. For documentation purpose, the entire surface of  
502 selected plates was imaged by sequentially capturing several fields of view, using a motorized  
503 stage piloted by Image Pro Plus (version 7.1; Media Cybernetics, Silver Spring, MD, USA) and then  
504 assembling a tiled image using the same software. The GFP and DsRed images of each plate were  
505 captured separately and then combined into one image.

506

507 To document the presence of colonies on the plates independently of fluorescence, the plates  
508 were subsequently imaged using the camera of a GelDocXR (Bio-Rad), operated in 'epiwhite'  
509 mode.

510

#### 511 **Dispersal potential of environmental communities.**

512 A soil sample was collected from the plow layer (5-15 cm depth) of a Danish agricultural field,  
513 included in the Danish Pesticide Risk Assessment Program (PLAP) (64) in March 2016 (Fårdrup,  
514 Sjælland). The soil is characterized by clay till and further details can be found at  
515 <http://pesticidvarsling.dk/>. The soil was stored at 4°C. For each experiment 25 g sieved (2 mm) soil  
516 was taken by composite sampling, i.e. as small subsamples taken from the original soil sample and  
517 then mixed. The soil bacteria were extracted using Nycodenz density gradient centrifugation as in  
518 (65), except for the final cell density determination, which was performed directly using a Thoma  
519 counting chamber. Cell density was adjusted to  $0.5 - 1 \times 10^6$  cells  $\mu\text{l}^{-1}$  in 0.9% NaCl solution and 10  
520  $\mu\text{l}$  inoculated as 1  $\mu\text{l}$  drops in the center of the ceramic discs. This inoculum corresponded to ca.



521 2000 to 8000 CFUs on R2A plates. All plates used for the environmental communities were  
522 amended with 100 mg l<sup>-1</sup> Delvolid to inhibit fungal growth (Natamycin, DSM food specialties,  
523 Delft, The Netherlands).

524 Lake water was sampled from the urban lake Sortedamssøen (Copenhagen), in September 2016.  
525 Four litre were collected from the surface water approximately 1.5 m from the shore. The water  
526 sample was filtrated first through a 2 µm glass fiber prefilter (Merck Millipore; Tullagreen, Ireland)  
527 and then through 0.2 µm polycarbonate filters (GVS Filter Technology; Morecambe, United  
528 Kingdom) on a filtration manifold (DHI Lab Products; Hørsholm, Denmark). The filters were  
529 transferred into a 15 ml falcon tube with 2.5 ml 0.9% NaCl solution and vortexed for 45 seconds.  
530 The filters were removed, and the cell density was adjusted by Thoma count to  $2 \times 10^6$  cells µl<sup>-1</sup>. 20  
531 µl of the suspension was inoculated as 1 µl drops, yielding 34125 CFUs per inoculum based on  
532 drop plate counts on R2A plates. Both the lake and soil inoculum were kept at 4°C overnight  
533 before inoculation on the ceramic discs. After inoculation, the discs were brought to matric  
534 potentials of -0.5 and -3.1 kPa and incubated at room temperature for 24 to 48 hours before  
535 sampling.

536

537 After sampling by pressing of the agar plate series on the PSMs at appropriate times, plates were  
538 incubated for 72 hours at 25°C. In addition to the presence / absence score, used in the pure  
539 culture studies, the coverage of bacterial growth on the individual agar plates were roughly  
540 estimated by eye using 4 categories; 1-25, 26-50, 51-75 and 76-100% coverage.

541

542 After scoring, for each pressed plate series, the plate with the fastest colonizers (bacteria present  
543 the furthest from center) and the Full Plate with the total cultivable community were chosen for

544 amplicon sequencing. In addition to these, for each separate experiment a “Reference Plate” was  
545 made, by drop plating 10  $\mu$ l of the inoculum onto the center of a small 25% R2A plate with 20 g  
546 agar l<sup>-1</sup>. This was meant as a motility-restricted control for the bacteria cultivable on the medium.

547

548 The bacteria were then washed from the agar plates by transferring the agar from the small petri  
549 dish into a standard size petri dish with a flamed spatula, adding 2 ml 0.9% NaCl solution for 10  
550 minutes and then gently rubbing the surface of the agar with a sterile inoculation loop and  
551 collecting the bacterial suspension by pipetting into an Eppendorf tube. The procedure was  
552 repeated twice with 1.5 ml 0.9% NaCl and the suspensions collected. The Eppendorf tubes were  
553 centrifuged for 5 minutes at 7500 x g before pooling into a single 1 ml sample suspension. The cell  
554 suspensions (plate wash) from the pressed plates, the Reference Plates, Nycodenz extracts, lake  
555 filtrate and leftover inoculums were all transferred to cryotubes and stored at -80°C.

556

#### 557 **DNA extraction and sequencing.**

558 DNA was extracted using the Powerlyzer Powersoil kit (MoBio; Carlsbad, USA) following the  
559 manufacturer’s protocol with a few changes. 500  $\mu$ l of the thawed plate wash was centrifuged for  
560 5 minutes at 10.000 x g. the supernatant was removed and the pellet dissolved by adding 750  $\mu$ l  
561 bead solution and vortexing. The suspensions were transferred to Glass Bead Tubes, 60  $\mu$ l C1  
562 solution was added and samples were placed in a Bead Beater for 5 minutes at 2000 RPM.  
563 Hereafter the manufacturer’s protocol was followed. DNA concentrations were measured on  
564 Qubit 2.0 (Life Technologies, Invitrogen; Carlsbad, USA) and stored at -80°C until sequencing.

565

566 The extracted DNA was PCR-amplified using the universal primer set PRK341F (5'-  
567 CCTAYGGGRBGCASCAG-3') and PRK806R (5'-GGACTACNNGGTATCTAAT-3') that amplify the V3-  
568 V4 hypervariable regions of the 16S rRNA genes (66). 2 x 300 bp Purified PCR products were  
569 sequenced on the Illumina MiSeq platform at the DTU Multi Assay Core Center (Lyngby, DK). All  
570 raw 16S rRNA gene amplicons were processed with the DADA2 pipeline (67) with default  
571 parameters. The sequences were classified based on the SILVA prokaryotic reference database  
572 version 123 (68). A total of 3.8 million sequences passed the filtering steps, representing an  
573 average of  $5.3 \times 10^4$  sequences per sample.

574

575 Shannon indices were computed in R software (version 3.3.1; R Core Team (2016)) using the  
576 "plot\_richness" function in the "phyloseq" package (69). Samples were rarefied to even depth  
577 (average of 10 iterations) with the "rarefy\_even\_depth" function in the "phyloseq" package before  
578 calculating Faith's Diversity with the "pd.query" function of the "PhyloMeasures" package (70).  
579 Heatmaps were plotted using the "amp\_heatmap" function of the "ampvis" package (71), while  
580 "ggplot2" (72) and "ggtree" (73) were used for plots and phylogenetic trees, respectively.

581 Type strains were identified using EZBioCloud ([www.ezbiocloud.net](http://www.ezbiocloud.net)) (74), and the closest match  
582 along with sequences for common *Pseudomonads* and *Aeromonads* were added to the trees for  
583 reference. For construction of phylogenetic trees with type strains, sequences were aligned with  
584 ClustalW in MEGA7 with the following parameters: Pairwise Alignment: Gap open: 1, extension:  
585 6.66, Multiple Alignment: Gap open: 15, extension: 6.66. Sequences were trimmed to even length.  
586 Tree was constructed with the "UPGMA" function in package "phangorn" (75).

587

588 All sequencing data have been deposited as a NCBI BioProject under accession number  
589 PRJNA400555.

590

591 **Statistical analysis.**

592 Mann-Whitney Rank Sum Test in Sigmaplot 13 (Systat Software Inc., San Jose, CA, USA) was used  
593 for the dispersal profile data. One-way ANOVA based on comparison of the differences between  
594 Full Plates and dispersed communities was used for 24 h Shannon Diversity Indices. Kruskal-Wallis  
595 Analysis of Variance on ranks (data was ranked due to unequal variance) and a paired t-test was  
596 used for 48 h Shannon Diversity Indices. *P* values < 0.05 were considered significant.

597

598 **Acknowledgements**

599 This study was funded by the Villum Kann Rasmussen Foundation through the Center for  
600 Environmental and Agricultural Microbiology (CREAM). The authors would like to thank Professor  
601 Mark McBride, University of Wisconsin for kindly supplying the *F. johnsoniae* strains used in this  
602 study.

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618 REFERENCES:

- 619 1. Holt RD. 2009. IJEE Soapbox. *Isr J Ecol Evol* 55:307–313.
- 620 2. Reichenbach T, Mobilia M, Frey E. 2007. Mobility promotes and jeopardizes biodiversity in  
621 rock–paper–scissors games. *Nature* 448:1046–1049.
- 622 3. Bonte D, Doherty M. 2017. Dispersal: a central and independent trait in life history. *Oikos*  
623 126:472–479.
- 624 4. Nemergut DR, Schmidt SK, Fukami T, O’Neill SP, Bilinski TM, Stanish LF, Knelman JE, Darcy  
625 JL, Lynch RC, Wickey P, Ferrenberg S. 2013. Patterns and Processes of Microbial Community  
626 Assembly. *Microbiol Mol Biol Rev* 77:342–356.
- 627 5. Bilton DT, Freeland JR, Okamura B. 2001. Dispersal in Freshwater Invertebrates. *Annu Rev*  
628 *Ecol Syst* 32:159–181.
- 629 6. Dechesne A, Badawi N, Aamand J, Smets BF. 2014. Fine scale spatial variability of microbial  
630 pesticide degradation in soil: Scales, controlling factors, and implications. *Front Microbiol*  
631 5:1–14.
- 632 7. Or D, Smets BF, Wraith JM, Dechesne A, Friedman SP. 2007. Physical constraints affecting  
633 bacterial habitats and activity in unsaturated porous media - a review. *Adv Water Resour*  
634 30:1505–1527.
- 635 8. Henrichsen J. 1972. Bacterial surface translocation: a survey and a classification. *Bacteriol*  
636 *Rev* 36:478–503.
- 637 9. Harshey R. 2003. Bacterial motility on a surface: many ways to a common goal. *Annu Rev*  
638 *Microbiol* 57:249–273.
- 639 10. Shrouf JD. 2015. A fantastic voyage for sliding bacteria. *Trends Microbiol* 23:244–246.

- 640 11. Hölscher T, Kovács ÁT. 2017. Sliding on the surface: Bacterial spreading without an active  
641 motor. *Environ Microbiol* 0:2537–2545.
- 642 12. McBride MJ. 2001. Bacterial Gliding Motility: Multiple Mechanisms for Cell Movement over  
643 Surfaces. *Annu Rev Microbiol* 55:49–75.
- 644 13. Snyder LAS, Loman NJ, Fütterer K, Pallen MJ. 2009. Bacterial flagellar diversity and  
645 evolution: seek simplicity and distrust it? *Trends Microbiol* 17:1–5.
- 646 14. Liu R, Ochman H. 2007. Stepwise formation of the bacterial flagellar system 104.
- 647 15. Luciano J, Agrebi R, Le Gall AV, Wartel M, Fiegna F, Ducret A, Brochier-Armanet C, Mignot T.  
648 2011. Emergence and modular evolution of a novel motility machinery in bacteria. *PLoS*  
649 *Genet* 7.
- 650 16. Whitman WB, Coleman DC, Wiebe WJ. 1998. Prokaryotes: The unseen majority. *Proc Natl*  
651 *Acad Sci* 95:6578–6583.
- 652 17. Ben-Jacob E, Finkelshtein A, Ariel G, Ingham C. 2016. Multispecies Swarms of Social  
653 Microorganisms as Moving Ecosystems. *Trends Microbiol* 24:257–269.
- 654 18. Dechesne A, Wang G, Gulez G, Or D, Smets BF. 2010. Hydration-controlled bacterial motility  
655 and dispersal on surfaces. *Proc Natl Acad Sci U S A* 107:14369–14372.
- 656 19. Venturi V, Bertani I, Kerényi Á, Netotea S, Pongor S. 2010. Co-swarms and local collapse:  
657 Quorum sensing conveys resilience to bacterial communities by localizing cheater mutants  
658 in *Pseudomonas aeruginosa*. *PLoS One* 5.
- 659 20. Zhou J, Ma Q, Yi H, Wang L, Song H, Yuan YJ. 2011. Metabolome profiling reveals metabolic  
660 cooperation between *Bacillus megaterium* and *Ketogulonicigenium vulgare* during induced  
661 swarm motility. *Appl Environ Microbiol* 77:7023–7030.
- 662 21. Finkelshtein A, Roth D, Ben Jacob E, Ingham CJ. 2015. Bacterial Swarms Recruit Cargo

- 663 Bacteria To Pave the Way in Toxic Environments. *MBio* 6:1–10.
- 664 22. Ellegaard-Jensen L, Knudsen BE, Johansen A, Albers CN, Aamand J, Rosendahl S. 2014.
- 665 Fungal-bacterial consortia increase diuron degradation in water-unsaturated systems. *Sci*
- 666 *Total Environ* 466–467:699–705.
- 667 23. Warmink JA, Nazir R, Corten B, van Elsas . JD. 2011. Hitchhikers on the fungal highway: The
- 668 helper effect for bacterial migration via fungal hyphae. *Soil Biol Biochem* 43:760–765.
- 669 24. Nazir R, Zhang M, de Boer W, van Elsas JD. 2012. The capacity to comigrate with *Lyophyllum*
- 670 *sp.* strain Karsten through different soils is spread among several phylogenetic groups
- 671 within the genus *Burkholderia*. *Soil Biol Biochem* 50:221–233.
- 672 25. Warmink JA, Van Elsas JD. 2009. Migratory response of soil bacteria to *Lyophyllum sp.* strain
- 673 Karsten In soil microcosms. *Appl Environ Microbiol* 75:2820–2830.
- 674 26. Brock DA, Read S, Bozhchenko A, Queller DC, Strassmann JE. 2013. Social amoeba farmers
- 675 carry defensive symbionts to protect and privatize their crops. *Nat Commun* 4:2385.
- 676 27. Dennis PG, Seymour J, Kumbun K, Tyson GW. 2013. Diverse populations of lake water
- 677 bacteria exhibit chemotaxis towards inorganic nutrients. *Isme J* 7:1661–1664.
- 678 28. Wolf AB, Rudnick M-BB, de Boer W, Kowalchuk GA. 2015. Early colonizers of unoccupied
- 679 habitats represent a minority of the soil bacterial community. *FEMS Microbiol Ecol*
- 680 91:fiv024.
- 681 29. Grossart HP, Riemann L, Azam F. 2001. Bacterial motility in the sea and its ecological
- 682 implications. *Aquat Microb Ecol* 25:247–258.
- 683 30. Fenchel T. 2001. Eppure si muove: Many water column bacteria are motile. *Aquat Microb*
- 684 *Ecol* 24:197–201.
- 685 31. Mitchell JG, Pearson L, Bonazinga A, Dillon S, Khouri H, Paxinos R. 1995. Long lag times and

- 686 high velocities in the motility of natural assemblages of marine bacteria. *Appl Environ*  
687 *Microbiol* 61:877–882.
- 688 32. Wolf AB, Vos M, de Boer W, Kowalchuk GA. 2013. Impact of Matric Potential and Pore Size  
689 Distribution on Growth Dynamics of Filamentous and Non-Filamentous Soil Bacteria. *PLoS*  
690 *One* 8:e83661.
- 691 33. Wang G, Or D. 2010. Aqueous films limit bacterial cell motility and colony expansion on  
692 partially saturated rough surfaces. *Environ Microbiol* 12:1363–1373.
- 693 34. Tecon R, Or D. 2016. Bacterial flagellar motility on hydrated rough surfaces controlled by  
694 aqueous film thickness and connectedness. *Sci Rep* 6:19409.
- 695 35. Dechesne A, Or D, Gulez G, Smets BF. 2008. The porous surface model, a novel  
696 experimental system for online quantitative observation of microbial processes under  
697 unsaturated conditions. *Appl Environ Microbiol* 74:5195–5200.
- 698 36. Jordan FL, Maier RM. 1999. Development of an agar lift-DNA/DNA hybridization technique  
699 for use in visualization of the spatial distribution of Eubacteria on soil surfaces. *J Microbiol*  
700 *Methods* 38:107–117.
- 701 37. Rhodes RG, Pucker HG, McBride MJ. 2011. Development and use of a gene deletion strategy  
702 for *Flavobacterium johnsoniae* to identify the redundant gliding motility genes *remF*, *remG*,  
703 *remH*, and *remI*. *J Bacteriol* 193:2418–2428.
- 704 38. Shrivastava A, Johnston JJ, Van Baaren JM, McBride MJ. 2013. *Flavobacterium johnsoniae*  
705 *GldK*, *GldL*, *GldM*, and *SprA* are required for secretion of the cell surface gliding motility  
706 adhesins *sprb* and *remA*. *J Bacteriol* 195:3201–3212.
- 707 39. Dechesne A, Smets BF. 2012. *Pseudomonad* Swarming Motility Is Restricted to a Narrow  
708 Range of High Matric Water Potentials. *Appl Environ Microbiol* 78:2936–2940.



- 709 40. Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should replace  
710 operational taxonomic units in marker-gene data analysis. *ISME J* 11:2639–2643.
- 711 41. Kleyer H, Tecon R, Or D. 2017. Hydration dynamics alters species composition of a synthetic  
712 soil bacterial community inhabiting unsaturated porous microcosms, p. 89, abstr P–44,. *In*  
713 BAGECO 14 Bacterial Genetics and Ecology, Aberdeen, UK.
- 714 42. Liebeke M, Brözel VS, Hecker M, Lalk M. 2009. Chemical characterization of soil extract as  
715 growth media for the ecophysiological study of bacteria. *Appl Microbiol Biotechnol* 83:161–  
716 173.
- 717 43. Holmsgaard PN, Norman A, Hede SC, Poulsen PHB, Al-Soud WA, Hansen LH, Sørensen SJ.  
718 2011. Bias in bacterial diversity as a result of Nycodenz extraction from bulk soil. *Soil Biol*  
719 *Biochem* 43:2152–2159.
- 720 44. Portillo MC, Leff JW, Lauber CL, Fierer N. 2013. Cell size distributions of soil bacterial and  
721 archaeal taxa. *Appl Environ Microbiol* 79:7610–7617.
- 722 45. Mctee MR, Gibbons SM, Feris K, Gordon NS, Gannon JE, Ramsey PW. 2013. Heavy metal  
723 tolerance genes alter cellular thermodynamics in *Pseudomonas putida* and river  
724 *Pseudomonas* spp. and influence ameбал predation. *FEMS Microbiol Lett* 347:97–106.
- 725 46. Bakken LR, Olsen R a. 1987. The relationship between cell size and viability of soil bacteria.  
726 *Microb Ecol* 13:103–14.
- 727 47. Wang Y, Hammes F, Boon N, Egli T. 2007. Quantification of the filterability of freshwater  
728 bacteria through 0.45, 0.22, and 0.1  $\mu\text{m}$  pore size filters and shape-dependent enrichment  
729 of filterable bacterial communities. *Environ Sci Technol* 41:7080–7086.
- 730 48. Hahn MW. 2004. Broad diversity of viable bacteria in “sterile” (0.2  $\mu\text{m}$ ) filtered water. *Res*  
731 *Microbiol* 155:688–691.

- 732 49. Givskov M, Givskov M, Eberl L, Eberl L, Moller S, Moller S, Poulsen LK, Poulsen LK, Molin S,  
733 Molin S. 1994. Responses To Nutrient Starvation In *Pseudomonas-putida* Kt2442 - Analysis  
734 Of General Cross-Protection, Cell-Shape, And Macromolecular Content 176:7–14.
- 735 50. Raaijmakers JM, De Bruijn I, Nybroe O, Ongena M. 2010. Natural functions of lipopeptides  
736 from *Bacillus* and *Pseudomonas*: more than surfactants and antibiotics. *FEMS Microbiol Rev*  
737 34:1037–1062.
- 738 51. Ndlovu T, Khan S, Khan W. 2016. Distribution and diversity of biosurfactant-producing  
739 bacteria in a wastewater treatment plant. *Environ Sci Pollut Res* 23:9993–10004.
- 740 52. Illori MO, Amobi CJ, Odocha AC. 2005. Factors affecting biosurfactant production by oil  
741 degrading *Aeromonas* spp. isolated from a tropical environment. *Chemosphere* 61:985–  
742 992.
- 743 53. Burch AY, Do PT, Sbodio A, Suslow T V., Lindow SE. 2016. High-level culturability of epiphytic  
744 bacteria and frequency of biosurfactant producers on leaves. *Appl Environ Microbiol*  
745 82:5997–6009.
- 746 54. Lindow SE, Brandl MT. 2003. Microbiology of the Phyllosphere MINIREVIEW Microbiology of  
747 the Phyllosphere. *Appl Environ Microbiol* 69:1875–1883.
- 748 55. Burch AY, Zeisler V, Yokota K, Schreiber L, Lindow SE. 2014. The hygroscopic biosurfactant  
749 syringafactin produced by *Pseudomonas syringae* enhances fitness on leaf surfaces during  
750 fluctuating humidity. *Environ Microbiol* 16:2086–98.
- 751 56. Burch AY, Shimada BK, Browne PJ, Lindow SE. 2010. Novel high-throughput detection  
752 method to assess bacterial surfactant production. *Appl Environ Microbiol* 76:5363–5372.
- 753 57. Shrivastava A, Lele PP, Berg HC. 2015. A rotary motor drives *Flavobacterium* gliding. *Curr*  
754 *Biol* 25:338–341.

- 755 58. Jarrell KF, McBride MJ. 2008. The surprisingly diverse ways that prokaryotes move. *Nat Rev*  
756 *Microbiol* 6:466–476.
- 757 59. McBride MJ. 2014. *The Prokaryotes*, 4th ed.
- 758 60. Sampedro I, Parales RE, Krell T, Hill JE. 2015. *Pseudomonas chemotaxis*. *FEMS Microbiol Rev*  
759 39:17–46.
- 760 61. Nakazawa T. 2002. Travels of a *Pseudomonas*, from Japan. *Environ Microbiol* 4:782–786.
- 761 62. Lin D, McBride MJ. 1996. Development of techniques for the genetic manipulation of the  
762 gliding bacteria *Lysobacter enzymogenes* and *Lysobacter brunescens*. *Can J Microbiol*  
763 42:896–902.
- 764 63. Liu J, McBride MJ, Subramaniam S. 2007. Cell surface filaments of the gliding bacterium  
765 *Flavobacterium johnsoniae* revealed by cryo-electron tomography. *J Bacteriol* 189:7503–  
766 7506.
- 767 64. Danish Pesticide Leaching Assessment Programme (PLAP). <http://pesticidvarsling.dk>.
- 768 65. Klümper U, Dechesne A, Smets B. 2014. Protocol for Evaluating the Permissiveness of  
769 Bacterial Communities Toward Conjugal Plasmids by Quantification and Isolation of  
770 Transconjugants. *Hydrocarb Lipid Microbiol Protoc Springer Protoc Handb* 1–14.
- 771 66. Yu Y, Lee C, Kim J, Hwang S. 2005. Group-specific primer and probe sets to detect  
772 methanogenic communities using quantitative real-time polymerase chain reaction.  
773 *Biotechnol Bioeng* 89:670–679.
- 774 67. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-  
775 resolution sample inference from Illumina amplicon data. *Nat Methods* 13:581–583.
- 776 68. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. 2007. SILVA: A  
777 comprehensive online resource for quality checked and aligned ribosomal RNA sequence

- 778 data compatible with ARB. *Nucleic Acids Res* 35:7188–7196.
- 779 69. McMurdie PJ, Holmes S. 2013. Phyloseq: An R Package for Reproducible Interactive Analysis  
780 and Graphics of Microbiome Census Data. *PLoS One* 8.
- 781 70. Tsirogiannis C, Sandel B. 2016. PhyloMeasures: a package for computing phylogenetic  
782 biodiversity measures and their statistical moments. *Ecography (Cop)* 39:709–714.
- 783 71. Albertsen M, Karst SM, Ziegler AS, Kirkegaard RH, Nielsen PH, Stokholm-Bjerregaard M.  
784 2015. Back to Basics – The Influence of DNA Extraction and Primer Choice on Phylogenetic  
785 Analysis of Activated Sludge Communities. *PLoS One* 10:e0132783.
- 786 72. Wickham H. 2009. *Ggplot2 : elegant graphics for data analysis*. Springer.
- 787 73. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree : an r package for visualization and  
788 annotation of phylogenetic trees with their covariates and other associated data. *Methods*  
789 *Ecol Evol* 8:28–36.
- 790 74. Yoon S-H, Ha S-M, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: A  
791 taxonomically united database of 16S rRNA and whole genome assemblies. *Int J Syst Evol*  
792 *Microbiol* 1613–1617.
- 793 75. Schliep KP. 2011. phangorn: Phylogenetic analysis in R. *Bioinformatics* 27:592–593.


794

795

796 **Figures**

797

798



799

800

801 **FIG 1 Proof of concept using pure cultures.**

802 A) Separation of the motile strain *P. putida* KT2440 GFP (green) and the non-motile *P. putida*  
803 KT2440 dsRed *fliM* (red) on agar plates pressed onto the ceramic disc as pictured with multiple  
804 fields of epifluorescence microscopy. The non-motile strain was only detected on the Full Plate  
805 press (red arrow), while the motile strain was detected on all of the pressed plates, including the  
806 one that captures the zone most distant from the inoculation point (green arrow). B) Dispersal  
807 assessed with a camera without fluorescence detection, which is the method used for  
808 environmental communities. Contrast has been digitally enhanced. The plates have been pressed  
809 after 40 h dispersal at -0.5 kPa.

810

811

812

813


814

815

816

817

818




819  
820  
821

822 **FIG 2 Dispersal dynamics of motile and non-motile strains as affected by the matric potential.**

823 The progressive dispersal of the motile strain *P. putida* KT2440 GFP was captured by our method,  
824 as well as the inability of the non-motile *P. putida* KT2440 *fliM* DsRed to disperse away from  
825 center of the ceramic disc. Both motile (green) and non-motile (red) strains were tested at three  
826 matric potentials (kPa). For the non-motile, only -0.5 kPa is depicted as the other values were  
827 similar, with bacteria solely present at the center. The distances shown are ranges, e.g. colonies  
828 have been observed on the agar ring at a distance between 11.5 to 15 mm from center. Numbers  
829 of replicate dispersal experiments vary from 2 to 5.

830  
831  
832  
833  
834  
835  
836

837



838

839

840 **FIG 3** Dispersal of a soil and a lake community over time at different matric potentials. Symbol

841 shading depicts bacterial coverage of the pressed agar plate, giving an indication of the extent of

842 colonization. The lake community was tested at two matric potentials vs four for the soil one; the

843 number of replication varied from two to five.

844

845

846


847

848

849

850

851



852

853 **FIG 4.** Heatmap of the relative abundance of the 20 most dominant genera across communities

854 derived from a soil extract and differing in their dispersal after being incubated at prescribed

855 matric potential for 24h or 48h. For 24 h, two additional matric potentials of 0.0 kPa and -4.2 kPa


856 (only one sample recovered at 20 mm) were added. Columns present the average of triplicate

857 communities, except for the motility restricted control (Reference Plate; n=4), the total

858 community on the Full Plate at -4.2 kPa (n= 1) and the fastest dispersed community at -1.2 for 24 h

859 (n=2) and at -3.1 kPa (n=2), and -4.2 kPa (n=1) for 48 h.






860

861 **FIG 5** Heatmap of the relative abundance of the 20 most dominant genera across communities  
 862 derived from a lake filtrate and differing in their dispersal after being incubated at prescribed  
 863 matric potential for 24 h or 48 h. Columns present the average of duplicate communities, except  
 864 for the motility restricted control (Reference Plate, n=1), the total community on the Full Plate at -  
 865 3.1 kPa (n= 3) for 48 h and the fastest dispersed community at -3.1 kPa for 24 h (n=1).

866

867



868

869 **FIG 6** Estimates of alpha-diversity (Shannon Diversity Index and Faith's Phylogenetic Diversity

870 index) for communities derived from soil or from a lake, after 48 hours incubation at prescribed

871 matric potentials. For each matric potential, the total community recovered from the full agar

872 plate (Full Plate) and the fastest dispersed community is presented. A motility restricted control

873 (Reference Plate) is also included. Replicates are depicted as separate dots. The Faith's

874 Phylogenetic Diversity indices reported are the average of values obtained for 10 random

875 rarefactions.

876