



Monge surfaces and planar geodesic foliations

Brander, David; Gravesen, Jens

Published in:
Journal of Geometry

Link to article, DOI:
[10.1007/s00022-018-0413-7](https://doi.org/10.1007/s00022-018-0413-7)

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Brander, D., & Gravesen, J. (2018). Monge surfaces and planar geodesic foliations. *Journal of Geometry*, 109(4). DOI: [10.1007/s00022-018-0413-7](https://doi.org/10.1007/s00022-018-0413-7)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Monge surfaces and planar geodesic foliations

David Brander and Jens Gravesen

Abstract. A Monge surface is a surface obtained by sweeping a generating plane curve along a trajectory that is orthogonal to the moving plane containing the curve. Locally, they are characterized as being foliated by a family of planar geodesic lines of curvature. We call surfaces with the latter property PGF surfaces, and investigate the global properties of these two naturally defined objects. The only compact orientable PGF surfaces are tori; these are globally Monge surfaces, and they have a simple characterization in terms of the directrix. We show how to produce many examples of Monge tori and Klein bottles, as well as tori that do not have a closed directrix.

Mathematics Subject Classification (2010). Primary 53A05; Secondary 53C12, 53C22.

Keywords. Monge surface, planar geodesic, developable surface.

1. Introduction

A *Monge surface* is the surface swept out by the motion of a plane curve (the *profile curve* or *generatrix*) whilst the plane is moved through space in such a way that the movement of the plane is always in the direction of the normal to the plane. They are also characterized as surfaces swept out by *parallels*, each of which is the orthogonal trajectory of a point on the initial plane curve, and all of which are parallel to each other. These surfaces were defined by Monge ([8], §XXV) and classically studied by, e.g. Darboux ([5], Section 85), and L. Raffy [9]. Each meridian (i.e., the intersection of the surface with the moving plane) is a normal section of the surface, and therefore a geodesic. Moreover, the meridians are all congruent. Hence a Monge surface is foliated by (congruent) planar geodesics.

Conversely, various practical applications, such as surfaces made from strips of wood [3], are modeled by surfaces foliated by planar geodesic lines of curvature. We call a surface with such a foliation a *PGF surface*. It was known classically (see [6], Section 129) that these surfaces are *locally* the

same objects as Monge surfaces. The purpose of this note is to consider the difference between the global objects given by these two natural definitions, and to investigate the problem of constructing compact examples.

To illustrate the difference at a global level between Monge surfaces and PGF surfaces, it is not hard to construct examples of PGF surfaces that are certainly not Monge surfaces (Figure 2). On the other hand, since a Monge surface is naturally defined by the arbitrary orthogonal motion of a plane through space, singularities naturally arise. For example, a sphere can be represented as a Monge surface with singularities at the poles, but a sphere does not have a global foliation by planar geodesics.

We investigate in a modern framework the basics of both the local and global theory that follows directly from the two natural definitions, focusing on PGF surfaces in Section 2 and Monge surfaces in Section 3.

In the last section we study the construction of *Monge tori*. We define a Monge torus to be a Monge surface that has the property that the profile curve and all of the parallels are closed curves. We show how to construct many examples of Monge tori and Klein bottles. We also answer a natural question that arises in the global theory of PGF surfaces: in Section 2 we observe (Proposition 2.6) that if $\mathbf{x} : M \rightarrow \mathbb{R}^3$ is a PGF immersion, and M is compact, then M is necessarily a torus. We also show (Theorem 2.4) that any complete immersed PGF surface $S \subset \mathbb{R}^3$ has a natural covering $\mathbb{R}^2 \rightarrow S$ consisting of lines of curvature, one family of which is the planar geodesic foliation, the other family being parallels of a Monge surface parameterization. If $S \subset \mathbb{R}^3$ is compact, one can ask whether or not the leaves of this double foliation are closed. It is easy to see that the planar geodesics are necessarily closed. However, one can show that the parallels need not be closed by constructing examples of Monge surfaces that are tori but not “Monge tori” – i.e., the parallels are not closed curves.

2. Planar Geodesic Foliations

If a plane curve is not a straight line, then the plane that it lies in is unique, and so a family of typical planar curves gives us a natural family of planes in \mathbb{R}^3 . For the case of a straight line, there are infinitely many planes that contain it. In order that there be a canonical choice, consider this property of the plane containing a non-rectilinear planar geodesic: the plane is orthogonal to the surface. This plane, determined by the tangent to the curve and the surface normal, is unique; and a straight line in a surface lies in such a plane if and only if the normal to the surface is constant along the line.

Definition 2.1. A *planar geodesic foliation* of an immersed surface $S \subset \mathbb{R}^3$ is a 1-dimensional foliation, the leaves of which are planar geodesic curves in S , with the additional property that, along each of the leaves, the normal to the surface is parallel to the plane containing the curve.

Note: it is not difficult to show that the conditions given on the leaves of the foliation are equivalent to requiring that they be planar geodesic *lines*

of curvature. The above definition includes all foliations by non-rectilinear planar geodesics. If the geodesics are straight lines, then the surface is developable, because the normal is required to be constant along the rulings. We will call a surface together with a planar geodesic foliation a *planar geodesically foliated (PGF) surface*.

2.1. Local planar geodesic foliations

Locally, a PGF surface is just a piece of a Monge surface: that is the statement of Theorem 2.2 below. A key ingredient of the proof is the existence, along any differentiable curve of a so-called *rotation minimizing frame*, or *relatively parallel adapted frame*, [2]. Given a differentiable curve \mathbf{r} , and any pair of vectors Q_1, Q_2 orthogonal to $\mathbf{r}'(v_0)$, there is a unique orthonormal frame field $(\mathbf{t}(v), \mathbf{q}_1(v), \mathbf{q}_2(v))$, where $\mathbf{t} = \mathbf{r}'/|\mathbf{r}'|$, $(\mathbf{q}_1, \mathbf{q}_2)(v_0) = (Q_1, Q_2)$, and such that the derivatives of \mathbf{q}_i are parallel to \mathbf{r}' , i.e., there is no rotation in the plane perpendicular to the curve. In Section 4 below, we use the fact that the torsion of the curve is the rate of rotation of such a rotation minimizing frame relative to the Frenet frame.

Theorem 2.2. *A planar geodesically foliated surface can, on a neighbourhood of any point, be locally parameterized in the form*

$$\mathbf{x}(u, v) = \mathbf{r}(v) + x(u) \mathbf{q}_1(v) + y(u) \mathbf{q}_2(v), \quad (2.1)$$

with \mathbf{r} a space curve with rotation minimizing frame $(\mathbf{t} = \mathbf{r}', \mathbf{q}_1, \mathbf{q}_2)$, and each isocurve $\gamma_v(u) = \mathbf{x}(u, v)$ a unit speed geodesic.

Proof. Without loss of generality, we can assume the surface is (locally) parameterized as $(u, v) \mapsto \mathbf{x}(u, v)$, where the isocurve $\gamma_{v_0}(u) := \mathbf{x}(u, v_0)$ is a planar unit-speed geodesic for each fixed v_0 . Now, varying v gives us a one parameter family $\Pi(v)$ of planes in \mathbb{R}^3 . Let $\mathcal{N}(v)$ be a family of normals to these planes. By integrating $\mathbf{r}'(v) = \mathcal{N}(v)$, with $\mathbf{r}(v_0) = \mathbf{x}(u_0, v_0)$, we obtain a curve $\mathbf{r}(v)$, the *spine curve* (or *directrix*), that intersects the planes orthogonally. Let $(\mathbf{t}, \mathbf{q}_1, \mathbf{q}_2)$ be the rotation minimizing frame along \mathbf{r} , with some choice of Q_1, Q_2 . Since each curve γ_v lies in the plane through $\mathbf{r}(v)$ spanned by $\mathbf{q}_i(v)$, we can write

$$\mathbf{x}(u, v) = \mathbf{r}(v) + x(u, v) \mathbf{q}_1(v) + y(u, v) \mathbf{q}_2(v). \quad (2.2)$$

The tangent to each γ_v is

$$\gamma'_v = \frac{\partial \mathbf{x}}{\partial u} = \frac{\partial x}{\partial u} \mathbf{q}_1 + \frac{\partial y}{\partial u} \mathbf{q}_2.$$

Set

$$\mathbf{n}_v = \mathbf{t} \times \gamma'_v = -\frac{\partial y}{\partial u} \mathbf{q}_1 + \frac{\partial x}{\partial u} \mathbf{q}_2.$$

At non-inflectional points, \mathbf{n}_v is the principal normal of the planar geodesic γ_v , but in any case it is perpendicular to both \mathbf{t} and γ'_v . By the assumption of Definition 2.1, the plane containing the curve γ_v is spanned by γ'_v and the

surface normal. Since \mathbf{n}_v also lies in this plane, it follows that \mathbf{n}_v is parallel to the surface normal or, equivalently,

$$\mathbf{n}_v \cdot \frac{\partial \mathbf{x}}{\partial v} = 0. \quad (2.3)$$

As \mathbf{t} is a unit vector field, there exist smooth functions κ_1 and κ_2 such that

$$\mathbf{t}'(v) = \kappa_1(v)\mathbf{q}_1(v) + \kappa_2(v)\mathbf{q}_2(v).$$

The rotation minimizing assumption means that

$$\frac{\partial \mathbf{x}}{\partial v} = (1 - x\kappa_1 - y\kappa_2)\mathbf{t} + \frac{\partial x}{\partial v}\mathbf{q}_1 + \frac{\partial y}{\partial v}\mathbf{q}_2.$$

So the condition (2.3) is

$$\frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = 0,$$

i.e., that the map $\phi(u, v) = (x(u, v), y(u, v))$ has rank less than two. Since, from (2.2), we must have $\partial\phi/\partial u \neq 0$, the image of ϕ is a curve. In other words, each curve $\alpha_v(u) = (x(u, v), y(u, v))$ is a reparameterization of a single curve $\alpha_{v_0}(u)$. From the assumption that $\gamma_v(u)$ is a unit speed geodesic, it follows easily that this reparameterization $u \mapsto \tilde{u}(u)$ is nothing but a translation $\tilde{u}(u) = u + c(v)$, and from this the stated result follows. \square

We will call a parameterization of the form (2.1) a *Monge surface parameterization*, as it follows from the definition of a Monge surface that a (regular) Monge surface has this form. We mention here that, given a Monge surface parameterization, where both the spine curve $\mathbf{r}(v)$ and the profile curve $(x(u), y(u))$ are parameterized by arc-length, the metric takes the form:

$$ds^2 = du^2 + (1 - \alpha)^2 dv^2, \quad \alpha(u, v) := x(u)\kappa_1(v) - y(u)\kappa_2(v). \quad (2.4)$$

Furthermore, one computes that the second fundamental form is diagonal, so (u, v) are principal coordinates, i.e., the parallels and meridians on a Monge surface are lines of curvature.

Let us call a surface *locally planar geodesically foliated* if there exists a neighbourhood around each point that admits a Monge surface parameterization. Using the fact that all geodesics on a sphere are planar, it is easy to construct complete examples of such surfaces with arbitrary topology by gluing in a C^∞ manner surfaces of revolution to pieces of spheres, as in Figure 1 (left), and then assembling multiple units.

Any surface such as those shown in Figure 1, where different choices of foliation are made for different regions of the surface, must, on the overlap of these regions, have more than one possible planar geodesic foliation. Obviously a piece of a plane, a sphere or a cylinder has this property. We now prove that these are essentially the only possibilities:

Theorem 2.3. *Let S be a connected surface that admits more than one distinct planar geodesic foliation. Then, either S is an open subset of a sphere, or every point of S has a neighbourhood U that is an open subset of a cylinder, $\gamma \times \mathbb{R}$, where γ is a plane curve.*




FIGURE 1. C^∞ immersed surfaces that are locally, but not globally, PGF surfaces.

Proof. Suppose S has two distinct foliations by planar geodesics, say \mathcal{F}_1 and \mathcal{F}_2 . Let p be a point of S , and suppose first that p is not an umbilical point. By Theorem 2.2, there is a neighbourhood U_1 of p such that U_1 is parameterized as a Monge surface $\mathbf{x}(u, v)$, where the geodesics are given by $v = \text{constant}$, and an analogous neighbourhood U_2 for \mathcal{F}_2 , parameterized by $\tilde{\mathbf{x}}(\tilde{u}, \tilde{v})$, where the planar geodesics are $\tilde{v} = \text{constant}$. Set $U = U_1 \cap U_2$. We can assume that U is small enough so that it contains no umbilic points. Then both (u, v) and (\tilde{u}, \tilde{v}) are curvature line coordinates on U . Since the lines of curvature through a non-umbilic point are unique, and the planar geodesic foliations are assumed to be distinct, it follows that the u isocurves are the \tilde{v} isocurves and vice versa, and we have $(\tilde{u}(u, v), \tilde{v}(u, v)) = (\tilde{u}(v), \tilde{v}(u))$. Moreover, all the u isocurves are congruent to each other, and all the v isocurves are congruent to each other. We can assume that both sets of coordinates are chosen such that both the profile curve and the spine curve are unit speed. From (2.4) and the relation between the variables we have

$$ds^2 = du^2 + (1 - \alpha(u, v))^2 dv^2 = (1 - \tilde{\alpha})^2 \left(\frac{\partial \tilde{v}}{\partial u} \right)^2 du^2 + \left(\frac{\partial \tilde{u}}{\partial v} \right)^2 dv^2,$$

where $\partial \tilde{u} / \partial v$ depends only on v . Hence $(1 - \alpha)^2$ depends only on v , and is constant equal to 1, given that the spine curve $\mathbf{r}(v)$ is unit speed. Thus, $ds^2 = du^2 + dv^2$, which implies that the surface is developable. A developable surface is locally a ruled surface, and, at non-umbilical points, the rulings are lines of curvature. Thus one of the families of planar geodesics consists of straight line segments. Since these line segments are also *parallel*, due to the second Monge surface representation, the surface U is a cylinder.

Now suppose that p is an umbilical point where the Gaussian curvature $K(p)$ at p is non-zero: then K must be non-zero on a neighbourhood W of p . By the discussion above, all of the points in W must be umbilic, because non-umbilical points have Gaussian curvature zero. Hence W is part of a sphere, with constant curvature K_0 . Moreover, the set of points at which the Gauss curvature is K_0 is clearly both open and closed, hence the whole of S .

The remaining case is that p is an umbilic and $K(p) = 0$. Since S is not part of a sphere, there can (by the above) be no umbilical points with positive

Gaussian curvature, and hence the Gauss curvature must be zero everywhere. Choose a neighbourhood U of p that has a Monge surface representation $\mathbf{x}(u, v)$. Since $K = 0$, we can assume that U is a part of a ruled surface. Then either U is a part of a plane and the proof is complete, or there exists a point q in U which is not an umbilic point. In the latter case, as previously discussed, the rulings, on a neighbourhood of q , are parallels of the Monge representation. Since a parallel curve to a straight line segment is also a straight line segment, it follows that all of the parallels on U are parallel straight line segments, and U is a cylinder. \square

2.2. Global planar geodesic foliations

We now consider global planar geodesically foliated surfaces. In general, a PGF surface does not have a global representation as a Monge surface. Figure 2 shows that it is easy to construct counterexamples: the developable surface to the left, although orientable, does not have a well-defined vector-field orthogonal to the foliation, whereas a Monge surface does, given by the direction of the parallels. The surface to the right has a different profile curve on one end of the surface from the other.




FIGURE 2. PGF surfaces that are not globally Monge surfaces..

Neither of the PGF surfaces in Figure 2 is *complete*. If we add this requirement, then such examples are ruled out:

Theorem 2.4. *Let S be a complete, connected, orientable PGF surface. Then S has a global parameterization in the form (2.1). In particular, this parameterization is a covering map $\mathbb{R}^2 \rightarrow S$ consisting of curvature line coordinates.*

Proof. Let γ_0 be a choice of one of the planar geodesics in the foliation, with a given arc-length parameterization. Denote the surface normal by N and let $\nu = N \times \gamma'_0$ denote the unit conormal along γ_0 . Such a conormal vector field is defined along any of the geodesics in the foliation, and we can always assume that, on neighbouring geodesics, the parameterizations are chosen so that ν varies continuously. Hence, given a point $p_0 = \gamma_0(s_0)$, we can integrate the vector field ν to obtain a curve \mathbf{r} in S , orthogonal to the foliation. Since S is complete and ν has unit length, \mathbf{r} is well-defined $\mathbb{R} \rightarrow S$. Let \tilde{S} be the Monge surface constructed from \mathbf{r} and γ_0 by the equation (2.1). This Monge surface is regular because a Monge surface is always regular on an

open set containing the spine curve, and the completeness of S , together with uniqueness of geodesics, means that \tilde{S} is a subset of the regular surface S . Now \tilde{S} is open in S , because any point on \tilde{S} has a neighbourhood parameterized in the form (2.1) with $(u, v) \in (u_0 - \varepsilon, u_0 + \varepsilon) \times (v_0 - \delta, v_0 + \delta)$. To show that \tilde{S} is the whole of S , consider an arbitrary point $q \in S$. Choosing the planar geodesic of the foliation that passes through q , we can similarly construct another Monge surface \hat{S} that contains q . If \hat{S} intersects \tilde{S} , then there is a planar geodesic that is contained in both \hat{S} and \tilde{S} . In this case, it follows by the construction of these sets that $\hat{S} = \tilde{S}$. Thus \hat{S} and \tilde{S} are either equal or disjoint. But both sets are open and S is connected, so they must be equal, and $q \in \tilde{S}$. By construction, the representation \tilde{S} is a regular immersion, hence a covering map $\mathbb{R}^2 \rightarrow S$. \square

Since the (u, v) coordinates of the covering map in Theorem 2.4 are canonically defined by the planar geodesic foliation, up to coordinate changes of the form $(u, v) \mapsto (\tilde{u}(u), \tilde{v}(v))$, this can be regarded as a canonical covering by the conformal Lorentz plane $\mathbb{R}^{1,1}$. Hence:

Corollary 2.5. *Every complete, connected, orientable PGF surface carries a Lorentzian conformal structure, the null lines of which are given by the planar geodesics and the family of parallels of a Monge surface parameterization.*

Note that not all planar geodesic foliations carry a global Lorentz structure (see Figure 2, left).

The Poincaré index theorem states that on a compact surface the Euler characteristic is equal to the sum of the indices of the zeros of any vector field. Since a surface satisfying the conditions of the above corollary carries a natural non-vanishing vector field, namely the tangent field to the foliation, this implies:

Proposition 2.6. *If M is a connected, compact orientable surface and $\mathbf{x} : M \rightarrow \mathbb{R}^3$ a PGF immersion, then M is a torus.*

3. Generalized Monge surfaces

We have shown that a PGF surface is not necessarily a Monge surface, unless it is complete. In this section we explore the converse direction, in order to clarify the precise difference between Monge surfaces and PGF surfaces, and find global conditions under which these objects coincide.

The idea of a Monge surface is defined kinematically by a plane $\Pi(t)$ moving through space such that the movement of any point on the surface is always orthogonal to the plane. We can describe such a moving plane by making an initial choice Q_1, Q_2 of orthonormal basis for the plane, and setting $\mathbf{q}_i(t)$ to be the corresponding vectors at time t . Similarly, making an initial choice of point p_0 on the plane, we let $p(t)$ be the corresponding point at time t . Finally, we set $\mathbf{t}(t)$ to be the normal to the plane at time t . By the definition of $\Pi(t)$, the derivatives of p, \mathbf{q}_1 and \mathbf{q}_2 have components only in

the direction of \mathbf{t} . Thus we have a map $F : (a, b) \rightarrow SE(3)$, from some open interval into the group of orientation preserving Euclidean motions, given by

$$F = \begin{pmatrix} \mathbf{q}_1 & \mathbf{q}_2 & \mathbf{t} & p \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

and the conditions on the derivatives are represented by

$$\omega = F^{-1} \frac{dF}{dt} = \begin{pmatrix} 0 & 0 & -\kappa_1 & 0 \\ 0 & 0 & -\kappa_2 & 0 \\ \kappa_1 & \kappa_2 & 0 & \lambda \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

We call the family of planes $\Pi(t)$, defined by F an *orthogonal family of planes*, and say that the family is *regular* if $(\kappa_1(t), \kappa_2(t), \lambda(t)) \neq (0, 0, 0)$, i.e., $\omega \neq 0$, for all $t \in (a, b)$. This definition of regularity does not depend on the choice of initial vectors Q_1 and Q_2 or the initial point p_0 . A different choice amounts to post-multiplying the frame F by a constant invertible matrix, and thus ω is conjugated by a constant invertible matrix, which does not change the condition $\omega \neq 0$.

Example 1. Given an arbitrary regular space curve \mathbf{r} , a regular orthogonal family of planes can be constructed by $p(t) = \mathbf{r}(t)$, $\mathbf{t} = \mathbf{r}'/|\mathbf{r}'|$ and $\mathbf{q}_1, \mathbf{q}_2$ a rotation minimizing frame. In that case λ is non-vanishing. More generally, there exist regular orthogonal families of planes where λ vanishes, regardless of the choice of initial point p_0 . For example, take

$$\begin{aligned} \mathbf{q}_1(t) &= (\cos t, \sin t, 0), & \mathbf{q}_2(t) &= (0, 0, 1), \\ \mathbf{t}(t) &= (-\sin t, \cos t, 0), & p(t) &= \mathbf{t}(t) - t\mathbf{t}'(t). \end{aligned}$$

This is a regular orthogonal family of planes, with $\kappa_1(t) = -1$, $\kappa_2 = 0$ and $\lambda(t) = t$. Here $\lambda(t)$ vanishes at $t = 0$. If we choose a different initial point \tilde{p}_0 on the plane, we have $\tilde{p}(t) = p(t) + c_1\mathbf{q}_1(t) + c_2\mathbf{q}_2(t)$ for some constants c_1 and c_2 . Then $\tilde{p}'(t) = (t + c_1)\mathbf{t}(t)$, so $\lambda(t)$ vanishes at $t = -c_1$.

Definition 3.1. Let $\gamma : (a, b) \rightarrow \mathbb{R}^2$ be a curve parameterized by arc-length, $\gamma(s) = (x(s), y(s))$, and let $F : (c, d) \rightarrow SE(3)$ be a regular orthogonal family of planes, with notation as above. Then the map $\mathbf{x} : (a, b) \times (c, d) \rightarrow \mathbb{R}^3$ given by

$$\mathbf{x}(u, v) = p(v) + x(u)\mathbf{q}_1(v) + y(u)\mathbf{q}_2(v),$$

is called a (*generalized*) *Monge surface*.

It should be noted that, if the initial point p_0 and the initial frame (Q_1, Q_2) are changed in the representation F for the family $\Pi(t)$, then the curve γ would need to be translated and rotated accordingly to get the same surface.

Since

$$\frac{\partial \mathbf{x}}{\partial u} = x'(u)\mathbf{q}_1(v) + y'(u)\mathbf{q}_2(v), \quad \frac{\partial \mathbf{x}}{\partial v} = (\lambda(v) - \kappa_1(v)x(u) - \kappa_2(v)y(u))\mathbf{t},$$

with $(x')^2 + (y')^2 = 1$, a Monge surface is regular at (u, v) if and only if

$$\lambda(v) \neq \kappa_1(v)x(u) + \kappa_2(v)y(u).$$

Geometrically, in each plane $\Pi(v)$, there is a (possibly empty) line, the instantaneous axis of rotation of the plane $\Pi(v)$,

$$L_v = \{(x, y) \mid \kappa_1(v)x + \kappa_2(v)y - \lambda(v) = 0\},$$

and the Monge surface is regular if and only if none of the lines L_v intersect the initial curve γ . Note that the spine curve p is a curve on the surface \mathbf{x} if and only if $(x(u), y(u)) = (0, 0)$ for some u . One can always arrange that p is a curve on the surface by choosing p_0 to be a point on the surface, and then translating γ accordingly.




FIGURE 3. Monge surfaces with the same singular spine curve, shown in red.

Example 2. Returning to the family of planes in Example 1, we have L_v given by the equation $x = v$. That is, the instantaneous axes of rotation are all vertical lines, and these lines sweep out the entire xy -plane as v runs over \mathbb{R} . Hence there is no choice of profile curve γ such that \mathbf{x} is everywhere a regular surface. Portions of the surfaces corresponding to three different lines as profile curve (the line $y = -x$, the x -axis and the y -axis) are shown in Figure 3. The first two have cuspidal edges, the last a fold singularity.

Note that the example of the sphere shows that the image of a generalized Monge surface can be a regular surface even if the parameterization \mathbf{x} is not regular. In general, however, there will be singularities that are not removable, as in Figure 3.

For a given generalized Monge surface, if the surface is regular then, by construction, it is a PGF surface. Combined with Theorem 2.4, we conclude:

Theorem 3.2. *Every regular Monge surface is a PGF surface, and every complete PGF surface has a covering by a Monge surface. In particular, complete, regular PGF surfaces are the same objects as complete regular Monge surfaces.*

4. Monge tori and Klein bottles

In this section we investigate the problem of constructing *Monge tori*. These are complete, regular surfaces, so can be regarded as either Monge surfaces or PGF surfaces. If the spine curve is a *closed* curve, i.e., a map $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^3$, satisfying $\mathbf{r}(v + P) = \mathbf{r}(v)$ for some P , it does not follow that the surface closes up to form a topological cylinder. Apart from very special cases such as a *tubular surface* (Figure 4, right), it is necessary that the surface normal, and hence the rotation minimizing frame, be periodic with the same period. This is a property of the geometry of the spine curve, and it has a very simple characterization for curves that have a well-defined Frenet frame.

4.1. Curves with closed rotation minimizing frames and Monge tori

Throughout this section, by a differentiable curve $\mathbf{r} : I \rightarrow \mathbb{R}^3$, we mean a regular unit-speed curve that has a well-defined differential binormal field \mathbf{b} along the curve. The principal normal is then defined by the formula $\mathbf{n} = \mathbf{b} \times \mathbf{t}$, where $\mathbf{t} = \mathbf{r}'(s)$. The signed curvature is given by $\mathbf{t}'(s) = \kappa(s)\mathbf{n}(s)$, and the torsion by $\mathbf{b}'(s) = \tau(s)\mathbf{n}(s)$. Given a closed curve with period L , denote the total torsion by $T = \int_0^L \tau(s)ds$. Note that the assumed binormal field is not automatically periodic: for example a closed curve could contain pieces of line segments, along which the binormal field could be anything at all. We therefore add the assumption of periodicity of the binormal field:

Lemma 4.1. *Let $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^3$ be a closed, differentiable curve with period L , and suppose that the binormal field \mathbf{b} is also periodic with the same period. Let $F = (\mathbf{t}, \mathbf{q}_1, \mathbf{q}_2)$ be any choice of rotation minimizing frame along \mathbf{r} . Then F is periodic with period L if and only if the total torsion T is an integer multiple of 2π .*

Proof. Because \mathbf{t} and \mathbf{b} are periodic with the same period, the Frenet frame $(\mathbf{t}, \mathbf{n}, \mathbf{b})$ also has this property. The torsion τ of the curve, given by $\mathbf{b}'(s) = \tau(s)\mathbf{n}(s)$, is the rate of rotation of the basis (\mathbf{n}, \mathbf{b}) for the plane $\Pi_s = \text{span}(\mathbf{n}, \mathbf{b})$ compared with the zero rotation of the rotation minimizing basis $(\mathbf{q}_1, \mathbf{q}_2)$. The latter basis (and hence F) thus closes up if and only if $2m\pi = \int \tau ds$, for some $m \in \mathbb{Z}$. \square

Let us define a *Monge cylinder* to be a regular Monge surface with closed spine curve and closed rotation minimizing frame. Note that this definition does not depend on the choice of spine curve: if \mathbf{r} is the spine curve, then any other spine curve is of the form $\tilde{\mathbf{r}}(s) = \mathbf{r}(s) + a\mathbf{q}_1(s) + b\mathbf{q}_2(s)$, which has the same properties. Given a Monge cylinder, it is a simple matter to construct a Monge torus by choosing as profile curve a small closed curve γ_0 in the plane $\Pi_0 = \text{span}(\mathbf{q}_1(0), \mathbf{q}_2(0))$, such that the origin lies on γ_0 . The Monge surface so constructed is regular on an open set containing the spine curve, which lies on the surface. Since the image of the spine curve is compact, it is always possible to choose γ_0 small enough so that the torus produced is regular.

4.2. Constructing Monge tori

Given the characterization in Lemma 4.1, it is not difficult to find examples of Monge cylinders, and hence tori. For example, a plane curve has zero torsion, but more generally, one can show [11] that a surface in \mathbb{R}^3 is a part of a sphere or a plane if and only if the total torsion of any closed curve in the surface is zero. Hence any close curve in a sphere can also be used to produce Monge cylinders.

The problem of constructing more general examples amounts to finding more general curves that satisfy the closing condition of Lemma 4.1. We are also interested in the case that the total torsion is an *odd* multiple of π , because this allows us to construct Möbius strips and Klein bottles. More generally still, since the total torsion is the total rotation angle of the rotation minimizing frame as the curve is traversed, by taking a profile curve γ_0 that has a rotational symmetry of order n around the origin, and using a spine curve with total torsion $2k\pi/n$ (where k is an integer), the Monge surface will close up after one period of the loop, although the Monge parameterization will be a covering of degree $d \leq n$, depending on the common divisors of n and k . If the total torsion is an irrational multiple of π , then a *circle* centered at the origin can be used as profile curve, to produce a *tubular* surface that is not a parameterized Monge torus, but the image of which is a torus. Note that the spine curve does not lie on the surface in this case, but at the center of the tube. Examples constructed using the method described in the next subsection are shown in Figure 4.




FIGURE 4. Monge coverings of embedded tori. Left: $T = L_S = 2\pi/4$, degree of covering is 4. The profile curve has order 4 symmetry about the origin of the plane. Right: $T = L_S = 2\pi/\pi$, degree of covering is uncountably infinite. The profile curve is a circle, the parallels do not close up.

It is thus interesting to know whether one can find closed curves with arbitrary values for the total torsion, or at least with arbitrary values in $[0, 2\pi)$. The answer to this question can be found in a survey by Fenchel [7], the relevant details of which we recall here: A convenient way to find curves with a specified value for the total torsion is to look for a curve with everywhere *positive* torsion, because if s is the arc-length on a curve \mathbf{r} and s_B the arc-length of the binormal \mathbf{b} then $ds_B = |\tau(s)|ds$. Hence, if $\tau > 0$,

and L_S is the length of the trace of \mathbf{b} on \mathbb{S}^2 , then

$$T = \int_0^L \tau(s) ds = \int_0^{L_S} dS_B = L_S.$$

In other words, if τ is non-vanishing, then the total torsion is just the length of the trace of the binormal curve on \mathbb{S}^2 . Fenchel also shows that a closed curve \mathbf{b} in \mathbb{S}^2 is the binormal to a *closed* curve \mathbf{r} in \mathbb{R}^3 if and only if the great circles tangent to \mathbf{b} sweep out the whole of the 2-sphere. Moreover, the torsion of \mathbf{r} is non-vanishing if \mathbf{b} has no cusps. There are many examples of such curves \mathbf{b} (see Figure 5), and curves such as those shown can clearly be scaled so that the length is anything between 0 and 2π , which solves the problem under discussion.




FIGURE 5. Examples of binormal images of closed space curves.

Note that we only obtain curves \mathbf{r} with non-vanishing torsion this way, and this does not include all interesting cases. For example if \mathbf{r} takes values in a sphere then the total torsion is zero. Hence the torsion must change sign, unless \mathbf{r} is a plane curve. It is possible to modify our discussion to include these by including curves \mathbf{b} that have cusps, but we omit this for the sake of simplicity.

4.3. Explicit constructions

Given a curve \mathbf{b} in the 2-sphere with the properties discussed above, there are infinitely many closed space-curves \mathbf{r} to choose from that have \mathbf{b} as the binormal field, so we would like to consider some natural choices. For example closed curves with non-vanishing curvature and *constant* torsion are found in work by J. Weiner [12], and Bates and Melko [1], however to construct these a special choice of \mathbf{b} is needed. Here we take \mathbf{b} to be arbitrary, subject to the conditions mentioned above.

Let $\mathbf{b}(t)$ be some parameterization of a regular closed curve in \mathbb{S}^2 , with period P , and the length of one period L_S . If \mathbf{b} is the binormal field to a space curve \mathbf{r} with unit tangent and normal fields \mathbf{t} and \mathbf{n} respectively, then \mathbf{b}' is proportional to \mathbf{n} , so $\mathbf{t} = \mathbf{n} \times \mathbf{b} = (\mathbf{b}'/|\mathbf{b}'|) \times \mathbf{b}$. If s is the arc-length parameter of \mathbf{r} , then

$$\mathbf{t} = \frac{1}{\tau} \frac{dt}{ds} \frac{d\mathbf{b}}{dt} \times \mathbf{b},$$

and

$$\mathbf{r} = \int \mathbf{t} \frac{ds}{dt} dt = \int \frac{1}{\tau} \frac{d\mathbf{b}}{dt} \times \mathbf{b} dt.$$

Given a choice of \mathbf{b} , a curve \mathbf{r} with binormal field \mathbf{b} is determined by any choice of function $\sigma(t) := 1/\tau(t)$. The closing condition for \mathbf{r} is

$$\int_0^P \sigma \frac{d\mathbf{b}}{dt} \times \mathbf{b} dt = 0, \quad (4.1)$$

and clearly many choices for σ will be available. To find a solution, we can set

$$\sigma(t) = \sum_{i=1}^n c_i B_i(t),$$

where B_i are some periodic basis functions, such as trigonometric functions or periodic B-splines. Then (4.1) is:

$$\sum_{i=1}^n c_i \mathbf{a}_i = 0, \quad \mathbf{a}_i = \int_0^T B_i(t) \mathbf{b}(t) \times \mathbf{b}'(t) dt.$$

Thus, given a choice of basis functions, the solutions are found by linear algebra. There can be many solutions for c_i , depending on the choice of basis functions. To choose a natural solution, we have used an optimization to minimize the elastic energy, $\int_0^L \kappa^2(s) ds$. To write this quantity in terms of \mathbf{b} and σ , let s_T denote the arc-length parameter of the curve \mathbf{t} and compute

$$\kappa = \tau \left| \mathbf{b} \times \frac{d^2 \mathbf{b}}{ds_B^2} \right| = \tau \left| \det \left(\mathbf{b} \frac{d\mathbf{b}}{ds_B} \frac{d^2 \mathbf{b}}{ds_B^2} \right) \right|,$$

and hence

$$\mathcal{E} = \int_0^L \kappa^2 ds = \int_0^T \frac{1}{\sigma} \frac{\det(\mathbf{b}\mathbf{b}'\mathbf{b}'')^2}{|\mathbf{b}'|^5} dt.$$

4.4. Concluding remarks

We have used the method just described to produce the examples shown in Figure 4. More examples can be found in the article [3]. If we scale \mathbf{b} to have length π , and use a profile curve that passes through the origin and is symmetric about the origin, then a Möbius strip is obtained. If the profile curve is a straight line the surface is developable. The problem of constructing developable Möbius strips has been considered, for example, in [13] and [4]. The method of [4] is, in essence, similar to that presented here, only the authors assume that the curvature of \mathbf{r} is non-vanishing. In [10], developable Möbius strips are characterized by a different method, using the so-called *center geodesic curve* instead of the spine curve.

If we take the spine curve of a developable Möbius strip, and take the profile curve to be a symmetric figure-8 curve, we obtain a Monge Klein bottle (see [3]). MATLAB functions that allow one to compute a Monge surface from a given spine curve and profile curve are currently available at <http://geometry.compute.dtu.dk/software>.

References

- [1] L Bates and O Melko. On curves of constant torsion I. *J. Geom.*, 104:213–227, 2013.
- [2] R Bishop. There is more than one way to frame a curve. *Amer. Math. Monthly*, 82:246–251, 1975.
- [3] D Brander and J Gravesen. Surfaces foliated by planar geodesics: A model for curved wood design. In D Swart, C Séquin, and K Fenyvesi, editors, *Proceedings of Bridges 2017: Mathematics, Art, Music, Architecture, Education, Culture*, pages 487–490, Phoenix, Arizona, 2017. Tessellations Publishing. Available on-line at <http://archive.bridgesmathart.org/2017/bridges2017-487.pdf>.
- [4] C Chicone and N Kalton. Flat embeddings of the Möbius strip in R^3 . *Comm. Appl. Nonlinear Anal.*, 9:31–50, 2002.
- [5] G Darboux. *Leçons sur la théorie générale des surfaces*, volume I. Gauthier-Villars, Paris, 1887.
- [6] L Eisenhart. *A treatise on the differential geometry of curves and surfaces*. The Atheneum Press, Boston, 1909.
- [7] W Fenchel. On the differential geometry of closed space curves. *Bull. Amer. Math. Soc.*, 57:44–54, 1951.
- [8] G Monge. *Application de l'analyse a la géométrie*. Bachelier, Paris, fifth edition, 1850.
- [9] L Raffy. Sur les surfaces à lignes de courbure planes, dont les plans enveloppent un cylindre. *Ann. de l'Éc. Norm. (3)*, 18:343–370, 1901.
- [10] T Randrup and P Røgen. Sides of the Möbius strip. *Arch. Math.*, 66:511–521, 1996.
- [11] W Scherrer. Eine Kennzeichnung der Kugel. *Vierteljschr. Naturforsch. Ges. Zürich*, 85:40–46, 1940.
- [12] J Weiner. Closed curves of constant torsion. II. *Proc. Amer. Math. Soc.*, 67:306–308, 1977.
- [13] W Wunderlich. Über ein abwickelbares Möbiusband. *Monatsh. Math.*, 66:276–289, 1962.

David Brander

Department of Applied Mathematics and Computer Science,
Technical University of Denmark
Matematiktorvet, Building 303 B
DK-2800 Kgs. Lyngby
Denmark
e-mail: dbra@dtu.dk

Jens Gravesen

Department of Applied Mathematics and Computer Science,
Technical University of Denmark
Matematiktorvet, Building 303 B
DK-2800 Kgs. Lyngby
Denmark
e-mail: jgra@dtu.dk