

Technical University of Denmark



## A new method for estimating transmission rates of mastitis-causing pathogens

Kirkeby, Carsten Thure; Hisham Beshara Halasa, Tariq; Gussmann, Maya Katrin; Græsbøll, Kaare

*Publication date:*  
2018

[Link back to DTU Orbit](#)

*Citation (APA):*

Kirkeby, C., Halasa, T., Gussmann, M. K., & Græsbøll, K. (2018). A new method for estimating transmission rates of mastitis-causing pathogens. Poster session presented at National Mastitis Council Annual Meeting, Tucson, United States.

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# A New Method for Estimating Transmission Rates of Mastitis-causing Pathogens

Carsten Kirkeby<sup>1</sup>, Tariq Halasa<sup>1</sup>, Maya Gussmann<sup>1</sup>, Kaare Græsbøll<sup>1</sup>

<sup>1</sup> Section for Epidemiology, National Veterinary Institute, Technical University of Denmark

## Background

- Epidemiological **simulation models** can predict effects of **control strategies for mastitis** if they rely on decent data
- The transmission rate controls the **spread of contagious pathogens** between animals in the herd.
- Transmission rates are estimated from **expensive** and **time-consuming** field studies testing all lactating cows repeatedly, often using large sampling intervals.
- Poisson regression** is often used to estimate the transmission rate, but may perform poorly at large sampling intervals

## Methods

- We derived a **new mathematical formula** for estimating the transmission rate. This new method is suitable for systems in equilibrium using the number of susceptible and infected animals directly **without regression analysis**. The new method allows for **multiple infection and recovery events between sampling points**, unlike Poisson regression.
- We used an existing simulation model to simulate transmission of ***S. aureus*** in a dairy herd using different transmission rates.
- We sampled the lactating cows in the simulated herd with different sampling intervals and **estimated the transmission rate** with traditionally used **Poisson regression (P)** and a **new method (N)**.
- The results were compared to determine whether one method performed **significantly better** than the other.





Figure 1: The optimal method for the true transmission rate and sampling interval. Letters show whether Poisson regression (P) or the New method (N) performed significantly better than or equally good as Poisson regression (P/N).



## Results

- The new method performed **equally well** to Poisson regression in most of the simulated scenarios, but **both models** showed areas where they were better than the other (Fig. 1).
- The new method proved more suitable for estimating transmission rates when **sampling with large intervals**, which can often be the case in field studies (green area).

## Take home message

- We developed a **new method to estimate transmission rates** from field studies of endemic contagious diseases like **mastitis**.
- The new method **performs better** than Poisson regression at **large sampling intervals**.
- Read more** in Scientific Reports 7, 9496 (2017) doi:10.1038/s41598-017-09209-x

