

HETEROCYCLES, Vol. , No. , , pp. -. © The Japan Institute of Heterocyclic Chemistry
Received, , Accepted, , Published online, .
DOI: 10.3987/COM- (Please do not delete.)

REGIOSELECTIVITY OF THE PALLADIUM-MEDIATED INTRAMOLECULAR COUPLING REACTION OF HIGHLY OXYGENATED PHENYL BENZOATE DERIVATIVES AND APPLICATION TO THE SYNTHESIS OF ALTERTENUOL

Takuya Matsukihira, Shumpei Saga, Yoshikazu Horino, and Hitoshi Abe*

Graduate School of Science and Engineering, University of Toyama, Gofuku,
Toyama 930-8555, Japan
abeh@eng.u-toyama.ac.jp

Abstract – The regioselectivity of the intramolecular biaryl coupling reaction of 3',4'-dialkoxyphenyl 2,4-dimethoxybenzoates was investigated using a palladium reagent, and transition state models of the reaction are proposed. As an application of this study, a short-step synthesis of alvertenuol is also performed.


INTRODUCTION

A palladium-mediated intramolecular biaryl coupling reaction of phenyl benzoate derivatives is a useful transformation for the preparation of the 6*H*-dibenzo[*b,d*]pyran-6-one skeleton.¹ There have been many studies of the total syntheses of natural products possessing this type of heterocyclic system, using the intramolecular biaryl coupling reaction.² In some cases, regioselectivity is a crucial concern in the intramolecular biaryl coupling, thus we recently reported several types of reactions involving the issue of regioselectivity.³ In this report, as a series of our study, regioselectivity is examined in the Pd-mediated intramolecular biaryl coupling reaction of 3',4'-dialkoxyphenyl 2,4-dimethoxybenzoates, and the synthesis of a natural polyphenolic compound, alvertenuol,⁴ is also described.

RESULTS AND DISCUSSION


As the substrates for the investigation, we prepared three kinds of phenyl benzoate derivatives **1a-1c**, which were easily obtained by a simple esterification between the corresponding phenols **2a-2c** and benzoic acid **3**. Initially, the regioselectivity was investigated using Ag₂CO₃ as the base (Table 1). For the ester **1a**, when Ph₃P was employed as the ligand, the *ortho*-product **5a** mainly generated. However, a poor regioselectivity for *para*- and *ortho*-products, **4a** and **5a**, was observed under no ligand condition (entries

1 and 2). On the contrary, for the substrates **1b** and **1c**, a dramatic change was observed in the product ratio. Namely, when using Ph_3P , the regioselectivity exhibited 1.4-0.7:1, although when using no ligand, the *para*-isomers **4b** and **4c** were predominantly generated (7.9-8.8:1, entries 3-6).



In contrast to the results above, employing K_2CO_3 as the base led to a different outcome for the product ratio. When several phosphine ligands⁵ were attempted with the ester **1a**, the regioselectivity was almost identical to the result in entry 1 (entries 7-9). On the other hand, a slightly increased ratio was observed when using no phosphine ligand (entry 10). In the case of both **1b** and **1c**, a moderate selectivity was observed in the range of 1.2-3.6:1 under the condition that the phosphine ligand was employed (entries 11, 13, 14, and 15). When no ligand was used for **1b** and **1c**, the reaction proceeded with the product ratio of 4.4-4.5:1 (entries 12 and 16).


Table 1. Reaction of Phenyl Benzoate Derivatives **1a-1c**



entry	substrate	ligand	yield (%)		ratio ^{b)}	entry	substrate	ligand	yield (%)		ratio ^{b)}
			4	5					4	5	
1	1a	Ph_3P	77		0.2 : 1	7	1a	XPhos ^{c)}	89		0.3 : 1
2	1a	none	79		1.8 : 1	8	1a	Ph_3P	81		0.2 : 1
3	1b	Ph_3P	92		1.4 : 1	9	1a	ⁿ Bu_3P	77		0.2 : 1
4	1b	none	89		8.8 : 1	10	1a	none	96		0.5 : 1
5	1c	Ph_3P	99		0.7 : 1	11	1b	Ph_3P	81		2.5 : 1
6	1c	none	80		7.9 : 1	12	1b	none	84		4.4 : 1
a) Ag_2CO_3 was used as the base in entries 1-6.											
b) Determined by ¹ H-NMR analysis.											
c) Ref. 5											
						13	1c	XPhos ^{c)}	79		3.6 : 1
						14	1c	Ph_3P	89		1.7 : 1
						15	1c	ⁿ Bu_3P	81		1.2 : 1
						16	1c	none	77		4.5 : 1


For the mechanistic aspect, we would be able to explain the regioselectivity by considering the transition state models.³ⁱ Based on Echavarren's intermolecular-assisted mechanism,⁶ the transition state of substrate **1a** can be illustrated as Figure 1. *A1* and *A2* describe the models in the case of using Ag_2CO_3 as the base. Strong affinity between the silver and the iodo atom causes a partial cationic character on the palladium. Therefore, the alkoxy oxygen can coordinate to the palladium atom (*A2*) leading to providing the *ortho*-products **5a-5c**, whereas the steric hindrance, which was generated between the alkoxy group and the ligand on the palladium atom, is also significant (*A2*; L = phosphine). Consequently, the product ratios are controlled by the balance of both the coordinating power and the steric factor. However, when using no phosphine ligand, the tight solvation of the DMA takes place to make the surrounding of the palladium to be bulkier (*A1* and *A2*; L = DMA). Thus, no phosphine ligand condition leads to the severe steric repulsion between the alkoxy group and DMA, as shown in *A2*, so that the ratio of the *para*-products **4a-4c** is increased.

Figure 1. Transition State Models under Ag_2CO_3 Condition




Contrary to the above, *B1-B3* illustrate the transition state models under the condition using K_2CO_3 as the base (Figure 2). In these cases, the cationic property of the palladium atom may not be considered because of the hard character of the potassium ion. The coordinating effect of the alkoxy group to the palladium is not an important factor for the substrates **1b** and **1c**, thus the *para*-isomers **4b** and **4c** are slightly predominant (*B1* and *B2*). Since the coordinating ability of the methylenedioxy group of the substrate **1a** is stronger than **1b** and **1c**, the attractive interaction between the oxygen and the palladium can control and mainly produce the *ortho*-isomer **5a** (*B3*).

Figure 2. Transition State Models under K_2CO_3 Condition



Since the coupling product **4c** was regioselectively obtained with the conditions of entry 6 in Table 1, we attempted a further transformation toward the natural product, alternenuol **6**, which was found from *Alternaria tenuis* in 1957.⁷ The selective demethylation of **4c** by the Node's method⁸ was successful to afford **7**, which was debenzylated by the conventional hydrogenolysis reaction leading to **6**. As an alternative way, partial dealkylation with BBr_3 proceeded effectively to produce **6** in a single step. The spectral data of synthetic **6** were in accordance with the reported data.⁴

Scheme 1. Transformation of **4c** into Altertenuol (**6**)



CONCLUSION

The regioselectivity of the Pd-mediated biaryl coupling reaction of 3',4'-dialkoxyphenyl 2,4-dimethoxybenzoates was intensively investigated. It was found that the combination of the ligand and the base influences the regioselectivity of the reaction. We also demonstrated the synthesis of altertenulol using the regioselective condition. This method would be utilized for further application to other syntheses of biaryl-type natural products.

EXPERIMENTAL

General Information

Melting points were measured using a Yanagimoto micro melting point hot-plate and are uncorrected. The IR spectra were recorded using a SHIMADZU FTIR-8400 spectrophotometer. The NMR spectra were obtained using a JEOL α -400 (400 MHz) instrument. The chemical shifts are given in δ ppm with TMS as an internal standard. The elemental analyses were performed using a Thermo Scientific FlashEA1112 analyzer. Silica gel column chromatography was carried out using a wakogel[®] C-200 or C-400HG (Wako).

3,4-Dimethoxyphenol (**2b**)⁹

A mixture of THF (2 mL), 31% H₂O₂ (0.13 mL), H₃BO₃ (189 mg, 3.05 mmol), and conc. H₂SO₄ (0.06 mL) was stirred for 30 min at rt. To the mixture was added a solution of 3,4-dimethoxybenzaldehyde (101 mg, 0.607 mmol) in THF (1 mL). The mixture was stirred for 19 h at rt, then a saturated NaHCO₃ aqueous solution was added. The mixture was extracted with CHCl₃ and the organic layer was washed with brine. After drying with MgSO₄ and evaporation, a residue was subjected to silica gel column chromatography with hexane-AcOEt (4:1) to give **2b** (73.1 mg, 78%) as colorless prisms, mp 83.9-85.2 °C (hexane-CHCl₃) [lit.⁹ mp 79 °C]. ¹H-NMR (400 MHz, CDCl₃) δ : 6.72 (1H, d, J = 8.8 Hz, ArH), 6.47 (1H, d, J = 2.7 Hz, ArH), 6.35 (1H, dd, J = 8.6, 2.7 Hz, ArH), 5.39 (1H, s, ArOH), 3.81 (3H, s, ArOMe), 3.79 (3H, s, ArOMe).

3,4-Dibenzyloxyphenol (**2c**)

A mixture of THF (2 mL), 31% H₂O₂ (0.07 mL), H₃BO₃ (100 mg, 1.62 mmol), and conc. H₂SO₄ (0.032 mL) was stirred for 30 min at rt. To the mixture was added a solution of 3,4-dibenzyloxybenzaldehyde (102 mg, 0.320 mmol) in THF (1.5 mL). The mixture was stirred for 19 h at rt, then a saturated NaHCO₃ aqueous solution was added. The mixture was extracted with CHCl₃ and the organic layer was washed with brine. After drying with MgSO₄ and evaporation, a residue was subjected to silica gel column chromatography with hexane-AcOEt (7:3) to give **2c** (86.8 mg, 88%) as a colorless solid, mp 117.4-118.8

°C (hexane-CHCl₃). ¹H-NMR (400 MHz, CDCl₃) δ: 7.44-7.28 (10H, m, ArH), 6.79 (1H, d, *J* = 8.6 Hz, ArH), 6.49 (1H, d, *J* = 3.0 Hz, ArH), 6.29 (1H, dd, *J* = 8.7, 2.8 Hz, ArH), 5.09 (2H, s, ArOCH₂), 5.07 (2H, s, ArOCH₂), 4.83 (1H, s, ArOH).

3,4-Methylenedioxyphenyl 2-iodo-4,6-dimethoxybenzoate (1a)

Under an N₂ atmosphere, a solution of 2-iodo-4,6-dimethoxybenzoic acid^{3d} (104 mg, 0.337 mmol), sesamol (50.7 mg, 0.367 mmol), EDC (90.9 mg, 0.474 mmol), and DMAP (40.7mg, 0.333 mmol) in dry CH₂Cl₂ (3 mL) was stirred for 1 h at rt. The mixture was washed with water and brine, then the organic layer was dried over MgSO₄. After evaporation, the resulting residue was subjected to silica gel column chromatography with chloroform to give **1** (90.2 mg, 63%). Colorless prisms, mp 110.2-110.9 °C (hexane-CH₂Cl₂). IR (KBr) cm⁻¹: 1742, 1593, 1562, 1503, 1478, 1459, 1438, 1408, 1309, 1287, 1243, 1220, 1188, 1169, 1144, 1114, 1097, 1068, 1023, 938, 926, 881, 841, 828, 807, 791, 597. ¹H-NMR (400 MHz, CDCl₃) δ: 6.95 (1H, d, *J* = 2.2 Hz, ArH), 6.84-6.81 (2H, t, ArH), 6.75 (1H, dd, *J* = 8.4, 2.3 Hz, ArH), 6.48 (1H, d, *J* = 1.9 Hz, ArH) 6.00 (2H, s, ArOCH₂OAr), 3.85 (3H, s, ArOMe), 3.82 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ: 166.3, 162.0, 158.2, 148.1, 145.7, 145.2, 122.4, 115.4, 114.2, 108.1, 103.9, 101.8, 99.1, 92.9, 56.2, 55.9. *Anal.* Calcd for C₁₆H₁₃IO₆: C, 44.88; H, 3.06. Found: C, 44.74; H, 2.94.

3,4-Dimethoxyphenyl 2-iodo-4,6-dimethoxybenzoate (1b)

In the same procedure mentioned above using **2b** (52.8 mg, 0.343 mmol), **1b** (63.4 mg, 44%) was obtained. Colorless prisms, mp 80.2-80.5 °C (hexane-CH₂Cl₂). IR (KBr) cm⁻¹: 1749, 1598, 1562, 1510, 1455, 1255, 1228, 1182, 1151, 1124, 1076, 1030. ¹H-NMR (400 MHz, CDCl₃) δ: 6.95 (1H, d, *J* = 2.2 Hz, ArH), 6.87 (2H, d, *J* = 1.5 Hz, ArH), 6.84 (1H, t, *J* = 1.5 Hz), 6.47 (1H, d, *J* = 2.2 Hz, ArH) 3.88 (3H, s, ArOMe), 3.87 (3H, s, ArOMe), 3.83 (3H, s, ArOMe), 3.80 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ: 166.3, 161.9, 158.1, 149.5, 147.2, 144.6, 122.5, 115.4, 113.0, 111.3, 105.9, 99.0, 92.9, 56.3, 56.2, 56.1, 55.8. *Anal.* Calcd for C₁₇H₁₇IO₆: C, 45.96; H, 3.86. Found: C, 45.84; H, 3.72.

3,4-Dibenzyloxyphenyl 2-iodo-4,6-dimethoxybenzoate (1c)

In the same procedure mentioned above using **2c** (4.04 g, 13.2 mmol), **1c** (4.87 g, 63%) was obtained. Colorless solid, mp 101.4-102.0 °C (Et₂O-CH₂Cl₂). IR (KBr) cm⁻¹: 1751, 1603, 1560, 1510, 1454, 1435, 1421, 1400, 1387, 1331, 1315, 1277, 1250, 1236, 1215, 1184, 1153, 1119, 1074, 1030, 1001, 989, 922, 883, 818, 804, 787, 770, 750, 731, 702, 594. ¹H-NMR (400 MHz, CDCl₃) δ: 7.47-7.29 (10H, m, ArH), 6.97-6.93 (3H, m, ArH), 6.84 (1H, dd, *J* = 8.7, 2.6 Hz, ArH), 6.47 (1H, d, *J* = 2.2 Hz, ArH), 5.18 (2H, s, ArOCH₂), 5.16 (2H, s, ArOCH₂), 3.82 (3H, s, ArOMe), 3.81 (3H, s, ArOMe). ¹³C-NMR (100 MHz,

CDCl₃) δ : 166.1, 161.9, 158.2, 149.6, 147.0, 145.1, 137.3, 137.0, 128.6, 128.6, 128.0, 127.9, 127.5, 127.5, 122.5, 115.6, 115.4, 114.0, 109.1, 99.1, 93.0, 72.0, 71.5, 56.2, 55.8. *Anal.* Calcd for C₂₉H₂₅IO₆: C, 58.40; H, 4.23. Found: C, 58.62; H, 4.15.

Typical Procedure for the Coupling Reaction (Table 1)

Under an N₂ atmosphere, a mixture of **1a** (100 mg, 0.234 mmol), Pd(OAc)₂ (0.023 mmol), ligand (0.047 mmol), base (0.023 mmol), and DMA (7 mL) was heated under reflux for 1 h with stirring. Unsolved materials were filtered off, then water was added to the filtrate. After extraction with CH₂Cl₂, the organic layer was washed with brine and dried over MgSO₄. Evaporation of the solvent gave a residue which was subjected to silica gel column chromatography with hexane-AcOEt-CH₂Cl₂ (1:1:3).

2,3-Methylenedioxy-7,9-dimethoxy-6H-benzo[*c*]chromen-6-one (4a)

Colorless needles, mp 240.5-241.0 °C (hexane-CH₂Cl₂). ¹H-NMR (400 MHz, CDCl₃) δ : 7.25 (1H, s, ArH), 6.83 (1H, d, *J* = 2.2 Hz, ArH), 6.75 (1H, s, ArH), 6.48 (1H, d, *J* = 2.2 Hz, ArH), 6.04 (2H, s, ArOCH₂OAr), 3.98 (3H, s, ArOMe), 3.96 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ : 165.5, 164.2, 158.1, 150.0, 148.1, 145.0, 139.7, 111.1, 102.9, 102.2, 101.2, 98.6, 98.3, 96.8, 56.5, 55.8. IR (KBr) cm⁻¹: 1719, 1603, 1574, 1504, 1472, 1454, 1433, 1381, 1344, 1254, 1211, 1180, 1146, 1092, 1063, 1030, 1005, 827. *Anal.* Calcd for C₁₆H₁₂O₆: C, 64.00; H, 4.03. Found: C, 63.85; H, 3.84.

1,2-Methylenedioxy-7,9-dimethoxy-6H-benzo[*c*]chromen-6-one (5a)

Colorless needles, mp 254.8-256.5 °C (hexane-CH₂Cl₂). ¹H-NMR (400 MHz, CDCl₃) δ : 7.58 (1H, d, *J* = 2.2 Hz, ArH), 6.91 (1H, d, *J* = 8.6 Hz, ArH), 6.80 (1H, d, *J* = 8.8 Hz, ArH), 6.56 (1H, d, *J* = 2.4 Hz, ArH), 6.16 (2H, s, ArOCH₂OAr), 4.00 (3H, s, ArOMe), 3.95 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ : 165.5, 163.8, 157.9, 146.8, 144.0, 143.7, 137.2, 109.8, 109.0, 105.3, 103.4, 102.3, 101.8, 99.4, 56.5, 55.8. IR (KBr) cm⁻¹: 1722, 1641, 1612, 1597, 1574, 1504, 1489, 1456, 1445, 1418, 1346, 1265, 1246, 1205, 1165, 1080, 1057, 1043, 1007, 986, 920, 839, 916. *Anal.* Calcd for C₁₆H₁₂O₆: C, 64.00; H, 4.03. Found: C, 63.92; H, 3.96.

2,3,7,9- Tetramethoxy-6H-benzo[*c*]chromen-6-one (4b)

Colorless solid, mp 180.8-181.7 °C (hexane-CH₂Cl₂). ¹H-NMR (400 MHz, CDCl₃) δ : 7.21 (1H, s, ArH), 6.87 (1H, d, *J* = 2.2 Hz, ArH), 6.77 (1H, s, ArH), 6.46 (1H, d, *J* = 2.2 Hz, ArH), 3.98 (3H, s, ArOMe), 3.98 (3H, s, ArOMe), 3.97 (3H, s, ArOMe), 3.91 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ : 165.1, 163.8, 157.9, 151.4, 146.5, 145.7, 139.0, 109.3, 104.0, 102.5, 99.8, 97.5, 96.2, 56.3, 56.0, 56.0, 55.5. IR (KBr) cm⁻¹: 3576, 1686, 1618, 1597, 1570, 1520, 1458, 1439, 1425, 1408, 1273, 1238, 1211, 1198, 1173, 1159, 1123, 1070, 995, 824. *Anal.* Calcd for C₁₇H₁₆O₆: C, 64.55; H, 5.10. Found: C, 64.33; H, 4.98.

1,2,7,9- Tetramethoxy-6H-benzo[*c*]chromen-6-one (5b)

Colorless solid, mp 202.2-203.5 °C (hexane-CH₂Cl₂). ¹H-NMR (400 MHz, CDCl₃) δ: 8.33 (1H, d, *J* = 2.2 Hz, ArH), 7.07 (2H, s, ArH), 6.61 (1H, d, *J* = 2.4 Hz, ArH), 3.99 (3H, s, ArOMe), 3.96 (3H, s, ArOMe), 3.92 (3H, s, ArOMe), 3.91 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ: 165.4, 163.7, 157.9, 149.5, 147.3, 146.4, 138.5, 114.7, 112.7, 112.4, 104.1, 102.3, 99.6, 60.4, 56.7, 56.5, 55.6. IR (KBr) cm⁻¹: 1724, 1601, 1587, 1570, 1497, 1474, 1456, 1443, 1414, 1337, 1298, 1267, 1246, 1229, 1213, 1204, 1163, 1090, 1072, 1059, 1024, 962, 837, 804. *Anal.* Calcd for C₁₇H₁₆O₆: C, 64.55; H, 5.10. Found: C, 64.45 ; H, 5.05.

2,3-Dibenzyloxy-7,9-dimethoxy-6H-benzo[*c*]chromen-6-one (4c)

Colorless solid, mp 166.5-167.3 °C (hexane-CH₂Cl₂). ¹H-NMR (400 MHz, CDCl₃) δ: 7.49-7.30 (11H, m, ArH), 6.81 (1H, s, ArH), 6.72 (1H, d, *J* = 2.2 Hz, ArH), 6.42 (1H, d, *J* = 2.2 Hz, ArH), 5.20 (2H, s, ArOCH₂), 5.18 (2H, s, ArOCH₂), 3.95 (3H, s, ArOMe), 3.92 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ: 165.4, 164.2, 158.2, 152.0, 147.4, 145.4, 139.4, 137.2, 136.2, 128.8, 128.7, 128.3, 128.1, 127.6, 127.3, 110.2, 109.9, 103.0, 102.5, 98.1, 96.6, 72.8, 71.0, 56.4, 55.7. IR (KBr) cm⁻¹: 1738, 1614, 1593, 1568, 1518, 1499, 1448, 1431, 1391, 1333, 1244, 1211, 1196, 1159, 1126, 1061, 1020, 1003, 924, 839, 824, 733, 694. *Anal.* Calcd for C₂₉H₂₄O₆: C, 74.35; H, 5.16. Found: C, 74.47; H, 5.22.

1,2-Dibenzyloxy-7,9-dimethoxy-6H-benzo[*c*]chromen-6-one (5c)

Colorless solid, mp 184.4-186.1 °C (hexane-CH₂Cl₂). ¹H-NMR (400 MHz, CDCl₃) δ: 8.29 (1H, d, *J* = 2.4 Hz, ArH), 7.47-7.31 (10H, m, ArH), 7.14 (1H, d, *J* = 9.0 Hz, ArH), 7.06 (1H, d, *J* = 9.0 Hz, ArH), 6.55 (1H, d, *J* = 2.2 Hz, ArH), 5.16 (2H, s, ArOCH₂), 5.08 (2H, s, ArOCH₂), 3.96 (3H, s, ArOMe), 3.43 (3H, s, ArOMe). ¹³C-NMR (100 MHz, CDCl₃) δ: 165.4, 163.5, 157.9, 148.7, 146.8, 138.7, 137.0, 136.7, 128.7, 128.6, 128.5, 128.4, 128.3, 127.8, 117.0, 113.2, 112.6, 104.1, 102.2, 100.2, 75.2, 72.1, 56.4, 55.3. IR (KBr) cm⁻¹: 1742, 1601, 1572, 1501, 1474, 1456, 1443, 1427, 1371, 1337, 1290, 1261, 1252, 1215, 1196, 1167, 1049, 1022, 959, 800, 752, 731, 696. *Anal.* Calcd for C₂₉H₂₄O₆: C, 74.35; H, 5.16. Found: C, 74.26; H, 5.13.

Altertenuol (2,3,7-Trihydroxy-9-methoxy-6H-benzo[*c*]chromen-6-one) (6)

Under N₂ atmosphere, to a solution of **4c** (50.1 mg, 0.107 mmol) in CH₂Cl₂ (3.5 mL) was added BBr₃ (1M CH₂Cl₂ solution, 0.320 mL, 0.320 mmol) at 0 °C. The mixture was stirred for 1 h at rt, and quenched with water. After extraction with AcOEt, the organic layer was washed with brine and dried over MgSO₄ to give a residue which was subjected to silica gel column chromatography with AcOEt. A colorless solid of **1** (27.9 mg, 95%) was obtained, mp 283.9-285.8 °C (acetone) [lit.⁷ mp 284-285 °C]. ¹H-NMR (400 MHz, CD₃OD) δ: 7.38 (1H, s, ArH), 6.94 (1H, d, *J* = 2.4 Hz, ArH), 6.74 (1H, s, ArH), 6.50 (1H, d, *J* = 2.2 Hz, ArH), 3.93 (3H, s, ArOMe). ¹³C-NMR (100 MHz, DMSO-d₆) δ: 166.8, 165.1, 163.8, 149.3, 144.4, 143.8, 137.6, 109.1, 109.0, 103.5, 100.0, 98.6, 98.0, 56.1. IR (KBr) cm⁻¹: 3535, 3285, 3105, 1647, 1607, 1568, 1528, 1510, 1456, 1431, 1279, 1227, 1202, 1165. *Anal.* Calcd for C₁₄H₁₀O₆: C, 61.32; H, 3.68.

Found: C, 61.26; H, 3.69.

2,3-Dibenzyloxy-7-hydroxy-9-methoxy-6H-benzo[*c*]chromen-6-one (7)^{4a}

Under N₂ atmosphere, to a solution of **4c** (102 mg, 0.218 mmol) in a mixed solvent of CH₂Cl₂ (4.7 mL) and CH₃CN (1.4 mL) was added AlCl₃ (69.6 mg, 0.522 mmol) and NaI (83.3 mg, 0.556 mmol) at 0 °C. The mixture was stirred for 4.5 h at rt, and quenched with water. After extraction with CH₂Cl₂, the organic layer was washed with brine and dried over MgSO₄ to give a residue which was subjected to silica gel column chromatography with hexane-AcOEt (4:3). Colorless solid of **7** (69.9 mg, 71%) was obtained, mp 211.3-212.5 °C (AcOEt) [lit.^{4a} mp 199-200 °C]. ¹H-NMR (400 MHz, CDCl₃) δ: 11.5 (1H, s, ArOH), 7.49-7.32 (11H, m, ArH), 6.88 (1H, s, ArH), 6.75 (1H, d, *J* = 2.2 Hz, ArH), 6.50 (1H, d, *J* = 2.2 Hz, ArH), 5.22 (4H, s, ArOCH₂), 3.91 (3H, s, ArOMe).

Conversion of 7 into 6

Under H₂ atmosphere, a mixture of **7** (53.6 mg, 0.118 mmol), 10% Pd/C (12.1 mg), and THF (4 mL) was vigorously stirred for 19 h at rt. After filtration and washing the filter paper with CHCl₃-MeOH (10:1), the filtrate was concentrated. An obtained residue was purified by silica gel column chromatography with AcOEt to give **6** (31.6 mg, 97%).

ACKNOWLEDGMENT

A part of this work was financially supported by JSPS KAKENHI Grant No. 22590003 and the Tamura Science and Technology Foundation.

REFERENCES AND NOTES

1. (a) D. E. Ames and A. Opalko, *Tetrahedron*, 1984, **40**, 1919; For reviews, see: (b) G. Bringmann, M. Breuning, and S. Tasler, *Synthesis*, 1999, 525; (c) J. Hassan, M. Sévignon, C. Gozzi, E. Schulz, and M. Lemaire, *Chem. Rev.*, 2002, **102**, 1359; (d) G. Bringmann, A. J. P. Mortimer, P. A. Keller, M. J. Gresser, J. Garner, and M. Breuning, *Angew. Chem. Int. Ed.*, 2005, **44**, 5384; (e) D. Alberico, M. E. Scott, and M. Lautens, *Chem. Rev.*, 2007, **107**, 174; (f) G. Bringmann, T. Gulder, T. A. M. Gulder, and M. Breuning, *Chem. Rev.*, 2011, **111**, 563; (g) J. Yamaguchi, A. D. Yamaguchi, and K. Itami, *Angew. Chem. Int. Ed.*, 2012, **51**, 8960.
2. (a) T. Matsumoto, T. Hosoya, and K. Suzuki, *J. Am. Chem. Soc.*, 1992, **114**, 3568; (b) T. Hosoya, E. Takashiro, T. Matsumoto, and K. Suzuki, *J. Am. Chem. Soc.*, 1994, **116**, 1004; (c) T. Harayama, H. Yasuda, T. Akiyama, Y. Takeuchi, and H. Abe, *Chem. Pharm. Bull.*, 2000, **48**, 861; (d) G. Bringmann and D. Menche, *Acc. Chem. Res.*, 2001, **34**, 615; (e) G. Bringmann, M. Breuning, R.-M. Pfeifer, W. A. Schenk, K. Kamikawa, and M. Uemura, *J. Organomet. Chem.*, 2002, **661**, 31; (f) G.

- Bringmann, S. Tasler, R.-M. Pfeifer, and M. Breuning, *J. Organomet. Chem.*, 2002, **661**, 49; (g) G. A. Molander, K. M. George, and L. G. Monovich, *J. Org. Chem.*, 2003, **68**, 9533; (h) H. Abe, S. Takeda, T. Fujita, K. Nishioka, Y. Takeuchi, and T. Harayama, *Tetrahedron Lett.*, 2004, **45**, 2327; (i) H. Abe, K. Nishioka, S. Takeda, M. Arai, Y. Takeuchi, and T. Harayama, *Tetrahedron Lett.*, 2005, **46**, 3197; (j) S. Takeda, H. Abe, Y. Takeuchi, and T. Harayama, *Tetrahedron*, 2007, **63**, 396; (k) H. Abe, Y. Sahara, Y. Matsuzaki, Y. Takeuchi, and H. Abe, *Tetrahedron Lett.*, 2008, **49**, 605; (l) H. Abe and T. Harayama, *Heterocycles*, 2008, **75**, 1305; (m) K. Shioe, Y. Takeuchi, T. Harayama, and H. Abe, *Chem. Pharm. Bull.*, 2010, **58**, 435; (n) K. Shioe, Y. Sahara, Y. Horino, T. Harayama, Y. Takeuchi, and H. Abe, *Tetrahedron*, 2011, **67**, 1960.
3. (a) T. Harayama, Y. Kawata, C. Nagura, T. Sato, T. Miyagoe, H. Abe, and Y. Takeuchi, *Tetrahedron Lett.*, 2005, **46**, 6091; (b) T. Harayama, A. Hori, H. Abe, and Y. Takeuchi, *Heterocycles*, 2006, **67**, 385; (c) H. Nishioka, C. Nagura, H. Abe, Y. Takeuchi, and T. Harayama, *Heterocycles*, 2006, **70**, 549; (d) H. Abe, T. Fukumoto, Y. Takeuchi, and T. Harayama, *Heterocycles*, 2007, **74**, 265; (e) T. Harayama, M. Asai, T. Miyagoe, H. Abe, Y. Takeuchi, A. Yamaguchi, and S. Fujii, *Heterocycles*, 2010, **81**, 1881; (f) T. Harayama, C. Nagura, T. Miyagoe, Y. Kawata, H. Abe, and Y. Takeuchi, *Heterocycles*, 2010, **81**, 2609; (g) H. Abe, T. Fukumoto, Y. Horino, T. Harayama, and Y. Takeuchi, *Heterocycles*, 2010, **82**, 851; (h) H. Abe, T. Matsukihira, T. Fukumoto, Y. Horino, Y. Takeuchi, and T. Harayama, *Heterocycles*, 2012, **84**, 323; (i) K. Maeda, T. Matsukihira, S. Saga, Y. Takeuchi, T. Harayama, Y. Horino, and H. Abe, *Heterocycles*, in press. DOI: 10.3987/COM-13-S(S)10.
4. (a) K. Kuramochi, K. Fukudome, I. Kuriyama, T. Takeuchi, Y. Sato, S. Kamisuki, K. Tsubaki, S. Sugawara, H. Yoshida, and Y. Mizushina, *Bioorg. Med. Chem.*, 2009, **17**, 7227; (b) G. Nemecek, J. Cudaj, and J. Podkech, *Eur. J. Org. Chem.*, 2012, 3863.
5. For Xphos, see: K. Billingsley and S. L. Buchwald, *J. Am. Chem. Soc.*, 2007, **129**, 3358.
6. (a) D. Garcia-Cuadrado, A. A. C. Braga, F. Maseras, and A. M. Echavarren, *J. Am. Chem. Soc.*, 2006, **128**, 1066; (b) D. Garcia-Cuadrado, P. de Mendoza, A. A. C. Braga, F. Maseras, and A. M. Echavarren, *J. Am. Chem. Soc.*, 2007, **129**, 6880.
7. (a) T. Rosett, R. H. Sankhala, C. E. Stickings, M. E. U. Taylor, and R. Thomas, *Biochem. J.*, 1957, **67**, 390; (b) R. Thomas, *Biochem. J.*, 1961, **80**, 234.
8. M. Node, T. Kajimoto, K. Nishide, E. Fujita, and K. Fuji, *Bull. Inst. Chem. Res., Kyoto Univ.*, 1992, **70**, 308.
9. A. Roy, K. R. Reddy, P. K. Mohanta, H. Ila, and H. Junjappa, *Synth. Commun.*, 1999, **29**, 3781.