

1 Effects of group sports on health-related physical fitness of overweight youth: A systematic
2 review and meta-analysis

3

4 Ruining head: Group sports and fitness in overweight youth

5

6 Ana Oliveira¹⁻³ MSc, alao@ua.pt; Ângela Monteiro², angela.monteiro@ua.pt; Cristina Jácome²
7 PhD, cristinajacome@ua.pt; Vera Afreixo^{3, 4} PhD, vera@ua.pt; Alda Marques^{2,3} PhD,
8 amarques@ua.pt

9

10 1 Faculty of Sports, University of Porto, Portugal

11 2 Lab 3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences
12 (ESSUA), University of Aveiro, Aveiro, Portugal

13 3 Institute for Research in Biomedicine (iBiMED), University of Aveiro, Aveiro, Portugal

14 4 Department of Mathematics, University of Aveiro, Aveiro, Portugal

15

16 Corresponding author: Alda Marques, PT, MSc, PhD, Senior Lecturer, School of Health Sciences,
17 University of Aveiro (ESSUA), Agrad do Crasto - Campus Universitário de Santiago, Edifício 30,
18 3810-193 Aveiro, Portugal. Email: amarques@ua.pt

19

1 **ABSTRACT**

2 Group sports interventions have been developed to improve the health-related physical fitness
3 of overweight/obese youth. However, its benefits are not systematically documented. This study
4 synthesize the evidence about the effects of group sports on health-related physical fitness of
5 overweight/obese youth.

6 Pubmed, Web of Knowledge, Scopus, Medline, CINAHL, SportDiscus and Academic Search
7 Complete were searched in February 2016. Studies assessing the effects of group sports on body
8 composition, cardiorespiratory endurance, muscle strength, flexibility and neuromotor fitness
9 of overweight/obese youth (aged <18 yrs) were included. Effect sizes (ES) were calculated with
10 Cohens' *d* and its 95% confidence intervals (CI).

11 Improvements were found in i) body composition - percentage of fat body mass (pooled
12 ES=0.67; 95% CI=0.24 to 1.10) and waist circumference (ES=0.69; p=0.004); ii) cardiorespiratory
13 endurance - peak oxygen consumption (pooled ES=0.53; 95% CI=0.13 to 0.92) and iii) muscle
14 strength – hand grip strength (ES=0.72; p=0.003). No significant effects were found for body
15 mass index (pooled ES=0.27; 95% CI=-0.14 to 0.69), percentage of lean body mass (ES=0.01;
16 p>0.05), maximal power output (ES from 0 to 0.06; p>0.05), sit-and-reach test (pooled ES=0.26;
17 95% CI=-0.16 to 0.68) and agility test (ES=0; p=0.48).

18 Group sports improve body composition, cardiorespiratory endurance and hand grip strength
19 of overweight/obese youth. Flexibility and neuromotor fitness do not seem to change following
20 group sports.

21 **Keywords:** team sports; physical fitness; children; adolescents; obese

22

1 Introduction

2 The number of overweight/obese youth has reached epidemic proportions (Daniels et al. 2009)
3 and it is now considered one of the main public health challenges of the 21st century (Karnik &
4 Kanekar 2012). Globally, the prevalence of overweight and obesity among youth in 2013 was
5 23% in developed countries and 13% in developing countries (Ng et al. 2014), which represents
6 an approximate 47% increase in the worldwide prevalence between 1980 and 2013 (Ng et al.
7 2014).

8 This phenomenon has raised attention of health policy makers and organisations as it is
9 associated with a range of adverse financial and health consequences. Currently, it is estimated
10 that around 7% of annual national health budgets across the European Union are spent on
11 diseases linked to obesity (European Union 2014), and this number is expected to increase if last
12 years' trend of increasing obesity is maintained. The rise in overweight and obesity in youth is
13 also distressing from a health related perspective due to the strong link between excess
14 adiposity and detrimental health and psychosocial outcomes in later life. These include, but are
15 not limited to, cardiovascular diseases, type 2 diabetes, as well as social stigmatisation and
16 mental health problems (World Health Organization 2004). Therefore, currently there is a major
17 research concern in developing effective and feasible interventions for the management of
18 overweight and obesity in youth (Aranceta et al. 2009; World Health Organization 2000).

19 Exercise has been recommended for the prevention and treatment of overweight and obesity
20 in youth (Kelley & Kelley 2013; Kelley et al. 2014), as weight gain is often caused by decreased
21 daily physical activity. Interventions showing the largest effects are often strictly structured and
22 comprise large components of educative and supportive programs about nutrition, healthy
23 lifestyle and exercise. However, these interventions result in large dropout rates due to lack of
24 enjoyment, perceptions of competence, social pressures, competing priorities and physical
25 factors (Crane & Temple 2015). For these reasons, group sports have emerged as potential
26 interventions for use with young populations as they link a variety of aerobic and resistance
27 activities with youths' interests of entertainment, improvement in skills and being with friends
28 (Salvy et al. 2012; Weintraub et al. 2008).

29 Several group sports have been studied in the adult population, such as tennis and football, and
30 positive associations with health benefits have been reported, including improved
31 cardiorespiratory endurance, lipid profile and leaner body (Oja et al. 2015; Pluim et al. 2007;
32 Randers et al. 2010). However, much less is known about the benefits of group sports on the
33 main components of health-related physical fitness (i.e., body composition, cardiorespiratory
34 endurance, muscle strength, flexibility and neuromotor) in overweight/obese youth
35 populations. A systematic review would be valuable to provide health and sports professionals
36 with the best evidence available on the benefits of group sports programs and guide
37 interventions for improving health-related physical fitness of overweight/obese youth.

38 Thus, the present review aimed to systematically summarise experimental and quasi-
39 experimental studies which reported the effects of group sports on health-related physical
40 fitness of overweight/obese youth when compared to absence of an intervention or to standard
41 interventions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

METHODS

Data sources and searches

A systematic literature search, limited to articles published in English, French, Portuguese and Spanish was conducted in February 2016 on Pubmed, ISI Web of Knowledge, Scopus, Medline, CINAHL, SportDiscus and Academic Search Complete. A specific search was also conducted in the Cochrane Library and in the International database of prospectively registered systematic reviews (PROSPERO) to exclude the existence of reviews with the same purpose as the present one. For the purpose of this review, “group sports” has been defined as “a collection of two or more individuals who possess a common identity, have common goals and objectives (...) and consider themselves to be a group”, following the definition proposed by Carron et al. (2005). Health-related physical fitness was composed of five components: body composition, cardiorespiratory endurance, muscle strength, flexibility and neuromotor fitness (American College of Sports Medicine 2013; Garber et al. 2011). Search terms were based on a combination of the following keywords: (obes* OR overweight) AND (youth OR child* OR adolescen* OR teen*) AND (“group sports” OR “team sports” OR “collective sports” OR football OR handball OR volleyball OR rugby OR basketball OR baseball OR soccer OR “water polo” OR hockey OR badminton OR tennis). The search terms were limited to titles and abstracts. Additionally, the reference lists of the selected articles were scanned for other potential eligible studies.

Eligibility criteria and study selection

Articles were included if they: i) involved overweight and/or obese (body mass index-BMI \geq 25Kg/m² or BMI \geq 85th percentile) (CDC/NDHS 2000) youth (<18 years old); ii) assessed the effects of group sports interventions; iii) were experimental or quasi-experimental (Centre for Evidence-Based Medicine 2016); iv) were full-text articles published in scientific journals or conference proceedings; and vi) were written in English, French, Portuguese or Spanish. Articles were excluded if the study was conducted only in adults or if it did not provide quantitative data on health-related physical fitness. Book chapters, review articles, abstracts of communications or meetings, letters to the editor, commentaries to articles, unpublished work and study protocols were also excluded from this study. After removing duplicates, two reviewers assessed all the potential studies identified. The studies were selected based on their titles and abstracts; when the title and abstract were relevant to the purpose of the review, the full-text article was read carefully to decide its inclusion. A third reviewer was consulted to solve any disagreements. This systematic review was reported using the systematic review method proposed by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) (Moher et al. 2009).

Quality assessment and data extraction

The quality of the included studies was assessed independently by two reviewers with a checklist adapted by Petticrew and colleagues based on the ‘Crombie criteria’ (Petticrew & Roberts 2006)

1 (Crombie 1996), according to previous systematic reviews (Barnard et al. 2010; Oliveira et al.
2 2014). This checklist provides a list of 8 questions about the research design, recruitment
3 strategy, response rate, sample representativeness, measures and statistics used and power, to
4 assess study quality.

5 Data from the included studies were extracted in a structured table-format comprising the
6 following topics: publication details (first author, year of publication, country); study design;
7 characteristics of participants (total number, age, gender; BMI), intervention (type, frequency,
8 duration and length of intervention), measures, outcome measures and findings.

9 10 **Data analysis and synthesis**

11
12 The consistency of the studies quality assessment, performed by two reviewers, was explored
13 with an inter-rater agreement analysis using Cohen's kappa. The value of Cohen's kappa ranges
14 from 0 to 1 and can be categorised as slight (<0.2), fair (0.21–0.4), moderate (0.41–0.6),
15 substantial (0.61–0.8) or almost perfect (>0.81) agreement (Landis & Koch 1977). The statistical
16 analysis was performed using PASW Statistics (version 18.0, SPSS Inc., Chicago, IL).

17 Meta-analysis was only possible to conduct with a limited number of variables due to the large
18 diversity of outcomes and units of measurement used in the included studies. For variables that
19 did not fit in the meta-analysis, effect sizes were computed. MetaXL 2.0 was used to calculate
20 the individual and pooled effect sizes. The input data were the Cohen's *d* value of each study
21 and the respective standard error. The output was the pooled Cohen's *d* value and
22 corresponding confidence intervals. Cohen's *d* effect sizes (ES) were interpreted as a small
23 (≥ 0.2), medium (≥ 0.5) or large (≥ 0.8) (Cohen 1988). Although comparisons between studies were
24 possible, some differences in study designs and methodologies were observed. Therefore,
25 statistical heterogeneity was considered and the random effect model was used to calculate the
26 pooled effect estimates. Additionally, the quality of individual studies was incorporated in meta-
27 analysis weights using the quality scores attributed (Doi & Thalib 2008). A sensitivity analysis, to
28 evaluate the robustness of the meta-analysis, was also performed by recalculating the pooled
29 results of the primary analysis and sequentially excluding each single study (Haidich 2010).

30 **RESULTS**

31 **Study selection**

32
33 The databases search identified 503 records. After duplicates removal, 263 records were
34 screened for relevant content. During the title and abstract screening, 223 articles were
35 excluded. Thus, the full-text of 40 potentially relevant articles was assessed and 34 articles were
36 excluded due to the following reasons: i) were not original scientific articles ($n=5$); ii) were
37 observational studies ($n=1$); iii) were written in Chinese, German or Serbian ($n=15$); iv) did not
38 include group sports interventions ($n=10$); v) did not include overweight/obese participants
39 ($n=1$); vi) did not assess health-related physical fitness components ($n=1$); vii) participants
40 presented other significant comorbidities than obesity ($n=1$, i.e., participants were also smokers
41 and alcohol consumers). Therefore, 6 original studies were included. Details of the search
42 process are shown in Supporting Information Fig. S1.

43 44 **Quality assessment**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

18

19
20

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

The results of the methodological quality assessment are presented in Supporting Information Table S1. All studies used objective measures and appropriate statistical analysis (Calcaterra et al. 2013; Faude et al. 2010; Hansen et al. 2013; Seabra et al. 2014; Vasconcellos et al. 2015; Weintraub et al. 2008). Sample representativeness was absent in all studies due to small sample sizes (Calcaterra et al. 2013; Faude et al. 2010; Hansen et al. 2013; Seabra et al. 2014; Vasconcellos et al. 2015; Weintraub et al. 2008), gender bias (i.e., samples were mainly or completely composed by males) (Hansen et al. 2013; Vasconcellos et al. 2015; Weintraub et al. 2008), lack of randomization processes (Hansen et al. 2013; Seabra et al. 2014) and no reasons given for selecting sub-samples for statistical analysis (Vasconcellos et al. 2015). Four studies did not justify their sample size (Calcaterra et al. 2013; Faude et al. 2010; Hansen et al. 2013; Vasconcellos et al. 2015), two presented inappropriate research designs (Calcaterra et al. 2013; Vasconcellos et al. 2015) to address the research question, two failed to report the recruitment strategy used (Calcaterra et al. 2013; Vasconcellos et al. 2015) and two failed to report the response rate (Faude et al. 2010; Vasconcellos et al. 2015). The agreement between the two reviewers was substantial ($k=0.702$; 95% CI=0.498–0.906; $p<0.001$; percentage of agreement of 85.4%).

Study characteristics

Characteristics of the studies included are shown in Supporting Information Table S2. Studies were published between 2008 and 2015 and were experimental (randomized controlled trial) (Faude et al. 2010; Weintraub et al. 2008) and quasi experimental (two groups pre-post design and two-stage cluster sample design) (Calcaterra et al. 2013; Hansen et al. 2013; Seabra et al. 2014; Vasconcellos et al. 2015) studies. All studies provided information of the effects of group sports interventions on body composition (Calcaterra et al. 2013; Faude et al. 2010; Hansen et al. 2013; Seabra et al. 2014; Vasconcellos et al. 2015; Weintraub et al. 2008), three on cardiorespiratory endurance (Calcaterra et al. 2013; Faude et al. 2010; Vasconcellos et al. 2015), two on flexibility (Calcaterra et al. 2013; Faude et al. 2010) and one on neuromotor fitness (Faude et al. 2010) and muscle strength (Calcaterra et al. 2013).

In total, 141 overweight/obese participants (45% male – data gathered from 4 studies, as Weintraub et al. (2008) and Vasconcellos et al. (2015) did not report the number of males/females enrolled), with age ranging from 8 to 17 years old were enrolled in the included studies. Group sports interventions, as well as control interventions, varied among studies. Studies conducting group sports interventions included different sports (football, rugby, volleyball and basketball), program lengths (3-6 months), session’s frequencies (3-4 days/week), durations (45-90 minutes) and intensities (50 % to higher than 80 % of the maximum heart rate). Five studies used football as their main intervention and presented similar sessions’ components and organisation (Faude et al. 2010; Hansen et al. 2013; Seabra et al. 2014; Vasconcellos et al. 2015; Weintraub et al. 2008). Hansen et al. (2013), Seabra et al. (2014) and Vasconcellos et al. (2015) sessions’ consisted of warm up (10-20 min), small sided football games (40-60 min) and cold down (10 min), whilst in the study of Faude et al. (2010), the sessions consisted of warm up (6 min), small sided football games (30 min), technique training (12 min) and fitness courses with

1 the ball (12 min). In the study of Weintraub et al. (2008), sessions began with teambuilding
2 exercises followed by 15 minutes of warm-up and stretching. The remainder practice was
3 devoted to learning football skills in the context of fun skill-building exercises and concluded
4 with a scrimmage. The number of players in each team (8 to 10 players) and the dimensions of
5 the court (55 x 24 m; length x width) were only reported by one study (Hansen et al. 2013). In
6 one study, participants included in the football group also performed the mandatory physical
7 education lessons (Seabra et al. 2014).

8 One study comprised a variety of group sports in their intervention including football, rugby,
9 volleyball, basketball and interactive video game exercises (Calcaterra et al. 2013). Each session
10 began and finished with a 10-min warm up/cool down period followed by a combination of
11 circuit-based aerobic exercises, strength and resistance exercises involving continuous work
12 bouts.

13 Adverse effects were reported in 2 studies (Hansen et al. 2013; Weintraub et al. 2008). Events
14 were not group sports-related and included foot injury, knee pain, eye pain, headaches, ingrown
15 toenail, ear infection, and skin rash in the control group and skin rash, car collision, and newly
16 diagnosed hypothyroidism in the group sports.

17

18 **Synthesis of results**

19 The effects of group sports interventions on the components of health-related physical fitness
20 of overweight and/or obese youth are summarised in the Supporting Information Table S2.

21 Body composition was analysed using BMI (Calcaterra et al. 2013; Faude et al. 2010; Hansen et
22 al. 2013; Seabra et al. 2014; Weintraub et al. 2008), waist circumference (Calcaterra et al. 2013),
23 percentage of fat body mass (Calcaterra et al. 2013; Seabra et al. 2014), and lean body mass
24 (Seabra et al. 2014).

25 All studies assessed BMI with a scale/stadiometer, however, only one found significant
26 differences favouring participants of group sports intervention (ES=0.75; $p=0.002$) (Calcaterra et
27 al. 2013). No significant differences were found between football based programs and standard
28 interventions (ES from 0.03 to 0.13; $p>0.05$) (Faude et al. 2010; Hansen et al. 2013; Seabra et al.
29 2014; Weintraub et al. 2008). The pooled Cohen's d estimate was 0.27 (95% CI=-0.14 to 0.69).
30 Medium amount of heterogeneity was found among studies ($I^2=40%$; $p=0.15$) (Supporting
31 Information Fig. S2).

32 Waist circumference was measured in one study using a measuring tape and significant
33 improvements, favouring group sports, were found (ES=0.69; $p=0.004$) (Calcaterra et al. 2013).

34 Percentage of fat body mass was measured in two studies, using a dual energy X-ray
35 absorptiometry (Seabra et al. 2014) and a bioelectrical impedance analysis (Calcaterra et al.
36 2013). Significant differences, favouring group sports interventions, were observed in one study
37 including a variety of group sports (ES=0.81; $p=0.001$) (Calcaterra et al. 2013). However, no
38 significant differences were found between participants enrolled in a football based program
39 and those in physical education lessons only (ES=0.28; $p>0.05$) (Seabra et al. 2014). The pooled

1 Cohen's d estimate for the percentage of fat body mass was 0.67 (95% CI=0.24 to 1.10). No
2 heterogeneity was found among studies ($I^2=10\%$; $p=0.29$) (Supporting Information Fig. S3).

3 The percentage of lean body mass was only assessed in one study which found no significant
4 differences between a football based exercise program and a standard exercise program
5 (ES=0.01; $p>0.05$) (Seabra et al. 2014).

6 Cardiorespiratory endurance was assessed with peak oxygen consumption (VO_{2peak})
7 (Vasconcellos et al. 2015) and maximal power output (PO_{max}) (Faude et al. 2010; Vasconcellos et
8 al. 2015) during maximal cardiopulmonary tests, on a cycle ergometer (Faude et al. 2010;
9 Vasconcellos et al. 2015) or on a treadmill (Calcaterra et al. 2013).

10 Conflicting findings were found for VO_{2peak} (Calcaterra et al. 2013; Faude et al. 2010;
11 Vasconcellos et al. 2015). Significant differences favouring a group sports intervention were
12 reported in two studies (ES=0.81 and ES=0.75; $p<0.05$) (Calcaterra et al. 2013; Vasconcellos et
13 al. 2015). No significant differences were found between participants whom enrolled in a
14 football based programe and those enrolled in physical education lessons only (ES=0; $p=0.98$)
15 (Faude et al. 2010). The pooled Cohen's d estimate for VO_{2peak} was 0.53 (95% CI=0.13 to 0.92),
16 however moderate evidence of heterogeneity was found between studies ($I^2=34\%$; $p=0.22$)
17 (Supporting Information Fig. S4).

18 No significant improvements were found for PO_{max} (Faude et al. 2010; Vasconcellos et al. 2015)
19 (ES=0 and ES=0.06; $p>0.05$) following group sports interventions. Due to the different units of
20 measurement used to report PO_{max} (i.e., Watts vs. Watts/Kg), it was not possible to calculate the
21 pooled Cohen's d.

22 Flexibility was assessed in two studies using the sit-and-reach test. Both studies reported no
23 significant improvements following group sports interventions (ES=0.25 and ES=0.28; $p>0.05$)
24 (Calcaterra et al. 2013; Faude et al. 2010). The pooled Cohen's d estimate for the sit-and-reach
25 test was 0.26 (95% CI=-0.16 to 0.68). No heterogeneity was found among studies ($I^2=0\%$; $p=0.94$)
26 (Supporting Information Fig. S5).

27 Neuromuscular fitness and muscle strength were assessed by Faude et al. (2010) and Calcaterra
28 et al. (2013), respectively. Faude et al. (2010), using the agility test, found no differences
29 between a group sports intervention and a standard intervention (ES=0; $p=0.48$). Calcaterra et
30 al. (2013), using a hand grip dynamometer, found significant muscle strength differences on a
31 pre-post intervention (ES=0.72; $p=0.003$).

32 The sensitivity analysis showed robustness for all variables included in the meta-analysis.

33

34 **DISCUSSION**

35 This systematic review and meta-analysis found that group sports interventions were effective
36 in improving body composition, cardiorespiratory endurance and muscle strength health-
37 related physical fitness components of overweight/obese youth Flexibility and neuromotor
38 fitness did not change after group sports interventions. Significant adverse effects were not
39 reported in control or group sports interventions.

1 Body composition was mainly assessed using BMI and percentage of body fat. Robust and
2 moderate evidence of improvement was observed for percentage of body fat and no
3 improvements were found for BMI. These results are similar to what has been found in adults
4 (Krustrup et al. 2010) and in other well established programs for weight loss in youth (Brown &
5 Summerbell 2009; Kelley & Kelley 2013; Kelley et al. 2014; Korsten-Reck et al. 2005). Four
6 reasons may explain the lack of BMI change. Firstly, the age range examined in the studies is
7 associated with major growing developments, in which significant gains in weight and height
8 may result in negligible changes in BMI (Malina et al. 2004). Secondly, BMI reflects both lean
9 and fat mass, thus similar increases in muscle mass and losses in fat mass following an
10 intervention may also result in negligible changes in BMI (Aasheim et al. 2011). Thirdly, the
11 interventions assessed were limited to group sports exercise programs, neglecting educational
12 and psychoeducational components. It has been suggested that more marked effects of activity
13 interventions on body composition would be achieved with multicomponent interventions,
14 including diet and behaviour modification (Nemet et al. 2005). Fourthly, it has been reported
15 that BMI lacks sensitivity to change following exercise programs (Krustrup et al. 2010; Watts et
16 al. 2005) and that percentage of body fat and H₂O, fat mass index and skinfolds are more
17 sensitive measures of body composition (Aasheim et al. 2011; Krustrup et al. 2010; Watts et al.
18 2005). Finally, these results should be interpreted with caution, as moderate heterogeneity was
19 found among studies. Such heterogeneity may be due to methodological differences (i.e., study
20 designs and sample sizes) and different intervention designs among studies (i.e., different
21 sports, program's lengths and intensities). Therefore, results from BMI should not be
22 overvalued. In fact it has been found that exercise improves health even if no weight is lost
23 (Shaw et al. 2006). Moreover, improved health and fitness may increase daily physical activity
24 and compliance with exercise programs which ultimately will lead to long-term weight
25 reductions.

26 Significant and medium effects were found following group sports interventions on
27 cardiorespiratory endurance of overweight and obese youth assessed with VO₂peak. Most
28 studies comprised group sports mainly focus on aerobic activities (e.g., football, basketball) with
29 moderate to high intensity, which are beneficial in increasing energy expenditure and weight
30 loss (Goran et al. 2000), hence improving VO₂peak (American College of Sports Medicine 2013).
31 Although these results are in line with other exercise programs developed for untrained adults
32 (Milanović et al. 2015; Oja et al. 2015) and overweight and obese youth (Kelley & Kelley 2013;
33 Kelley et al. 2015), their interpretation should be carefully assessed as, similar to body
34 composition, moderate heterogeneity was found among studies. Additionally, whilst VO₂peak
35 (i.e., the highest value of VO₂ attained on an incremental or high-intensity test, designed to take
36 an individual to his/her limit of tolerance) was the most used measure to assess
37 cardiorespiratory endurance (probably due to its easiness to define and determine), maximum
38 oxygen consumption (VO₂max -*"the oxygen intake during an exercise intensity at which actual
39 oxygen intake reaches a maximum beyond which no increase in effort can raise it"*(Hill & Lupton
40 1923)) is the single best indicator of cardiorespiratory endurance according to the World Health
41 Organisation (Shephard et al. 1968). Thus, it would be valuable if future studies could focus in
42 this variable to produce more secure and reliable information about the benefits of group sports
43 on the cardiorespiratory endurance of overweight and obese youth.

1 No evidence of improvements in flexibility and neuromotor fitness following group sports
2 interventions were found. This was expected, as the group sports reviewed in this meta-analysis
3 did not incorporate specific training modules for these fitness components. Also, recent studies
4 have been reporting no differences among normal weight, overweight and obese youth in
5 flexibility and neuromotor fitness (Đokić & Međedović 2013; Dumith et al. 2010; Fogelholm et
6 al. 2008). Thus, although aerobic activities are important for cardiorespiratory and body
7 composition health (American College of Sports Medicine 2013), it may be valuable to also
8 include flexibility and neuromotor components in exercise programmes. This may encourage
9 overweight/obese youth to develop positive experiences from physical activities that are not as
10 affected by excess weight as aerobic activities.

11 Only one study assessed the effects of group sports interventions on muscle strength, and
12 reported significant and large improvements in this fitness component. Strength is of particular
13 importance as the ability of overweight/obese youth to produce strength during field tests does
14 not solely reflect muscle capacity, but also relies on other musculoskeletal health factors such
15 as range of motion, flexibility, balance and coordination (Thivel et al. 2016). Additionally, it has
16 been reported that muscle strength is often the only health-related physical fitness component
17 that is improved in overweight youth (Dumith et al. 2010). Therefore, similarly to flexibility and
18 neuromotor activities, strength activities must be integrated in group sports exercise programs,
19 not only to preserve the strength component but also to highlight youths' abilities in these
20 activities when compared to normal weight peers (Dumith et al. 2010). Such strategy may be of
21 value to further motivate and commit overweight/obese children and adolescents to the
22 intervention program.

23 Health professionals often raise the objection that some group sports, namely European and
24 American football, are not appropriate for overweight/obese youth as the relative risk for
25 injuries in obese football players are 2.5 times higher than in healthy peers (Kaplan et al. 1995).
26 In the present review only two studies, which included European football as the main
27 intervention, reported incidence of adverse effects, and no significant football-related injuries
28 were reported. Conversely to professional football trainings, the interventions of the included
29 studies were not focused in improving sports technical gestures, thus the high mechanical
30 stresses and repetition movements, which often led to injuries (Caine et al. 2006), were reduced.
31 Nevertheless, future studies should investigate the incidence of adverse events in
32 overweight/obese youth per group sport, so that health and sports professionals could choose
33 and adapt their group sports interventions attending to injury prevention.

34 Finally, it should be noted that only six studies with different research designs, small sample
35 sizes, and interventions mainly based on football were included, demonstrating that the
36 evidence about the effects of group sports on health-related physical fitness of
37 overweight/obese youth is still limited. This was unexpected, considering that group sports are
38 often popular among children/adolescents; and that health institutions and policy makers
39 worldwide are continuously calling attention for the need to develop effective interventions for
40 the management of overweight and obesity in youth (Aranceta et al. 2009; World Health
41 Organization 2000). This review has presented the positive results that can be achieved with
42 group sports interventions and may serve as motivation and rationale for further studies and
43 funding opportunities currently emerging in this field. This would largely contribute not only to

1 increase the evidence of group sports interventions but, together with nutritional and
2 psychoeducational programs, may help to develop an effective and feasible intervention to
3 tackle youth obesity.

4

5 **Limitations**

6 This systematic review and meta-analysis has a number of limitations that should be considered.
7 Firstly, during the databases search, only titles and abstracts were searched for the selected
8 keywords, and no specific health-related fitness terms, such as cardiorespiratory endurance or
9 muscle strength, were included. Although this could have influenced the studies found, the
10 broad terms used allowed to find a larger number of studies in a preliminary phase and thus,
11 reduce the searching bias. Additionally to the search in the databases, the reference lists from
12 the studies included were reviewed. Thus, it is believed that this systematic review contains the
13 most relevant studies on the proposed topic. Secondly, only six studies with different research
14 designs and methodologies were found. Therefore, the extent to which the conclusions of this
15 review can be generalised to different group sports remains unclear. Lastly, it was not possible
16 to categorise children/adolescents into subgroups (e.g. infants, children and adolescents) due
17 to the limited number of studies found and lack of detailed information about age in the
18 included studies. Future studies should further investigate the effect sizes by age groups and in
19 adult and senior populations, to capture the developmental differences in performing group
20 sports.

21

22 **Perspectives**

23 It is well known that overweight/obese youth have reduced health-related physical fitness
24 compared with healthy peers. These findings are alarming as it is also known that poor health-
25 related physical fitness leads to cardiovascular diseases and obesity, increasing morbidity and
26 mortality risks. Therefore, it is imperative to design effective interventions to increase physical
27 fitness among youth.

28 Group sports interventions are often popular among youth, do not require specialised
29 equipment and or/infrastructures and have shown to be effective in improving body
30 composition, cardiorespiratory and muscle strength in overweight/obese youth. Nevertheless,
31 current studies using this intervention have overlooked health-related physical fitness
32 components in which overweight/obese youth have similar or better performances than their
33 normal weight peers (i.e., muscle strength, flexibility and neuromotor components). Therefore,
34 it is suggested that health and sports professionals adapt group sports interventions to include
35 these fitness components in order to match youth's expectations without compromising the
36 desired outcomes.

37 **Acknowledges**

38 AO is supported by Fundação para a Ciência e Tecnologia (PhD grant – Ref.
39 SFRH/BD/101951/2014), Portugal. This work was also partially supported by FEDER through
40 COMPETE and FCT under project UID/BIM/04501/2013 and by National Funds through FCT, in

1 the context of the projects UID/CEC/00127/2013 and Incentivo/EEI/UI0127/2014. The authors
2 would also like to acknowledge to Marta João Oliveira for her assistance in the edition of images.

3 REFERENCES

4 Aasheim ET, Aylwin SJ, Radhakrishnan ST, Sood AS, Jovanovic A, Olbers T, le Roux CW.
5 Assessment of obesity beyond body mass index to determine benefit of treatment. *Clin Obes.*
6 2011; **1**: 77-84.

7 American College of Sports Medicine. *ACSM's guidelines for exercise testing and prescription.*
8 Philadelphia: American College of Sports Medicine. 2013.

9 Aranceta J, Moreno B, Moya M, Anadon A. Prevention of overweight and obesity from a public
10 health perspective. *Nutr Rev.* 2009; **67 Suppl 1**: S83-88.

11 Barnard K, Thomas S, Royle P, Noyes K, Waugh N. Fear of hypoglycaemia in parents of young
12 children with type 1 diabetes: a systematic review. *BMC Pediatr.* 2010; **10**: 50.

13 Brown T, Summerbell C. Systematic review of school-based interventions that focus on changing
14 dietary intake and physical activity levels to prevent childhood obesity: an update to the obesity
15 guidance produced by the National Institute for Health and Clinical Excellence. *Obes Rev.* 2009;
16 **10**: 110-141.

17 Caine D, DiFiori J, Maffulli N. Physeal injuries in children's and youth sports: reasons for concern?
18 *British journal of sports medicine.* 2006; **40**: 749-760.

19 Calcaterra V, Larizza D, Codrons E, De Silvestri A, Brambilla P, Abela S, Arpesella M, Vandoni M.
20 Improved metabolic and cardiorespiratory fitness during a recreational training program in
21 obese children. *Journal of pediatric endocrinology & metabolism : JPEM.* 2013; **26**: 271-276.

22 Carron AV, Hausenblas HA, Eys MA. *Group Dynamics in Sport.* 3rd ed: Fitness Information
23 Technology 2005.

24 CDC/NDHS. CDC Growth charts: United State. . Centers for Disease Control and
25 Prevention/National Center for Health Statistics (CDC/NDHS), 2000.

26 Centre for Evidence-Based Medicine. Study Designs. 2016.

27 Cohen J. *Statistical Power Analysis for the Behavioral Sciences.* 2nd ed. Hellsdale, New Jersey:
28 Lawrence Earlbaum Associates 1988.

29 Crane J, Temple V. A systematic review of dropout from organized sport among children and
30 youth. *Eur Phys Educ Rev.* 2015; **21**: 114-131.

31 Crombie I. *The pocket guide to critical appraisal.* London: BMJ Publishing Group 1996.

32 Daniels SR, Jacobson MS, McCrindle BW, Eckel RH, Sanner BM. American Heart Association
33 Childhood Obesity Research Summit: executive summary. *Circulation.* 2009; **119**: 2114-2123.

34 Doi SA, Thalib L. A quality-effects model for meta-analysis. *Epidemiology (Cambridge, Mass).*
35 2008; **19**: 94-100.

36 Đokić Z, Međedović B. Relationship Between Overweight, Obesity and the Motor Abilities of 9-12
37 Year Old School Children. *Physical Culture.* 2013; **67**: 81-102.

- 1 Dumith SC, Ramires VV, Souza MA, Moraes DS, Petry FG, Oliveira ES, Ramires SV, Hallal PC.
2 Overweight/obesity and physical fitness among children and adolescents. *J Phys Act Health*.
3 2010; **7**: 641-648.
- 4 European Union. EU Action Plan on Childhood Obesity 2014-2020. 2014.
- 5 Faude O, Kerper O, Mulhaupt M, Winter C, Beziel K, Junge A, Meyer T. Football to tackle
6 overweight in children. *Scand J Med Sci Sports*. 2010; **20 Suppl 1**: 103-110.
- 7 Fogelholm M, Stigman S, Huisman T, Metsamuuronen J. Physical fitness in adolescents with
8 normal weight and overweight. *Scand J Med Sci Sports*. 2008; **18**: 162-170.
- 9 Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain DP.
10 American College of Sports Medicine position stand. Quantity and quality of exercise for
11 developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in
12 apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011; **43**:
13 1334-1359.
- 14 Goran M, Fields DA, Hunter GR, Herd SL, Weinsier RL. Total body fat does not influence maximal
15 aerobic capacity. *International journal of obesity and related metabolic disorders : journal of the
16 International Association for the Study of Obesity*. 2000; **24**: 841-848.
- 17 Haidich AB. Meta-analysis in medical research. *Hippokratia*. 2010; **14**: 29-37.
- 18 Hansen PR, Ndersen LJ, Ebelo AN, Rito J, Hornstrup T, Schmidt JF, Jackman SR, Mota J, Rego C,
19 Olweira J, Seabra A, Krstrup P. Cardiovascular effects of 3 months of football training in
20 overweight children examined by comprehensive echocardiography: a pilot study. *J Sports Sci*.
21 2013; **31**: 1432-1440.
- 22 Hill AV, Lupton H. Muscular Exercise, Lactic Acid, and the Supply and Utilization of Oxygen. *QJM*.
23 1923; **os-16**: 135-171.
- 24 Kaplan TA, Digel SL, Scavo VA, Arellana SB. Effect of obesity on injury risk in high school football
25 players. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport
26 Medicine*. 1995; **5**: 43-47.
- 27 Karnik S, Kanekar A. Childhood Obesity: A Global Public Health Crisis. *Int J Prev Med*. 2012; **3**: 1-
28 7.
- 29 Kelley GA, Kelley KS. Effects of Exercise in the Treatment of Overweight and Obese Children and
30 Adolescents: A Systematic Review of Meta-Analyses. *Journal of Obesity*. 2013; **2013**: 783103.
- 31 Kelley GA, Kelley KS, Pate RR. Effects of exercise on BMI z-score in overweight and obese children
32 and adolescents: a systematic review with meta-analysis. *BMC Pediatrics*. 2014; **14**: 1-16.
- 33 Kelley GA, Kelley KS, Pate RR. Exercise and BMI in Overweight and Obese Children and
34 Adolescents: A Systematic Review and Trial Sequential Meta-Analysis. *BioMed research
35 international*. 2015; **2015**: 704539.
- 36 Korsten-Reck U, Kromeyer-Hauschild K, Wolfarth B, Dickhuth HH, Berg A. Freiburg Intervention
37 Trial for Obese Children (FITOC): results of a clinical observation study. *Int J Obes (Lond)*. 2005;
38 **29**: 356-361.
- 39 Krstrup P, Aagaard P, Nybo L, Petersen J, Mohr M, Bangsbo J. Recreational football as a health
40 promoting activity: a topical review. *Scand J Med Sci Sports*. 2010; **20 Suppl 1**: 1-13.

- 1 Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*.
2 1977; **33**: 159-174.
- 3 Malina RM, Bouchard C, Bar-Or O. *Growth, maturation and physical activity*. Champaign: Human
4 Kinetics 2004.
- 5 Milanović Z, Pantelić S, Čović N, Sporiš G, Krstrup P. Is Recreational Soccer Effective for
6 Improving VO₂ max? A Systematic Review and Meta-Analysis. *Sports Medicine (Auckland, Nz)*.
7 2015; **45**: 1339-1353.
- 8 Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews
9 and Meta-Analyses: The PRISMA StatementThe PRISMA Statement. *Ann Intern Med*. 2009; **151**:
10 264-269.
- 11 Nemet D, Barkan S, Epstein Y, Friedland O, Kowen G, Eliakim A. Short- and long-term beneficial
12 effects of a combined dietary-behavioral-physical activity intervention for the treatment of
13 childhood obesity. *Pediatrics*. 2005; **115**: e443-449.
- 14 Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, Mullany EC, Biryukov S,
15 Abbafati C, Abera SF, Abraham JP, Abu-Rmeileh NM, Achoki T, AlBuhairan FS, Alemu ZA, Alfonso
16 R, Ali MK, Ali R, Guzman NA, Ammar W, Anwari P, Banerjee A, Barquera S, Basu S, Bennett DA,
17 Bhutta Z, Blore J, Cabral N, Nonato IC, Chang JC, Chowdhury R, Courville KJ, Criqui MH, Cundiff
18 DK, Dabhadkar KC, Dandona L, Davis A, Dayama A, Dharmaratne SD, Ding EL, Durrani AM,
19 Esteghamati A, Farzadfar F, Fay DF, Feigin VL, Flaxman A, Forouzanfar MH, Goto A, Green MA,
20 Gupta R, Hafezi-Nejad N, Hankey GJ, Harewood HC, Havmoeller R, Hay S, Hernandez L, Hussein
21 A, Idrisov BT, Ikeda N, Islami F, Jahangir E, Jassal SK, Jee SH, Jeffreys M, Jonas JB, Kabagambe EK,
22 Khalifa SE, Kengne AP, Khader YS, Khang YH, Kim D, Kimokoti RW, Kinge JM, Kokubo Y, Kosen S,
23 Kwan G, Lai T, Leinsalu M, Li Y, Liang X, Liu S, Logroscino G, Lotufo PA, Lu Y, Ma J, Mainoo NK,
24 Mensah GA, Merriman TR, Mokdad AH, Moschandreas J, Naghavi M, Naheed A, Nand D,
25 Narayan KM, Nelson EL, Neuhouser ML, Nisar MI, Ohkubo T, Oti SO, Pedroza A, Prabhakaran D,
26 Roy N, Sampson U, Seo H, Sepanlou SG, Shibuya K, Shiri R, Shiue I, Singh GM, Singh JA, Skirbekk
27 V, Stapelberg NJ, Sturua L, Sykes BL, Tobias M, Tran BX, Trasande L, Toyoshima H, van de Vijver
28 S, Vasankari TJ, Veerman JL, Velasquez-Melendez G, Vlassov VV, Vollset SE, Vos T, Wang C, Wang
29 X, Weiderpass E, Werdecker A, Wright JL, Yang YC, Yatsuya H, Yoon J, Yoon SJ, Zhao Y, Zhou M,
30 Zhu S, Lopez AD, Murray CJ, Gakidou E. Global, regional, and national prevalence of overweight
31 and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden
32 of Disease Study 2013. *Lancet*. 2014; **384**: 766-781.
- 33 Oja P, Titze S, Kokko S, Kujala UM, Heinonen A, Kelly P, Koski P, Foster C. Health benefits of
34 different sport disciplines for adults: systematic review of observational and intervention studies
35 with meta-analysis. *British journal of sports medicine*. 2015.
- 36 Oliveira A, Jacome C, Marques A. Physical fitness and exercise training on individuals with spina
37 bifida: a systematic review. *Res Dev Disabil*. 2014; **35**: 1119-1136.
- 38 Petticrew M, Roberts H. *Systematic reviews in the social sciences: a practical guide*. Oxford:
39 Blackwell 2006.
- 40 Pluim BM, Staal JB, Marks BL, Miller S, Miley D. Health benefits of tennis. *British journal of sports
41 medicine*. 2007; **41**: 760-768.
- 42 Randers MB, Nybo L, Petersen J, Nielsen JJ, Christiansen L, Bendiksen M, Brito J, Bangsbo J,
43 Krstrup P. Activity profile and physiological response to football training for untrained males

- 1 and females, elderly and youngsters: influence of the number of players. *Scand J Med Sci Sports*.
2 2010; **20 Suppl 1**: 14-23.
- 3 Salvy SJ, Bowker JC, Germeroth L, Barkley J. Influence of peers and friends on overweight/obese
4 youths' physical activity. *Exerc Sport Sci Rev*. 2012; **40**: 127-132.
- 5 Seabra AC, Seabra AF, Brito J, Krstrup P, Hansen PR, Mota J, Rebelo A, Rêgo C, Malina RM.
6 Effects of a 5-month football program on perceived psychological status and body composition
7 of overweight boys. *Scandinavian Journal of Medicine and Science in Sports*. 2014; **24**: 10-16.
- 8 Shaw K, Gennat H, O'Rourke P, Del Mar C. Exercise for overweight or obesity. *Cochrane Database*
9 *Syst Rev*. 2006: Cd003817.
- 10 Shephard RJ, Allen C, Benade AJ, Davies CT, Di Prampero PE, Hedman R, Merriman JE, Myhre K,
11 Simmons R. The maximum oxygen intake. An international reference standard of
12 cardiorespiratory fitness. *Bulletin of the World Health Organization*. 1968; **38**: 757-764.
- 13 Thivel D, Ring-Dimitriou S, Weghuber D, Frelut ML, O'Malley G. Muscle Strength and Fitness in
14 Pediatric Obesity: a Systematic Review from the European Childhood Obesity Group. *Obesity*
15 *facts*. 2016; **9**: 52-63.
- 16 Vasconcellos F, Seabra A, Montenegro R, Cunha F, Bouskela E, Farinatti P. Can Heart Rate
17 Variability be used to Estimate Gas Exchange Threshold in Obese Adolescents? *Int J Sports Med*.
18 2015; **36**: 654-660.
- 19 Watts K, Jones T, Davis E, Green D. Exercise Training in Obese Children and Adolescents. *Sports*
20 *Med*. 2005; **35**: 375-392.
- 21 Weintraub DL, Tirumalai EC, Haydel KF, Fujimoto M, Fulton JE, Robinson TN. Team sports for
22 overweight children: the Stanford Sports to Prevent Obesity Randomized Trial (SPORT). *Arch*
23 *Pediatr Adolesc Med*. 2008; **162**: 232-237.
- 24 World Health Organization. Obesity: preventing and managing the global epidemic. Geneva:
25 World Health Organization, 2000.
- 26 World Health Organization. Global Strategy on Diet, Physical Activity and Health. France: World
27 Health Organization, 2004.
- 28