

Florida International University FIU Digital Commons

Department of Biostatistics Faculty Publications

Robert Stempel College of Public Health & Social
Work

9-21-2015

The benefits of inositol-stabilized arginine silicate as a workout ingredient

S. Rood-Ojalvo
Nutrition 21, LLC

D. Sandler
StrengthPro Inc

Emir Veledar
Emory University; Department of Biostatistics, Florida International University, eveledar@fiu.edu

J. Komorowski

Follow this and additional works at: https://digitalcommons.fiu.edu/biostatistics_fac

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Rood-Ojalvo et al.: The benefits of inositol-stabilized arginine silicate as a workout ingredient. *Journal of the International Society of Sports Nutrition* 2015 12(Suppl 1):P14.

This work is brought to you for free and open access by the Robert Stempel College of Public Health & Social Work at FIU Digital Commons. It has been accepted for inclusion in Department of Biostatistics Faculty Publications by an authorized administrator of FIU Digital Commons. For more information, please contact dcc@fiu.edu.

POSTER PRESENTATION

Open Access

The benefits of inositol-stabilized arginine silicate as a workout ingredient

S Rood-Ojalvo^{1*}, D Sandler², E Veledar³, J Komorowski¹

From The Twelfth International Society of Sports Nutrition (ISSN) Conference and Expo
Austin, TX, USA. 11-13 June 2015

Background

The purpose of this study was to examine the benefits of inositol-stabilized arginine silicate (ASI; Nitrosigine[®]) as a workout ingredient in healthy adults. ASI has been previously shown to significantly enhance blood levels of arginine up to six hours post-dose and increase nitric oxide levels. To investigate reports of enhanced energy, increased muscle pump and stamina during workouts, and faster muscle recovery post-workout, ASI (1,500mg/day) was tested in a double-blind placebo-controlled crossover-design (DBPC-X) study using the POMS vigor-activity and fatigue-inertia sub-scores, blood flow measurements, leg circumference measurements, and biomarkers of muscle recovery (creatine kinase (CK) and lactate dehydrogenase (LDH)) as outcome measures.

Methods

The DBPC-X study was conducted in male subjects (N = 16 per group) who had limited exercise routines prior to participating in the study. These subjects took ASI daily for 4 days. Subjects had baseline measurements drawn at the hour 0 visit, took the study product, and completed an intense leg extension exercise protocol to induce muscle soreness. Subjects returned after 24, 48, and 72 hours for additional study measurements. After 72 hours, subjects repeated the leg extension exercise protocol. There was a seven-day washout between test products. Between-product assessments were primary endpoints and within-product assessments secondary endpoints.

Results

Sixteen (16) healthy male subjects (19-33 years of age) completed the study. Perceived energy, measured using the POMS vigor-activity sub-scores, significantly increased

after 72 hours compared to placebo ($p = 0.039$). At 72 hours, perceived fatigue, measured using the POMS fatigue-inertia sub-scores, significantly decreased in the ASI group ($p = 0.041$) from pre-dose, compared to a non-significant change in the placebo group ($p = 0.580$); $p = 0.055$ between groups.

Hyperemia, measured using leg circumference, increased significantly in the ASI group by 1.8cm ($p = 0.001$) at 72 hours from pre-dose, compared to a non-significant increase in the placebo group by 0.8cm ($p = 0.091$); $p = 0.070$ between groups.

Blood flow, measured by blood velocity through the femoral artery using a Doppler Ultrasound, increased 59.9 cm/s in the ASI group ($p < 0.005$) and 49.9cm/s in the placebo group ($p < 0.005$) after exercise on Day 3; $p = 0.2$ between groups.

CK levels significantly decreased in the ASI group at 24 ($p = 0.040$), 48 ($p = 0.017$) and 72 ($p = 0.034$) hours post-exercise compared to the placebo group. Immediately post-exercise at the hour 0 visit, ASI supplementation led to 44% less muscle damage, measured by CK levels, than placebo ($p = 0.057$). LDH levels significantly increased from baseline immediately after exercise in the placebo group ($p = 0.015$), but not in the ASI group ($p = 0.366$); $p = 0.133$ between groups. No safety concerns were raised by this study.

Conclusion

Both primary and secondary endpoints show that daily doses of ASI prior to workout significantly increased pre-workout energy levels, increased muscle pump immediately following a workout, and decreased biomarkers of muscle damage immediately after a workout and during recovery. These results demonstrate multiple benefits of ASI as a functional workout ingredient.

* Correspondence: sroodojalvo@nutrition21.com

¹Nutrition 21, LLC, Purchase, Harrison, NY, USA

Full list of author information is available at the end of the article

Authors' details

¹Nutrition 21, LLC, Purchase, Harrison, NY, USA. ²StrengthPro Inc, Golden, CO, USA. ³Emory University, Atlanta, GA, USA.

Published: 21 September 2015

doi:10.1186/1550-2783-12-S1-P14

Cite this article as: Rood-Ojalvo et al.: The benefits of inositol-stabilized arginine silicate as a workout ingredient. *Journal of the International Society of Sports Nutrition* 2015 **12**(Suppl 1):P14.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

