



A Combined Linkage and Exome Sequencing Analysis for Electrocardiogram Parameters in the Erasmus Rucphen Family Study

Claudia T. Silva^{1,2,3}, Irina V. Zorkoltseva⁴, Najaf Amin¹, Ayşe Demirkan^{1,5}, Elisabeth M. van Leeuwen¹, Jan A. Kors⁶, Marten van den Berg⁶, Bruno H. Stricker^{7,8,9}, André G. Uitterlinden⁸, Anatoly V. Kirichenko⁴, Jacqueline C. M. Witteman⁷, Rob Willemssen¹⁰, Ben A. Oostra^{1,11}, Tatiana I. Axenovich⁴, Cornelia M. van Duijn^{1,11†} and Aaron Isaacs^{1,11*†}

OPEN ACCESS

Edited by:

Gudrun A. Brockmann,
Humboldt University of Berlin,
Germany

Reviewed by:

Catherine Stein,
Case Western Reserve University,
USA
Anke Hinney,
University of Duisburg-Essen,
Germany

*Correspondence:

Aaron Isaacs
aaron.isaacs@gmail.com

†These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Applied Genetic Epidemiology,
a section of the journal
Frontiers in Genetics

Received: 02 June 2016

Accepted: 11 October 2016

Published: 08 November 2016

Citation:

Silva CT, Zorkoltseva IV, Amin N, Demirkan A, van Leeuwen EM, Kors JA, van den Berg M, Stricker BH, Uitterlinden AG, Kirichenko AV, Witteman JCM, Willemssen R, Oostra BA, Axenovich TI, van Duijn CM and Isaacs A (2016) A Combined Linkage and Exome Sequencing Analysis for Electrocardiogram Parameters in the Erasmus Rucphen Family Study. *Front. Genet.* 7:190. doi: 10.3389/fgene.2016.00190

¹ Genetic Epidemiology Unit, Department of Epidemiology, Erasmus University Medical Center, Rotterdam, Netherlands, ² Doctoral Program in Biomedical Sciences, Universidad del Rosario, Bogotá, Colombia, ³ GENIUIROS Group, Genetics and Genomics Research Center CIGGUR, School of Medicine and Health Sciences, Universidad del Rosario, Bogotá, Colombia, ⁴ Institute of Cytology and Genetics, Siberian Branch of Russian Academy of Sciences, Novosibirsk, Russia, ⁵ Department of Human Genetics, Leiden University Medical Center, Leiden, Netherlands, ⁶ Department of Medical Informatics, Erasmus University Medical Center, Rotterdam, Netherlands, ⁷ Department of Epidemiology, Erasmus University Medical Center, Rotterdam, Netherlands, ⁸ Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, Netherlands, ⁹ Inspectorate of Health Care, The Hague, Netherlands, ¹⁰ Department of Clinical Genetics, Erasmus University Medical Center, Rotterdam, Netherlands, ¹¹ Center for Medical Systems Biology, Leiden, Netherlands

Electrocardiogram (ECG) measurements play a key role in the diagnosis and prediction of cardiac arrhythmias and sudden cardiac death. ECG parameters, such as the PR, QRS, and QT intervals, are known to be heritable and genome-wide association studies of these phenotypes have been successful in identifying common variants; however, a large proportion of the genetic variability of these traits remains to be elucidated. The aim of this study was to discover loci potentially harboring rare variants utilizing variance component linkage analysis in 1547 individuals from a large family-based study, the Erasmus Rucphen Family Study (ERF). Linked regions were further explored using exome sequencing. Five suggestive linkage peaks were identified: two for QT interval (1q24, LOD = 2.63; 2q34, LOD = 2.05), one for QRS interval (1p35, LOD = 2.52) and two for PR interval (9p22, LOD = 2.20; 14q11, LOD = 2.29). Fine-mapping using exome sequence data identified a C > G missense variant (c.713C > G, p.Ser238Cys) in the *FCRL2* gene associated with QT (rs74608430; $P = 2.8 \times 10^{-4}$, minor allele frequency = 0.019). Heritability analysis demonstrated that the SNP explained 2.42% of the trait's genetic variability in ERF ($P = 0.02$). Pathway analysis suggested that the gene is involved in cytosolic Ca^{2+} levels ($P = 3.3 \times 10^{-3}$) and AMPK stimulated fatty acid oxidation in muscle ($P = 4.1 \times 10^{-3}$). Look-ups in bioinformatics resources showed that expression of *FCRL2* is associated with *ARHGAP24* and *SETBP1* expression. This finding was not replicated in the Rotterdam study. Combining the bioinformatics information with the association and linkage analyses, *FCRL2* emerges as a strong candidate gene for QT interval.

Keywords: genetics, epidemiology, electrocardiography, linkage, exome

INTRODUCTION

The electrocardiogram (ECG) is an important tool for diagnosing, monitoring and evaluating risk in patients with cardiovascular disease (CVD; Lin et al., 2013; Pelto et al., 2013). ECG measurements, such as PR interval, QRS complex duration, and QT interval, are used for the diagnosis and prediction of cardiac arrhythmias and sudden cardiac death (SCD; Kolder et al., 2012). Myocardial depolarization and repolarization time are measured by the QT interval: the time between the onset of the QRS complex and the end of the T wave. QT shortening or prolongation has been associated with an increased risk for arrhythmias and SCD (Newton-Cheh and Shah, 2007). PR interval and QRS duration are measures of cardiac conduction time; QRS duration reflects conduction through the ventricular myocardium, while PR interval measures atrial and atrioventricular conduction from the sinoatrial node to the ventricular myocardium, primarily through the atrioventricular node (Cheng et al., 2015; Mozos and Caraba, 2015).

There are significant genetic contributions to ECG measurements; genome-wide association studies (GWAS) identified at least 71 common variants associated with their variability (Arking et al., 2006, 2014; Newton-Cheh et al., 2007, 2009; Pfeufer et al., 2009, 2010; Holm et al., 2010; Sotoodehnia et al., 2010). A number of these associations were established in loci containing genes that encode proteins with previously known roles in heart development and function, such as cardiac transcription factors; sodium, calcium, and potassium ion channels; genes with a role in myocardial electrophysiology; and others involved in the conduction of electrical impulses (Kolder et al., 2012). These include *ARHGAP24*, *SETBP1*, *LRIG1*, *CREBBP*, *MEIS1*, *TBX20*, and *TBX5*. Some ion channel encoding genes, such as *SCN5A*, *HERG*, *KCNE1*, and *KCNE2*, have been associated with long QT syndrome (LQTS; Tristani-Firouzi et al., 2001), atrial fibrillation (AF) and Brugada Syndrome (Hedley et al., 2011). Collectively, however, these loci explain only modest proportions of phenotypic variability; GWAS SNPs specific for each trait account for limited trait heritability (17% for QRS, 4% for QT, and 2% for PR) (Silva et al., 2015).

Genome-wide association studies generally interrogate only common variants, typically of small effect. Families, in addition to being robust against population stratification, may be enriched for less frequent variants, which can potentially be identified by linkage and fine mapping. The aim of this study, therefore, was to discover less frequent variants using linkage analysis in a large family-based study, the Erasmus Rucphen Family Study (ERF).

MATERIALS AND METHODS

Study Population

The ERF study, which is a part of the Genetic Research in Isolated Populations (GRIP) Program, is a family-based study including over 3000 participants descendant from 22 couples that lived in the Rucphen region in the southwest Netherlands in the 19th century (Pardo et al., 2005). All descendants of those couples were invited to visit the clinical research center in the

region, where they were examined in person (Aulchenko et al., 2004). Interviews at the time of blood sampling were performed by medical practitioners and included questions on a broad range of topics, including current medication use and medical history (Sayed-Tabatabaei et al., 2005). Height and weight were measured with the participant in light underclothing and body mass index (kg/m^2) was computed. Blood pressure (BP) was measured twice on the right arm in a sitting position after at least five minutes rest, using an automated device (OMRON 711, Omron Healthcare, Bannockburn, IL, USA). The average of the two measures was used for analysis. Hypertension was defined through the use of antihypertensive medication and/or through the assessment of BP measurements according to the World Health Organization (1999) guidelines (individuals with $\text{BP} \geq 140/90$ mmHg should be regarded as hypertensive). The Medical Ethics Committee of the Erasmus University Medical Center approved the ERF study protocol and all participants, or their legal representatives, provided written informed consent.

ECG Measurement and Interpretation

Examinations included 10 s 12-lead ECG measurements, recorded with an ACTA-ECG (Esaote, Florence, Italy) with a sampling frequency of 500 Hz. Digital measurements of the ECG parameters were made using the Modular ECG Analysis System (MEANS; van Bommel et al., 1990). Briefly, MEANS operates on multiple simultaneously recorded leads, which are transformed to a detection function that brings out the QRS complex and the other parts of the signal. MEANS determines common onsets and offsets for all 12 leads together on one representative averaged beat, with the use of template matching techniques. The measurement and diagnostic performance of MEANS have been extensively evaluated, both by the developers and by others (Willems et al., 1987, 1991; van Bommel et al., 1990; de Bruyne et al., 1997; Eijgelsheim et al., 2009). The MEANS criteria for MI are mainly based on pathological Q waves, QR ratio, and R-wave progression (Leening et al., 2010). A cardiologist, specialized in ECG methodology, ascertained the final diagnosis of MI. QT interval was corrected for heart rate using Bazett's formula in all analyses (Funck-Brentano and Jaillon, 1993).

Genotyping and Statistical Analyses of the Linkage Study

Illumina's HumanHap6k Genotyping BeadChip (6K Illumina Linkage IV Panels[®]) was used for genotyping for the linkage analyses. All genotyping procedures were performed according to the manufacturer's protocols. Only markers with minor allele frequency (MAF) > 0.05 were selected for further analysis. Genotyping errors leading to Mendelian inconsistencies were detected using PedCheck (O'Connell and Weeks, 1998). Unlikely double recombination events were detected using MERLIN (Abecasis et al., 2002). All observed Mendelian errors were eliminated from the data. A total of 5250 autosomal SNPs with a call rate greater than 95% were included in the linkage analyses. All traits were adjusted for age, sex, BMI and height and inverse-normal transformation of ranks was applied before analysis. One thousand five hundred and forty-seven people with

complete ECG, covariate, and genotype data were included in the initial analysis. Variance component multipoint linkage was performed using the `-vc` option in the MERLIN v.1.0.1 software (Gudbjartsson et al., 2000; Abecasis et al., 2002). This program calculates exact IBD sharing probabilities using the Lander-Green algorithm, requiring restriction of pedigree size. Because of this, the large single ERF pedigree with multiple loops was split into non-overlapping fragments of no more than 18 bits with the help of the PedSTR program (Kirichenko et al., 2009). Final variance component two-point linkage analysis for the identified *FCRL2* variant (rs74608430) was performed using Merlin in one large, single pedigree.

Regions of interest with LOD > 1.9 were selected for further study (Lander and Kruglyak, 1995). Borders of the linkage areas were defined as LOD score minus 2 support intervals (LOD-2 SI) around the linkage peaks. Genes within the LOD-2 SI were annotated using SCAN (SNP and CNV Annotation Database¹).

Exome Sequencing

Exomes for 1336 individuals from ERF were sequenced at the Center for Biomics, Department of Cell Biology, Erasmus MC, the Netherlands, using the Agilent V4 capture kit on an Illumina HiSeq2000 sequencer using the TruSeq Version 3 protocol. Mean depth base was 74.23x (median = 57x) and mean depth region was 65.26x (median = 52.87x). The sequence reads were aligned to the human genome build 19 (hg19) using BWA and the NARWHAL pipeline (Li and Durbin, 2009; Brouwer et al., 2012). The aligned reads were processed further using the IndelRealigner, MarkDuplicates, and TableRecalibration tools from the Genome Analysis Toolkit (GATK) and Picard² to remove systematic biases and to recalibrate the PHRED quality scores in the alignments. Genetic variants were called using the Unified Genotyper tool of the GATK. About 1.4 million Single Nucleotide Variants (SNVs) were called and, after removing the low quality variants (QUAL < 150), we retrieved 577,703 SNVs in 1,309 individuals. Linear regression analyses, with SNVs in an additive model, were conducted on ECG measures, adjusted for age, sex, BMI, and height. To reduce the burden of multiple testing, we assessed only damaging variants in the LOD-2 SI; we found 324 such variants for QT, 52 for QRS and 61 for PR. We employed a Bonferroni correction for the number of deleterious mutations selected for each trait (QT: $P = 1.5 \times 10^{-4}$, QRS: $P = 9.6 \times 10^{-4}$, and PR: $P = 8.2 \times 10^{-4}$). The proportion of trait variance explained by the SNP was calculated using the Merlin software (Abecasis et al., 2002).

Replication

We sought to replicate our findings in the Rotterdam Study (RS) cohort. The RS is an ongoing prospective cohort study conducted since 1990 in the city of Rotterdam in The Netherlands (Hofman et al., 2013). The Illumina Exome BeadChip array (“exome chip”) was developed through a large international initiative to efficiently study coding variants spanning the genome. The v1.0 array contains 247,870 variants, which were genotyped in 3,183

individuals from the RS population. Calling for this sample, and numerous others, was done centrally (in total, 62,267 samples). After rigorous quality control and exclusion of variants that were monomorphic or too rare to analyze, the final dataset consisted of 108,678 polymorphic variants in 3,163 individuals.

Bioinformatics Analysis

To predict the functionality of genetic variants, and for comparison to BWA and NARWHAL, annotations were also performed using the dbNSFP (database of human non-synonymous SNPs and their functional predictions³ and Seattle⁴ databases. These databases gave functional prediction results from four different programs (PolyPhen-2, SIFT, MutationTaster, and LRT) (Chun and Fay, 2009; Adzhubei et al., 2010; Schwarz et al., 2010; Vaser et al., 2016), in addition to gene and variant annotations. Genes containing nominally significant variants (Table 2) were analyzed using Ingenuity Pathway Analysis (IPA; Ingenuity systems Inc, Redwood city, CA, USA). Several IPA modules were implemented: the “core analysis” was used to assess pathways, relationships, and mechanisms relevant to the dataset; the “upstream regulator analysis” was implemented to identify molecules (including microRNA and transcription factors) that may affect expression levels; and the “downstream effects analysis” was utilized to predict downstream biological processes that are increased or decreased.⁵ The GEO2R⁶ tool was used to analyse microarray-based expression data in the GEO database (GEO Accession numbers: GSE2240 and GSE41177). The Gene Network tool⁷ was used to describe co-expression networks and to assess potential functional effects of identified genes.

RESULTS

Table 1 shows the characteristics of the participants included in the discovery linkage analyses and exome sequencing, as well as the exome chip replication sample. There were no significant differences between the largely overlapping linkage and exome sequence groups. The replication sample was considerably older, and was characterized by increased frequency of hypertension (and BP differences), increased PR interval and decreased QT interval compared to the discovery samples. The three ECG traits studied (the QT, QRS, and PR intervals) demonstrated only modest pair-wise correlations in the discovery dataset (Supplementary Table 1).

Supplementary Table 2 shows the linkage results for the ECG traits, which yielded a total of five regions with suggestive LOD scores (LOD > 1.9). QT was suggestively linked to two regions, on chromosome 1 (LOD = 2.63) and on chromosome 2 (LOD = 2.05). A suggestive LOD score for QRS was observed on chromosome 1 (LOD = 2.52) and, for PR, two suggestive regions

³<http://varianttools.sourceforge.net/Annotation/DbNSFP>

⁴<http://snp.gs.washington.edu/SeattleSeqAnnotation/>

⁵https://www.ingenuity.com/wp-content/themes/ingenuity-qiagen/pdf/ipa/ipa_datasheet.pdf

⁶<http://www.ncbi.nlm.nih.gov/geo/geo2r/>

⁷<http://genenetwork.nl:8080/GeneNetwork>

¹<http://www.scandb.org>

²<http://picard.sourceforge.net>

TABLE 1 | Descriptive statistics of the linkage, exome sequence, and replication populations.

	Linkage Studies			Exome sequence			Exome array		
	ERF = 1860			ERF = 1212			Rotterdam Study = 2300		
	Mean (SD)	Minimum	Maximum	Mean (SD)	Minimum	Maximum	Mean (SD)	Minimum	Maximum
Males (n, %)	775 (42%)			488 (40%)			1012 (44%)		
Age (years)	46.4 (13.8)	16.6	85.3	47.4 (14.0)	18.2	86.1	67.9 (8.1)	55.0	101.0
BMI (kg/m ²)	26.6 (4.6)	15.5	61.8	26.6 (4.4)	15.5	61.8	26.2 (3.6)	14.3	44.2
Height (cm)	167.4 (9.1)	143.6	196.5	166.8 (9.1)	141.0	196.5	167.4 (9.4)	137.0	198.0
Weight (kg)	75.9 (15.1)	41.9	161.0	74.2 (14.5)	42.1	161.0	73.6 (11.9)	40.0	130.8
SBP (mm Hg)	137.7 (19.1)	85.5	217.0	138.0 (19.3)	85.5	239.0	139.5 (21.7)	82.0	231.0
DBP (mm Hg)	79.6 (9.7)	54.5	120.0	79.3 (9.7)	53.5	127.5	73.8 (11.0)	36.0	129.0
Hypertension	766 (41%)			517 (43%)			1133 (49%)		
PR	152.0 (22.4)	92.0	308.0	152.8 (22.4)	96.0	308.0	165.5 (20.8)	102.0	220.0
QT	403.1 (22.4)	336.0	531.0	403.6 (22.0)	336.0	531.0	396.6 (28.5)	282.0	516.0
QRS	96.8 (9.9)	68.0	120.0	96.8 (9.9)	68.0	120.0	96.9 (10.6)	68.0	120.0

Mean and standard deviation (SD) are given for the continuous measurements and number (percentage) is given for the categorical variables. BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; QT, Bazett's corrected QT interval; SD, standard deviation.




TABLE 2 | Selection of the coding variants.

Trait	Locus	Variants in the coding region				Observations ≤ 5%				Predicted to be damaging				Genes
		Synonymous	Missense	Stop	Splice	Total	Missense	Stop	Splice	Missense	Stop	Splice		
QT	1	3110	5089	117	36	8353	660	0	4	207	2	0	DENVD2C, RWDD3, FCER1A, GPR25, CDT1C, OMA1, LXL1L, LRRCEB, TPR, HOOK1, GIF2B, TXNIP, DDR2, CNN3, RBM15, BCL9, INVS1ABP, TNN, CEPT1, ACOT11, SARS, VAV3, TOMM40L, GABPB2, RFX5, ETV3L, APOBEC4, KIAA1614, ASPM, SPRR3, CEP350, C1orf168, COL24A1, SEMA6C, C1orf49, CACNA1S, IVL, VSG8, EDEM3, HMCN1, TBX19, GLRX2, IFI16, PODN, INADL, MPL, HY1, CAPZA1, AMIGO1, HCN3, RITCD1, OR10J1, FLG, DMRTB1, SPTA1, HFMT1, CFHR2, FCRL2, NCF2, CHIA, RBMXL1, C8A, SGIP1, FMO4, GBP1, CELSR2, ODF2L, PEAR1, FCRL1, SLC44A5, UROD, MOBKL2C, LRRC7, LRRCEC, IPO9, PRPF38B, MSH4, KIFAP3, LAMC2, PAQR6, ZNF687, MIER1, SMG7, TMEM61, ALX3, FAM189B, PDE4DIR, ATPAF1, C1orf50, PRRC2C, ZNF281, IGSF3, CRCT1, UOGRH, SLC27A3, NPHS2, PKLR, ATP1A4, TMEM125, TNR, OVGPI, SHCBP1L, UHMK1, B4GALT2, RNF220, PIAS3, KIF2C, TARS2, TMEM59, PI3K, CMPK1, PIK3R3, METTL11B, CITED4, EFCAB7, TTF2, AXDND1, DDX20, IGSF9, LEPRE1, ADAMTSL4, WDR77, GNAT2, GFSM2, PPM1J, ABCA4, EXTL2, AP4B1, HIVEP3, UBQLN4, POLR3C, NEGR1, TBX15, GBR6, KIAA1324, DPYD, F5, GJA5, CYP4A22, HENMT1, MIRPL37, TDRD5, ZBTB7B, SPATA6, FORLB, ABL2, ZFYVE9, LAMC1, RHBG, DUSP12, ZYG11A, WDR3, FAAH, C1orf106, HSD3B1, CTSS, TRIM45, ALG6, ACP6, PRUNE, TRIM46, AGL, MAG3, C1orf27, AL359075.1, SLC5A9, EBNA1BP2, COL11A1, FGGY, AIMP1, FAM63A, GLT25D2, DIMRTA2, EVI5, DPT, OR6P1, CRYGA, TTN, ARMC9, GIF3C3, ADAM23, ZFAND2B, PER2, COL6A3, TNS1, PAX3, HDAC4, OBSL1, CAPN10, IGFBP5, TMEM198, ESPNL, SPAG16, COL4A3, ANKAR, NEUROD1, NOP58, DNAH7, IQCA1, CCDC141, KIF1A, CASP10, SSFA2, CRYGC, ECEL1, AP1S3, COL5A2, NDUFS1, ATF2, STK36, UNG80, ABCB6, KIAA1486, ANKMY1, C2orf67, PLEKHM3, CNPPD1, ALPP, EFHD1, ZSWIM2, C2orf62, AQP12B, WIPF1, PDE11A, GLB1L, CCDC150, DGKD, SERPINE2, ABCA12, ITGAV, IDH1, SPHKAP, FNI, CDK15, GPR35, WNT10A, CYP27A1, ACSL3, ANKZF1, DNAJC10, FBXO36, STK16, MYO1B, KLHL30, PIKFYVE, DES, ASNSD1	
	2	1662	2444	40	17	4165	328	0	2	113	2	0	OTUD3, PHC2, SYF2, DHDDS, EPB41, NBP33, ZBTB40, COL16A1, RAP1GAP, C1orf38, EPHA10, MACEF1, PAD14, LDLRAP1, FCC2, AK2, SEPN1, TMC02, HSPG2, MAP3K6, TMC04, CCDC28B, TMEM234, GRHL3, ALDH4A1, GJB4, MAN1C1, SERINC2, E2F2, MUL1, PHACTR4, MYOM3, SRRM1, RLF, TINAGL1, KIAA0319L, C1orf94, C1orf63, UBXN11, USP48	
QRS	1	1057	1446	25	17	2546	152	1	4	51	1	0		

(Continued)

TABLE 2 | Continued

Trait	Locus	Variants in the coding region				Observations ≤ 5%				Predicted to be damaging			
		Synonymous	Missense	Stop	Splice	Total	Missense	Stop	Splice	Missense	Stop	Splice	Genes
PR	9	375	656	8	5	1053	96	0	0	29	0	0	DENND4C, CA9, FRMPD1, PLIN2, COIN, IFT74, UBAP1, IFNA10, RECK, UNC13B, GRHPR, KIAA1045, FREM1, OR2S2, IFNA14, FAM154A, KIAA1797, RGP1, ALDH1B1, NOL6, (GALT; GALT; RP11-195F19.29), PTPPLAD2, DDX58
	14	440	792	24	6	1276	86	0	1	31	1	0	HEATR5A, RABGGTA, LRRC16B, RBM23, CMA1, SUPT16H, MMP14, PARP2, CEBPE, OR4K1, PRKD1, LRRC16B, MYH6, PSMB11, HEATR5A, LRPT0, LRRC16B, TTC5, OR10G3, OR4N5, MYH6, TEPI, SDR39U1, TEPI, SLC7A7, LRPT0, TEPI, ADCY4, (AL163636.2;AL163636.2;RNASE4;RNASE4;RNASE4), PCK2, ARHGGEF40, KLHL33

Overview of variants predicted to be damaging for each trait with a frequency or less than 5%.

were located on chromosomes 9 and 14 with LOD scores of 2.20 and 2.29, respectively (Supplementary Table 2). Plots of the linked regions are shown in **Figure 1**.

Our analysis of coding variants in these linked regions revealed 55,050 variants in coding regions of genes under the peaks, as described in **Table 2**. Of these mutations, 1334 had a frequency less than or equal to 5%, 437 were predicted to be damaging by at least two of the prediction software packages used, and six were nonsense variants. By linkage peak, there were 207 missense damaging mutations and two nonsense mutations on 1q24 and 113 missense damaging mutations and two nonsense mutations on 2q32 for QT; 51 missense mutations and one nonsense mutation on 1p36 for QRS; and 29 missense mutations on 9q21 and 31 missense mutations and one nonsense mutation on 14q12 for PR. In total, 21 variants had nominal regression *P*-values less than 0.05 (the smallest *P*-values under each linkage peak were $P = 2.8 \times 10^{-4}$ for QT on chromosome 1, $P = 2.3 \times 10^{-2}$ for QT on chromosome 2, $P = 2.6 \times 10^{-2}$ for QRS on chromosome 1, $P = 1.9 \times 10^{-2}$ for PR on chromosome 9, and $P = 1.9 \times 10^{-2}$ for PR on chromosome 14) without reaching the significance levels needed to account for multiple comparisons (Supplementary Table 3). Looking for known genes under the linkage peaks (Supplementary Table 4), we found two variants previously related to heart failure, *TTN* (rs72648923; $P = 5.5 \times 10^{-2}$, MAF = 1.4×10^{-2}) and *HSD3B1* ($P = 3.9 \times 10^{-2}$ MAF = 1.1×10^{-2}). Neither achieved statistical significance after Bonferroni correction, although both genes were marginally associated with QT. Only a single variant, a C > G (Ser > Cys) variant in *FCRL2* (rs74608430; $P = 2.8 \times 10^{-4}$, MAF = 1.9×10^{-2}), approached the Bonferroni threshold for multiple-testing ($P = 1.5 \times 10^{-4}$). This variant, under the linkage peak on chromosome 1q23.1 for QT, is highly conserved (scorePhastCons = 0.998) and also predicted by PolyPhen-2 to be damaging (0.999). In the whole ERF population, rs74608430 explained 2.42% of the heritability of QT (reducing the LOD to 1.1; $h^2 = 0.87\%$; $P = 0.02$). This finding was not replicated in the RS ($P = 0.12$, $\beta = 0.14$). A sequence kernel association test analysis of the gene also failed to achieve significance in the replication sample ($P = 0.44$).

Not much is known about the function of *FCRL2*. Among the functions predicted by Gene Network are the regulation of cytosolic Ca^{2+} levels ($P = 3.3 \times 10^{-3}$) and AMPK stimulated fatty acid oxidation in muscle ($P = 4.1 \times 10^{-3}$). In the GEO database, *FCRL2* expression was higher in AF (Barth et al., 2005; Yeh et al., 2013). Supplementary Figure 1A shows the genes co-expressed with *FCRL2*, according to Gene Network. Two genes that have been associated with ECG outcomes by GWAS emerge: *ARHGAP24*, associated with PR, and *SETBP1*, associated with QRS (Holm et al., 2010; Pfeufer et al., 2010; Sotoodehnia et al., 2010). In the chromosome 1 region linked to QT, looking for co-expression, we found correlations between *DMRTA2*, *CEP350*, and *MPL* with genes previously associated with ECG traits: *DMRTA2* is co-expressed with *LRIG1*, a QRS associated gene (Supplementary Figure 1B); *MPL* is in a module with *MEIS1*, associated with PR (Supplementary Figure 1C); and *CEP350* interacts with *CREBBP*, associated with QT (Supplementary Figure 1D). These three genes are not in linkage disequilibrium

with each other. At the chromosome 2q34 locus linked with QT, a heart failure gene, *TTN*, was under the linkage peak. According to Gene Network analysis, expression of *TTN* is related to expression of three previously known QT genes (*ATP1B*, *TCEA3*, and *PLN*) and two QRS and PR associated genes (*TBX20* and *TBX5*) (Supplementary Figure 1E) (Holm et al., 2010; Pfeufer et al., 2010; Sotoodehnia et al., 2010; Arking et al., 2014). Additionally, *SPHKAP*, on chromosome 2 under the QT linkage peak, is co-expressed with *TBX5* (Supplementary Figure 1F).

DISCUSSION

Linkage analysis is an important tool for the identification of genomic regions influencing trait variability. The role of *TPM1* mutations with sudden death is a clear example of a locus discovered by linkage analysis (Ott et al., 2011; Mango et al., 2016). The advantages of family studies include control of heterogeneity and population stratification (Reich et al., 2005; Ott et al., 2011). We performed a linkage study on ECG measurements and identified five suggestive regions (1p35.1, 1q24.2, 2q34, 9p22.2, 14q11.2). Rare variant analysis in these regions uncovered two genes related to heart failure, *TTN* ($P = 5.5 \times 10^{-2}$) and *HSD3B1* ($P = 3.9 \times 10^{-2}$) and one gene with unknown cardiac function *FCRL2* ($P = 2.8 \times 10^{-4}$). None of them reaches statistical significance level after correction for multiple comparisons.

This study was conducted in a large, well-characterized family-based cohort, ascertained on the basis of genealogy and not phenotype. Multiple levels of genetic data, including a linkage panel and exome sequence data, provided a powerful dataset for identifying variants that may not be easily discovered with GWAS. Unfortunately, exome data was not available in the whole cohort, which could limit our ability to identify causal variants. Additionally, the sequence data did not include extra-genic or intronic variants that may be responsible for the observed linkage peaks.

Our analysis of rare coding variants in these linkage regions revealed 55,050 variants in coding regions. One thousand three hundred and thirty-four of these mutations had a frequency less than or equal to 5% and 437 were predicted to be damaging; none reached the significance threshold accounting for multiple comparisons. These variants spanned genes, including *TTN* and *HSD3B1*, which have been previously related to CVDs. *HSD3B*, a gene on chromosome 1 (1p13.1), has two isoforms (*HSD3B1* and *HSD3B2*) that were found to be associated with an increase in plasma aldosterone (Shimodaira et al., 2010). Changes in circulating aldosterone levels can modulate BP and hypertrophy (HT). A genome wide linkage analysis revealed that *HSD3B1* is a locus for BP variation (Shimodaira et al., 2010).

Another interesting gene covered by these variants was *TTN*; this gene encodes a sarcomeric protein named Titin, with a crucial role in sarcomeric structural integrity and muscle elasticity. Mutations in *TTN* have been shown to cause heart failure in humans. Additionally, mouse models with *TTN* mutations exhibit weak heart contractility and heart failure (Gerull et al., 2002; Xu et al., 2002; May et al., 2004) and hearts of

mutant embryos displayed weak spontaneous contraction (May et al., 2004). Additionally, the *TTN* network includes three QT associated genes, *ATP1B*, *TCEA3*, and *PLN*; *TBX320*, a QRS associated gene; and *TBX5* (a QRS and QT associated gene).

We also identified a less frequent C > G missense variant (rs74608430) in the *FCRL2* gene under the linkage peak on chromosome 1p23.1. This variant explains 2.42% ($h^2 = 0.87\%$, $P = 0.02$) of the total genetic variance of QT ($h^2 = 36\%$) in the ERF population. *FCRL2* has not been previously described with respect to cardiac function. Bioinformatics resources, however, showed that *FCRL2* expression is associated with *ARHGAP24* and *SETBP1* expression, two genes implicated in ECG variability by GWAS. This suggests that *FCRL2* may be relevant for heart function. *FCRL2* is expressed mostly in liver, heart, testis and kidney⁸. Gene Network predicts that it may be relevant for cytosolic Ca^{2+} levels and AMPK stimulated fatty acid oxidation in muscle. These are plausible pathways for QT function. This finding for rs74608430, however, was not replicated in the RS, in which the MAF was 2.9×10^{-2} . The absence of replication could be related to environmental differences influencing complex gene-environment interactions between these two study groups (Teo et al., 2009). Another plausible explanation is that, due to longer stretches of linkage disequilibrium in the family-based ERF sample, rs74608430 is tagging another variant in ERF and this is not the case in the general population.

Further, Ingenuity analysis revealed that *FCRL2* is correlated with some microRNAs (such as miR-1263, miR337-5p, miR-4699-3p, miR518e-3p, miR-507, miR3689a-5p, miR-507, miR-3622a-5p, miR-450b-5p, miR-4720-3p, and miR-1253). Among these, miR-337-5p is known to be differentially expressed in patients with valvular heart disease and patients with chronic AF (Cooley et al., 2012). This is consistent with the GEO database at NCBI⁹, which suggests that *FCRL2* is upregulated in patients with AF and dilated cardiomyopathy. In summary, the bioinformatics data available for this gene supports the hypothesis that *FCRL2* may be involved in heart function, and, specifically, related to ECG variability.

Additional interesting genes have been uncovered under the linkage peaks. First, the PR linkage peak on chromosome 14 contains damaging variants in the alpha and beta subunits of cardiac myosin *MYH6* and *MYH7*. Previous studies showed that genetic variants in these two genes have been found in hypertrophic cardiomyopathy (Anan et al., 1994; Lankford et al., 1995; Rayment et al., 1995; Niimura et al., 2002; Carniel et al., 2005), dilated cardiomyopathy (Kamisago et al., 2000; Carniel et al., 2005) and atrial septal defect (Ching et al., 2005). Second, we found *TNNT2* under the linkage peak on chromosome 1 for QT, which harbors known mutations underlying hypertrophic cardiomyopathy (Thierfelder et al., 1994) and familial dilated cardiomyopathy (Kamisago et al., 2000).

No explanatory variants were found for the other loci, for which there are a number of potential explanations. Linkage peaks are not precise in highlighting the location of the causal variant; even the region of interest cannot be easily pinpointed.

⁸<http://www.bioinfo.mochsl.org.br/miriad/gene/FCRL2/>

⁹<http://www.ncbi.nlm.nih.gov/geo/>

Additionally, we did not take into account alternative forms of genetic variation, such as structural and copy number variations (CNVs) or repeats in the linkage regions. Lastly, causal rare variants may be located outside the coding sequence, which we did not include in our sequencing analyses.

CONCLUSION

Although the combination of linkage and exome sequencing did not lead to the identification of a causal variant, suggestive linkage regions contain a number of plausible candidate genes, including *FCRL2*, *TTN*, *MYH6*, *MYH7*, *TNNT2*, and *HSD321*. Further analysis will need to be performed to demonstrate the involvement of these proteins in ECG measurements. We could not explain these with exonic sequence variants, so they will require more extensive follow-up, but provide potentially important indicators of the location of variation influencing ECG.

AUTHOR CONTRIBUTIONS

CS: Formal analysis, writing – original draft preparation; IZ: Formal analysis; NA: Formal analysis; AD: Formal analysis; EvL: Formal analysis; JK: Formal analysis, investigation, software; MvB: Formal analysis; BS: Investigation, resources; AU: Investigation, resources; AK: Formal analysis, software; JW: Investigation, resources; RW: Writing – original draft preparation, supervision; BO: Investigation, resources; TA: Formal analysis, supervision; CvD: Conceptualization, formal analysis, investigation, resources, writing – original draft preparation, supervision; AI: Conceptualization, formal analysis, writing – original draft preparation, supervision.

FUNDING

The Erasmus Rucphen Family (ERF) study as a part of EUROSPAN (European Special Populations Research Network)

REFERENCES

- Abecasis, G. R., Cherny, S. S., Cookson, W. O., and Cardon, L. R. (2002). Merlin-rapid analysis of dense genetic maps using sparse gene flow trees. *Nat. Genet.* 30, 97–101. doi: 10.1038/ng786
- Adzhubei, I. A., Schmidt, S., Peshkin, L., Ramensky, V. E., Gerasimova, A., Bork, P., et al. (2010). A method and server for predicting damaging missense mutations. *Nat. Methods* 7, 248–249. doi: 10.1038/nmeth0410-248
- Anan, R., Greve, G., Thierfelder, L., Watkins, H., McKenna, W. J., Solomon, S., et al. (1994). Prognostic implications of novel beta cardiac myosin heavy chain gene mutations that cause familial hypertrophic cardiomyopathy. *J. Clin. Invest.* 93, 280–285. doi: 10.1172/JCI116957
- Arking, D. E., Pfeufer, A., Post, W., Kao, W. H., Newton-Cheh, C., Ikeda, M., et al. (2006). A common genetic variant in the NOS1 regulator NOS1AP modulates cardiac repolarization. *Nat. Genet.* 38, 644–651. doi: 10.1038/ng1790
- Arking, D. E., Pulit, S. L., Crotti, L., van der Harst, P., Munroe, P. B., Koopmann, T. T., et al. (2014). Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat. Genet.* 46, 826–836. doi: 10.1038/ng.3014

was supported by European Commission FP6 STRP Grant No. 018947 (LSHG-CT-2006- 01947) and also received funding from the European Community's Seventh Framework Programme (FP7/2007–2013)/Grant Agreement HEALTH-F4- 2007-201413 by the European Commission under the programme “Quality of Life and Management of the Living Resources” of 5th Framework Programme (No. QLG2-CT-2002-01254). High-throughput analysis of the ERF data was supported by joint grant from Netherlands Organization for Scientific Research and the Russian Foundation for Basic Research (NWO-RFBR 047.017.043), and Russian Federal Agency of Scientific Organizations projects VI.53.2.2 and 0324-2015-0003. Exome sequencing analysis in ERF was supported by the Netherlands Organization for the Health Research and Development grant for Project No. 91111025. This work was also supported by the Federal Agency of Scientific Organizations #0324-2015-0003 (IZ, AK and TA). The Rotterdam Study is funded by Erasmus Medical Center (MC) and Erasmus University, Rotterdam; Netherlands Organization for the Health Research and Development; the Research Institute for Diseases in the Elderly; the Ministry of Education, Culture and Science; the Ministry for Health, Welfare and Sports; the European Commission; and the Municipality of Rotterdam.

ACKNOWLEDGMENTS

We thank all study participants and their relatives, general practitioners, and neurologists for their contributions and P. Veraart for help in genealogy, J. Vergeer for the supervision of the laboratory work, and P. Snijders for help in data collection.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fgene.2016.00190/full#supplementary-material>

- Aulchenko, Y. S., Heutink, P., Mackay, I., Bertoli-Avella, A. M., Pullen, J., Vaessen, N., et al. (2004). Linkage disequilibrium in young genetically isolated Dutch population. *Eur. J. Hum. Genet.* 12, 527–534. doi: 10.1038/sj.ejhg.5201188
- Barth, A. S., Merk, S., Arnoldi, E., Zwermann, L., Kloos, P., Gebauer, M., et al. (2005). Reprogramming of the human atrial transcriptome in permanent atrial fibrillation: expression of a ventricular-like genomic signature. *Circ. Res.* 96, 1022–1029. doi: 10.1161/01.RES.0000165480.82737.33
- Brouwer, R. W., van den Hout, M. C., Grosveld, F. G., and van Ijcken, W. F. (2012). NARWHAL, a primary analysis pipeline for NGS data. *Bioinformatics* 28, 284–285. doi: 10.1093/bioinformatics/btr613
- Carniel, E., Taylor, M. R., Sinagra, G., Di Lenarda, A., Ku, L., Fain, P. R., et al. (2005). Alpha-myosin heavy chain: a sarcomeric gene associated with dilated and hypertrophic phenotypes of cardiomyopathy. *Circulation* 112, 54–59. doi: 10.1161/CIRCULATIONAHA.104.507699
- Cheng, M., Lu, X., Huang, J., Zhang, S., and Gu, D. (2015). Electrocardiographic PR prolongation and atrial fibrillation risk: a meta-analysis of prospective cohort studies. *J. Cardiovasc. Electrophysiol.* 26, 36–41. doi: 10.1111/jce.12539

- Ching, Y. H., Ghosh, T. K., Cross, S. J., Packham, E. A., Honeyman, L., Loughna, S., et al. (2005). Mutation in myosin heavy chain 6 causes atrial septal defect. *Nat. Genet.* 37, 423–428. doi: 10.1038/ng1526
- Chun, S., and Fay, J. C. (2009). Identification of deleterious mutations within three human genomes. *Genome Res.* 19, 1553–1561. doi: 10.1101/gr.092619.109
- Cooley, N., Cowley, M. J., Lin, R. C., Marasco, S., Wong, C., Kaye, D. M., et al. (2012). Influence of atrial fibrillation on microRNA expression profiles in left and right atria from patients with valvular heart disease. *Physiol. Genomics* 44, 211–219. doi: 10.1152/physiolgenomics.00111.2011
- de Bruyne, M. C., Kors, J. A., Hoes, A. W., Kruijssen, D. A., Deckers, J. W., Grosfeld, M., et al. (1997). Diagnostic interpretation of electrocardiograms in population-based research: computer program research physicians, or cardiologists? *J. Clin. Epidemiol.* 50, 947–952. doi: 10.1016/S0895-4356(97)00100-5
- Eijgelsheim, M., Aarnoudse, A. L., Rivadeneira, F., Kors, J. A., Witteman, J. C., Hofman, A., et al. (2009). Identification of a common variant at the NOS1AP locus strongly associated to QT-interval duration. *Hum. Mol. Genet.* 18, 347–357. doi: 10.1093/hmg/ddn341
- Funck-Brentano, C., and Jaillon, P. (1993). Rate-corrected QT interval: techniques and limitations. *Am. J. Cardiol.* 72, 17B–22B. doi: 10.1016/0002-9149(93)90035-B
- Gerull, B., Gramlich, M., Atherton, J., McNabb, M., Trombitás, K., Sasse-Klaassen, S., et al. (2002). Mutations of TTN, encoding the giant muscle filament titin, cause familial dilated cardiomyopathy. *Nat. Genet.* 30, 201–204. doi: 10.1038/ng815
- Gudbjartsson, D. F., Jonasson, K., Frigge, M. L., and Kong, A. (2000). Allegro, a new computer program for multipoint linkage analysis. *Nat. Genet.* 25, 12–13. doi: 10.1038/75514
- Hedley, P. L., Haundrup, O., Andersen, P. S., Aidt, F. H., Jensen, M., Moolman-Smook, J. C., et al. (2011). The KCNE genes in hypertrophic cardiomyopathy: a candidate gene study. *J. Negat. Results Biomed.* 10:12. doi: 10.1186/1477-5751-10-12
- Hofman, A., Darwish Murad, S., van Duijn, C. M., Franco, O. H., Goedegebure, A., Ikram, M. A., et al. (2013). The rotterdam study: 2014 objectives and design update. *Eur. J. Epidemiol.* 28, 889–926. doi: 10.1007/s10654-013-9866-z
- Holm, H., Gudbjartsson, D. F., Arnar, D. O., Thorleifsson, G., Thorgeirsson, G., Stefansdottir, H., et al. (2010). Several common variants modulate heart rate, PR interval and QRS duration. *Nat. Genet.* 42, 117–122. doi: 10.1038/ng.511
- Kamisago, M., Sharma, S. D., DePalma, S. R., Solomon, S., Sharma, P., McDonough, B., et al. (2000). Mutations in sarcomere protein genes as a cause of dilated cardiomyopathy. *N. Engl. J. Med.* 343, 1688–1696. doi: 10.1056/NEJM200012073432304
- Kirichenko, A. V., Belonogova, N. M., Aulchenko, Y. S., and Axenovich, T. I. (2009). PedStr software for cutting large pedigrees for haplotyping, IBD computation and multipoint linkage analysis. *Ann. Hum. Genet.* 73(Pt 5), 527–531. doi: 10.1111/j.1469-1809.2009.00531.x
- Kolder, I. C., Tanck, M. W., and Bezzina, C. R. (2012). Common genetic variation modulating cardiac ECG parameters and susceptibility to sudden cardiac death. *J. Mol. Cell Cardiol.* 52, 620–629. doi: 10.1016/j.yjmcc.2011.12.014
- Lander, E., and Kruglyak, L. (1995). Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nat. Genet.* 11, 241–247. doi: 10.1038/ng1195-241
- Lankford, E. B., Epstein, N. D., Fananapazir, L., and Sweeney, H. L. (1995). Abnormal contractile properties of muscle fibers expressing beta-myosin heavy chain gene mutations in patients with hypertrophic cardiomyopathy. *J. Clin. Invest.* 95, 1409–1414. doi: 10.1172/JCI117795
- Leening, M. J., Elias-Smale, S. E., Felix, J. F., Kors, J. A., Deckers, J. W., Hofman, A., et al. (2010). Unrecognised myocardial infarction and long-term risk of heart failure in the elderly: the Rotterdam Study. *Heart* 96, 1458–1462. doi: 10.1136/hrt.2009.191742
- Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25, 1754–1760. doi: 10.1093/bioinformatics/btp324
- Lin, W. H., Zhang, H., and Zhang, Y. T. (2013). Investigation on cardiovascular risk prediction using physiological parameters. *Comput Math. Methods Med.* 2013:272691. doi: 10.1155/2013/272691
- Mango, R., Luchetti, A., Sanguolo, R., Ferradini, V., Briglia, N., Giardina, E., et al. (2016). Next generation sequencing and linkage analysis for the molecular diagnosis of a novel overlapping syndrome characterized by hypertrophic cardiomyopathy and typical electrical instability of Brugada syndrome. *Circ. J.* 80, 938–949. doi: 10.1253/circj.CJ-15-0685
- May, S. R., Stewart, N. J., Chang, W., and Peterson, A. S. (2004). A Titin mutation defines roles for circulation in endothelial morphogenesis. *Dev. Biol.* 270, 31–46. doi: 10.1016/j.ydbio.2004.02.006
- Mozos, I., and Caraba, A. (2015). Electrocardiographic predictors of Cardiovascular mortality. *Dis. Markers* 2015, 727401. doi: 10.1155/2015/727401
- Newton-Cheh, C., Eijgelsheim, M., Rice, K. M., de Bakker, P. I., Yin, X., Estrada, K., et al. (2009). Common variants at ten loci influence QT interval duration in the QTGEN Study. *Nat. Genet.* 41, 399–406. doi: 10.1038/ng.364
- Newton-Cheh, C., Guo, C. Y., Wang, T. J., O'Donnell, C. J., Levy, D., and Larson, M. G. (2007). Genome-wide association study of electrocardiographic and heart rate variability traits: the Framingham Heart Study. *BMC Med. Genet.* 8(Suppl. 1):S7. doi: 10.1186/1471-2350-8-S1-S7
- Newton-Cheh, C., and Shah, R. (2007). Genetic determinants of QT interval variation and sudden cardiac death. *Curr. Opin. Genet. Dev.* 17, 213–221. doi: 10.1016/j.gde.2007.04.010
- Niimura, H., Patton, K. K., McKenna, W. J., Soultis, J., Maron, B. J., Seidman, J. G., et al. (2002). Sarcomere protein gene mutations in hypertrophic cardiomyopathy of the elderly. *Circulation* 105, 446–451. doi: 10.1161/hc0402.102990
- O'Connell, J. R., and Weeks, D. E. (1998). PedCheck: a program for identification of genotype incompatibilities in linkage analysis. *Am. J. Hum. Genet.* 63, 259–266. doi: 10.1086/301904
- Ott, J., Kamatani, Y., and Lathrop, M. (2011). Family-based designs for genome-wide association studies. *Nat. Rev. Genet.* 12, 465–474. doi: 10.1038/nrg2989
- Pardo, L. M., MacKay, I., Oostra, B., van Duijn, C. M., and Aulchenko, Y. S. (2005). The effect of genetic drift in a young genetically isolated population. *Ann. Hum. Genet.* 69(Pt. 3), 288–295. doi: 10.1046/J.1469-1809.2005.00162.x
- Pelto, H., Owens, D., and Drezner, J. (2013). Electrocardiographic findings suggestive of cardiomyopathy: what to look for and what to do next. *Curr. Sports Med. Rep.* 12, 77–85. doi: 10.1249/JSR.0b013e3182874abb
- Pfeufer, A., Sanna, S., Arking, D. E., Muller, M., Gateva, V., Fuchsberger, C., et al. (2009). Common variants at ten loci modulate the QT interval duration in the QTSCD Study. *Nat. Genet.* 41, 407–414. doi: 10.1038/ng.362
- Pfeufer, A., van Noord, C., Marcianti, K. D., Arking, D. E., Larson, M. G., Smith, A. V., et al. (2010). Genome-wide association study of PR interval. *Nat. Genet.* 42, 153–159. doi: 10.1038/ng.517
- Rayment, I., Holden, H. M., Sellers, J. R., Fananapazir, L., and Epstein, N. D. (1995). Structural interpretation of the mutations in the beta-cardiac myosin that have been implicated in familial hypertrophic cardiomyopathy. *Proc. Natl. Acad. Sci. U.S.A.* 92, 3864–3868. doi: 10.1073/pnas.92.9.3864
- Reich, D., Patterson, N., De Jager, P. L., McDonald, G. J., Waliszewska, A., Tandon, A., et al. (2005). A whole-genome admixture scan finds a candidate locus for multiple sclerosis susceptibility. *Nat. Genet.* 37, 1113–1118. doi: 10.1038/ng1646
- Sayed-Tabatabaei, F. A., van Rijn, M. J., Schut, A. F., Aulchenko, Y. S., Croes, E. A., Zillikens, M. C., et al. (2005). Heritability of the function and structure of the arterial wall: findings of the Erasmus Rucphen Family (ERF) study. *Stroke* 36, 2351–2356. doi: 10.1161/01.STR.0000185719.66735.dd
- Schwarz, J. M., Rodelsperger, C., Schuelke, M., and Seelow, D. (2010). MutationTaster evaluates disease-causing potential of sequence alterations. *Nat. Methods* 7, 575–576. doi: 10.1038/nmeth0810-575
- Shimodaira, M., Nakayama, T., Sato, N., Aoi, N., Sato, M., Izumi, Y., et al. (2010). Association of HSD3B1 and HSD3B2 gene polymorphisms with essential hypertension, aldosterone level, and left ventricular structure. *Eur. J. Endocrinol.* 163, 671–680. doi: 10.1530/EJE-10-0428
- Silva, C. T., Kors, J. A., Amin, N., Dehghan, A., Witteman, J. C., Willemsen, R., et al. (2015). Heritabilities, proportions of heritabilities explained by GWAS findings, and implications of cross-phenotype effects on PR interval. *Hum. Genet.* 134, 1211–1219. doi: 10.1007/s00439-015-1595-9
- Sotoodehnia, N., Isaacs, A., de Bakker, P. I., Dorr, M., Newton-Cheh, C., Nolte, I. M., et al. (2010). Common variants in 22 loci are associated with QRS duration and cardiac ventricular conduction. *Nat. Genet.* 42, 1068–1076. doi: 10.1038/ng.716

- Teo, Y. Y., Fry, A. E., Bhattacharya, K., Small, K. S., Kwiatkowski, D. P., and Clark, T. G. (2009). Genome-wide comparisons of variation in linkage disequilibrium. *Genome Res.* 19, 1849–1860. doi: 10.1101/gr.092189.109
- Thierfelder, L., Watkins, H., MacRae, C., Lamas, R., McKenna, W., Vosberg, H. P., et al. (1994). Alpha-tropomyosin and cardiac troponin T mutations cause familial hypertrophic cardiomyopathy: a disease of the sarcomere. *Cell* 77, 701–712. doi: 10.1016/0092-8674(94)90054-X
- Tristani-Firouzi, M., Chen, J., Mitcheson, J. S., and Sanguinetti, M. C. (2001). Molecular biology of K(+) channels and their role in cardiac arrhythmias. *Am. J. Med.* 110, 50–59. doi: 10.1016/S0002-9343(00)00623-9
- van Bommel, J. H., Kors, J. A., and van Herpen, G. (1990). Methodology of the modular ECG analysis system MEANS. *Methods Inf. Med.* 29, 346–353.
- Vaser, R., Adusumalli, S., Leng, S. N., Sikic, M., and Ng, P. C. (2016). SIFT missense predictions for genomes. *Nat. Protoc.* 11, 1–9. doi: 10.1038/nprot.2015.123
- Willems, J. L., Abreu-Lima, C., Arnaud, P., van Bommel, J. H., Brohet, C., Degani, R., et al. (1991). The diagnostic performance of computer programs for the interpretation of electrocardiograms. *N. Engl. J. Med.* 325, 1767–1773. doi: 10.1056/NEJM199112193252503
- Willems, J. L., Arnaud, P., van Bommel, J. H., Bourdillon, P. J., Degani, R., Denis, B., et al. (1987). A reference data base for multilead electrocardiographic computer measurement programs. *J. Am. Coll. Cardiol.* 10, 1313–1321. doi: 10.1016/S0735-1097(87)80136-5
- World Health Organization (1999). World health organization-international society of hypertension guidelines for the management of hypertension. Guidelines Subcommittee. *J. Hypertens* 17, 151–183.
- Xu, X., Meiler, S. E., Zhong, T. P., Mohideen, M., Crossley, D. A., Burggren, W. W., et al. (2002). Cardiomyopathy in zebrafish due to mutation in an alternatively spliced exon of titin. *Nat. Genet.* 30, 205–209. doi: 10.1038/ng816
- Yeh, Y. H., Kuo, C. T., Lee, Y. S., Lin, Y. M., Nattel, S., Tsai, F. C., et al. (2013). Region-specific gene expression profiles in the left atria of patients with valvular atrial fibrillation. *Heart Rhythm* 10, 383–391. doi: 10.1016/j.hrthm.2012.11.013

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Silva, Zorkoltseva, Amin, Demirkan, van Leeuwen, Kors, van den Berg, Stricker, Uitterlinden, Kirichenko, Witteman, Willemsen, Oostra, Axenovich, van Duijn and Isaacs. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.