

## Original Paper

## Neonatology

Neonatology 2016;109:14–21

DOI: 10.1159/000438978

Received: March 24, 2015

Accepted after revision: July 27, 2015

Published online: September 23, 2015

# Neurodevelopmental Outcome in High-Risk Congenital Diaphragmatic Hernia Patients: An Appeal for International Standardization

Kitty G. Snoek<sup>a</sup> Irma Capolupo<sup>c</sup> Annabella Braguglia<sup>c</sup> Lucia Aite<sup>c</sup>  
 Joost van Rosmalen<sup>b</sup> Laura Valfrè<sup>c</sup> René M. Wijnen<sup>a</sup> Pietro Bagolan<sup>c</sup>  
 Dick Tibboel<sup>a</sup> Hanneke IJsselstijn<sup>a</sup>

<sup>a</sup>Intensive Care and Department of Pediatric Surgery, Erasmus MC-Sophia Children's Hospital, and <sup>b</sup>Department of Biostatistics, Erasmus MC, Rotterdam, The Netherlands; <sup>c</sup>Department of Medical and Surgical Neonatology, Bambino Gesù Children's Hospital, Rome, Italy

## Key Words

Congenital diaphragmatic hernia · Follow-up program · Neurodevelopmental outcome

## Abstract

**Background:** Since mortality in congenital diaphragmatic hernia (CDH) is decreasing, morbidity such as neurodevelopmental outcome is becoming increasingly important. **Objectives:** We evaluated neurodevelopmental outcome in high-risk CDH patients treated according to the CDH EURO Consortium standardized treatment protocol. **Methods:** This observational, prospective cohort study was conducted in two European centers. Neurodevelopment of 88 patients (Rotterdam n = 49; Rome n = 39) was assessed at 12 and 24 months with the Bayley Scales of Infant Development (BSID)-II-NL (Rotterdam) or BSID-III (Rome). Data of the centers were analyzed separately. **Results:** Cognition was normal in 77.8% of children from Rotterdam and in 94.8% from Rome at 12 months, and in 70.7 and 97.4%, respectively, at 24 months. Motor function was normal in 64.3% from Rotterdam and in 81.6% from Rome at 12 months and in 45.7 and 89.8%, respectively, at 24 months. Longer length of hospital stay (LoS) was associated with worse cognitive outcome and motor function; LoS, low socioeconomic sta-

tus, and ethnicity were associated with lower cognition. **Conclusions:** At 2 years, most CDH patients have normal cognition, but are at risk for motor function delay. Due to differences in outcomes between centers, careful interpretation is needed before conclusions can be drawn for other centers. Future multicenter collaboration should not only focus on standardization of postnatal care, but also on international standardization of follow-up to identify risk factors and thereby reduce morbidity. © 2015 S. Karger AG, Basel

## Introduction


Congenital diaphragmatic hernia (CDH) affects approximately 1 in 3,000 newborns [1]. It is a life-threatening disease caused mainly by persistent pulmonary hypertension and pulmonary hypoplasia [1]. Severely ill CDH patients may receive extracorporeal membrane oxygenation (ECMO) [1]. In 2008, a standardized neonatal treatment protocol was developed at a consensus meeting of the CDH EURO Consortium [1]. The mortality rate has decreased from 33 to 12% since this protocol was implemented [2]. Consequently, worldwide the focus of interest has shifted to morbidity [3–5].

KARGER

© 2015 S. Karger AG, Basel  
 1661–7800/15/1091–0014\$39.50/0

E-Mail karger@karger.com  
 www.karger.com/neo

Hanneke IJsselstijn, MD, PhD  
 Intensive Care and Department of Pediatric Surgery, Room Sk-1280  
 Erasmus MC-Sophia Children's Hospital, PO Box 2060  
 NL-3000 CB, Rotterdam (The Netherlands)  
 E-Mail h.ijsselstijn@erasmusmc.nl



**Fig. 1.** Flowchart of inclusion of all patients. NA = Not applicable.

Previous studies found associated morbidity in about 87% of CDH survivors, mainly related to the lungs or the gastrointestinal tract [6, 7]. While most follow-up studies have focused on morbidity, few have examined neurodevelopment. Most of those studies were cross-sectionally performed in small series in single centers [8–10]. Recently, longitudinal studies have shown neurodevelopmental delays in CDH patients [3, 4, 11]. To date, no published studies have examined neurodevelopmental outcome after implementation of a standardized treatment protocol in a multicenter setting.

We longitudinally investigated neurodevelopmental outcome in high-risk patients managed according to the CDH EURO Consortium standardized treatment protocol [1] in two high-volume CDH centers. Secondly, we identified determinants of neurodevelopmental outcome.

### Materials and Methods

This is an observational, prospective cohort study in patients born between January 2009 and May 2012 with high-risk CDH, i.e. antenatal diagnosis/respiratory insufficiency within 6 h postnatal-

ly. All patients were treated according to a standardized treatment protocol [1]; however, ECMO was only offered in Rotterdam. Rotterdam and Rome are two of the largest European CDH centers. As standard of care, survivors were offered a structured, longitudinal follow-up program, initiated in Rotterdam in 1999 [11] and in Rome in 2004 [12]. We evaluated prospectively collected data of repeated measurements at corrected ages of 12 and 24 months. Since subjects were not submitted to any handling and no rules of human behavior were imposed, institutional review board approval was waived. Parents were informed that data were used for research purposes.


Patient characteristics were retrieved from medical records. Neurodevelopmental outcome was assessed with the Bayley Scales of Infant Development (BSID). In Rotterdam, the Dutch-language version of the BSID-II (BSID-II-NL) was administered with Dutch normative data [13]. In both centers cognitive development was assessed at both time points (by two developmental psychologists per center). Psychomotor development was assessed by developmental psychologists at both time points in Rome and at 12 months in Rotterdam. In Rotterdam, psychomotor development was assessed by one physiotherapist at 24 months. In Rome, the Italian translation of the BSID-III [14] with American normative data was administered. Scores were grouped as normal ( $>-1$  SD), mildly delayed ( $-2 < SD < -1$ ) and severely delayed ( $<-2$  SD). For further details, see the supplementary file Methods (for all suppl. material, see [www.karger.com/doi/10.1159/000438978](http://www.karger.com/doi/10.1159/000438978)).

#### Data Analysis

Descriptive statistics are presented as number (%), mean (SD) or median (range). Independent samples *t* tests,  $\chi^2$  tests and Mann-Whitney *U* tests, where appropriate, were used to compare characteristics between centers. One-sample *t* tests served to compare mean scores to the normalized population mean. Linear mixed models were used to estimate neurodevelopmental outcome over time. These models included only the time point (12 or 24 months) as the independent variable, and a random intercept to account for the within-subject correlations. Perinatal factors associated with adverse neurodevelopment were determined using multivariate linear regression. Cognition and motor outcome were chosen as dependent variables in two different models; the data were analyzed separately for each assessment, age and each center. Based on clinical experience and the literature, the following independent variables were included in the linear regression analysis: tube feeding at discharge, ECMO support, ethnicity and SES as categorical variables, and SNAP-II score and length of hospital stay (LoS; log-transformed due to skewed distribution) as continuous variables. Analyses were performed using SPSS version 20.

## Results

In the study period, 141 CDH patients were admitted (Rotterdam  $n = 76$ ; Rome  $n = 65$ ). Thirty-eight children died (Rotterdam  $n = 19$ ; Rome  $n = 19$ ) and 5 were not high-risk CDH patients; thus, 98 children were eligible for follow-up. Ultimately, 88 patients were assessed (fig. 1).



**Fig. 2.** Neurodevelopmental outcome in CDH survivors at 12 and 24 months. Black: Rotterdam; grey: Rome.

Baseline characteristics of participants (table 1) were not significantly different from nonparticipants. Four patients (Rotterdam  $n = 1$ ; Rome  $n = 3$ ) underwent fetoscopic endotracheal occlusion. In Rotterdam, 28 patients underwent thoracoscopic repair and 7 were converted to laparotomy. In Rome, all children underwent laparotomy. None of the children received anticonvulsants. Six patients from Rotterdam had a genetic or chromosomal abnormality that was not associated with neurodevelopment. Patch repair, episodes of general anesthesia 0–24 months, and involvement of a physical therapist and speech-language pathologist were significantly different

**Table 1.** Background characteristics

	Rotterdam (n = 49)	Rome (n = 39)	p value <sup>1</sup>
<i>Perinatal and postnatal characteristics</i>			
LHR	1.9 (0.8)	2.2 (0.8)	0.32
Missing	15	13	
O/E LHR	51.6 (17.4)	53.3 (15.8)	0.71
Missing	21	13	
Male sex	28 (57.1%)	25 (64.1%)	0.51
Birth weight, kg	3.07 (0.56)	2.99 (0.41)	0.45
Born before 37 weeks of gestation	8 (16.3%)	4 (10.3%)	0.41
SNAP-II score	18 (12)	13 (9)	0.07
Age at repair, days	3.8 (2.7)	3.6 (2.1)	0.77
Left-sided CDH	42 (85.7%)	38 (97.4%)	0.06
Liver: intrathoracic	19 (38.8%)	12 (30.8%)	0.44
Defect size			0.24
A	5 (10.2%)	4 (10.3%)	
B	18 (26.7%)	22 (56.4%)	
C	24 (49.0%)	11 (28.2%)	
D	2 (4.1%)	2 (5.1%)	
Patch repair	35 (71.4%)	11 (28.2%)	<0.001
Defect size A	n = 1	n = 0	
Defect size B	n = 8	n = 1	
Defect size C	n = 24	n = 8	
Defect size D	n = 2	n = 2	
Initial ventilation			0.83
CMV	24 (49.0%)	20 (51.3%)	
HFO	25 (51.0%)	19 (48.7%)	
Length of ventilation, days	8 (1–271)	8 (2–70)	0.55
Length of initial hospital stay, days	28 (6–387)	30 (15–161)	0.54
CLD			0.12
No CLD	29 (59.2%)	32 (82.1%)	
Mild CLD	8 (16.3%)	2 (5.1%)	
Moderate CLD	1 (2.0%)	0 (0%)	
Severe CLD	11 (22.4%)	5 (12.8%)	

between Rotterdam and Rome (table 1). After exclusion of ECMO-treated patients (online suppl. table 1a), episodes of general anesthesia 0–24 months and involvement of physical therapy were not significantly different between the two centers (online supplementary table 1b).

Based on the cutoff scores from the BSID manuals, cognition was normal in 71 children (85.5%) and motor

	Rotterdam (n = 49)	Rome (n = 39)	p value <sup>1</sup>
<i>Length of morphinomimetics/sedatives</i>			
<1 week	25 (51.0%)	14 (35.9%)	0.28
1 week to 1 month	17 (34.7%)	20 (51.3%)	
>1 month	7 (14.3%)	5 (12.8%)	
<i>Episodes of general anesthesia 0–24 months</i>			
	2 (1–13)	1 (1–3)	0.01
<i>At discharge</i>			
Tube feeding	29 (59.2%)	20 (51.3%)	0.46
Physical therapy at home	20 (40.8%)	7 (17.9%)	0.02
Speech language pathologist involved	19 (38.8%)	3 (7.7%)	0.001
<i>Demographic variables</i>			
<i>Ethnicity</i>			
Native	34 (69.4%)	33 (84.6%)	0.13
Nonnative	14 (28.6%)	6 (15.4%)	
Unknown	1 (2.0%)	0 (0%)	
<i>SES mother</i>			
Low	13 (26.5%)	9 (23.1%)	0.12
Medium	15 (30.6%)	21 (53.8%)	
High	18 (36.7%)	9 (23.1%)	
Unknown	3 (6.1%)	0 (0%)	

Categorical variables are shown as n (%) and compared between centers using  $\chi^2$  tests, normally distributed variables are shown as means (SD) and compared between groups using independent samples t tests, and continuous variables that are not normally distributed are shown as medians (range) and compared between centers using Mann-Whitney U tests. LHR = Lung-head ratio [15]; O/E LHR = observed to expected lung-head ratio; SNAP-II score = Score for Neonatal Acute Physiology-II [16]; CMV = conventional mechanical ventilation; HFO = high-frequency oscillation; CLD = chronic lung disease [17]; SES = socioeconomic status [18]. <sup>1</sup> Comparison between Rotterdam and Rome.

function was normal in 58 children (72.5%) at 12 months; this held true for 67 patients (83.8%) and 56 patients (65.9%), respectively, at 24 months. Proportions of mild and severe neurodevelopmental delay are shown in figure 2. Table 2 shows means (SD) of neurodevelopmental outcome. In online supplementary figure 2a and online supplementary table 2a, data were presented after exclusion of ECMO-treated patients.

In Rotterdam, cognition and motor function remained stable over time. In Rome, cognition and motor function improved over time [mean differences 4.1 (p = 0.02) and 5.8 (p = 0.003), respectively].

**Table 2.** Neurodevelopmental outcome scores at 12 and 24 months (corrected for prematurity)

	12 months				24 months			
	Rotterdam <sup>1</sup>	p value <sup>2</sup>	Rome (n = 38)	p value <sup>2</sup>	Rotterdam <sup>3</sup>	p value <sup>2</sup>	Rome (n = 39)	p value <sup>2</sup>
Cognitive outcome <sup>4</sup>	97.8 (19.8)	0.46	97.9 (11.8)	0.28	96.0 (18.4)	0.17	102.1 (13.9)	0.36
Motor outcome <sup>4</sup>	87.7 (18.8)	<0.001	93.2 (12.2)	0.002	82.9 (16.7)	<0.001	98.2 (14.8)	0.45
Language outcome <sup>4</sup>	–	–	97.7 (8.6)	0.10	–	–	97.7 (12.6)	0.26
Receptive language <sup>5</sup>	–	–	9.4 (1.7)	0.04	–	–	10.2 (2.4)	0.60
Expressive language <sup>5</sup>	–	–	9.8 (1.9)	0.51	–	–	8.9 (2.5)	0.01
Fine motor skills <sup>5</sup>	–	–	10.1 (2.1)	0.76	–	–	10.5 (2.6)	0.23
Gross motor skills <sup>5</sup>	–	–	7.7 (2.5)	<0.001	–	–	8.9 (2.7)	0.01

<sup>1</sup> Cognitive outcome: n = 45; motor outcome: n = 42. <sup>2</sup> Outcome scores were compared to the expected normal score of reference population; Rotterdam: BSID-II-NL; Rome: BSID-III. <sup>3</sup> Cognitive outcome: n = 41; motor outcome: n = 46. <sup>4</sup> Mean (SD) of expected normal score of reference population is 100 (15). <sup>5</sup> Mean (SD) of expected normal score of reference population is 10 (3).

In Rotterdam, cognitive outcome at 24 months was negatively associated with low socioeconomic status (SES) (B -36.2; 95% CI: -58.1 to -14.3) and non-Dutch ethnicity (B 25.1; 95% CI: 5.9 to 44.2; table 3). Motor function was negatively associated with longer LoS at 24 months (B -7.8; 95% CI: -15.3 to -0.3). For example, taking into account the logarithmic transformation of LoS, children who were hospitalized twice as long scored 5.4 points lower (calculation:  $B \times \ln 2 = -7.8 \times \ln 2$ ) on motor function at 24 months. In non-ECMO-treated patients, the SNAP-II score was significantly associated with cognition and motor function at 12 months. At 24 months, low SES and non-Dutch ethnicity were associated with poor cognition, and low SES and LoS were associated with poor motor function (online suppl. table 3a). In Rome, longer LoS was significantly associated with cognition at 12 months (B -13.7; 95% CI: -21.2 to -6.2) and 24 months (B -10.2; 95% CI: -20.0 to -0.4), and longer LoS and motor function at 12 months (B -9.5; 95% CI: -17.7 to -1.4) and 24 months (B -16.7; 95% CI: -25.1 to -8.2).

## Discussion

This longitudinal study was performed in two European high-volume centers that use the same neonatal treatment protocol [1] in high-risk CDH patients. We found normal cognition in 78% of children aged 12 months in Rotterdam and 95% in Rome. At 24 months this was 71% in Rotterdam and 97% in Rome. Normal development of motor function occurred in 64% in Rot-

terdam and 82% in Rome at 12 months. At 24 months, this was 46% in Rotterdam and 90% in Rome.

A few cross-sectional studies have reported neurodevelopmental outcome in CDH patients [8–10, 19]. Danzer et al. [9] reported delayed cognition in 32% of 41 children, assessed with BSID-II or BSID-III at a median age of 24 months, and 24% of 42 prospectively enrolled patients at 2–4 years assessed with BSID-III [8]. The prevalence of cognitive problems using similar outcome categories was comparable for Rotterdam, whilst fewer patients in Rome had cognitive delays. It is possible that patients in the study by Danzer et al. [8] were more severely ill since they were ventilated longer and more infants received ECMO. In a multicenter study by Wynn et al. [19], mean cognitive scores – obtained with BSID-III in 48 CDH patients at 24 months – were significantly below normal (mean 93; SD 15). We found the same trend; although different BSID versions were applied, results from Rotterdam are comparable and results from Rome are better. Many factors may contribute to cognitive delay in CDH. Consistent with previous studies, low SES was a significant determinant of cognitive delay [19]. Previously reported predictive factors, such as tube feeding at discharge and ECMO need [19], were not confirmed in our study. However, failure to reach statistical significance for ECMO need may have been due to the fact that 6 survivors needed ECMO treatment.

In general, impaired motor function is more frequent than cognitive problems in CDH. Our results on motor function were more favorable than those of Danzer et al. [9], who reported mild (23%) and severe (31%) delay on



**Table 3.** Determinants of impaired neurodevelopmental outcome; results of linear regression models

	Independent variables	Rotterdam		Rome	
		Parameter estimate	95% CI	Parameter estimate	95% CI
Cognition at 12 months	SNAP-II score	0.5	-0.2 to 1.2	0.1	-0.0 to 0.33
	ECMO	2.3	-25.8 to 30.3	-	-
	Length of hospital stay, days <sup>1</sup>	-9.6	-20.0 to 0.7	-13.7	-21.2 to -6.2
	Tube feeding at discharge	-0.2	-19.2 to 19.0	-1.9	-10.6 to 6.8
	Ethnicity	13.3	-7.7 to 34.3	1.5	-8.1 to 11.1
	SES mother				
	Low	-11.5	-34.9 to 11.9	1.9	-8.4 to 12.1
	Medium	4.6	-16.1 to 25.2	1.2	-7.7 to 10.1
High <sup>2</sup>	-	-	-	-	
Cognition at 24 months	SNAP-II score	0.1	-0.4 to 0.5	0.0	-0.2 to 0.3
	ECMO	20.8	-9.0 to 50.6	-	-
	Length of hospital stay, days <sup>1</sup>	-5.8	-14.1 to 2.6	-10.2	-20.0 to -0.4
	Tube feeding at discharge	-3.5	-18.4 to 11.3	-4.1	-16.2 to 8.0
	Ethnicity	25.1	5.9 to 44.2	-2.1	-15.5 to 11.4
	SES mother				
	Low	-36.2	-58.1 to -14.3	-7.0	-21.3 to 7.4
	Medium	-11.5	-27.9 to 4.8	-3.6	-16.0 to 8.7
High <sup>2</sup>	-	-	-	-	
Motor function at 12 months	SNAP-II score	0.6	-0.0 to 1.2	0.0	-0.2 to 0.2
	ECMO	5.6	-22.3 to 33.5	-	-
	Length of hospital stay, days <sup>1</sup>	-10.5	-21.6 to 0.5	-9.5	-17.7 to -1.4
	Tube feeding at discharge	-0.2	-18.4 to 17.9	-6.6	-16.1 to 2.8
	Ethnicity	15.3	-5.6 to 36.3	3.6	-6.8 to 14.1
	SES mother				
	Low	-0.8	-23.2 to 21.6	-3.1	-14.2 to 8.1
	Medium	1.2	-18.8 to 21.2	-2.4	-12.1 to 7.3
High <sup>2</sup>	-	-	-	-	
Motor function at 24 months	SNAP-II score	0.3	-0.2 to 0.8	0.1	-0.2 to 0.3
	ECMO	14.4	-7.8 to 36.6	-	-
	Length of hospital stay, days <sup>1</sup>	-7.8	-15.3 to -0.3	-16.7	-25.1 to -8.2
	Tube feeding at discharge	3.0	-11.0 to 17.0	0.8	-9.7 to 11.3
	Ethnicity	6.2	-7.4 to 19.9	2.1	-9.5 to 13.7
	SES mother				
	Low	-15.4	-32.1 to 1.4	-0.2	-12.7 to 12.2
	Medium	-2.7	-17.2 to 11.8	5.5	-5.2 to 16.2
High <sup>2</sup>	-	-	-	-	

SNAP-II score = Score for Neonatal Acute Physiology II. <sup>1</sup> Because of skewed distribution, these variables were log-transformed before including them into the linear regression model. <sup>2</sup> Reference category.

motor function at 24 months. Friedman et al. [3] reported motor problems in 60% at 1 year and 73% at 3 years. They retrospectively evaluated medical records documenting motor function. It is conceivable that standardized developmental assessment would have given other outcomes. On the other hand, Wynn et al. [19] found that motor function at 24 months was sig-

nificantly lower than the population norm (mean 95; SD 11), which is comparable to Rome. Scores in Rotterdam, however, were on average 12 points lower. We could not confirm their results of adverse motor function in patients with low SES, but could confirm the result of another study [4] that LoS was associated with poorer motor function.

The strengths of our study were that children were treated according to a standardized protocol [1] and followed in a prospective, longitudinal, standardized follow-up program. We collected data on ethnicity and SES, which are well-known modifiers of neurodevelopmental outcome [19].

Neurodevelopmental outcomes were different between the centers, and changes between the two measurement moments were only seen in Rome. We suggest three possible reasons. The first is the use of different assessment instruments. Since recent studies reported that BSID-II and BSID-III are not comparable [20], we refrained from pooling the data [1]. In the transition from BSID-II to BSID-III, 23 items from the BSID-II mental scale were moved to the BSID-III fine motor scale. Thus, with a higher proportion of items in the fine motor function domain in BSID-III compared to BSID-II (48 vs. 28%, respectively) the fine motor score contributes relatively much to the total motor composite score in BSID-III. This is reflected by the fact that in Rome gross motor scores are lower than fine motor scores at both ages. We assume that the outcome for patients who have a developmental delay in both cognition and motor function will not be very different for the two BSID versions. However, results may differ for patients with CDH, who will typically show impaired motor function. Moreover, recent studies showed differences in reference scores of BSID-III between populations [21, 22]. These considerations possibly explain the underestimation of neurodevelopmental delay in Rome. We have few standardized neurodevelopmental tests with population-matched reference data and the best option seems to compare neurodevelopmental outcome with healthy reference populations in each country. The use of American reference data in Rome might also have caused bias. In another study, neurodevelopmental outcomes were not different from those of healthy matched controls aged 12 and 36 months [10]. Disease-determining parameters were not reported, which precludes comparison with our study.

Second, professional background may have played a role. In Rotterdam, motor function at 24 months was assessed by a pediatric physical therapist, whereas in Rome assessments were performed by developmental psychologists. Since results of our study did not significantly differ from the PDI assessed by developmental psychologists in a previous study of our group [11], we assume that this may not fully explain differences in motor function. Moreover, this contradicts our assumption that neurodevelopmental outcome has worsened due to survival of severely ill patients [2].

Third, center-specific differences may have played a role. Although both tertiary intensive care centers have a comparable referral area, patients from Rotterdam may have been more critically ill than those from Rome (higher proportion of large diaphragmatic defects and higher patch use). However, SNAP-II scores and observed to expected lung-head ratio were not significantly different. The surgical approach differed, too. Primarily, a thoracoscopic approach was performed in more than 50% in Rotterdam versus open surgery in all patients in Rome. A randomized controlled trial of open versus thoracoscopic repair in CDH concluded that thoracoscopic repair was associated with more prolonged and severe intraoperative hypercapnia and acidosis than open surgery [23]. Moreover, only in Rotterdam was ECMO available. Since only 6 survivors received ECMO and patient characteristics did not significantly change after exclusion of these children, we assume that this hardly contributes to the outcome differences. The influence of center-specific differences such as surgical care on neurodevelopmental outcome remains speculative, but these observations emphasize the need for multicenter studies with standardized protocols. Since scores in Rotterdam slowly deteriorate at 24 months, and it is known that early adverse development in children with major congenital anomalies is predictive of development at 5 years [5], follow-up was planned for patients from Rotterdam.

Next to the differences in study design, the fact that not every child was tested at both time points could be considered a limitation. However, the numbers of drop-outs were relatively low and the use of linear mixed models in the analysis over time accounted for missing data, provided that these outcomes were missing at random [24].

Interpreting outcomes in the context of international multicenter studies is difficult if the follow-up program is not standardized. International guidelines on standardization of follow-up programs in CDH, however, are still lacking. So far, the international CDH registry [25] collects only prenatal, perinatal and early postnatal data. We recommend setting up standardized follow-up programs using population-appropriate reference data and similar assessment instruments as well as assessments performed by professionals of the same background. In multicenter collaboration, video assessments should be included in the training sessions. We like to make an appeal for not only international standardization of postnatal care, but also for follow-up care beyond the neonatal period, as this is essential to improve outcome for CDH patients. This would require standardization of treatment protocols

and decision flowcharts of referring criteria to pediatric physical therapists/speech-language pathologists should be involved. Moreover, children should be followed up to adolescence since deficits may worsen.

In conclusion, although most CDH patients have normal neurodevelopment within the first 2 years of life, they

are at risk for impaired motor function. Standardization of multicenter long-term follow-up programs using standardized assessment instruments and stratification to illness severity is necessary to compare neurodevelopmental outcomes between centers and to evaluate long-term effects of interventions.

## References

- 1 Reiss I, Schaible T, van den Hout L, Capolupo I, Allegaert K, van Heijst A, Gorett Silva M, Greenough A, Tibboel D; CHE EURO Consortium: Standardized postnatal management of infants with congenital diaphragmatic hernia in Europe: The CDH EURO Consortium consensus. *Neonatology* 2010;98:354–364.
- 2 van den Hout L, Schaible T, Cohen-Overbeek TE, Hop W, Siemer J, van de Ven K, Wessel L, Tibboel D, Reiss I: Actual outcome in infants with congenital diaphragmatic hernia: the role of a standardized postnatal treatment protocol. *Fetal Diagn Ther* 2011;29:55–63.
- 3 Friedman S, Chen C, Chapman JS, Jeruss S, Terrin N, Tighiouart H, Parsons SK, Wilson JM: Neurodevelopmental outcomes of congenital diaphragmatic hernia survivors followed in a multidisciplinary clinic at ages 1 and 3. *J Pediatr Surg* 2008;43:1035–1043.
- 4 Danzer E, Gerdes M, D'Agostino JA, Hoffman C, Bernbaum J, Bebbington MW, Siegle J, Sulkowski J, Rintoul NE, Flake AW, Scott Adzick N, Hedrick HL: Longitudinal neurodevelopmental and neuromotor outcome in congenital diaphragmatic hernia patients in the first 3 years of life. *J Perinatol* 2013;33:893–898.
- 5 Mazer P, Gischler SJ, Van der Cammen-Van Zijp MH, Tibboel D, Bax NM, IJsselstijn H, Van Dijk M, Duivenvoorden HJ: Early developmental assessment of children with major non-cardiac congenital anomalies predicts development at the age of 5 years. *Dev Med Child Neurol* 2010;52:1154–1159.
- 6 Spoel M, van der Cammen-van Zijp MH, Hop WC, Tibboel D, de Jongste JC, IJsselstijn H: Lung function in young adults with congenital diaphragmatic hernia; a longitudinal evaluation. *Pediatr Pulmonol* 2013;48:130–137.
- 7 Peetsold MG, Kneepkens CM, Heij HA, IJsselstijn H, Tibboel D, Gemke RJ: Congenital diaphragmatic hernia: long-term risk of gastroesophageal reflux disease. *J Pediatr Gastroenterol Nutr* 2010;51:448–453.
- 8 Danzer E, Gerdes M, D'Agostino JA, Partridge EA, Hoffman-Craven CH, Bernbaum J, Rintoul NE, Flake AW, Adzick NS, Hedrick HL: Preschool neurological assessment in congenital diaphragmatic hernia survivors: outcome and perinatal factors associated with neurodevelopmental impairment. *Early Hum Dev* 2013;89:393–400.
- 9 Danzer E, Gerdes M, Bernbaum J, D'Agostino J, Bebbington MW, Siegle J, Hoffman C, Rintoul NE, Flake AW, Adzick NS, Hedrick HL: Neurodevelopmental outcome of infants with congenital diaphragmatic hernia prospectively enrolled in an interdisciplinary follow-up program. *J Pediatr Surg* 2010;45:1759–1766.
- 10 Leeuwen L, Walker K, Halliday R, Fitzgerald DA: Neurodevelopmental outcome in congenital diaphragmatic hernia survivors during the first three years. *Early Hum Dev* 2014;90:413–415.
- 11 Gischler SJ, Mazer P, Duivenvoorden HJ, van Dijk M, Bax NM, Hazebroek FW, Tibboel D: Interdisciplinary structural follow-up of surgical newborns: a prospective evaluation. *J Pediatr Surg* 2009;44:1382–1389.
- 12 Valfre L, Braguglia A, Conforti A, Morini F, Trucchi A, Iacobelli BD, Nahom A, Chukhlantseva N, Dotta A, Corchia C, Bagolan P: Long term follow-up in high-risk congenital diaphragmatic hernia survivors: patching the diaphragm affects the outcome. *J Pediatr Surg* 2011;46:52–56.
- 13 Ruiter SSH, van der Meulen B, Nakken H: The BSID-II NL: construction, standardisation, and instrumental utility. *Neth J Psychol* 2008;64:15–40.
- 14 Bayley N: Bayley Scales of Infant and Toddler Development (BSID-III), ed 3. San Antonio, Harcourt Assessment, 2006.
- 15 Jani J, Nicolaides KH, Keller RL, Benachi A, Peralta CF, Favre R, Moreno O, Tibboel D, Lipitz S, Eggink A, Vaast P, Allegaert K, Harrison M, Deprest J; Antenatal-CDH-Registry Group: Observed to expected lung area to head circumference ratio in the prediction of survival in fetuses with isolated diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2007;30:67–71.
- 16 Richardson DK, Corcoran JD, Escobar GJ, Lee SK: SNAP-II and SNAPPE-II: simplified newborn illness severity and mortality risk scores. *J Pediatr* 2001;138:92–100.
- 17 Jobe AH, Bancalari E: Bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 2001;163:1723–1729.
- 18 Centraal Bureau voor de Statistiek (Statistics Netherlands) Standaard Onderwijsindeling 2006 (The Dutch Standard Classification of Education). [www.cbs.nl/nl-NL/menu/methoden/classificaties/overzicht/soi/2003/default.htm](http://www.cbs.nl/nl-NL/menu/methoden/classificaties/overzicht/soi/2003/default.htm) (accessed on January 6, 2015).
- 19 Wynn J, Aspelund G, Zygmunt A, Stolar CJ, Mychaliska G, Butcher J, Lim FY, Gratton T, Potoka D, Brennan K, Azarow K, Jackson B, Needelman H, Crombleholme T, Zhang Y, Duong J, Arkovitz MS, Chung WK, Farkouh C: Developmental outcomes of children with congenital diaphragmatic hernia: a multicenter prospective study. *J Pediatr Surg* 2013;48:1995–2004.
- 20 Lowe JR, Erickson SJ, Schrader R, Duncan AF: Comparison of the Bayley II Mental Developmental Index and the Bayley III Cognitive Scale: are we measuring the same thing? *Acta Paediatr* 2012;101:e55–e58.
- 21 Anderson PJ, De Luca CR, Hutchinson E, Roberts G, Doyle LW; Victorian Infant Collaborative Group: Underestimation of developmental delay by the new BAYLEY-III Scale. *Arch Pediatr Adolesc Med* 2010;164:352–356.
- 22 Chinta S, Walker K, Halliday R, Loughran-Fowlds A, Badawi N: A comparison of the performance of healthy Australian 3-year-olds with the standardised norms of the Bayley Scales of Infant and Toddler Development (version-III). *Arch Dis Child* 2014;99:621–624.
- 23 Bishay M, Giacomello L, Retrosi G, Thyoka M, Garriboli M, Brierley J, Harding L, Scuplak S, Cross KM, Curry JI, Kiely EM, De Coppi P, Eaton S, Pierro A: Hypercapnia and acidosis during open and thoracoscopic repair of congenital diaphragmatic hernia and esophageal atresia: results of a pilot randomized controlled trial. *Ann Surg* 2013;258:895–900.
- 24 Cnaan A, Laird NM, Slasor P: Using the general linear mixed model to analyse unbalanced repeated measures and longitudinal data. *Stat Med* 1997;16:2349–2380.
- 25 Harting MT, Lally KP: The congenital diaphragmatic hernia study group registry update. *Semin Fetal Neonatal Med* 2014;19:370–375.