

Effect of prednisolone on inflammatory markers in pericardial tuberculosis: A pilot study

Justin Shenje^a, Rachel P. Lai^b, Ian L. Ross^a, Bongani M. Mayosi^a, Robert J. Wilkinson^{a,b,c,d}, Mpiko Ntsekhe^a, Katalin A. Wilkinson^{a,b,c,*}

^a The Cardiac Clinic, Department of Medicine, University of Cape Town, Grootte Schuur Hospital, Observatory 7925, South Africa

^b The Francis Crick Institute, NW1 2AT London, United Kingdom

^c Wellcome Centre for Infectious Diseases Research in Africa, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Observatory 7925, South Africa

^d Imperial College London, W2 1PG, United Kingdom

ARTICLE INFO

Article history:

Received 6 June 2017

Received in revised form 27 September 2017

Accepted 22 October 2017

Available online 3 November 2017

Keywords:

Tuberculosis

HIV

Pericarditis

Steroids

Treatment monitoring

ABSTRACT

Background: Pericardial disorders are a common cause of heart disease, and the most common cause of pericarditis in developing countries is tuberculous (TB) pericarditis. It has been shown that prednisolone added to standard anti-TB therapy leads to a lower rate of constrictive pericarditis. We conducted a pilot study to evaluate the effect of adjunctive prednisolone treatment on the concentration of inflammatory markers in pericardial tuberculosis, in order to inform immunological mechanisms at the disease site.

Methods: Pericardial fluid, plasma and saliva samples were collected from fourteen patients with pericardial tuberculosis, at multiple time points. Inflammatory markers were measured using multiplex luminex analysis and ELISA.

Results: In samples from 14 patients we confirmed a strongly compartmentalized immune response at the disease site and found that prednisolone significantly reduced IL-6 concentrations in plasma by 8 hours of treatment, IL-1beta concentrations in saliva, as well as IL-8 concentrations in both pericardial fluid and saliva by 24 hours.

Conclusion: Monitoring the early effect of adjunctive immunotherapy in plasma or saliva is a possibility in pericarditis.

© 2017 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Tuberculosis (TB) is the most common opportunistic infection in HIV-1 infected persons in Sub-Saharan Africa. Tuberculous pericarditis is an inflammation of the pericardium caused by *Mycobacterium tuberculosis*. It is a life-threatening extrapulmonary form of TB that results in accumulation of fluid around the heart, potentially leading to constriction. In developed countries it accounts for ~5% of all cases of acute pericarditis, compared to up to 90% in Sub-Saharan Africa, where it is the most common cause of pericardial effusions in HIV-1 co-infected patients [1]. Mortality in these patients is up to 40% in the absence of anti-retroviral treatment, despite antituberculosis therapy, pericardial drainage, or pericardiectomy [2].

The recently completed Investigation of the Management of Pericarditis (IMPI), a large clinical trial examining the effect of prednisolone or *Mycobacterium indicus pranii* or both added to the standard regimen of isoniazid, rifampin, ethambutol and pyrazinamide showed, that neither

the standard anti-TB therapy alone nor the addition of prednisolone to chemotherapy resulted in a clinically satisfactory mortality reduction, however the prednisolone group had a lower rate of constrictive pericarditis and fewer hospitalisations compared to placebo [3]. Since we have recently shown that pericardial tuberculosis is characterised by a compartmentalized profibrotic immune response, we hypothesised that prednisolone had a suppressive effect on the concentration of inflammatory and potentially profibrotic cytokines in the pericardium [4].

In this pilot study we evaluated the effect of prednisolone on inflammatory markers in pericardial fluid, plasma and saliva, in a subset of patients from the above clinical trial in order to improve our understanding of immunological mechanisms at the disease site, which could inform development of more targeted interventions. Specific analytes were selected based on our recent analysis of differentially abundant inflammatory markers (at both RNA and protein level) between the blood and pericardial fluid compartments [4]. Additionally, we also evaluated the hypothesis that inflammation induced cell death in the pericardial compartment would be in part due to apoptosis, which is initiated by two major pathways: the extrinsic (through ligation of death receptors or TNF receptors) and intrinsic pathways (mitochondria mediated) [5]. Caspase 8 activation is an essential early step in the induction of the

* Corresponding author at: Institute of Infectious Disease and Molecular Medicine, Room N2.09.B3 Wernher Beit North Building, Observatory 7925, South Africa.
E-mail address: katalin.wilkinson@uct.ac.za (K.A. Wilkinson).

apoptosis by the extrinsic pathway, while Caspase 9 activation is part of the intrinsic pathway, both leading to activation of Caspase 3 [6]. We therefore also assessed the effect of prednisolone on the concentration of Caspases 3, 8 and 9 in the pericardial fluid and plasma.

2. Materials and methods

2.1. Patient population

Patients were recruited from Groote Schuur Hospital, Cape Town, South Africa as part of an intensive pharmacokinetic sampling study of the Investigation of the Management of Pericarditis (IMPI) trial, with a computer-generated randomisation list as described previously [3,7]. Ethical approval for these studies was obtained from the Faculty of Health Sciences Human Research Ethical Committee at the University of Cape Town (Reference numbers 102/2003, 402/2005, 289/2007) and as published [3,7]. Briefly, patients were eligible for inclusion in the trial if they were 18 years of age or older, had a pericardial effusion requiring pericardiocentesis confirmed by echocardiography, had evidence of definite or probable tuberculous pericarditis and started antituberculosis treatment less than a week before enrolment into the comparison of prednisolone versus placebo arm of the IMPI trial and provided written informed consent for inclusion in the study. Pericardial fluid, plasma and saliva were obtained from 14 patients at multiple time points following randomisation to prednisolone/placebo for pharmacokinetic studies to assess antibiotic penetration as described [7]. Pericardial samples were collected via a catheter which was left in the pericardial space for 24 h post-pericardiocentesis. Not all samples were available from all patients at all time points, thus the present analysis evaluated the concentration of cytokines, chemokines and caspases at 0, 8 and 24 h only.

2.2. Luminex multiplex assay for cytokines and chemokines

Mediators analysed in undiluted plasma (P), pericardial fluid (PF) and saliva (S) samples included IFN- γ , IL-1 α , IL-1 β , IL-6, IL-10, IL-12p40, TNF, CXCL8 (IL-8) and CXCL10 (IP-10), using customized Milliplex™ kits (HCYTOMAG-60 K, Millipore, St Charles, MO, USA) on the Bio-Plex platform (Bio-Rad Laboratories, Hercules, CA, USA) as described [8]. Caspases 3, 8 and 9 in undiluted pericardial fluid and plasma were measured using human in vitro ELISA kits from Abcam (Cambridge, UK), following the manufacturer's recommendations.

2.3. Statistical analysis

Statistical analysis was performed using GraphPad Prism Version 7.0c for Mac. The normality of data was assessed using the D'Agostino

and Pearson normality test. As not all patients were sampled at all time points, some data were not paired. Between group comparisons of unpaired non-normally distributed data were analysed using the Mann-Whitney *U* test. No correction for multiple comparisons was performed.

3. Results

Of the 14 patients nine were on placebo and five on prednisolone (at a dose of 120 mg per day in the first week). Patient characteristics were previously described as part of the main study [7], with the subset included in this pilot study being detailed in Table 1. Nine patients had definite tuberculous pericarditis (3 on prednisolone and 6 on placebo) and five had probable tuberculous pericarditis (2 on prednisolone and 3 on placebo). The pericardial protein measured in pericardial fluid was median 61 g/l (IQR 55–67.5) in the definite TB patients and median 66 g/l (IQR 56–76) in the probable TB patients respectively, while the ADA was median 87.3 U/l (IQR 43–127) and 51 U/l (32.8–67.6) in the definite and probable patients respectively. There was no significant difference between the groups with respect to either parameter. There was no difference between age and weight between the groups, gender composition was male/female 5/4 and 2/3 for the placebo and prednisolone groups respectively. Seven out of nine patients in the placebo and three out of five patients in the prednisolone group were HIV infected. Only two patients were receiving antiretroviral treatment at the time of the study, both in the placebo group (Table 1). Median CD4 counts were not different between the groups (240 cells/ml for the placebo and 139 cells/ml for the prednisolone group). Since our previous findings indicated that neither HIV-1 coinfection status, nor CD4 count or pericardial fluid *Mycobacterium tuberculosis* culture result affected the compartmentalized profibrotic immune response [4], we combined patients with and without HIV infection for analysis, as well as patients with definite and probable pericarditis.

Results are summarised in Table 2, indicating the median (IQR) of all analytes measured in all samples at the tested time points. Since samples were collected before prednisolone administration, we combined the baseline data to analyse the effect of compartmentalization: higher concentrations of IFN- γ , IL-10, IL-1 β , IL-6, IL-8, IP-10 and TNF were found at the disease site compared to plasma at day 0, supporting our previous findings [4]. Interestingly, IL-1 α and IL-12p40 was mostly detectable in the saliva samples, which also contained elevated concentrations of IL-1 β , IL-8 and IP-10 compared to plasma. Only IL-6 concentrations were higher in the plasma, compared to saliva ($p = 0.03$) at baseline. Caspase 3, 8 and 9 were only detectable in the pericardial fluid as opposed to plasma ($p = 0.02, 0.0001$ and 0.03 respectively for the three caspases measured).

Table 1
Patient characteristics.

Patient number	Age (years)	Wt (kg)	Gender	HIV	HAART	ART therapy	CD4 (cells/ μ l)	Creatinine (μ mol/l)	Globulin (g/l)	Steroid allocation	Pericardial protein (g/l)	ADA (units/l)	TB microscopy
1	28	53	male	Positive	No	N/A	115	76	46	Placebo	58	Hemolyzed	Positive
2	29	40	female	Positive	No	N/A	50	20	45	Placebo	54	83	Positive
3	24	66	female	Positive	No	N/A	42	43	56	Prednisolone	68	57	Positive
4	56	82	male	Negative	N/A	N/A	485	121	30	Placebo	62	133	Positive
5	31	72	male	Negative	N/A	N/A	319	82	36	Prednisolone	55	119	Positive
6	51	53	female	Positive	Yes	Tdf/FTC/EFV	159	80	56	Placebo	50	53	Negative
7	24	45	female	Positive	No	N/A	321	257	56	Placebo	67	92	Positive
8	27	66	female	Positive	No	N/A	135	45	55	Prednisolone	70	26	Positive
9	44	52	female	Positive	No	N/A	139	65	56	Prednisolone	66	51	Negative
10	59	66	male	Negative	N/A	N/A	874	97	38	Prednisolone	62	33	Negative
11	45	70	male	Negative	N/A	N/A	721	109	47	Placebo	76	32	Negative
12	33	47	male	Positive	No	N/A	116	71	49	Placebo	61	130	Positive
13	25	73	male	Positive	No	N/A	^a	73	^a	Placebo	55	38	Positive
14	27	^a	female	Positive	Yes	Tdf/FTC/EFV	255	41	74	Placebo	76	68	Negative

^a Data not available.

Table 2
Median (IQR) of all analytes measured in all samples.

Site	N	Time	Group	IFN- γ pg/ml	IL-1 α pg/ml	IL-1 β pg/ml	IL-6 pg/ml	IL-10 pg/ml	IL-12p40 pg/ml	TNF pg/ml	IL-8 pg/ml	IP-10 pg/ml	Caspase 3 pg/ml	Caspase 8 ng/ml	Caspase 9 ng/ml	
Pericardial fluid	14	D0	Combined	2061	0	15	9064	26	0	197	7670	1414	93	7	35	
				648–3285	0–62	0–114	8225–10,000	21–65	0–5	100–354	1841–10,434	1013–1646	0–313	5–16	30–73	
				1585	0	0	9277	28	0	180	3725	1435	134	6	31	
	5	D0	Prednisolone	533–6363	0–149	0–109	8533–9969	22–47	0–3	107–469	1512–9987	846–1537	0–250	5–14	24–82	
				1436	0	0	9166	37	0	218	4635	1495	193	8	38	
				383–2903	0–213	0–177	8269–9390	25–56	0–0	142–483	2876–9677	1050–1701	0–193	6–13	26–61	
	3	8 h	Placebo	838	0	0	8797	24	0	265	1834	1245	48	6	32	
				278–2696	0–20	0–42	8102–9755	13–31	0–0	141–394	1576–2890	865–1335	0–137	4–9	24–42	
				2536	0	18	8850	22	0	213	9329	1348	88	7	36	
	9	D0	Placebo	553–3687	0–111	7–292	7243–10,254	17–83	0–10	99–291	1903–10,855	790–1887	0–403	5–20	30–91	
				1347	2	14	9097	29	0	182	2603	1462	72	10	41	
				298–3485	0–18	8–74	8058–9483	11–125	0–3	96–353	2334–9106	657–1972	0–216	8–17	31–70	
	6	24 h	Placebo	927	8	58	8695	16	1	175	7067	1758	168	11	48	
				482–2619	0–124	14–185	6377–9394	8–46	0–18	83–299	4180–9957	908–1944	0–684	4–21	32–61	
				28	0	0	23	4	0	33	23	82	0	0	24	
	Plasma	12	D0	Combined	10–116	0–0	0–0	16–61	0–7	0–0	21–53	11–33	40–158	0–0	0–2	19–40
					84	0	0	20	3	0	35	21	77	0	0	29
					28–231	0–0	0–5	13–23	0–12	0–0	28–67	16–30	26–127	0–56	0–6	23–55
3		8 h	Placebo	72	0	0	4	0	0	23	10	52	0	2	49	
				14–102	0–0	0–5	0–15	0–8	0–0	15–25	10–25	33–72	0–0	1–4	17–55	
				37	0	0	6	0	0	19	28	49	0	2	25	
3		24 h	Placebo	9–259	0–0	0–5	6–73	0–7	0–0	19–57	16–54	21–59	0–470	0–2	14–34	
				13	0	0	35	4	0	26	25	86	0	0	20	
				5–118	0–0	0–0	19–88	0–7	0–0	12–55	5–72	36–305	0–0	0–0	18–34	
5		8 h	Placebo	16	0	0	49	6	0	22	20	68	0	2	23	
				4–117	0–0	0–0	19–67	0–10	0–0	14–94	8–99	36–214	0–109	0–3	19–45	
				11	0	0	44	1	0	20	19	47	0	0	20	
6		24 h	Placebo	3–83	0–0	0–0	17–74	0–3	0–0	17–68	16–30	24–288	0–0	0–2	15–22	
				17	3393	28	11	15	18	15	295	1123	nd	nd	nd	
				10–23	949–11,180	18–144	2–28	1–25	0–39	9–80	134–961	167–2316	nd	nd	nd	
Saliva		5	D0	Prednisolone	18	1758	26	10	21	31	13	325	1061	nd	nd	nd
					10–30	980–8567	15–663	5–29	15–57	0–48	10–47	88–1257	201–2420	nd	nd	nd
					17	341	12	5	37	37	14	52	139	nd	nd	nd
	3	8 h	Placebo	17–35	336–5752	5–20	1–10	4–40	10–86	7–20	32–55	45–762	nd	nd	nd	
				10	390	11	6	28	23	12	44	92	nd	nd	nd	
				1–36	59–3341	2–14	1–11	2–63	0–88	3–15	3–117	30–552	nd	nd	nd	
	7	D0	Placebo	15	6473	30	11	3	11	59	265	1184	nd	nd	nd	
				0–23	722–12,416	17–146	0–29	0–22	0–39	6–210	156–486	67–2361	nd	nd	nd	
				16	1399	14	8	4	0	22	140	345	nd	nd	nd	
	6	8 h	Placebo	13–17	401–13,354	4–56	4–30	1–37	0–2	4–110	22–234	90–1095	nd	nd	nd	
				15	2890	26	10	5	0	26	163	376	nd	nd	nd	
				11–23	531–7632	17–105	2–12	0–77	0–32	5–56	94–361	240–2661	nd	nd	nd	

N: number of patient samples available for analysis at specific time point; D0: day 0; nd: not done.

Prednisolone significantly decreased the concentration of IL-6 by 8 h in plasma, compared to the patients who received placebo (Fig. 1A, $p = 0.036$, Mann Whitney *U* test). In pericardial fluid, prednisolone significantly reduced IL-8 concentrations by 24 h (Fig. 1B, $p = 0.03$, Mann Whitney *U* test), compared to the placebo treated patients. There was a trend towards decreased IL-1beta concentrations, however significance was not reached due to low numbers ($p = 0.06$, not shown). Finally in saliva, we found a significant reduction in concentration of both IL-8 and IL-1beta by 24 h of treatment in the prednisolone group (Fig. 1C, $p = 0.04$, Mann Whitney *U* test; and Fig. 1D, $p = 0.027$, Mann Whitney *U* test), compared to the placebo group. The concentration of caspase 3, 8 and 9 did not appear to be affected by prednisolone in any of the compartments evaluated.

4. Discussion

The largest clinical trial evaluating the effect of adjunctive prednisolone therapy showed no effect on the primary composite of death, cardiac tamponade requiring pericardiocentesis or constrictive pericarditis,

however, with respect to the secondary outcomes, the IMPI trial demonstrated that prednisolone reduced the incidence of constrictive pericarditis and the incidence of hospitalisations [3]. Here we hypothesised that the antiinflammatory effects of prednisolone would also manifest in reduced measurable concentrations of inflammatory cytokines and chemokines. Our findings suggest that the beneficial effect of prednisolone is associated with the suppression of inflammatory mediators IL-6, IL-8 and IL-1beta.

These results support our previous findings described during the treatment of tuberculosis associated immune reconstitution inflammatory syndrome (TB-IRIS) with prednisone vs placebo [9], where prednisone reduced the duration of hospitalisation and the number of outpatient therapeutic procedures. At the same time, IL-6, IL-10, IL-12 p40, TNF, IFN-gamma, and IFN-gamma-induced protein-10 (IP-10, CXCL10) concentrations significantly decreased in the serum of prednisone, but not placebo, treated patients [10]. In a separate study we found that adjunct corticosteroid therapy modifies the inflammatory profile of those who develop TB-IRIS, with lower concentrations of IFN-gamma, IP-10, TNF, IL-6, IL-8, IL-10, IL-12p40, and IL-18 [11]. Thus, the beneficial




Fig. 1. Panel A. Concentration of IL-6 (pg/ml) in plasma at baseline, 8 and 24 h after initiation of prednisolone treatment, which resulted in significant reduction by 8 h ($p = 0.036$, Mann Whitney test). Concentration of IL-8 (pg/ml) in pericardial fluid (Panel B) and saliva (Panel C) at baseline, 8 and 24 h after initiation of prednisolone treatment, which led to significant decrease by 24 h in both pericardial fluid and saliva ($p = 0.03$ and 0.04 respectively, Mann Whitney test). Panel D. Concentration of IL-1beta (pg/ml) in saliva at baseline, 8 and 24 h after initiation of prednisolone treatment, leading to significant decrease by 24 h ($p = 0.027$, Mann Whitney test).

effects of prednisone appear to be mediated via suppression of predominantly proinflammatory cytokine responses of innate immune origin, similar to our current findings.

We reported earlier that cell-death enrichment factors were elevated in pericardial fluid [4] and hypothesised that TB antigen specific activated T cells that enter the pericardium and release INF-gamma die, thus potentially activating the inflammasome pathway resulting in pyroptosis and release of IL-1beta, ultimately leading to more cell death. Here we show a trend towards decreased IL-1beta concentrations in the pericardial fluid of prednisolone treated patients, as well as higher concentrations of caspases 3, 8 and 9 in the pericardial fluid, thereby supporting our hypothesis for compartmentalized inflammation induced apoptosis in the pericardial fluid.

This report is limited by the small sample size therefore the statistical significance of the findings is weak. The small sample size further limited our capacity to analyse the results based on the HIV status of the patients, since it has been described that HIV infection is associated with a lower incidence of pericardial constriction in patients with presumed tuberculous pericarditis [12], and IL-13-secreting CD4 T cells, which are reduced by HIV, regulate fibrogenesis directly, through stimulating collagen synthesis by fibroblasts and indirectly by promoting TGF-beta1 production by macrophages [13]. However, we based our analysis strategy on our previous findings, that showed neither HIV-1 coinfection status, nor CD4 count or pericardial fluid *Mycobacterium tuberculosis* culture result affected the compartmentalized profibrotic immune response with respect to the selected analytes we measured [4]. Additional limitation of this pilot study is that pericardial fluid sampling was restricted within 24 h of initiation of prednisolone therapy due to safety concerns over leaving the pericardial catheter in the pericardial space for a prolonged period of time.

These findings however are of interest because TB remains the leading cause of constrictive pericarditis in Africa. The treatment for chronic pericardial tuberculosis leading to constriction is pericardiectomy, which is associated with high mortality and morbidity but also problematic due to the fact that cardiac surgery is not widely available in Africa [14]. Thus adjunctive immunotherapy that reduces the incidence of constrictive pericarditis, and which could be monitored by measuring inflammatory markers in easily obtainable samples such as plasma or saliva, might be beneficial during early patient follow up in reducing morbidity and possibly even mortality.

Author contribution

JS, ILR, BMM, RJW, MN and KAW conceived and designed the study. JS, ILR, NP, BMM recruited, sampled and collected data from patients. KAW performed the experiments with input into experimental design from

RPL, KAW and RJW analysed the data. KAW, RJW wrote the manuscript, which was revised by all authors.

Funding

This work was supported by the a South African Medical Research Council Self-Initiated Award; The Francis Crick Institute (FC00110218); The Wellcome Trust (104803, 203135); National Research Foundation of South Africa (96841); European Union TBVAC2020 (643381).

Conflict of interest

The authors report no relationships that could be construed as a conflict of interest.

References

- [1] G. Lazaros, D. Tousoulis, Tuberculous pericarditis: a complex puzzle to put together, *EBioMedicine* 2 (2015) 1570–1571.
- [2] B.M. Mayosi, C.S. Wiysonge, M. Ntsekhe, et al., Mortality in patients treated for tuberculous pericarditis in sub-Saharan Africa, *S. Afr. Med. J.* 98 (2008) 36–40.
- [3] B.M. Mayosi, M. Ntsekhe, J. Bosch, et al., Prednisolone and *Mycobacterium indicus pranii* in tuberculous pericarditis, *N. Engl. J. Med.* 371 (2014) 1121–1130.
- [4] K. Matthews, A. Deffur, M. Ntsekhe, et al., A compartmentalized profibrotic immune response characterizes pericardial tuberculosis, irrespective of HIV-1 infection, *Am. J. Respir. Crit. Care Med.* 192 (2015) 1518–1521.
- [5] M. Divangahi, S.M. Behar, H. Remold, Dying to live: how the death modality of the infected macrophage affects immunity to tuberculosis, *Adv. Exp. Med. Biol.* 783 (2013) 103–120.
- [6] E.M. Creagh, Caspase crosstalk: integration of apoptotic and innate immune signalling pathways, *Trends Immunol.* 35 (2014) 631–640.
- [7] J. Shenje, F. Ifeoma Adimora-Nweke, I.L. Ross, et al., Poor penetration of antibiotics into pericardium in pericardial tuberculosis, *EBioMedicine* 2 (2015) 1640–1649.
- [8] S. Marais, K.A. Wilkinson, M. Lesosky, et al., Neutrophil-associated central nervous system inflammation in tuberculous meningitis immune reconstitution inflammatory syndrome, *Clin. Infect. Dis.* 59 (2014) 1638–1647.
- [9] G. Meintjes, R.J. Wilkinson, C. Morroni, et al., Randomized placebo-controlled trial of prednisone for paradoxical tuberculosis-associated immune reconstitution inflammatory syndrome, *AIDS* 24 (2010) 2381–2390.
- [10] G. Meintjes, K.H. Skolimowska, K.A. Wilkinson, et al., Corticosteroid-modulated immune activation in the tuberculosis immune reconstitution inflammatory syndrome, *Am. J. Respir. Crit. Care Med.* 186 (2012) 369–377.
- [11] A. Conesa-Botella, G. Meintjes, A.K. Coussens, et al., Corticosteroid therapy, vitamin D status, and inflammatory cytokine profile in the HIV-tuberculosis immune reconstitution inflammatory syndrome, *Clin. Infect. Dis.* 55 (2012) 1004–1011.
- [12] M. Ntsekhe, C.S. Wiysonge, F. Gumedze, et al., HIV infection is associated with a lower incidence of constriction in presumed tuberculous pericarditis: a prospective observational study, *PLoS One* 3 (2008), e2253.
- [13] T.A. Wynn, Fibrotic disease and the T(H)1/T(H)2 paradigm, *Nat. Rev. Immunol.* 4 (2004) 583–594.
- [14] A.K. Mutyaba, S. Balkaran, R. Cloete, et al., Constrictive pericarditis requiring pericardiectomy at Groote Schuur Hospital, Cape Town, South Africa: causes and perioperative outcomes in the HIV era (1990–2012), *J. Thorac. Cardiovasc. Surg.* 148 (2014) 3058–3065.e1.