

Coalgebraic completeness-via-canonicity for distributive substructural logics

Fredrik Dahlqvist, David Pym

University College London

Abstract

We prove strong completeness of a range of substructural logics with respect to a natural poset-based relational semantics using a coalgebraic version of completeness-via-canonicity. By formalizing the problem in the language of coalgebraic logics, we develop a modular theory which covers a wide variety of different logics under a single framework, and lends itself to further extensions. Moreover, we believe that the coalgebraic framework provides a systematic and principled way to study the relationship between *resource models* on the semantics side, and *substructural logics* on the syntactic side.

Keywords: Completeness, canonicity, coalgebraic logic, substructural logic, resource modelling, separation logic.

1. Introduction

This work lies at the intersection of resource semantics/modelling, substructural logics, and the theory of canonical extensions and canonicity. These three areas respectively correspond to the semantic, proof-theoretic, and algebraic sides of the problem we tackle: to give a systematic, modular account of the relation between resource semantics and logical structure. Our approach will mostly be semantically driven, guided by the resource models of separation logic. We will therefore not delve into the proof theory of substructural logics, but rather deal with the equivalent algebraic formulations in terms of residuated lattices ([1] and [2] give an overview of the correspondence between classes of residuated lattices and substructural logics).

Resource semantics and modelling. Resource interpretations of substructural logics — see, for example, [3, 4, 5, 6, 7] — are well-known and exemplified in the context of program verification and semantics by Ishtiaq and O’Hearn’s pointer logic [8] and Reynolds’ separation logic [9], each of which amounts to a model of a specific theory in Boolean BI. Resource semantics and

Email addresses: f.dahlqvist@ucl.ac.uk (Fredrik Dahlqvist), d.pym@ucl.ac.uk (David Pym)

modelling with resources has become an active field of investigation in itself (see, for example, [10]). Certain requirements, discussed below, seem natural (and useful in practice) in order to model naturally arising examples of resource.

1. We need to be able to compare at least some resources. Indeed, in a completely discrete model of resource (i.e., where no two resources are comparable) it is impossible to model key concepts such as ‘having enough resources’. On the other hand, there is no reason to assume that *any two* resources be comparable (e.g., heaps). This suggests at least a pre-order structure on models. In fact, we take the view that comparing two resources is fundamental and, in particular, if two resources cannot be distinguished in this way then they can be identified. We thus add antisymmetry and work with posets.
2. We need to be able to combine (some) resources to form new resources (e.g., union of heaps with disjoint domains [8]). We denote the combination operation by \otimes . An alternative, relational, point of view is that we should be able to specify how resources can be ‘split up’ into pairs of constituent resources. From this perspective, it makes sense to be able to list for a given resource r , the pairs (s_1, s_2) of resources which combine to form a resource $s_1 \otimes s_2 \leq r$.
3. All reasonable examples of resources possess ‘unit’ resources with respect to the combination operation \otimes ; that is, special resources that leave other resources unchanged under the combination operation.
4. The last requirement is crucial, but slightly less intuitive. In the most well-behaved examples of resource models (e.g., heaps or \mathbb{N}), if we are given a resource r and a ‘part’ s of r , there exists a resource s' that ‘completes’ s to make r ; that is, we can find a resource s' such that $s \otimes s' = r$. More generally, given two resources r, s , we want to be able to find the the best s' such that $s \otimes s' \leq r$. In a model of resource without this feature, it is impossible to provide an answer to legitimate questions such as ‘how much additional resource is needed to make statement ϕ hold?’. Mathematically, this requirement says that the resource composition is a residuated mapping in both its arguments.

The literature on resource modelling, and on separation logic in particular, is vast, but two publications – [11] and [12] – are strongly related to this work. Both show completeness of ‘resource logics’ by using Sahlqvist formulas, which amounts to using completeness-via-canonicity ([13, 14]).

Completeness-via-canonicity and substructural logics. The logical side of resource modelling is the world of substructural logics, such as BI, and of their algebraic formulations; that is, residuated lattices, residuated monoids, and related structures. The past decade has seen a fair amount of research into proving the completeness of relational semantics for these logics (for BI, for example, [5, 6]), using, among other approaches, techniques from the duality

theory of lattices. In [15], Dunn *et al.* prove completeness of the full Lambek calculus and several other well-known substructural logics with respect to a special type of Kripke semantics by using duality theory. This type of Kripke semantics, which is two-sorted in the non-distributive case, was studied in detail by Gehrke in [16]. The same techniques have been applied to prove Kripke completeness of fragments of linear logic in [17]. Finally, the work of Suzuki [18] explores in much detail completeness-via-canonicity for substructural logics. Our work follows in the same vein but with some important differences. Firstly, we use a dual adjunction rather than a dual equivalence to connect syntax and semantics. This is akin to working with Kripke frames rather than descriptive general frames in modal logics: the models are simpler and more intuitive, but the tightness of the fit between syntax and semantics is not as strong. Secondly, we use the topological approach to canonicity of [19, 20, 21] because we feel it is the most flexible and modular approach to building canonical (in)equations. Thirdly, we only consider distributive structures. This is to some extent a matter of taste. Our choice is driven by the desire to keep the theory relatively simple (the non-distributive case is more involved), by the fact that, from a resource-modelling perspective, the non-distributive case does not seem to occur ‘in the wild’ and, finally, because we place ourselves in the framework of coalgebraic logic, where the category of distributive lattices forms a particularly nice ‘base category’.

Completeness-via-canonicity, coalgebraically. The coalgebraic perspective brings many advantages to the study of completeness-via-canonicity. First, it greatly clarifies the connection between canonicity as an algebraic method and the existence of ‘canonical models’; that is, strong completeness. Second, it provides a generic framework in which to prove completeness-via-canonicity for a vast range of logics ([22]). Third, it is intrinsically modular; that is, it provides theorems about complicated logics by combining results for simpler ones ([23, 24]). We will return to the advantages of working coalgebraically throughout the paper.

2. A coalgebraic perspective on substructural logics

We use the ‘abstract’ version of coalgebraic logic developed in, for example, [25], [26] and [27]; that is, we require the following basic situation:

$$\begin{array}{ccc}
 L & & T^{\text{op}} \\
 \downarrow & \xrightarrow{F} & \downarrow \\
 \mathcal{C} & \perp & \mathcal{D}^{\text{op}} \\
 \uparrow & \xleftarrow{G} & \uparrow
 \end{array}
 \quad (1)$$

The left hand-side of the diagram is the syntactic side, and the right-hand side the semantic one. The category \mathcal{C} represents a choice of ‘reasoning kernel’; that is, of logical operations which we consider to be fundamental, whilst L is a syntax constructing functor which builds terms over the reasoning kernel. Objects in \mathcal{D} are the carriers of models and T specifies the coalgebras on these

carriers in which the operations defined by L are interpreted. The functors F and G relate the syntax and the semantics, and F is left adjoint to G . We will denote such an adjunction by $F \dashv G : \mathcal{C} \rightarrow \mathcal{D}$. Note, as mentioned in the introduction, that we only need a dual adjunction, not a full duality.

2.1. Syntax

Reasoning kernels. There are three choices for the category \mathcal{C} which are particularly suited to our purpose, the category **DL** of distributive lattices, the category **BDL** of bounded distributive lattices, and the category **BA** of boolean algebras. The categories **DL**, **BDL** and **BA** have a very nice technical feature from the perspective of coalgebraic logic: each category is locally finite; that is, finitely generated objects are finite. This is a very desirable technical property for the presentation of endofunctors on this category and for coalgebraic strong completeness theorems. We denote by $F \dashv U$ the usual free-forgetful adjunction between **DL** (resp. **BDL**, resp. **BA**) and **Set**.

True and false. The choice of including (or not) \top and \perp to the logic is clearly provided by the choice of reasoning kernel.

Algebras. Recall that an algebra for an endofunctor $L : \mathcal{C} \rightarrow \mathcal{C}$ is an object A of \mathcal{C} together with a morphism $\alpha : LA \rightarrow A$. We refer to endofunctors $L : \mathcal{C} \rightarrow \mathcal{C}$ as *syntax constructors*.

Resource operations. The operations on resources specified in the introduction; that is, a combination operation and its left and right residuals, are introduced via the following syntax constructor:

$$L_{\text{RL}} : \mathcal{C} \rightarrow \mathcal{C}, \begin{cases} L_{\text{RL}} A = F\{I, a \otimes b, a \multimap b, a \otimes\!-\! b \mid a, b \in \mathsf{UA}\} / \equiv \\ L_{\text{RL}} f : L_{\text{RL}} A \rightarrow L_{\text{RL}} B, [a]_{\equiv} \mapsto [f(a)]_{\equiv}, \end{cases}$$

where \equiv is the fully invariant equivalence relation in \mathcal{C} generated by the following Distribution Laws for non-empty finite subsets X of A :

$$\begin{array}{ll} \text{DL1. } \bigvee X \otimes a = \bigvee [X \otimes a] & \text{DL4. } \bigvee X \multimap a = \bigwedge [X \multimap a] \\ \text{DL2. } a \otimes \bigvee X = \bigvee [a \otimes X] & \text{DL5. } \bigwedge X \otimes\!-\! a = \bigwedge [X \otimes\!-\! a] \\ \text{DL3. } a \multimap \bigwedge X = \bigwedge [a \multimap X] & \text{DL6. } a \otimes\!-\! \bigvee X = \bigwedge [a \otimes\!-\! X]. \end{array}$$

where $\bigvee [X \otimes a] = \bigvee \{x \otimes a \mid x \in X\}$ and similarly for the other operations. For the categories **BDL** and **BA** we allow X to be the empty set and use the usual convention that $\bigvee \emptyset = \perp$ and $\bigwedge \emptyset = \top$. The language defined by L_{RL} is the free L_{RL} -algebra over FV which we shall denote by $\mathcal{L}(L_{\text{RL}}, V)$ or simply $\mathcal{L}(L_{\text{RL}})$ when a choice of propositional variables V has been established. It is not difficult to see that $\mathcal{L}(L_{\text{RL}})$ is the language of the distributive full Lambek calculus (or residuated lattices) quotiented under the axioms of \mathcal{C} and DL1-6. An L_{RL} -algebra is simply an object of \mathcal{C} endowed with a nullary operation I and binary operations \otimes , \multimap and $\otimes\!-\!$ satisfying the distribution laws above. Note that an L_{RL} -algebra is *not* a distributive residuated lattice. Only some

features of this structure have been captured by the axioms above. But several are still missing, and will be added subsequently as *canonical frame conditions*. L_{RL} -algebras are an example of *Distributive Lattice Expansions*, or DLEs; that is, distributive lattices endowed with a collection of maps of finite arities. When $\mathcal{C} = \mathbf{BA}$, L_{RL} -algebras are an example *Boolean Algebra Expansions*, or BAEs.

Modularity. The syntax developed above is completely modular in two respects. First, it is modular in the choice of ‘reasoning kernel’ since the same formal functor can be overloaded to be used on several different choices of base categories. Second, and most importantly, it allows for a very concise definition and construction of the *fusion* of logics ([24]); that is, the free combination of two logics defined on the same base categories. If $L_1, L_2 : \mathcal{C} \rightarrow \mathcal{C}$ are functors defining languages $\mathcal{L}(L_1), \mathcal{L}(L_2)$, then the fusion $\mathcal{L}(L_1) \oplus \mathcal{L}(L_2)$ of these languages is simply given by $\mathcal{L}(L_1 + L_2)$ where $+$ is the object-wise coproduct in \mathcal{C} . As an example, consider modal substructural logics; for instance, the ‘relevant modal logic’ of [18] or the modal resource logics of [10, 28, 29]. The language of positive modal logic ([30]) is given by the functor

$$L_{\text{ML}} : \mathcal{C} \rightarrow \mathcal{C}, \begin{cases} L_{\text{ML}}A = \mathbf{F}\{\Diamond a, \Box a \mid a \in \mathbf{UA}\} / \equiv \\ L_{\text{ML}}f : L_{\text{ML}}A \rightarrow L_{\text{ML}}B, [a]_{\equiv} \mapsto [f(a)]_{\equiv}, \end{cases}$$

where \equiv is the fully invariant equivalence relation in \mathcal{C} generated by the following Distribution Laws for finite subsets X of A :

$$\text{ML1. } \Diamond(\bigvee X) = \bigvee[\Diamond X] \qquad \text{ML2. } \Box(\bigwedge X) = \bigwedge[\Box X].$$

where $\bigvee[\Diamond X] = \bigvee\{\Diamond x \mid x \in X\}$ and similarly for \Box . When $\mathcal{C} = \mathbf{BA}$ one can of course use a single modality and define its dual in the usual fashion, but nothing is lost by considering the full signature, so we will consider L_{ML} to be the functor defining modal logics across all our reasoning kernels, and the *language* of modal logics is then given in our framework by the free L_{ML} -algebra over \mathbf{FV} ; that is, $\mathcal{L}(L_{\text{ML}})$. The language of the various substructural modal logics cited above, which is the fusion $\mathcal{L}(L_{\text{RL}}) \oplus \mathcal{L}(L_{\text{ML}})$, is thus simply given by $\mathcal{L}(L_{\text{RL}} + L_{\text{ML}})$. Similarly, we can consider bi-substructural languages as is done in [31]; that is, languages which allow resources to be combined in two different ways. In this case the language is simply given by $\mathcal{L}(L_{\text{RL}} + L_{\text{RL}})$.

2.2. Coalgebraic semantics

Semantic domain. As we mentioned in the introduction, it is reasonable to assume that a model of resources should be a poset, and thus taking $\mathcal{D} = \mathbf{Pos}$ is intuitively justified. This is a particularly attractive choice of ‘semantic domain’ given that the category \mathbf{Pos} is related to \mathbf{DL} by the dual adjunction $\mathbf{Pf} \dashv \mathcal{U} : \mathbf{DL} \rightarrow \mathbf{Pos}^{\text{op}}$, where \mathbf{Pf} is the functor sending a distributive lattice to its poset of prime filters, and \mathbf{DL} -morphisms to their inverse images, and \mathcal{U} is the functor sending a poset to the distributive lattice of its upsets and monotone maps to their inverse images. When a distributive lattice is a boolean

algebra, it is well-known that prime filters are maximal (i.e., ultrafilters) and the partial order on the set of ultrafilter is thus discrete; that is, ultrafilters are only related to themselves. Thus the dual adjunction $\mathbf{Pf} \dashv \mathcal{U}$ becomes the well-known adjunction $\mathbf{Uf} \dashv \mathcal{P}$ between \mathbf{BA} and \mathbf{Set}^{op} .

Coalgebras. Recall that a coalgebra for an endofunctor $T : \mathcal{D} \rightarrow \mathcal{D}$, is an object W of \mathcal{D} together with a morphism $\gamma : W \rightarrow TW$. The endofunctors that we will consider are built from products and ‘powersets’ and will be referred to as *model constructors*. Note that \mathbf{Pos} has products, which are simply the \mathbf{Set} products with the obvious partial order on pairs of elements. The ‘powerset’ functor which we will consider is the *convex powerset* functor: $P_c : \mathbf{Pos} \rightarrow \mathbf{Pos}$, sending a poset to its set of convex subsets, where a subset U of a poset (X, \leq) is convex if $x, z \in U$ and $x \leq y \leq z$ implies $y \in U$. The set $P_c X$ is given a poset structure via the *Egli-Milner* order (see [32, 33]). Note that if X is a set, it can be seen as a trivial poset where any element is only related to itself, and in this case it is not difficult to see that *any* subset $U \subseteq X$ is convex. Thus in the case of sets, the convex subset functor P_c is simply the usual covariant powerset functor. It therefore makes sense to consider P_c over all our ‘semantic domains’.

Coalgebras for the resource operations. We define the following model constructor, which is used to interpret $I, \otimes, -\otimes$ and $\otimes-$:

$$T_{\text{RL}} : \mathcal{D} \rightarrow \mathcal{D}, \begin{cases} T_{\text{RL}} W = 2 \times P_c(W \times W) \times P_c(W^{\text{op}} \times W) \times P_c(W \times W^{\text{op}}) \\ T_{\text{RL}} f : T_{\text{RL}} W \rightarrow T_{\text{RL}} W', U \mapsto (\text{Id}_2 \times (f \times f)^3)[U]. \end{cases}$$

The intuition is that the first component of the structure map of a T_{RL} -coalgebra (to the poset 2) separates states into units and non-units. The second component sends each ‘state’ $w \in W$ to the pairs of states which it ‘contains’, the next two components are used to interpret $-\otimes$ and $\otimes-$, respectively, and will turn out to be very closely related to the second component. Note that if $\mathcal{D} = \mathbf{Pos}$, the structure map of coalgebras are monotone, intuitively this means bigger resources can be split up in more ways.

The semantic transformations. In the abstract flavour of coalgebraic logic, the semantics is provided by a natural transformation $\delta : LG \rightarrow GT^{\text{op}}$ called the *semantic transformation*. We show below how this defines an interpretation map, but we first define our semantic transformation at every poset W by its action on the generators of $L_{\text{RL}}GW$:

$$\begin{aligned} \delta_W^{\text{RL}}(I) &= \{t \in T_{\text{RL}} W \mid \pi_1(t) = 0 \in 2\} \\ \delta_W^{\text{RL}}(u \otimes v) &= \{t \in T_{\text{RL}} W \mid \exists (x, y) \in \pi_2(t), x \in u, y \in v\} \\ \delta_W^{\text{RL}}(u -\otimes w) &= \{t \in T_{\text{RL}} W \mid \forall (x, y) \in \pi_3(t), x \in u \Rightarrow y \in w\} \\ \delta_W^{\text{RL}}(w \otimes - v) &= \{t \in T_{\text{RL}} W \mid \forall (x, y) \in \pi_4(t), x \in v \Rightarrow y \in w\} \end{aligned}$$

where $\pi_i, 1 \leq i \leq 4$ are the usual projections maps, and $u, v \in GW$.

Proposition 1. *The natural transformation δ^{RL} is well-defined.*

Proof. Let us first check that for any $u, v \in \mathcal{U}W$, $\delta_W^{\text{RL}}(u \otimes v)$, $\delta_W^{\text{RL}}(u \multimap v)$ and $\delta_W^{\text{RL}}(u \otimes -v)$ are upsets in $T_{\text{RL}}W$. Assume first that $t \in \delta_W^{\text{RL}}(u \otimes v)$ and that $t \leq t' \in T_{\text{RL}}W$, we want to show that $t' \in \delta_W^{\text{RL}}(u \otimes v)$ too. By definition of the partial order on $T_{\text{RL}}W$, we have that $\pi_2(t) \leq \pi_2(t')$ for the (component-wise) Egli-Milner order; that is, for each $(x, y) \in \pi_2(t)$ there exists $(x', y') \in \pi_2(t')$ such that $x \leq x'$ and $y \leq y'$. But by definition of $\delta_W^{\text{RL}}(u \otimes v)$ we know that there exists $(x, y) \in \pi_2(t)$ such that $x \in u, y \in v$, and since u, v are upsets it follows that $x' \in u, y' \in v$ and thus $t' \in \delta_W^{\text{RL}}(u \otimes v)$ as desired. Assume now that $t \in \delta_W^{\text{RL}}(u \multimap v)$ and that $t \leq t' \in T_{\text{RL}}W$, we want to show that $t' \in \delta_W^{\text{RL}}(u \multimap v)$. To see that this is the case, take any $(x', y') \in \pi_3(t')$ and assume that $x' \in u$, we need to show that $y' \in v$. By definition of the Egli-Milner order we know that there exists $(x, y) \in \pi_3(t)$ such that $x' \leq x$ and $y \leq y'$ (note the inequality reversal due to the presence of $(-)^{\text{op}}$ in the definition of T_{RL}). Since u is an upset, it follows that $x \in u$ and since $t \in \delta_W^{\text{RL}}(u \multimap v)$, it follows that $y \in v$, and thus $y' \in v$ as v is an upset. The proof is identical for $\delta_W^{\text{RL}}(u \otimes -v)$.

Let us now show that δ_W^{RL} satisfies the distributivity laws DL1-6. For any $u_1, u_2, v \in \mathcal{U}W$ we have

$$\begin{aligned} \delta_X^{\text{RL}}(u_1 \cup u_2, v) &= \{t \in T_{\text{RL}}W \mid \exists (x, y) \in \pi_2(t), x \in u_1 \cup u_2, y \in v\} \\ &= \{t \in T_{\text{RL}}W \mid \exists (x, y) \in \pi_2(t), x \in u_1, y \in v\} \cup \\ &\quad \{t \in T_{\text{RL}}W \mid \exists (x, y) \in \pi_2(t), x \in u_2, y \in v\} \\ &= \delta_X^{\text{RL}}(u_1, v) \cup \delta_X^{\text{RL}}(u_2, v), \end{aligned}$$

and the proof is clearly identical for the second argument. The meet preservation in the second argument of \multimap is easy:

$$\begin{aligned} \delta_X^{\text{RL}}(v \multimap (u_1 \cap u_2)) &= \{t \in T_{\text{RL}}W \mid \forall (x, y) \in \pi_3(t), x \in v \Rightarrow y \in (u_1 \cap u_2)\} \\ &= \{t \in T_{\text{RL}}W \mid \forall (x, y) \in \pi_3(t), x \in v \Rightarrow y \in u_1\} \cap \\ &\quad \{t \in T_{\text{RL}}W \mid \forall (x, y) \in \pi_3(t), x \in v \Rightarrow y \in u_2\} \\ &= \delta_X^{\text{RL}}(v \multimap u_1) \cap \delta_X^{\text{RL}}(v \multimap u_2). \end{aligned}$$

For the anti-preservation of joins in the first argument we have

$$\begin{aligned} \delta_X^{\text{RL}}((u_1 \cup u_2) \multimap v) &= \{t \in T_{\text{RL}}W \mid \forall (x, y) \in \pi_3(t), x \in (u_1 \cup u_2) \Rightarrow y \in v\} \\ &= \{t \in T_{\text{RL}}W \mid \forall (x, y) \in \pi_3(t), x \in u_1 \Rightarrow y \in v\} \cap \\ &\quad \{t \in T_{\text{RL}}W \mid \forall (x, y) \in \pi_3(t), x \in u_2 \Rightarrow y \in v\} \\ &= \delta_X^{\text{RL}}(u_1 \multimap v) \cap \delta_X^{\text{RL}}(u_2 \multimap v), \end{aligned}$$

where we use the classical equality $x \in (u_1 \cup u_2) \Rightarrow y \in v$ iff $x \notin (u_1 \cup u_2)$ or $y \in v$ iff $(x \notin u_1 \text{ or } x \in v)$ and $(x \notin u_2 \text{ or } x \in v)$. The proof for \otimes is identical. \square

The semantic transformations are thus well-defined. We now show how the interpretation map arises from the semantic transformation. Recall that, for a given syntax constructor $L : \mathcal{C} \rightarrow \mathcal{C}$, the language $\mathcal{L}(L)$ of L is the free L -algebra over FV . This is equivalent to saying that it is the initial $L(-) + \text{FV}$ -algebra.

We use initiality to define the interpretation map by putting an $L(-) + FV$ -algebra structure on the ‘predicates’ of a T -coalgebra $\gamma : W \rightarrow TW$; that is, on the carrier set GW . By definition of the coproduct, this means defining a morphism $LGW \rightarrow GW$ and a morphism $FV \rightarrow GW$. By adjointness it is easy to see that the latter is simply a valuation $v : V \rightarrow UGW$. For the former we simply use the semantic transformation and G applied to the coalgebra. The interpretation map $\llbracket - \rrbracket_\gamma$ is thus given by the catamorphism:

$$\begin{array}{ccc}
L\mathcal{L}(L) + FV & \xrightarrow{L\llbracket - \rrbracket_\gamma + \text{Id}_{FV}} & LGW + FV \\
\downarrow & & \downarrow \delta_W + \text{Id}_{FV} \\
& & GTW + FV \\
& & \downarrow G\gamma + v \\
\mathcal{L}(L) & \xrightarrow{\llbracket - \rrbracket_\gamma} & GW
\end{array}$$

Modularity. Following our point on the modularity of the syntax, we highlight the modularity of the coalgebraic semantics too. Modal logic will be interpreted in T_{ML} -coalgebra for the functor

$$T_{\text{ML}} : \mathcal{D} \rightarrow \mathcal{D}, \begin{cases} T_{\text{RL}}W = P_c(W) \times P_c(W) \\ T_{\text{ML}}f : T_{\text{ML}}W \rightarrow T_{\text{ML}}W', U \mapsto (f)^2[U]. \end{cases}$$

Note that we are interpreting \Diamond and \Box using different relations. Modulo Dunn’s *interaction axioms* ([30]) one can show that these two relations must be equal in the case of boolean modal logic, and that they can be assumed to be equal in the case of positive modal logic (although models where they are not equal will exist too). The semantics is given as usual by the transformation $\delta^{\text{ML}} : L_{\text{ML}}G \rightarrow GT_{\text{ML}}$ defined at every poset W by its action on the generators of $L_{\text{ML}}GW$:

$$\begin{aligned}
\delta_W^{\text{ML}}(\Diamond u) &= \{(x, y) \in T_{\text{ML}}W \mid x \cap u \neq \emptyset\} \\
\delta_W^{\text{ML}}(\Box u) &= \{(x, y) \in T_{\text{ML}}W \mid y \subseteq u\}
\end{aligned}$$

Model constructors and semantic transformations can be assembled in a way that is dual to the the syntax constructors; that is, using products rather than co-products. Formally, for languages defined by functors $L_1, L_2; \mathcal{C} \rightarrow \mathcal{C}$ interpreted in coalgebras for the functors $T_1, T_2; \mathcal{D} \rightarrow \mathcal{D}$ by semantic transformations δ^1, δ^2 respectively, the fusion $\mathcal{L}(L_1 + L_2)$ is interpreted in $T_1 \times T_2$ -coalgebra, where the product is taken object-wise in \mathcal{D} , via the semantic transformation

$$(G\pi_1 + G\pi_2) \circ \delta^1 + \delta^2 : L_1G + L_2G \rightarrow GT_1 + GT_2 \rightarrow GT_1 \times GT_2$$

In particular, the semantics of the modal substructural logics defined above is

given by the following interpretation maps:

$$\begin{array}{ccc}
& L_{\text{ML}} + L_{\text{RL}} \llbracket - \rrbracket_{(\gamma_1 \times \gamma_2)} + \text{Id}_{\text{FV}} & \\
(L_{\text{ML}} + L_{\text{RL}}) \mathcal{L}(L_{\text{ML}} + L_{\text{RL}}(-)) + \text{FV} & \dashrightarrow & L_{\text{ML}} G W + L_{\text{RL}} G W + \text{FV} \\
\downarrow & & \downarrow \delta_W^{\text{ML}} + \delta_W^{\text{RL}} + \text{Id}_{\text{FV}} \\
& & G T_{\text{ML}} W + G T_{\text{RL}} W + \text{FV} \\
& & \downarrow G(\gamma_1 \times \gamma_2) \circ (G\pi_1 + G\pi_2) + v \\
\mathcal{L}(L_{\text{ML}} + L_{\text{RL}}) & \dashrightarrow & \llbracket - \rrbracket_{(\gamma_1 \times \gamma_2)} \dashrightarrow G W
\end{array}$$

2.3. Advantages of the coalgebraic approach.

Before we move on to the technical part of this paper, we return to the advantages of our set up. From the perspective of studying the relation between resource semantics and logics, the fundamental situation described by Diagram 1 is particularly promising. Going from substructural logics to resource models, the coalgebraic approach allows us to ‘guess’ and generate appropriate resource models. Indeed, starting from a ‘reasoning kernel’ \mathcal{C} , the existence of a dual adjunction $F \dashv G$ with a category \mathcal{D} restricts the kind of model carriers we should consider. Moreover, as we will see later, objects of the type GFA for $A \in \mathcal{C}$ will play a crucial role and should be *canonical extensions*. This extra requirement determines to a great extent the useful structure(s) one ought to consider for the carriers of resource models. For example when $\mathcal{C} = \mathbf{DL}$, we cannot take $\mathcal{D} = \mathbf{Set}$, because GFA is then given by $\mathcal{PPf}A$ which is not the canonical extension of A . It is therefore the framework itself which suggests that non-boolean substructural logics should have posets of resources as their models. Similarly, as we have shown above, the choice of T , that is to say of relational structure on the carrier, can be guessed from that of L in a systematic fashion – at least for the languages we consider here.

Conversely, if we start from requirements on resource models, such as the natural conditions listed in the introduction, we can work from resource model to logic via the existence of a dual adjunction $F \dashv G$ and the constraint that GFA should be the canonical model of A . In this way, the ‘natural’ logics to reason about partially ordered models of resources are positive; that is, based on \mathbf{DL} . Moreover, the relational structure suggested in the introduction suggests adding binary modalities, in other words functors $L : \mathbf{DL} \rightarrow \mathbf{DL}$ building binary ‘modal’ formulas over \mathbf{DL} , as was done in this section. Thus we see that in either direction the categorical clarity of coalgebraic logics provides us with a natural and principled methodology for building resource models from substructural logics and vice versa.

Finally, we note that recent work on *positive* coalgebraic logics ([32, 33]) suggests that what is known of boolean modal logics with relational semantics can be adapted in a systematic and principled way to the positive modal logics that we are considering here. Indeed, our choice of semantics in terms

of convex powerset coalgebras is dictated by the fact that this functor is a universal extension to **Pos** of the usual powerset functor on **Set**. In this sense, the coalgebraic perspective also suggests what a ‘correct’ relational model on a poset of resources should be.

3. Jónsson-Tarski extensions

The languages $\mathcal{L}(L_{\text{RL}})$ and $\mathcal{L}(L_{\text{ML}})$ which we have introduced earlier are part a class of logics with a very strong property: they are strongly complete with respect to their semantics. This is what we will now establish, and it is the first step is showing strong completeness of more complex logics based on the languages $\mathcal{L}(L_{\text{RL}})$ and $\mathcal{L}(L_{\text{ML}})$. The proof is an application of the coalgebraic Jónsson-Tarski theorem.

Theorem 2 (Coalgebraic Jónsson-Tarski theorem, [26]). *Assuming the basic situation of Diagram (1) and a semantic transformation $\delta : LG \rightarrow GT$, if its adjoint transpose $\hat{\delta} : TF \rightarrow FL$ has a right-inverse $\zeta : FL \rightarrow TF$, then for every L -algebra $\alpha : LA \rightarrow A$, the embedding $\eta_A : A \rightarrow GFA$ of A into its canonical extension can be lifted to the following L -algebra embedding:*

$$\begin{array}{ccccc} LA & \xrightarrow{\alpha} & & & A \\ \downarrow L\eta_A & & & & \downarrow \eta_A \\ LGFA & \xrightarrow{\delta_{FA}} & GTFA & \xrightarrow{G\zeta_A} & GFLA & \xrightarrow{GF\alpha} & GFA \end{array} \quad (2)$$

We call the coalgebra $\zeta \circ F\alpha : FA \rightarrow TFA$ a *canonical model* of (the L -algebra) A . If A is the free L -algebra over FV we recover the usual notion of canonical model. The ‘truth lemma’ follows from the definition of η . We will call the L -algebra $LGFA \rightarrow GFA$ defined by Diagram (2) a *Jónsson-Tarski extension* of the $\alpha : LA \rightarrow A$.

We now prove the existence of canonical models for the logics defined by L_{ML} and L_{RL} . The result generalizes Lemma 5.1 of [30], which builds canonical models for countable DLs with a unary operator, and Lemma 4.26 of [13], which builds canonical models for countable BAs with n -ary operators. We essentially show how to build canonical models for arbitrary DLs with n -ary expansions all of whose arguments either (1) preserve joins or anti-preserve meets, or (2) preserve meets or anti-preserve joins. The proof is rather involved and is detailed in the appendix.

Theorem 3. *The adjoint transpose of the transformation $\delta^{\text{RL}} : L_{\text{RL}}G \rightarrow GT_{\text{RL}}$ (resp. $\delta^{\text{ML}} : L_{\text{ML}}G \rightarrow GT_{\text{ML}}$) has right inverses at every distributive lattice.*

Strong completeness: Let us now define what we exactly mean by strong completeness. Let \mathcal{C} be **DL**, **BDL** or **BA**, $L : \mathcal{C} \rightarrow \mathcal{C}$, V be a set of propositional variables, $q : \mathcal{L}(L) \rightarrow \mathcal{Q}$ be a regular epi, and let $\Phi, \Psi \subseteq \mathcal{Q}$ be two families of ‘formulas’ such that $\Phi \not\vdash \Psi$; that is, such that no finite set Φ_0 of elements of Φ

and no finite set Ψ_0 of elements of Ψ can be found such that $\bigwedge \Phi_0 \leq \bigvee \Psi_0$. The statement that \mathcal{Q} is strongly complete w.r.t. to a class \mathcal{T} of T -coalgebras means that for any such choice of Φ, Ψ there exists a T -coalgebra $\gamma : X \rightarrow TX$ in \mathcal{T} , a valuation $v : FV \rightarrow GX$, and a point $x \in X$ such that $x \in \llbracket a \rrbracket_{(\gamma, v)}$ for all $a \in \Phi$ and $x \notin \llbracket b \rrbracket_{(\gamma, v)}$ for all $b \in \Psi$.

Theorem 4. *The logic defined by L_{RL} (resp. L_{ML}) is sound and strongly complete with respect to the class of all T_{RL} - (resp. T_{ML} -) coalgebras.*

Proof. We show the case of $\mathcal{L}(L_{\text{RL}})$. Let $\Phi, \Psi \subseteq \mathcal{L}(L_{\text{RL}})$ and $\Phi \not\subseteq \Psi$. Then the filter $\langle \Phi \rangle^\uparrow$ generated by Φ and the ideal $\langle \Psi \rangle^\downarrow$ generated by Ψ obey $\langle \Phi \rangle^\uparrow \cap \langle \Psi \rangle^\downarrow = \emptyset$. By the PIT there exists a prime filter w_Φ extending $\langle \Phi \rangle^\uparrow$ such that $w_\Phi \cap \langle \Psi \rangle^\downarrow = \emptyset$. By Theorems 2 and 3, the L -algebra $\mathcal{L}(L_{\text{RL}})$ has a Jónsson-Tarski extension which provides an interpretation in the T -coalgebra $\text{Pf}\mathcal{L}(L_{\text{RL}}) \rightarrow \text{Pf}L_{\text{RL}}\mathcal{L}(L_{\text{RL}}) \xrightarrow{\zeta_{\mathcal{L}(L_{\text{RL}})}} \text{TPf}\mathcal{L}(L_{\text{RL}})$ coinciding with $\eta_{\mathcal{L}(L_{\text{RL}})}$. In this interpretation $w_\Phi \in \llbracket a \rrbracket$ for all $a \in \Phi$ and $w_\Phi \notin \llbracket b \rrbracket$ for all $b \in \Psi$. \square

4. Canonical extensions and canonical equations

In the previous section we have shown how to embed an L -algebra with carrier A into an L -algebra with carrier GFA . When $\mathcal{C} = \mathbf{DL}, \mathbf{BDL}$ or \mathbf{BA} carriers of this shape are known as *canonical extensions* (and denoted A^σ) and a great deal is known about them. The theory of canonical extensions in \mathbf{DL} has been extended to boolean algebras with operators (BAOs) ([34]) and to distributive lattice expansions (DLEs) ([35, 20]) and forms the basis of the theory of canonicity which consists in determining when the validity of an equation in a DLEs transfer to its canonical extension; that is, when $A \models s = t$ implies $A^\sigma \models s = t$. Note that the canonical and Jónsson-Tarski extensions are in general not equal. This section deals only with canonical extension, but we will see in the next section how these results can be combined with the Jónsson-Tarski construction of Theorem 2.

4.1. Canonical extension of distributive lattices

We now briefly present the salient facts about canonical extensions for distributive lattices. For any A in \mathbf{DL} , $\mathcal{UPf}A$ is known as the *canonical extension* of A and denoted A^σ . It can be characterised uniquely up to isomorphism through purely algebraic properties, namely that A is *dense* and *compact* in A^σ . For our purpose however, *defining* the canonical extension of A as $\mathcal{UPf}A$ will be sufficient. The canonical extension A^σ of a distributive lattice A is always *completely distributive* (see [20]). The following terminology will be important: A^σ is a completion of A and all joins of elements of A therefore exist in A^σ , such elements are called *open* and their set is denoted by $O(A)$. Dually, meets in A^σ of elements of A will be called *closed* and their set denoted $K(A)$. Elements of $A = K(A) \cap O(A)$ are therefore called *clopens*.

4.2. Canonical extension of distributive lattice expansions

We now sketch the theory of canonical extensions for Distributive Lattice Expansions (DLE) — for the details, see [35, 20]. Each map $f : \mathsf{UA}^n \rightarrow \mathsf{UA}$ can be extended to a map $(\mathsf{UA}^\sigma)^n \rightarrow \mathsf{UA}^\sigma$ in two canonical ways:

$$\begin{aligned} f^\sigma(x) &= \bigvee \{ \bigwedge f[d, u] \mid K^n \ni d \leq x \leq u \in O^n \} \\ f^\pi(x) &= \bigwedge \{ \bigvee f[d, u] \mid K^n \ni d \leq x \leq u \in O^n \}, \end{aligned}$$

where $f[d, u] = \{f(a) \mid a \in A^n, d \leq a \leq u\}$. Note that since A is compact in A^σ the intervals $[d, u]$ are never empty, which justifies these definitions. For a signature Σ , the *canonical extension* of a Σ -DLE $(A, (f_s : \mathsf{UA}^{\text{ar}(n)} \rightarrow \mathsf{UA})_{s \in \Sigma})$ is defined to be the Σ -DLE $(A^\sigma, (f_s^\sigma : \mathsf{U}(A^\sigma)^{\text{ar}(n)} \rightarrow \mathsf{UA}^\sigma)_{s \in \Sigma})$, and similarly for BAEs. We summarize some important facts about canonical extensions of maps in the following proposition, proofs can be found in, for example, [19, 20, 21]:

Proposition 5. *Let A be a distributive lattice, and $f : \mathsf{UA}^n \rightarrow \mathsf{UA}$.*

1. $f^\sigma \upharpoonright A^n = f^\pi \upharpoonright A^n = f$.
2. $f^\sigma \leq f^\pi$ under pointwise ordering.
3. If f is monotone in each argument, then $f^\sigma \upharpoonright (K \cup O)^n = f^\pi \upharpoonright (K \cup O)^n$.

We call a monotone map $f : \mathsf{UA}^n \rightarrow \mathsf{UA}$ *smooth in its i^{th} argument* ($1 \leq i \leq n$) if, for every $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n \in K \cup O$,

$$f^\sigma(x_1, \dots, x_{i-1}, x_i, x_{i+1}, \dots, x_n) = f^\pi(x_1, \dots, x_{i-1}, x_i, x_{i+1}, \dots, x_n),$$

for every $x_i \in A^\sigma$. A map $f : \mathsf{UA}^n \rightarrow \mathsf{UA}$ is called *smooth* if it is smooth in each of its arguments.

In order to study effectively the canonical extension of maps, we need to define six topologies on A^σ . First, we define $\sigma^\uparrow, \sigma^\downarrow$ and σ as the topologies generated by the bases $\{\uparrow p \mid p \in K\}$, $\{\downarrow u \mid u \in O\}$ and $\{\uparrow p \cap \downarrow u \mid K \ni p \leq u \in O\}$. The next set of topologies is well-known to domain theorists: a *Scott open* in A^σ is a subset $U \subseteq A^\sigma$ such that (1) U is an upset and (2) for any up-directed set D such that $\bigvee D \in U$, $D \cap U \neq \emptyset$. The collection of Scott opens forms a topology called the *Scott topology*, which we denote γ^\uparrow . The dual topology will be denoted by γ^\downarrow , and their join by γ . It is not too hard to show (see [19, 21]) that $\gamma^\uparrow \subseteq \sigma^\uparrow$, $\gamma^\downarrow \subseteq \sigma^\downarrow$, and $\gamma \subseteq \sigma$. We denote the product of topologies by \times , and the n -fold product of a topology τ by τ^n . The following result shows why these topologies are important: they essentially characterize the canonical extensions of maps:

Proposition 6 ([19]). *For any DL A and any map $f : \mathsf{UA}^n \rightarrow \mathsf{UA}$,*

1. f^σ is the largest $(\sigma^n, \gamma^\uparrow)$ -continuous extension of f ,
2. f^π is the smallest $(\sigma^n, \gamma^\downarrow)$ -continuous extension of f

3. f is smooth iff it has a unique (σ^n, γ) -continuous extension.

From this important result, it is not hard to get the following key theorem, sometimes known as *Principle of Matching Topologies*, which underlies the basic ‘algorithm’ for canonicity:

Theorem 7 (Principle of Matching Topologies, [19, 21]). *Let A be a distributive lattice, and $f : \mathbb{U}A^n \rightarrow \mathbb{U}A$ and $g_i : \mathbb{U}A^{m_i} \rightarrow \mathbb{U}A$, $1 \leq i \leq n$ be arbitrary maps. Assume that there exist topologies τ_i on A , $1 \leq i \leq n$ such that each g_i^σ is (σ^{m_i}, τ_i) -continuous, then*

1. if f^σ is $(\tau_1 \times \dots \times \tau_n, \gamma^\uparrow)$ -continuous, then $f^\sigma(g_1^\sigma, \dots, g_n^\sigma) \leq (f(g_1, \dots, g_n))^\sigma$
2. if f^σ is $(\tau_1 \times \dots \times \tau_n, \gamma^\downarrow)$ -continuous, then $f^\sigma(g_1^\sigma, \dots, g_n^\sigma) \geq (f(g_1, \dots, g_n))^\sigma$
3. if f^σ is $(\tau_1 \times \dots \times \tau_n, \gamma)$ -continuous, then $f^\sigma(g_1^\sigma, \dots, g_n^\sigma) = (f(g_1, \dots, g_n))^\sigma$.

The last piece of information we need to effectively use the Principle of Matching Topologies is to determine when maps are continuous for a certain topology, based on the distributivity laws they satisfy. For our purpose the following results will be sufficient:

Proposition 8 ([35, 19, 20, 21]). *Let A be a distributive lattice, and let $f : \mathbb{U}A^n \rightarrow \mathbb{U}A$ be a map. For every $(n-1)$ -tuple $(a_i)_{1 \leq i \leq n-1}$, we denote by $f_a^k : A \rightarrow A$ the map defined by $x \mapsto f(a_1, \dots, a_{k-1}, x, a_k, \dots, a_{n-1})$.*

1. If f_a^k preserves binary joins, then $(f^\sigma)_a^k$ preserve all non-empty joins and is $(\sigma^\downarrow, \sigma^\downarrow)$ -continuous.
2. If f_a^k preserves binary meets, then $(f^\sigma)_a^k$ preserve all non-empty meets and is $(\sigma^\uparrow, \sigma^\uparrow)$ -continuous.
3. If f_a^k anti-preserved binary joins (i.e., turns them into meets), then $(f^\sigma)_a^k$ anti-preserve all non-empty joins and is $(\sigma^\downarrow, \sigma^\uparrow)$ -continuous.
4. If f_a^k anti-preserved binary meets (i.e., turns them into joins), then $(f^\sigma)_a^k$ anti-preserve all non-empty meets and is $(\sigma^\uparrow, \sigma^\downarrow)$ -continuous.
5. In each case f is smooth in its k^{th} argument.

4.3. Canonical (in)equations

To say anything about the canonicity of equations, we need to compare interpretations in A with interpretations in A^σ . It is natural to try to use the extension $(\cdot)^\sigma$ to mediate between these interpretations, but $(\cdot)^\sigma$ is defined on maps, not on terms. Moreover, not every valuation on A^σ originates from valuation on A . We would therefore like to recast the problem in such a way that (1) terms are viewed as maps, and (2) we do not need to worry about valuations.

Term functions. The solution is to adopt the language of *term functions* (as first suggested in [14]). Given a signature Σ , let $\mathsf{T}(V)$ denote the language of Σ -DLEs (or Σ -BAEs) over a set V of propositional variables. We view each term $t \in \mathsf{T}(V)$ as defining, for each Σ -DLE A , a map $t^A : A^n \rightarrow A$. This allows us to consider its canonical extension $(t^A)^\sigma$, and also allows us to reason without having to worry about specifying valuations. Formally, given a signature Σ and a set V a propositional variables, we inductively define the term function associated with an element t built from variables $x_1, \dots, x_n \in V$ as follows:

- $x_i^A = \pi_i^n : A^n \rightarrow A, 1 \leq i \leq n$;
- $(f(t_1, \dots, t_m))^A = f^A \circ \langle t_1^A, \dots, t_m^A \rangle$.

where π_i is the usual projection on the i^{th} component, f^A is the interpretation of the symbol f in A and $\langle t_1^A, \dots, t_m^A \rangle$ is usual the product of m maps. Note that in this definition we work in **Set**, and the building blocks of term functions are thus the variables in V (interpreted as projections) and all operation symbols, including \vee, \wedge and possibly \neg .

Proposition 9. *Let s, t be terms in the language defined by a signature Σ and A be a Σ -DLE,*

$$A \models s = t \text{ iff } s^A = t^A.$$

Canonical (in)equations. An equation $s = t$ where $s, t \in \mathsf{T}(V)$ is called *canonical* if $A \models s = t$ implies $A^\sigma \models s = t$, and similarly for inequations. Following [14], we say that $t \in \mathsf{T}(V)$ is *stable* if $(t^A)^\sigma = t^{A^\sigma}$, that t is *expanding* if $(t^A)^\sigma \leq t^{A^\sigma}$, and that t is *contracting* if $(t^A)^\sigma \geq t^{A^\sigma}$, for any A . The inequality between maps is taken pointwise. The following proposition illustrates the usefulness of these notions:

Proposition 10 ([14]). *If $s, t \in \mathsf{T}(V)$ are stable then the equation $s = t$ is canonical. Similarly, let $s, t \in \mathsf{T}(V)$ such that s is contracting and t is expanding, then the inequality $s \leq t$ is canonical.*

Proof. Let A be an arbitrary Σ -DLE. If $A \models s = t$, then $s^A = t^A$ by Proposition 9. Therefore $(s^A)^\sigma = (t^A)^\sigma$ and thus $s^{A^\sigma} = t^{A^\sigma}$, by stability, and it follows that $A^\sigma \models s = t$ by Proposition 9.

Similarly, if $A \models s \leq t$ then $s^A \leq t^A$ by Proposition 9 and thus $(s^A)^\sigma \leq (t^A)^\sigma$. By the assumptions on s and t , this means that we also have $s^{A^\sigma} \leq t^{A^\sigma}$, and thus $A^\sigma \models s \leq t$ by Proposition 9. \square

5. Coalgebraic Completeness via-canonicity

In this section we will combine the results of Sections 2 and 3. We will first exhibit a set of canonical axioms which complete the definition of L_{RL} and completely axiomatize the distributive full Lambek calculus. This will prove that the variety defined by these axioms is canonical; that is, closed under canonical extension. We will then show that the canonical and Jónsson-Tarski extensions defined by Theorems 3 and 2 coincide. This will allow us to conclude strong completeness of the distributive full Lambek calculus.

5.1. Axiomatizing distributive residuated lattices

So far we have only captured part of the structure of distributive residuated lattices, namely we have enforced the distribution properties of \rightarrow , \otimes , \multimap and \boxtimes by our definition of the syntax constructor L_{RL} . In order to capture the rest of the structures we now add axioms which, when added to DL1-6 fully axiomatize distributive residuated lattices. Due to the constraints that these axioms must be canonical, we choose the following Frame Conditions:

$$\begin{array}{ll}
\text{FC1. } a \otimes I = a, I \otimes a = a & \text{FC4. } (c \boxtimes b) \otimes a \leq c \boxtimes (a \otimes b) \\
\text{FC2. } I \leq a \multimap a, I \leq a \boxtimes a & \text{FC5. } (a \boxtimes b) \otimes b \leq a \\
\text{FC3. } a \otimes (b \multimap c) \leq (a \otimes b) \multimap c & \text{FC6. } b \otimes (b \multimap a) \leq a.
\end{array}$$

Proposition 11. *The axioms DL1-6 and FC1-6 axiomatize distributive residuated lattices.*

Proof. It is straightforward to check that axioms DL1-6 and FC1-6 hold in any residuated lattice. Conversely, we show that if FC1-6 hold in an L_{RL} -algebra, then this L_{RL} -algebra is a residuated lattice. It is clear from FC1 that \otimes defines a monoid on the carrier set. It remains to show that the residuation conditions are satisfied. Assume that $a \otimes b \leq c$. We have

$$\begin{array}{ll}
I \leq b \multimap b & \text{FC2} \\
a \leq a \otimes (b \multimap b) & \text{FC1 and monotonicity of } \otimes \text{ from } L_{\text{RL}} \\
\leq (a \otimes b) \multimap b & \text{FC3} \\
\leq c \multimap b & \text{Monotonicity of } \multimap \text{ from } L_{\text{RL}}
\end{array}$$

Now assume that $a \leq b \multimap c$. Then we have

$$\begin{array}{ll}
a \otimes b \leq (b \multimap c) \otimes b & \text{Monotonicity of } \otimes \text{ from } L_{\text{RL}} \\
\leq c & \text{FC5.}
\end{array}$$

The proof for the left residual \boxtimes is identical. Note that the monotonicity of the operators are consequences of DL1-6. \square

We now show one of the crucial steps.

Proposition 12. *The axioms FC1-6 are canonical.*

Proof. The proof is an application of Theorem 7 and Proposition 10.

FC1: Since \otimes preserves binary joins in each argument, it is smooth by Prop. 8, and it follows that it is (σ^2, γ) -continuous. Since π_1^σ and I^σ are trivially (σ, σ) -continuous, it follows from Theorem 7 that $(\otimes \circ \langle \pi_1, I \rangle)^\sigma = \otimes^\sigma \circ \langle \pi_1, 1 \rangle^\sigma$. Each side of the equation is thus stable and the result follows from Prop. 10.

FC2: I is stable and thus contracting, and $(\multimap \circ \langle \pi_1, \pi_1 \rangle)^\sigma = \multimap^\sigma \circ \langle \pi_1, \pi_1 \rangle^\sigma$, since π_1^σ is (σ, σ) -continuous and \multimap^σ is smooth. The RHS of the inequality is thus stable, and a fortiori expanding, and the inequality is thus canonical.

FC3-4: Since \otimes^σ preserve joins in each argument, it preserves up-directed ones, and is thus $((\gamma^\uparrow)^2, \gamma^\uparrow)$ -continuous. Since $\neg\otimes^\sigma$ is smooth it is in particular $(\sigma^2, \gamma^\uparrow)$ -continuous. Since π_1^σ is $(\sigma, \gamma^\uparrow)$ -continuous, we get that $\otimes^\sigma \circ \langle \pi_1^\sigma, \neg\otimes^\sigma \circ \langle \pi_2^\sigma, \pi_3^\sigma \rangle \rangle$ is $(\sigma^3, \gamma^\uparrow)$ -continuous and thus contracting. For the RHS, note that since $\neg\otimes^\sigma$ preserves meets in its first argument, it must in particular preserve down-directed ones, thus $\neg\otimes^\sigma$ is $(\gamma^\downarrow, \gamma^\downarrow)$ -continuous in its first argument. Similarly, since $\neg\otimes^\sigma$ anti-preserve joins in its second argument, it must in particular anti-preserve up-directed ones, and is thus $(\gamma^\uparrow, \gamma^\downarrow)$ -continuous in its second argument. This means that $\neg\otimes^\sigma$ is $(\gamma^2, \gamma^\downarrow)$ -continuous. We thus have that the full term is $(\sigma^3, \gamma^\downarrow)$ continuous, and thus expanding. The inequation is therefore canonical.

FC5-6: The LHS is contracting by the same reasoning as above, and the RHS is stable and thus expanding. \square

5.2. Jónsson-Tarski vs canonical extensions

We have just shown that the variety of L_{RL} -algebras defined by the equations FC1-6 is canonical; that is, closed under canonical extension. However, since we want to exhibit models, what we really need to show is that the variety defined by FC1-6 is closed under *Jónsson-Tarski extensions*. Fortunately, for the logics of interest to us here the two extensions in fact coincide. This is what we will now show. The proof is not difficult but rather long, and can be found in the appendix.

Proposition 13. *The structure map of the Jónsson-Tarski extension of an L_{RL} -algebra is equal to the canonical extension of its structure map (in the sense of Section 4.2).*

5.3. Strong completeness

We are now ready to combine all our results and to state and prove our main completeness theorem.

Theorem 14 (Strong completeness theorem). *The Distributive Full Lambek Calculus is strongly complete with respect to the class of T_{RL} -coalgebras validating FC1-6.*

Theorem 14. Let Φ, Ψ be (not necessarily finite) subsets of $\mathcal{L}(L_{\text{RL}})$; that is, elements of the free L_{RL} -algebra over FV , such that

$$\text{FC1} - 6 + \Phi \not\models \Psi$$

We need to find a T_{RL} -model validating the axioms FC1-6 such that each $a \in \Phi$ and no $b \in \Psi$ is satisfied in this model. Now consider the Lindenbaum-Tarski L_{RL} -algebra $\alpha : L_{\text{RL}}\mathcal{L} \rightarrow \mathcal{L}$ defined by axioms FC1-6; that is,

$$\mathcal{L} = \mathcal{L}(L_{\text{RL}})/(\text{FC1} - 6),$$

where the quotient is under the fully invariant equivalence relation in \mathcal{C} generated by the frame conditions FC1-6. Note that this algebra comes equipped with a canonical valuation $v : FV \rightarrow \mathcal{L}$. By construction, \mathcal{L} validates FC1-6, and since we've established, in Proposition 12, that they are canonical, the L_{RL} -algebra

$$L_{RL}\mathcal{UPf}\mathcal{L} \xrightarrow{\alpha^\sigma} \mathcal{UPf}\mathcal{L}$$

also validates these axioms. By Proposition 13 we know that this L_{RL} -algebra is the Jónsson-Tarski extension of \mathcal{L} , and as a consequence

$$L_{RL}\mathcal{UPf}\mathcal{L} \xrightarrow{\delta_{Pf\mathcal{L}}^{RL}} \mathcal{UT}_{RL}Pf\mathcal{L} \xrightarrow{\mathcal{U}\zeta_{\mathcal{L}}} \mathcal{UPf}L_{RL}\mathcal{L} \xrightarrow{\mathcal{UPf}\alpha} \mathcal{UPf}\mathcal{L}$$

validates FC1-6. As a direct consequence of the definition of coalgebraic semantics, we have the following commutative diagram:

$$\begin{array}{ccccc}
& & L_{RL}[-]_{\mathcal{UPf}\mathcal{L}} + \text{Id}_{FV} & & \\
& \nearrow & & \searrow & \\
L_{RL}\mathcal{L}(L_{RL}) + FV & \xrightarrow{L_{RL}[-]_{\mathcal{L}} + \text{Id}_{FV}} & L_{RL}\mathcal{L} + FV & \xrightarrow{L_{RL}\eta_{\mathcal{L}} + \text{Id}_{FV}} & L_{RL}\mathcal{UPf}\mathcal{L} + FV \\
\downarrow & & \downarrow \alpha & & \downarrow \delta_{Pf\mathcal{L}}^{RL} + \text{Id}_{FV} \\
& & & & \mathcal{UT}_{RL}Pf\mathcal{L} + FV \\
& & & & \downarrow \mathcal{U}\zeta_{Pf\mathcal{L}} + \text{Id}_{FV} \\
& & & & \mathcal{UPf}\mathcal{L} + FV \\
& & & & \downarrow \mathcal{UPf}\alpha + v \\
\mathcal{L}(L_{RL}) & \xrightarrow{[-]_{\mathcal{L}}} & \mathcal{L} & \xrightarrow{\eta_{\mathcal{L}}} & \mathcal{UPf}\mathcal{L} \\
& \searrow & & \nearrow & \\
& & [-]_{\mathcal{UPf}\mathcal{L}} & &
\end{array}$$

It follows easily that at every prime filter $w \in Pf\mathcal{L}$, $w \models \text{FC1-6}$, since $[-]_{\mathcal{UPf}\mathcal{L}}$ must factor through $[-]_{\mathcal{L}}$ which ensures precisely that FC1-6 are valid. Thus $Pf\mathcal{L}$ is a model validating the axioms. We now need to find a point in $w_\Phi \in Pf\mathcal{L}$ such that $w \models \Phi$ but $w \not\models \Psi$. For this we start by considering the filter-ideal pair

$$(\langle \Phi \rangle^\uparrow, \langle \Psi \rangle^\downarrow)$$

where $\langle \Phi \rangle^\uparrow$ is the filter generated by the equivalence classes in the Lindenbaum-Tarski algebra \mathcal{L} of formulas in Φ , and similarly for the ideal generated by Ψ . It is clear that $\langle \Phi \rangle^\uparrow$ is proper, or else we would necessarily have $\text{FC1-6} + \Phi \vdash \Psi$, a contradiction. For the same reason it is clear that $\langle \Phi \rangle^\uparrow \cap \langle \Psi \rangle^\downarrow = \emptyset$. By the PIT, we can find a prime filter $w_\Phi \supseteq \langle \Phi \rangle^\uparrow$ in $Pf\mathcal{L}$ such that $w_\Phi \cap \langle \Psi \rangle^\downarrow = \emptyset$. It follows immediately that

$$w_\Phi \models \Phi \text{ and } w_\Phi \not\models \Psi$$

which is what we wanted to show. \square

5.4. Modularity.

The coalgebraic setting allows us to combine completeness-via-canoncity results from simple logics to get results for more complicated logics. It can be shown that the coalgebraic Jónsson-Tarski theorem is modular in the following sense.

Theorem 15 (Strong completeness transfers under fusion). *Let $L_i : \mathcal{C} \rightarrow \mathcal{C}, T_i : \mathcal{D} \rightarrow \mathcal{D}, \delta^i : L_i G \rightarrow GT_i, i = 1, 2$. For any $(L_1 + L_2)$ -algebra (A, α) , if $\hat{\delta}_A^i$ has a right inverse $\zeta_A^i, i = 1, 2$, then $\eta_A : A \rightarrow GFA$ lifts to an $L_1 + L_2$ -algebra morphism.*

Proof. We show that the following diagram commutes:

$$\begin{array}{ccc}
 L_1 A + L_2 A & \xrightarrow{L_1 \eta_A + L_2 \eta_A} & L_1 GFA + L_2 GFA \\
 \downarrow \alpha & \searrow \eta_{L_1 A + L_2 A} & \downarrow \delta_A^1 + \delta_A^2 \\
 & & GT_1 FA + GT_2 FA \\
 & & \downarrow G\pi_1 + G\pi_2 \\
 & & G(T_1 FA \times T_2 FA) \\
 & & \downarrow G(\zeta_A^1 \times \zeta_A^2) \\
 & & G(FL_1 A \times FL_2 A) \simeq GF(L_1 A + L_2 A) \\
 & & \downarrow GF\alpha \\
 A & \xrightarrow{\eta_A} & GFA
 \end{array}$$

F being left adjoint preserves colimits, and thus turns coproduct in \mathcal{C} into products in \mathcal{D} . The bottom left-hand corner trapezium thus commutes by naturality of η . So we must show the commutativity of top-right-hand corner triangle. For this we first show that

$$(G\pi_1 + G\pi_2) \circ (\delta^1 + \delta^2) \circ (L_1 \eta_A + L_2 \eta_A) = G(\hat{\delta}_A^1 \times \hat{\delta}_A^2) \circ \eta_{L_1 A + L_2 A}$$

This is easily seen from the following diagram, which unravels the definition of adjoint transposes and uses the fact that F preserves colimits:

$$\begin{array}{ccc}
 L_1 A + L_2 A & \xrightarrow{\quad} & GF(L_1 A + L_2 A) \simeq G(FL_1 A \times FL_2 A) \\
 \downarrow L_1 \eta_A + L_2 \eta_A & & \downarrow GF(L_1 \eta_A + L_2 \eta_A) \simeq G(FL_1 \eta_A \times FL_2 \eta_A) \\
 L_1 GFA + L_2 GFA & \xrightarrow{\quad} & GF(L_1 GFA + L_2 GFA) \simeq G(FL_1 GFA \times FL_2 GFA) \\
 \downarrow (\delta^1)_{FA} + (\delta^2)_{FA} & & \downarrow GF((\delta^1)_{FA} + (\delta^2)_{FA}) \simeq G(F(\delta^1)_{FA} \times G(F(\delta^2)_{FA})) \\
 GT_1 FA + GT_2 FA & \xrightarrow{\quad} & GF(GT_1 FA + GT_2 FA) \simeq G(FGT_1 FA \times FGT_2 FA) \\
 \downarrow G\pi_1 + G\pi_2 & \swarrow G(\epsilon_{T_1 FA} \circ \pi_1 \times \epsilon_{T_2 FA} \circ \pi_2) & \\
 G(T_1 FA \times T_2 FA) & &
 \end{array}$$

All the horizontal arrows are simply given by the unit $\eta : \mathbf{Id} \rightarrow GF$ (we have omitted the labels to keep the diagram readable), and thus the two top rectangles commute by naturality. Finally, we are left to deal with the bottom triangle which can be seen to commute from the following commutative diagram:

$$\begin{array}{ccccc}
GT_1FA & \xrightarrow{i_1} & G(T_1FA \times T_2FA) \simeq GT_1FA + GT_2FA & \xleftarrow{i_2} & GT_2FA \\
\downarrow \eta_{GT_1FA} & & \downarrow \eta_{GT_1FA + GT_2FA} & & \downarrow \eta_{GT_2FA} \\
GFGT_1FA & \xrightarrow[\simeq G\pi_1]{GF i_1} & G(FGT_1FA \times FGT_2FA) \simeq GF(GT_1FA + GT_2FA) & \xleftarrow[\simeq G\pi_2]{GF i_2} & GFGT_1FA \\
\downarrow G\epsilon_{T_1FA} & & \downarrow G(\epsilon_{T_1FA} \circ \pi_1 \times \epsilon_{T_2FA} \circ \pi_2) & & \downarrow G\epsilon_{T_2FA} \\
GT_1FA & \xrightarrow{G\pi_1} & G(T_1FA \times T_2FA) & \xleftarrow{G\pi_2} & GT_2FA
\end{array}$$

The top squares commute by naturality of η , the bottom squares commute by naturality of ϵ and the two squares can be joined by the fact that F turns coproducts into products. Note also that $G\epsilon_{T_1FA} \circ \eta_{GT_1FA} = \mathbf{Id}_{GT_1FA}$ by the fact that $F \dashv G$, and the desired result follows from the unicity of the coproduct map $G\pi_1 + G\pi_2$. It is now easy to see that

$$\begin{aligned}
& G(\zeta_A^1 \times \zeta_A^2) \circ (G\pi_1 + G\pi_2) \circ (\delta^1 + \delta^2) \circ (L_1\eta_A + L_2\eta_A) \\
&= G(\zeta_A^1 \times \zeta_A^2) \circ G(\hat{\delta}_A^1 \times \hat{\delta}_A^2) \circ \eta_{L_1A + L_2A} \\
&= G((\hat{\delta}_A^1 \times \hat{\delta}_A^2) \circ (\zeta_A^1 \times \zeta_A^2)) \circ \eta_{L_1A + L_2A} \\
&= \eta_{L_1A + L_2A},
\end{aligned}$$

by the assumption that ζ_A^1 and ζ_A^2 are right inverses. \square

Note that we can extract a *model* of the right type from the proof above, namely

$$\zeta_A^1 \times \zeta_A^2 \circ F\alpha : FA \rightarrow T_1FA \times T_2FA.$$

6. Applications to distributive substructural logics.

6.1. Describing T_{RL} -coalgebras validating FC1-6

The axioms FC1-6 translate as two simple frame conditions on the relational (resource) models interpreting the logic defined by L_{RL} and these axioms; one dealing with the unit I of the language, and the other with the residuation of \multimap and \otimes with respect to \otimes . The class of T_{RL} -coalgebras satisfying the second frame condition is described in Theorem 19. It is intuitive and indeed corresponds to the usual relational semantics of distributive substructural logics (see, e.g., [36]) or separation logic/BI. However, proving it ‘from first principles’ as we do here is much more intricate than might be expected and, indeed, much more so than is clear in [37].

Let $\gamma : W \rightarrow T_{\text{RL}}W$ be a T_{RL} -coalgebra validating the axioms FC1-6. Axioms FC1 means that every world $w \in W$ must have amongst its successors pairs

(w, x) and (y, w) such that x, y are ‘unit states’, viz. $x, y \models I$, moreover these are the only successors of w containing a unit state. This condition can be found in, for example, [11]. The other axioms are simply designed to capture the residuation condition in such a way that canonicity can be used; so a model in which FC2-6 are valid is simply a model in which the residuation conditions hold, viz. $a \otimes b \leq c$ iff $b \leq a - \otimes c$ iff $a \leq c \otimes b$.

To see what this means for T_{RL} -coalgebras we need the following lemma. For any T_{RL} -coalgebra $\gamma : W \rightarrow 2 \times \text{P}_c(W \times W) \times \text{P}_c(W^{\text{op}} \times W) \times \text{P}_c(W \times W^{\text{op}})$, let $\gamma_I, \gamma_{\otimes}, \gamma_{-\otimes}$ and $\gamma_{\otimes-}$ define the four components of the structure map. We define

$$\begin{aligned}\gamma_{\otimes}^{(\downarrow \times \downarrow)}(w) &= \{(x, y) \mid \exists (x', y') \in \gamma_{\otimes}(w), x \leq x', y \leq y'\} \\ \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(w) &= \{(x, y) \mid \exists (x', y') \in \gamma_{-\otimes}(w), x \leq x', y' \leq y\} \\ \gamma_{\otimes-}^{(\uparrow \times \downarrow)}(w) &= \{(x, y) \mid \exists (x', y') \in \gamma_{\otimes-}(w), x' \leq x, y \leq y'\}\end{aligned}$$

Lemma 16. *Let $\gamma : W \rightarrow T_{\text{RL}}W$ be a coalgebra and let $\hat{\gamma} : W \rightarrow T_{\text{RL}}W$ be the T_{RL} -coalgebra defined by $\gamma_I, \gamma_{\otimes}^{(\downarrow \times \downarrow)}, \gamma_{-\otimes}^{(\downarrow \times \uparrow)}, \gamma_{\otimes-}^{(\uparrow \times \downarrow)}$ defined as above, then for any valuation $v : FW \rightarrow \mathcal{U}W$ and any $w \in W$*

$$(w, \gamma, v) \models a \text{ iff } (w, \hat{\gamma}, v) \models a$$

Proof. This is an easy consequence of the the definition of T_{RL} and of the fact that the denotation of any formula is an upset. \square

We can now formulate the residuation condition.

Lemma 17. *Equations FC2-6 are valid in $\gamma : W \rightarrow T_{\text{RL}}W$ iff*

$$(y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x) \text{ iff } (y, x) \in \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z) \text{ iff } (x, z) \in \gamma_{\otimes-}^{(\uparrow \times \downarrow)}(y)$$

Proof. The ‘if’ direction follows easily by unravelling the definition of the semantics. So let us turn to the ‘only if’ direction. Assume that $a \otimes b \leq c$ iff $b \leq a - \otimes c$ and let $\gamma : W \rightarrow T_{\text{RL}}W$. We will show that $(y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)$ iff $(y, x) \in \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z)$, the case of $(x, z) \in \gamma_{\otimes-}^{(\uparrow \times \downarrow)}(y)$ is treated identically.

We start by showing that if $(y, x) \in \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z)$, then $(y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)$. Assume $(y, x) \in \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z)$ — that is, that there exist $(y', x') \in \gamma_{-\otimes}(z)$ such that $y \leq y'$ and $x' \leq x$. Consider a valuation such that $\llbracket a \rrbracket = \uparrow y$ and $\llbracket b \rrbracket = \uparrow z$. We’re assuming that $a \otimes b \leq c$ iff $b \leq a - \otimes c$, which means in particular that $b \leq a - \otimes (a \otimes b)$. Since $z \models b$, it must therefore also be the case that $z \models a - \otimes (a \otimes b)$. Since $\llbracket a \rrbracket = y$ and $y \leq y'$ we have $y' \models a$ and it must therefore also be the case that $x' \models a \otimes b$, and hence $x \models a \otimes b$. It follows that there exist $(y'', z'') \in \gamma_{\otimes}(x)$ with $y'' \geq y$ and $z'' \geq z$; that is, $(y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)$.

For the converse, we show that if $(y, x) \notin \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z)$ then $(y, z) \notin \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)$. Let $x, y, z \in W$ with $(y, x) \notin \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z)$ and consider a valuation such that $\llbracket a \rrbracket = \uparrow y$ and $\llbracket c \rrbracket = (\downarrow x)^c$. It follows that $z \models a - \otimes c$. Indeed, assume the opposite; that is,

that there exists $(y', x') \in \gamma_{-\otimes}(z)$ such that $y' \models a$ — that is, $y \leq y'$ — and $x' \not\models c$ — that is, $x' \notin (\downarrow x)^c$; that is, $x' \leq x$. This means exactly that $(y, x) \in \gamma_{-\otimes}^{(\downarrow \times \uparrow)}(z)$, a contradiction. Thus $z \models a - \otimes c$. Now assume that $(y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)$; that is, there exist $(y'', z'') \in \gamma_{\otimes}(x)$ such that $y'' \geq y, z'' \geq z$. Since $y \models a$ and $z \models a - \otimes c$ we have $y'' \models a$ and $z'' \models a - \otimes c$ and therefore we have $x \models a \otimes (a - \otimes c)$. Since we're assuming that $a \otimes b \leq c$ iff $b \leq a - \otimes c$, we have in particular that $a \otimes (a - \otimes c) \leq c$, and thus $x \models c$. But this is impossible since $\llbracket c \rrbracket = (\downarrow x)^c$. Thus we cannot have $(y, z) \in \gamma_{\otimes}(x)^{(\downarrow \times \downarrow)}$; that is, $(y, z) \notin \gamma_{\otimes}(x)^{(\downarrow \times \downarrow)}$. \square

The entire information required to encode a T_{RL} -coalgebra validating FC2-6 is therefore entirely contained in $\gamma_{\otimes}^{(\downarrow \times \downarrow)}$ (or $\gamma_{-\otimes}^{(\downarrow \times \uparrow)}$ or $\gamma_{\otimes-}^{(\uparrow \times \downarrow)}$). We will now show that we can in fact simply consider γ_{\otimes} (or $\gamma_{-\otimes}$ or $\gamma_{\otimes-}$). Note first that Lemma 17 enforces a strict constraint on $\gamma_{\otimes}^{(\downarrow \times \downarrow)}$: since $\gamma_{-\otimes}^{(\downarrow \times \uparrow)}, \gamma_{\otimes-}^{(\uparrow \times \downarrow)}$ are monotone it follows that if $z \leq z'$ we must have

$$\{(y, x) \mid (y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)\} \leq_{W^{\text{op}} \times W} \{(y', x') \mid (y', z') \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x')\} \quad (3)$$

and similarly, if $y \leq y'$

$$\{(x, z) \mid (y, z) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)\} \leq_{W \times W^{\text{op}}} \{(x', z') \mid (y', z') \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x')\} \quad (4)$$

for the Egli-Milner order on $W^{\text{op}} \times W$. The inequations (3) and (4) are quite strong and must be satisfied by any map $\gamma_{\otimes}^{(\downarrow \times \downarrow)}$ capable of reconstructing the entire T_{RL} -coalgebra. More generally, we say that a monotone map $\gamma : W \rightarrow P_c(W \times W)$ obeys the *Residuation Compatibility Condition* (RCC) if

$$\begin{aligned} \{(y, x) \mid (y, z) \in \gamma(x)\} &\leq_{W^{\text{op}} \times W} \{(y', x') \mid (y', z') \in \gamma(x')\} \text{ and} \\ \{(x, z) \mid (y, z) \in \gamma(x)\} &\leq_{W \times W^{\text{op}}} \{(x', z') \mid (y', z') \in \gamma(x')\} \end{aligned} \quad (\text{RCC})$$

We now generalize the statement of Lemma 17 to an arbitrary monotone map $\gamma : W \rightarrow P_c(W \times W)$ by defining

$$\overline{\gamma} : W \rightarrow P_c(W^{\text{op}} \times W), z \mapsto \{(y, x) \mid (y, z) \in \gamma(x)\} \quad (5)$$

$$\underline{\gamma} : W \rightarrow P_c(W \times W^{\text{op}}), y \mapsto \{(x, z) \mid (y, z) \in \gamma(x)\} \quad (6)$$

Lemma 18. *If $\gamma : W \rightarrow P_c(W \times W)$ satisfies (RCC), then $\overline{\gamma}$ and $\underline{\gamma}$ are monotone maps.*

Proof. Immediate from the definitions. \square

With this notation in place we can now state the main result of this Section, which relies on showing that we can impose (RCC) on γ_{\otimes} rather than $\gamma_{\otimes}^{(\downarrow \times \downarrow)}$.

Theorem 19. *The logic defined by L_{RL} and the axioms FC2-6 is strongly complete with respect the class of T_{RL} -coalgebras of the shape*

$$\gamma_I \times \gamma_{\otimes} \times \overline{\gamma}_{\otimes} \times \underline{\gamma}_{\otimes} : W \rightarrow \mathbb{2} \times P_c(W \times W) \times P_c(W^{\text{op}} \times W) \times P_c(W \times W^{\text{op}})$$

such that γ_{\otimes} satisfies (RCC).

Proof. Let $\gamma = \gamma_I \times \gamma_{\otimes} \times \gamma_{-\otimes} \times \gamma_{\otimes-} : W \rightarrow T_{\text{RL}}W$ be a coalgebra in which the frame conditions FC2-6 are valid. From Lemma 16 it follows that the frame conditions are valid in the coalgebra

$$\gamma_I \times \gamma_{\otimes}^{(\downarrow \times \downarrow)} \times \gamma_{-\otimes}^{(\downarrow \times \uparrow)} \times \gamma_{\otimes-}^{(\uparrow \times \downarrow)} : W \rightarrow T_{\text{RL}}W$$

and from Lemma 17 this coalgebra is in fact of the shape

$$\gamma_I \times \gamma_{\otimes}^{(\downarrow \times \downarrow)} \times \overline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}} \times \underline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}} : W \rightarrow T_{\text{RL}}W$$

If we can show that $\overline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}} = \overline{\gamma_{\otimes}}^{(\downarrow \times \downarrow)}$ and $\underline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}} = \underline{\gamma_{\otimes}}^{(\downarrow \times \downarrow)}$, then our claim will follow from Lemma 16. We show the first equality, the second one being similar.

$$\begin{aligned} (y, x) &\in \overline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}}(z) \\ \Leftrightarrow (y, z) &\in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x) && \text{Def. of } \overline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}} \\ \Leftrightarrow \exists y', z' \text{ s.th. } &y \leq y', z \leq z' \text{ and } (y', z') \in \gamma_{\otimes}(x) && \text{Def. of } \gamma_{\otimes}^{(\downarrow \times \downarrow)} \\ \Leftrightarrow \exists y', z' \text{ s.th. } &y \leq y', z \leq z' \text{ and } (y', x) \in \overline{\gamma_{\otimes}}(z') && \text{Def. of } \overline{\gamma_{\otimes}} \\ \Rightarrow \exists x', y' \text{ s.th. } &x' \leq x, y \leq y' \text{ and } (y', x') \in \overline{\gamma_{\otimes}}(z) && (RCC) \text{ and } z \leq z' \\ \Leftrightarrow (y, x) &\in \overline{\gamma_{\otimes}}^{(\downarrow \times \uparrow)}(z) && \text{Def. of } \overline{\gamma_{\otimes}}^{(\downarrow \times \uparrow)} \end{aligned}$$

Conversely, we have

$$\begin{aligned} (y, x) &\in \overline{\gamma_{\otimes}}^{(\downarrow \times \uparrow)}(z) \\ \Leftrightarrow \exists x', y' \text{ s.th. } &y \leq y', x' \leq x \text{ and } (y', x') \in \overline{\gamma_{\otimes}}(z) && \text{Def. of } \overline{\gamma_{\otimes}}^{(\downarrow \times \uparrow)} \\ \Leftrightarrow \exists x', y' \text{ s.th. } &y \leq y', x' \leq x \text{ and } (y', z) \in \gamma_{\otimes}(x') && \text{Def. of } \overline{\gamma_{\otimes}} \\ \Rightarrow \exists y', z' \text{ s.th. } &y \leq y', z \leq z' \text{ and } (y', z') \in \gamma_{\otimes}(x) && \gamma_{\otimes} \text{ monotone and } x' \leq x \\ \Leftrightarrow (y, z) &\in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x) && \text{Def. of } \gamma_{\otimes}^{(\downarrow \times \downarrow)} \\ \Leftrightarrow (y, x) &\in \overline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}}(z) && \text{Def. of } \overline{\gamma_{\otimes}^{(\downarrow \times \downarrow)}} \end{aligned}$$

□

Example 20. *Heaps, which for the sake of brevity and convenience we shall define as partial maps on \mathbb{N}_+ with finite domain, form a model satisfying Theorem 19. To show this, we first define the set of heaps*

$$\mathcal{H} = \{f : \mathbb{N}_+ \rightarrow \mathbb{N}_+ \mid \text{dom}(f) \in \mathcal{P}_{\omega}(\mathbb{N}_+)\}$$

It forms a poset under the order

$$f \leq g \text{ if } f = g \upharpoonright \text{dom}(f)$$

We now define a T_{RL} -coalgebra structure on \mathcal{H} as follows:

$$\begin{aligned} \gamma_I : \mathcal{H} &\rightarrow \mathbb{2}, & f &\mapsto \begin{cases} 0 & \text{if } \text{dom}(f) = \emptyset \\ 1 & \text{else} \end{cases} \\ \gamma_{\otimes} : \mathcal{H} &\rightarrow \text{P}_c(\mathcal{H} \times \mathcal{H}), & f &\mapsto \{(g, h) \mid \text{dom}(g) \cap \text{dom}(h) = \emptyset, \\ & & & f \upharpoonright \text{dom}(g) = g, f \upharpoonright \text{dom}(h) = h\} \end{aligned}$$

Clearly γ_I is trivially monotone, and validates the axioms FC1. To see that γ_{\otimes} is well-typed, note first that $\gamma_{\otimes}(f)$ is a down-set, and therefore also convex. Moreover, it is not hard to see that if f' extends f ; that is, $f \leq f'$ then $\gamma_{\otimes}(f) \subseteq \gamma_{\otimes}(f')$. The first half of the Egli-Milner definition is therefore trivially satisfied. For the second half, if $(g', h') \in \gamma_{\otimes}(f')$, then $(g' \upharpoonright \text{dom}(f), h' \upharpoonright \text{dom}(f)) \in \gamma_{\otimes}(f)$ provides the witness we need. It follows that γ_{\otimes} is also monotone and thus well-typed. Finally, we need to check that it satisfies (RCC); that is, if $h \leq h'$

$$\{(g, f) \mid (g, h) \in \gamma_{\otimes}(f)\} \leq_{\mathcal{H}^{\text{op}} \times \mathcal{H}} \{(g', f') \mid (g', h') \in \gamma_{\otimes}(f')\}$$

Starting from (g, f) in the first set, we define f' by $f' = f$ on $\text{dom}(f)$ and $f' = h'$ on $\text{dom}(h') \cap (\text{dom}(f))^c$ and get an element (g, f') such that $(g, h') \in \gamma_{\otimes}(f')$, $g \leq g$ and $f \leq f'$, which shows that the first direction of the Egli-Milner order holds. For the second, start with (g', f') in the second set and define f by $f = h$ on $\text{dom}(h)$ and $f = g'$ on $\text{dom}(g')$. It is easy to see that $(g', h) \in \gamma_{\otimes}(f)$, $g' \leq g'$ and $f \leq f'$, providing a witness for the second direction of the Egli-Milner order. It follows that γ_{\otimes} satisfies (RCC), and heaps therefore provide a class of models for the distributive full Lambek calculus.

6.2. Additional frame conditions.

Structural rules can be added to the full distributive Lambek calculus to form new logics. These rules are the exchange rule (e), the contraction rule (c), the left weakening rule (lw), and the right weakening rule (rw). Relevance logic for example consists in adding (c) and (e) to the distributive Lambek calculus, adding only (c) defines the positive MALL⁺ fragment of linear logic ([36]), whilst the combination of (lw), (rw) and (e) defines affine logic. These structural rules correspond to (in)equations in the theory of residuated lattices (see [36, 1, 2]); that is, in the language of L_{RL} -algebras. Let us go through them in order.

Exchange. The exchange rule corresponds to the axiom (e) given by $a \otimes b = b \otimes a$. It is easy to see from the results of Section 4.2 that it is canonical. The class of T_{RL} -coalgebras in which this axiom is valid is characterised by

$$(y, z) \in \gamma_{\otimes}(x) \Rightarrow (z, y) \in \gamma_{\otimes}^{(\downarrow \times \downarrow)}(x)$$

in other words, the ternary relation defined by $\gamma_{\otimes}^{(\downarrow \times \downarrow)}$ is closed under permuting the successor states. In the case of the boolean Lambek calculus, that is to say in the classical case, we clearly have $(y, z) \in \gamma_{\otimes}(x)$ iff $(z, y) \in \gamma_{\otimes}(x)$.

Contraction. The contraction rule corresponds to the axiom (c) given by $a \leq a \otimes a$; that is, increasing idempotency (see [1, 2]). Once again, the canonicity of this axiom is almost immediate from Section 4.2. The class of T_{RL} -coalgebras in which (c) is valid is characterized by

$$\forall x \in W \exists (y, z) \in \gamma(x) \text{ s.th. } x \leq y, z$$

and in the classical case, this means that $(x, x) \in \gamma_{\otimes}(x)$.

Left weakening. The weakening rule corresponds to the axiom $a \leq I$, viz. every state is a unit state. Coalgebraically, this means that the component γ_I of a T_{RL} -coalgebra is the constant map $0 \in \mathbb{2}$. In the classical case, this amounts to saying that $I = \top$.

For the right weakening rule we need to introduce a new unit, which we will denote J . In fact we use this as an opportunity to introduce a whole new signature, dual to the signature defining L_{RL} . We define

$$L_{\text{RL}}^{\partial} : \mathcal{C} \rightarrow \mathcal{C}, \begin{cases} L_{\text{RL}}^{\partial} A = \mathbf{F}\{J, a \oplus b, a \oplus b, a \oplus b \mid a, b \in \mathbf{UA}\} / \equiv \\ L_{\text{RL}}^{\partial} f : L_{\text{RL}} A \rightarrow L_{\text{RL}} B, [a]_{\equiv} \mapsto [f(a)]_{\equiv}, \end{cases}$$

where \equiv is the fully invariant equivalence relation in \mathcal{C} generated by following the Distribution Laws for non-empty finite subsets X of A :

$$\begin{array}{ll} \text{DL}^{\partial}1. \ \wedge X \oplus a = \wedge[X \oplus a] & \text{DL}^{\partial}4. \ \wedge X \oplus a = \wedge[X \oplus a] \\ \text{DL}^{\partial}2. \ a \oplus \wedge X = \wedge[a \oplus X] & \text{DL}^{\partial}5. \ \vee X \oplus a = \vee[X \oplus a] \\ \text{DL}^{\partial}3. \ a \oplus \vee X = \vee[a \oplus X] & \text{DL}^{\partial}6. \ a \oplus \wedge X = \wedge[a \oplus X]. \end{array}$$

Note that the equations $\text{DL}^{\partial}1$ -6 are dual to the equations $\text{DL}1$ -6. In particular, \oplus is a binary \square , whilst \otimes is a binary \diamond . The language $\mathcal{L}(L_{\text{RL}}^{\partial})$ will also be interpreted in T_{RL} coalgebras, via the semantic transformation $\delta^{\text{RL}\partial} : L_{\text{RL}}^{\partial} G \rightarrow GT_{\text{RL}}$ defined at every poset W via its action on the generators:

$$\begin{aligned} \delta_W^{\text{RL}\partial}(J) &= \{t \in T_{\text{RL}} W \mid \pi_1(t) = 1 \in \mathbb{2}\} \\ \delta_W^{\text{RL}\partial}(u \oplus v) &= \{t \in T_{\text{RL}} W \mid \forall (x, y) \in \pi_2(t), x \in u \text{ or } y \in v\} \\ \delta_W^{\text{RL}\partial}(u \oplus w) &= \{t \in T_{\text{RL}} W \mid \exists (x, y) \in \pi_3(t), x \notin u \text{ and } y \in w\} \\ \delta_W^{\text{RL}\partial}(w \oplus v) &= \{t \in T_{\text{RL}} W \mid \exists (x, y) \in \pi_4(t), x \notin v \text{ and } y \in w\} \end{aligned}$$

A proof very similar to that of Proposition 1 shows that $\delta^{\text{RL}\partial}$ is well-defined, and using exactly the same technique as in Theorem 3, it can be shown that the adjoint transpose of $\delta^{\text{RL}\partial}$ has right inverses at every distributive lattice. It follows that $\mathcal{L}(L_{\text{RL}}^{\partial})$ is strongly complete with respect to the class of all T_{RL} -coalgebras. However, the intended interpretation of \oplus and \oplus is once again that they should be the left and right residuals of \oplus . We therefore introduce the following axioms.

$$\begin{array}{ll}
\text{FC}^\partial 1. & a \oplus J = a, \ J \oplus a = a, \\
\text{FC}^\partial 2. & I \leq a \multimap a, \ I \leq a \multimap a, \\
\text{FC}^\partial 3. & a \oplus (b \multimap c) \leq (a \oplus b) \multimap c, \\
\text{FC}^\partial 4. & (c \oplus b) \oplus a \leq c \oplus (a \oplus b), \\
\text{FC}^\partial 5. & (a \oplus b) \oplus b \leq a, \text{ and} \\
\text{FC}^\partial 6. & b \oplus (b \multimap a) \leq a.
\end{array}$$

These axioms capture identities and residuation, and are therefore of exactly the same shape as axioms FC1-6. A dual construction to that of Section 6.1, shows that equations FC[∂]1-6 are valid in a T_{RL} -coalgebra $\gamma : W \rightarrow T_{\text{RL}}$ iff it satisfies

$$(y, z) \in \gamma_{\oplus}^{(\uparrow \times \uparrow)}(x) \text{ iff } (y, x) \in \gamma_{\oplus}^{(\uparrow \times \downarrow)}(z) \text{ iff } (x, z) \in \gamma_{\oplus}^{(\downarrow \times \uparrow)}(y)$$

where $\bar{\gamma}_{\oplus}$ and $\underline{\gamma}_{\oplus}$ are defined by Eqs. (5), (6). The fact that $\gamma_{\oplus}^{(\uparrow \times \downarrow)}$ and $\gamma_{\oplus}^{(\downarrow \times \uparrow)}$ should be monotone means that $\gamma_{\oplus}^{(\uparrow \times \uparrow)}$ must satisfy (RCC). Finally, dualising Theorem 19 we get

Theorem 21. $\mathcal{L}(L_{\text{RL}}^\partial)$ quotiented by the axioms FC[∂]2-6 is strongly complete with respect to the class of T_{RL} -coalgebras of the shape

$$\gamma_J \times \gamma_{\oplus} \times \bar{\gamma}_{\oplus} \times \underline{\gamma}_{\oplus} : W \rightarrow \mathbb{2} \times \text{P}_c(W \times W) \times \text{P}_c(W^{\text{op}} \times W) \times \text{P}_c(W \times W^{\text{op}})$$

such that γ_{\oplus} satisfies (RCC).

Combining this theorem with our earlier result on the modularity of coalgebraic completeness-via-canonicity (Theorem 15) we get that the logic defined by $\mathcal{L}(L_{\text{RL}} + L_{\text{RL}}^\partial)$ and the axioms FC2-6 and FC[∂]2-6 is strongly complete with respect to $T_{\text{RL}} \times T_{\text{RL}}$ -coalgebras of the shape:

$$\begin{aligned}
& (\gamma_I \times \gamma_{\otimes} \times \bar{\gamma}_{\otimes} \times \underline{\gamma}_{\otimes}) \times (\gamma_J \times \gamma_{\oplus} \times \bar{\gamma}_{\oplus} \times \underline{\gamma}_{\oplus}) : \\
& W \rightarrow (\mathbb{2} \times \text{P}_c(W \times W) \times \text{P}_c(W^{\text{op}} \times W) \times \text{P}_c(W \times W^{\text{op}}))^2
\end{aligned} \tag{7}$$

where both γ_{\otimes} and γ_{\oplus} satisfy (RCC). We can now return to the structural rules which we described at the beginning of this section and account for right weakening.

Right weakening. Working in the signature of the fusion $\mathcal{L}(L_{\text{RL}} \oplus L_{\text{RL}}^\partial)$, right weakening corresponds to the axioms $J \leq a$ which is clearly canonical. Coalgebraically this axiom corresponds to saying that if $\gamma_J(w) = 1$ in a model, then $w \models a$ for any formula a and any valuation. This is clearly not possible: let $p \in V$ be a propositional variable and consider a valuation such that $\llbracket p \rrbracket = (\downarrow w)^c$ (a valid upset), clearly $w \not\models p$. It follows that γ_J must be the constant monotone map $0 \in \mathbb{2}$. In particular if left weakening is also allowed, then nothing distinguishes γ_I and γ_J and J holds precisely when I does not; that is, never. In the classical case we clearly have $J = \perp$.

The logic defined by $L_{\text{RL}} + L_{\text{RL}}^\partial$ and the axioms FC1-6, FC[∂]1-6 and the exchange axioms $a \otimes b = b \otimes a$ and $a \oplus b = b \oplus a$ is the positive fragment of the

Multiplicative-Additive Linear Logic (MALL^+ in [36]). Many additional features could be added to the quotient of $\mathcal{L}(L_{\text{RL}} + L_{\text{RL}}^\partial)$ under FC1-6 and FC^∂ 1-6, most notably one could define the ‘negation’ operations $\sim a \triangleq a - \otimes J$ and $\neg a \triangleq J \otimes - a$ and use them to connect the behaviour of the two halves of the signature. We refer the reader to [36],[1] and [2] for such considerations. To conclude we return to our heap model and show that it is a model for the logic we have just defined.

Example 22 ([12]). Recall from Example 20 that the poset of heaps \mathcal{H} can be equipped with a map $\gamma_\otimes : \mathcal{H} \rightarrow \text{P}_c(\mathcal{H}^{\text{op}} \times \mathcal{H})$ which satisfies (RCC), and can thus be used to reconstruct an entire T_{RL} -coalgebra structure (modulo a monotone map $\gamma_I : \mathcal{H} \rightarrow \mathbb{2}$). We will now define a second such map which will interpret the dual signature given by L_{RL}^∂ in the way suggested by [12]. We choose an arbitrary upset of heaps $U \subseteq \mathcal{H}$ and define

$$\begin{aligned} \gamma_J : \mathcal{H} &\rightarrow \mathbb{2}, & f &\mapsto \begin{cases} 1 & \text{if } f \in U \\ 0 & \text{else} \end{cases} \\ \gamma_\oplus : \mathcal{H} &\rightarrow \text{P}_c(\mathcal{H} \times \mathcal{H}), & f &\mapsto \{(g, h) \mid \text{dom}(f) = \text{dom}(g) \cap \text{dom}(h) \\ & & & g \upharpoonright \text{dom}(f) = h \upharpoonright \text{dom}(f)\} \end{aligned}$$

The map γ_J is well-typed by construction. To see that γ_\oplus is also well-typed, note first that $\gamma_\oplus(f)$ is this time an upset, and therefore convex. It is not difficult to check in the same way as in Example 20 that if $f \leq f'$ then $\gamma_\oplus(f) \leq \gamma_\oplus(f')$ for the Egli-Milner order; that is, γ_\oplus is monotone and thus well-typed. Simple set-theoretic considerations of the same type as in Example 20 also show that γ_\oplus defined as above satisfies (RCC). It follows that the data of $\gamma_\otimes, \gamma_\oplus$ and U endows the poset \mathcal{H} of heaps with a $(T_{\text{RL}})^2$ -coalgebra structure allowing the interpretation of $\mathcal{L}(L_{\text{RL}} + L_{\text{RL}}^\partial)$ -formulas. It is shown in [12] that several additional $\mathcal{L}(L_{\text{RL}} + L_{\text{RL}}^\partial)$ -axioms may be envisaged. Most notably the axioms $a \leq a \oplus 0$ and $a \oplus 0 \leq a$ (which combined give FC^∂ 1, an axiom we have chosen not to enforce as ‘standard’), the contraction axiom $a \oplus a \leq a$ (dual to axiom (c)) and the weak distribution axiom $a \otimes (b \oplus c) \leq (a \otimes b) \oplus a$. These axioms are canonical, and Theorems 14 and 15 therefore easily provides strong completeness for these axioms too, although as was is shown in [12], no heap model can validate all three of these axioms simultaneously.

7. Conclusion and future work

We have shown how distributive substructural logics can be formalized and given a semantics in the framework of coalgebraic logic, and highlighted the modularity of this approach. By choosing a syntax whose operators explicitly follow distribution rules, we can use the elegant topological theory of canonicity for DLs, and in particular the notion of smoothness and of topology matching, to build a set of canonical (in)equation capturing the distributive full Lambek calculus. The coalgebraic approach makes the connection between algebraic canonicity and canonical models explicit, categorical and generalizable.

The modularity provided by our approach is twofold. Firstly, we have a generic method for building canonical (in)equations by using the Principle of Matching Topologies. Getting completeness results with respect to simple Kripke models for variations of the distributive full Lambek calculus (e.g. relevant logic, MALL^+ , etc...) becomes very straightforward. Secondly, adding more operators to the fundamental language simply amounts to taking a *coproduct* of syntax constructors (e.g., $L_{\text{RL}} + L_{\text{ML}}$ to define modal substructural logics) and interpreting it with a *product* of model constructors (e.g., $T_{\text{RL}} \times T_{\text{ML}}$). This is particularly suited to logics which build on BI such as the bi-intuitionistic boolean BI of [12].

The operators \otimes , \multimap , \boxtimes all satisfy simple distribution laws, but our approach could also accommodate operators with more complicated distribution laws and non-relational semantics. For example, the theory presented in this work could perhaps be extended to cover a graded version of \otimes , say \otimes_k , whose interpretation would be ‘there are at least k ways to separate a resource such that...’ or ‘a resource can be split in two at a cost of k ...’, the semantics would be given by coalgebras of the type $2 \times \mathcal{B}(- \times -)$ where \mathcal{B} is the ‘bag’ or multiset functor. Similarly, a graded version \rightarrow_k of the intuitionistic implication whose meaning would be ‘... implies ... apart from at most k exceptions’ and interpreted by $\mathcal{B}(- \times -)$ -coalgebras could also be covered by our approach. Crucially, such operators do satisfy (more complicated) distribution laws which lead to generalizations of the results in Section 4.2, and the possibility of building canonical (in)equations. The coalgebraic infrastructure would then allow the rest of the theory to stay essentially unchanged. We are currently investigating these possibilities.

Acknowledgement

The first author is funded by EPSRC grant EP/H008373/2.

References

- [1] H. Ono, Substructural Logics and Residuated Lattices – an Introduction, Trends in Logic 20 (2003) 177–212.
- [2] N. Galatos, P. Jipsen, T. Kowalski, H. Ono, Residuated Lattices: An Algebraic Glimpse at Substructural Logics: An Algebraic Glimpse at Substructural Logics, vol. 151, Elsevier, 2007.
- [3] J.-Y. Girard, Linear logic., Theoret. Comp. Sci. 50 (1987) 1–102.
- [4] P. W. O’Hearn, D. J. Pym, The logic of bunched implications, Bulletin of Symbolic Logic 5 (02) (1999) 215–244.
- [5] D. Pym, P. W. O’Hearn, H. Yang, Possible Worlds and Resources: The Semantics of BI., Theoret. Comp. Sci. 315 (1) (2004) 257–305.

- [6] D. Galmiche, D. Méry, D. Pym, The Semantics of BI and Resource Tableaux., *Mathematical Structures in Computer Science* 15 (2005) 1033–1088.
- [7] M. Collinson, D. Pym, Algebra and logic for resource-based systems modelling, *Mathematical Structures in Computer Science* 5 (2009) 959–1027.
- [8] S. S. Ishtiaq, P. W. O’Hearn, BI as an assertion language for mutable data structures, in: *ACM SIGPLAN Notices*, vol. 36.3, ACM, 14–26, 2001.
- [9] J. C. Reynolds, Separation logic: A logic for shared mutable data structures, in: *Logic in Computer Science, 2002. Proceedings. 17th Annual IEEE Symposium on*, IEEE, 55–74, 2002.
- [10] M. Collinson, B. Monahan, D. J. Pym, *A discipline of mathematical systems modelling*, College Publications, 2012.
- [11] C. Calcagno, P. Gardner, U. Zarfaty, Context logic as modal logic: completeness and parametric inexpressivity, in: *ACM SIGPLAN Notices*, vol. 42.1, ACM, 123–134, 2007.
- [12] J. Brotherston, J. Villard, Sub-classical Boolean bunched logics and the meaning of par, in: *LIPICs-Leibniz International Proceedings in Informatics*, vol. 41, Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2015.
- [13] P. Blackburn, M. de Rijke, Y. Venema, *Modal Logic*, vol. 53 of *Cambridge Tracts in Theoretical Computer Science*, Cambridge University Press, 2001.
- [14] B. Jónsson, On the canonicity of Sahlqvist identities, *Studia Logica* 53 (4) (1994) 473–492.
- [15] J. M. Dunn, M. Gehrke, A. Palmigiano, Canonical Extensions and Relational Completeness of Some Substructural Logics, *The Journal of Symbolic Logic* 70 (3) (2005) pp. 713–740.
- [16] M. Gehrke, Generalized Kripke Frames, *Studia Logica* 84 (2) (2006) 241–275.
- [17] D. Coumans, M. Gehrke, L. van Rooijen, Relational semantics for a fragment of linear logic, *Proceedings of PhDs in Logic III*.
- [18] T. Suzuki, Canonicity results of substructural and lattice-based logics, *The Review of Symbolic Logic* 4 (2011) 1–42.
- [19] M. Gehrke, J. Harding, Bounded Lattice Expansions, *Journal of Algebra* 238 (1) (2001) 345 – 371.
- [20] M. Gehrke, B. Jónsson, Bounded distributive lattice expansions., *Mathematica Scandinavica* 94 (2004) 13–45.

- [21] Y. Venema, Algebras and Coalgebras, in: J. van Benthem, P. Blackburn, F. Wolter (Eds.), *Handbook of Modal Logic*, Elsevier, 2006.
- [22] F. Dahlqvist, D. Pattinson, Some Sahlqvist Completeness Results for Coalgebraic Logics, in: *FOSSACS 2013. Proceedings*, vol. 7794 of *LNCS*, Springer, 193–208, 2013.
- [23] C. Cirstea, D. Pattinson, Modular construction of complete coalgebraic logics, *Theor. Comput. Sci.* 388 (1-3) (2007) 83–108.
- [24] F. Dahlqvist, D. Pattinson, On the Fusion of Coalgebraic Logics, in: *CALCO 2011. Proceedings*, vol. 6859 of *LNCS*, Springer, 161–175, 2011.
- [25] C. Kupke, A. Kurz, D. Pattinson, Algebraic Semantics for Coalgebraic Logics, in: *CMCS 2004*, vol. 106 of *Electr. Notes in Theoret. Comp. Sci.*, 219–241, 2004.
- [26] C. Kupke, A. Kurz, D. Pattinson, Ultrafilter Extensions for Coalgebras, in: *CALCO*, vol. 3629 of *LNCS*, Springer, 263–277, 2005.
- [27] B. Jacobs, A. Sokolova, Exemplaric Expressivity of Modal Logics, *J. Log. and Comput.* 20 (2010) 1041–1068.
- [28] J.-R. Courtault, D. Galmiche, A modal BI logic for dynamic resource properties, in: *Logical Foundations of Computer Science*, Springer, 134–148, 2013.
- [29] J.-R. Courtault, D. Galmiche, A modal separation logic for resource dynamics, *Journal of Logic and Computation* (2015) exv031.
- [30] J. M. Dunn, Positive modal logic, *Studia Logica* 55 (2) (1995) 301–317.
- [31] M. Collinson, K. McDonald, D. Pym, Layered graph logic as an assertion language for access control policy models, *Journal of Logic and Computation* In Press.
- [32] M. Bílková, A. Kurz, D. Petrişan, J. Velebil, Relation liftings on preorders and posets, in: *Algebra and Coalgebra in Computer Science*, Springer, 115–129, 2011.
- [33] A. Balan, A. Kurz, J. Velebil, Positive fragments of coalgebraic logics, in: *Algebra and Coalgebra in Computer Science*, Springer, 51–65, 2013.
- [34] B. Jónsson, A. Tarski, Boolean Algebras with Operators. Part 1., *Amer. J. Math.* 33 (1951) 891–937.
- [35] M. Gehrke, B. Jónsson, Bounded distributive lattices with operators., *Mathematica Japonica* 40, No. 2 (1994) 207–215.
- [36] G. Restall, *An introduction to substructural logics*, Routledge, 2002.

- [37] F. Dahlqvist, D. Pym, Completeness-via-canonicity for distributive substructural logics, a coalgebraic perspective, in: Proc 15th Int. Conf. on Relational and Algebraic Methods in Computer Science (RAMiCS 2015), vol. LNCS 9348, 119–135, 2015.

Appendix

Proof of Theorem 3. We prove the result for L_{RL} and T_{RL} , the same technique can readily be applied to L_{ML} and T_{ML} . We need to prove that $\hat{\delta}^{RL}_A$ has a natural right inverse. By describing a prime filter of LA in terms of the ‘generators’ it contains we get the following characterization of $\hat{\delta}^{RL}_A : TPfA \rightarrow PfLA$:

$$\begin{aligned} I \in \hat{\delta}_A(U_1, U_2, U_3, x) &\text{ iff } x = 0 \\ a \otimes b \in \hat{\delta}_A(U_1, U_2, U_3, x) &\text{ iff } \exists (F_1, F_2) \in U_1, (a, b) \in (F_1, F_2) \\ a - \otimes b \in \hat{\delta}_A(U_1, U_2, U_3, x) &\text{ iff } \forall (F_1, F_2) \in U_2, a \in F_1 \Rightarrow b \in F_2 \\ a \otimes - b \in \hat{\delta}_A(U_1, U_2, U_3, x) &\text{ iff } \forall (F_1, F_2) \in U_3, b \in F_2 \Rightarrow a \in F_1 \end{aligned}$$

At every distributive lattice A , we now define $\gamma_A : PfLA \rightarrow TPfA$ by

$$F \mapsto \begin{cases} 0 & \text{if } I \in F, 1 \text{ else} \\ \{(F_1, F_2) \in (PfA)^2 \mid a \otimes b \in F \text{ whenever } (a, b) \in (F_1, F_2)\} \\ \{(G_1, G_2) \in (PfA)^2 \mid a \in G_1 \Rightarrow b \in G_2 \text{ whenever } a - \otimes b \in F\} \\ \{(H_1, H_2) \in (PfA)^2 \mid b \in H_2 \Rightarrow a \in H_1 \text{ whenever } a \otimes - b \in F\}. \end{cases}$$

By unravelling the definition of $\hat{\delta}^{RL}_A$, we get that for γ_A to be a right inverse of $\hat{\delta}^{RL}_A$ it must satisfy:

$$(a \otimes b) \in F \Leftrightarrow \exists (F_1, F_2) \in \pi_1(\gamma_A(F)) \text{ s.th. } (a, b) \in (F_1, F_2) \quad (8)$$

$$a - \otimes b \in F \Leftrightarrow \forall G_1, G_2 \in \pi_2(\gamma_A(F)) \quad a \in G_1 \Rightarrow b \in G_2 \quad (9)$$

$$a \otimes - b \in F \Leftrightarrow \forall H_1, H_2 \in \pi_2(\gamma_A(F)) \quad b \in H_2 \Rightarrow a \in H_1 \quad (10)$$

Note that the first component of γ_A poses no difficulty since

$$I \in \hat{\delta}_A(\gamma_A(F)) \Leftrightarrow \pi_1(\gamma_A(F)) = 0 \Leftrightarrow I \in F$$

Note also that the right-to-left direction of (8) and the left-to-right direction of (9,10) follows immediately from the definitions. The hard part of the proof are the opposite directions.

Left-to-right direction of (8): Assume that $a \otimes b \in F$, we need to build F_1, F_2 such that: (1) $a \in F_1$, (2) $b \in F_2$ and (3) $(F_1, F_2) \in \pi_1(\gamma_A(F))$; that is, $a' \in F_1$ and $b' \in F_2$ implies $a' \otimes b' \in F$, or equivalently, $a' \otimes b' \notin F$ implies $a' \notin F_1$ or $b' \notin F_2$. We will build F_1, F_2 , using a proof which is similar to the proof of the prime ideal theorem for filter-ideal pairs. Let us denote by $\mathcal{P}(a, b)$ the set of pairs $((F_1, I_1), (F_2, I_2))$ such that

1. $\uparrow a \subseteq F_1$
2. $\uparrow b \subseteq F_2$
3. $I_1 = \{c \mid \exists d \in F_2 \text{ s.th. } c \otimes d \notin F\}$
4. $I_2 = \{d \mid \exists c \in F_1 \text{ s.th. } c \otimes d \notin F\}$
5. $F_1 \subseteq \{c \mid \forall d \in F_2, c \otimes d \in F\}$
6. $F_2 \subseteq \{d \mid \forall c \in F_1, c \otimes d \in F\}$

We make the following observations about $\mathcal{P}(a, b)$

- it is non-empty: $((\uparrow a, \{c \mid c \otimes b \notin F\}), (\uparrow b, \{d \mid a \otimes d \notin F\})) \in \mathcal{P}(a, b)$
- it forms a poset under component-wise set inclusion.
- I_1, I_2 are ideals. It is clear that they are down-sets. Moreover, if $c, c' \in I_1$ then there exist $d, d' \in F_2$ s.th. $c \otimes d, c' \otimes d' \notin F$, and as a consequence $c \otimes (d \wedge d'), c' \otimes (d \wedge d') \notin F$ and $d \wedge d' \in F_2$ since F_2 is a filter. Since F is prime and \otimes distributes over joins it follows that $(c \vee c') \otimes (d \wedge d') \notin F$, and thus $d \wedge d'$ witnesses the fact that $c \vee c' \in I_1$. The proof is identical for I_2 .
- $F_1 \cap I_1 = F_2 \cap I_2 = \emptyset$ for each $((F_1, I_1), (F_2, I_2)) \in \mathcal{P}(a, b)$. Indeed assume that there exist $f \in F_1, i \in I_1$ such that $f \leq i$, then we have $f \otimes d \leq i \otimes d$ for some $d \in F_2$ such that $i \otimes d \notin F$. But by construction we must have $f \otimes d \in F$ which contradicts $i \otimes d \notin F$, since F is a filter.

Let us now check that $\mathcal{P}(a, b)$ has upper bounds of chains. Assume that $((F_1^i, I_1^i), (F_2^i, I_2^i)) \in \mathcal{P}(a, b), i \in \omega$ and define

$$F_i^\infty = \bigcup_j F_i^j, \quad I_i^\infty = \bigcup_j I_i^j, \quad i = 1, 2$$

It is well-known and easy to check that the union of a chain of filter (resp. ideals) is a filter (resp. ideals). Let us now check that conditions 1.-6. are satisfied too. The first two conditions are trivially satisfied. For 3.-4., let $x \in I_1^\infty$, by definition there exists $i \in \omega$ s.th. $c \in I_1^i$ and thus there exist $d \in F_1^i$ s.th. $c \otimes d \notin F$, but clearly $d \in F_1^\infty$ too, which shows that $I_1^\infty \subseteq \{c \mid d \in F_2^\infty \text{ s.th. } c \otimes d \notin F\}$. The opposite inclusion works in exactly the same way: let c be s.th. there exists $d \in F_2^\infty$ s.th. $c \otimes d \notin F$, then this d can be traced back to a certain F_2^i and thus $c \in I_1^i$. The proof for 5.-6. is very similar, let $c \in F_1^\infty$, then there exist $i \in \omega$ s.th. $c \in F_1^i$. Now let $d \in F_2^\infty$, then there exist $j \in \omega$ s.th. $d \in F_2^j$. By taking $k = \max(i, j)$ we get that $c \in F_1^k, d \in F_2^k$ from which it follows that $c \otimes d \in F$.

We can now apply Zorn's lemma to get the existence of a maximal element of $\mathcal{P}(a, b)$, which we will call

$$((\hat{F}_1, \hat{I}_1), (\hat{F}_2, \hat{I}_2))$$

and we claim that \hat{F}_1, \hat{F}_2 are two prime filters satisfying conditions (1)-(3) which we specified at the beginning of the proof. It is clear that $a \in \hat{F}_1$ and $b \in \hat{F}_2$, thus (1) and (2) are satisfied. For (3), assume that $a' \otimes b' \notin F$ and that $a' \in \hat{F}_1$, then by construction, $b' \in \hat{I}_2$, and since $\hat{F}_2 \cap \hat{I}_2 = \emptyset$ we get $b' \notin \hat{F}_2$ which is what we needed to show. The last step of the proof is to show that \hat{F}_1, \hat{F}_2 are prime. Assume that $c \vee c' \in \hat{F}_1$ but that $c, c' \notin \hat{F}_1$. It follows that

$$((\hat{F}_1, \hat{I}_1), (\hat{F}_2, \hat{I}_2)) \not\sqsubseteq (((\hat{F}_1 \cup \{c\}), \hat{I}_1), (\hat{F}_2, \{d \mid \exists c \in \langle \hat{F}_1 \cup \{c\} \rangle \text{ s.t. } c \otimes d \notin F\}))$$

where $\langle \hat{F}_1 \cup \{c\} \rangle$ is the filter generated by $\hat{F}_1 \cup \{c\}$. Since the left-hand side of the inequality is maximal, it cannot be the case that the right-hand side belongs to $\mathcal{P}(a, b)$ that is, one of the conditions 1.-6. cannot hold. Clearly 1. and 2. must hold, and 3. and 4. hold by construction, thus 5. or 6. cannot hold. In fact both conditions will not hold precisely if there exist $d \in \hat{F}_2$ and $f \in \hat{F}_1$ such that $(f \wedge c) \otimes d \notin F$; that is, $f \wedge c \in \hat{I}_1$; that is, there exist $i \in I_1$ s.th.

$$f \wedge c \leq i$$

A completely similar reasoning shows that there must exist $f' \in \hat{F}_1$ and $i' \in \hat{I}_1$ such that

$$f' \wedge c' \leq i'$$

It thus follows that

$$(f \vee f') \wedge (f \vee c') \wedge (c \vee f') \wedge (c \vee c') \leq i \vee i'$$

Since \hat{F}_1 is a filter, \hat{I}_1 is an ideal, and we've assumed $c \vee c' \in \hat{F}_1$, we get that $\hat{F}_1 \cap \hat{I}_1 \neq \emptyset$ which is a contradiction by virtue of the properties of elements of $\mathcal{P}(a, b)$. Thus either c or c' belongs to \hat{F}_1 which is thus prime as desired. A completely analogous argument shows that \hat{F}_2 is prime too.

Right-to-left direction of (9-10): We show the contrapositive; that is, that if $a - \otimes b \notin F$ there exists F_1, F_2 such that $a \in F_1$ and $b \notin F_2$. We proceed as in the case of \otimes by defining the set $\mathcal{P}(a, b)$ of filter-ideal pairs $((F_1, I_1), (F_2, I_2))$ such that

1. $\uparrow a \subseteq F_1$
2. $I_1 = \{c \mid \exists d \in I_2 (c - \otimes d) \in F\}$
3. $F_2 = \{d \mid \exists c \in F_1 (c - \otimes d) \in F\}$
4. $\downarrow a \subseteq I_2$
5. $F_1 \subseteq \{c \mid \forall d \in I_2 (c - \otimes d) \notin F\}$
6. $I_2 \subseteq \{d \mid \forall c \in F_1 (c - \otimes d) \notin F\}$

We make the following observations about $\mathcal{P}(a, b)$

- it is not empty: $((\uparrow a, \{c \mid \exists d \leq b(c \multimap d) \in F\}), (\{d \mid a \multimap d \in F\}, \downarrow b)) \in \mathcal{P}(a, b)$. We need only check that conditions 5. and 6. are satisfied. Let c be s.th. there exist $d \leq b$ with $c \multimap d \in F$, and assume $a \leq c$, it follows that

$$F \ni c \multimap d \leq a \multimap d \leq a \multimap b \notin F$$

a contradiction. Similarly, if there exist d such that $a \multimap d \in F$ and $d \leq b$, then $F \ni a \multimap d \leq a \multimap b \notin F$, a contradiction.

- it forms a poset under component-wise set inclusion
- I_1 is an ideal: assume $c \in I_1$ and $c' \leq c$, since \multimap is antitone in its first argument, we get $c \multimap d \leq c' \multimap d$ and thus $c' \multimap d \in F$. Moreover, if $c, c' \in I_1$ then there exist $d, d' \in I_2$ such that $c \multimap d, c' \multimap d' \in F$. Since F is a filter, and I_2 is an ideal, we get $c \multimap (d \vee d'), c' \multimap (d \vee d') \in F$ and $(d \vee d') \in I_2$. If we now consider $(c \vee c') \multimap (d \vee d')$ we get by the anti-join preservation law of \multimap , $c \multimap (d \vee d') \wedge c' \multimap (d \vee d')$ which is a meet of elements of F and thus an element of F . Thus $(d \vee d')$ witnesses that $c \vee c' \in I_1$
- For completely dual reasons, F_2 is a filter.
- $F_1 \cap I_2 = F_2 \cap I_1 = \emptyset$ for each $((F_1, I_1), (F_2, I_2)) \in \mathcal{P}(a, b)$: assume $f \in F_1$ and $i \in I_1$ s.th. $f \leq i$, then by definition of I_1 there exist d such that $i \multimap d \in F$ but since \multimap is antitone in its first argument, this would mean $f \multimap d \in F$ which contradicts property 5. of F_1 . Dually, assume that there exists $f \in F_2, i \in I_2$ s.th. $f \leq i$, then by definition of F_2 , there exist d such that $f \multimap d \in F$, but then we would also have $i \multimap d \in F$ which contradicts the property 6. of I_2 .

We now check that $\mathcal{P}(a, b)$ has upper bound of chains, let $((F_1^i, I_1^i), (F_2^i, I_2^i)) \in \mathcal{P}(a, b), i \in \omega$ and define

$$F_i^\infty = \bigcup_j F_i^j, \quad I_i^\infty = \bigcup_j I_i^j, \quad i = 1, 2$$

It is not difficult to see that $((F_1^\infty, I_1^\infty), (F_2^\infty, I_2^\infty)) \in \mathcal{P}(a, b)$ by proceeding as in the existence lemma for \otimes . We then apply Zorn's lemma to get a maximal element $((\hat{F}_1, \hat{I}_1), (\hat{F}_2, \hat{I}_2))$ of $\mathcal{P}(a, b)$. It is clear that $a \in \hat{F}_1, b \notin \hat{F}_2$. We need only check that they are prime filters. Assume $c \vee c' \in \hat{F}_1$ but $c, c' \notin \hat{F}_1$, it follows that

$$((\hat{F}_1, \hat{I}_1), (\hat{F}_2, \hat{I}_2)) \not\sqsubseteq ((\hat{F}_1 \cup \{c\}, \hat{I}_1), (\{c \mid \exists d \in \hat{F}_1 \cup \{c\}\}(c \multimap d) \in F), \hat{I}_2))$$

Since $((\hat{F}_1, \hat{I}_1), (\hat{F}_2, \hat{I}_2))$ is maximal, the right-hand side of the inequality must violate either 5. or 6., which in fact amounts to the same thing, namely the existence of $f \in \hat{F}_1, d \in \hat{I}_2$ s.th. $(f \vee c) \multimap d \in F$; that is, $(f \wedge c) \in \hat{I}_1$; that is, $(f \vee c) \leq i$, for some $i \in \hat{I}_1$. A similar argument implies the existence of $f' \in \hat{F}_1, i' \in \hat{I}_1$ such that $f' \wedge c \leq i'$ and a contradiction follows as in the proof for \otimes . To show that \hat{F}_2 is prime is equivalent to showing that \hat{I}_2 is prime, and a proof totally dual to the above proof shows just that. The proof for \otimes is clearly identical. \square

Proof of Proposition 13. Recall Diagram (2):

$$\begin{array}{ccc}
L_{\text{RL}}A & \xrightarrow{L_{\text{RL}}\eta_A} & L_{\text{RL}}\mathcal{UPf}A \\
\downarrow \alpha & \searrow \eta_{LA} & \downarrow \delta_{\text{Pf}A}^{\text{RL}} \\
& & \mathcal{UT}_{\text{RL}}\text{Pf}A \\
& & \downarrow \mathcal{U}(\zeta_A) \\
& & \mathcal{UPf}L_{\text{RL}}A \\
& & \downarrow \mathcal{UPf}\alpha \\
A & \xrightarrow{\eta_A} & \mathcal{UPf}A
\end{array}$$

where ζ_A is the canonical model structure map whose existence we have established in Theorem 3. Recall that by definition of L_{RL} , α is equivalent to a nullary and three binary maps on $\mathcal{U}A$, which we denote as $I, \alpha_{\otimes}, \alpha_{-\otimes}$ and $\alpha_{\otimes-}$ satisfying the distribution laws DL1-DL6.

Similarly, $\mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_{\text{Pf}A}^{\text{RL}}$ is equivalent to a nullary operator and three binary maps on $\mathcal{U}\mathcal{UPf}A = \mathcal{U}A^\sigma$ which we will denote by $\mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ I', \mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_{\otimes}, \mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_{-\otimes}$ and $\mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_{\otimes-}$ and satisfy DL1-DL6. By commutativity of the above diagram these operators are extensions of $I, \alpha_{\otimes}, \alpha_{-\otimes}$ and $\alpha_{\otimes-}$ respectively. We want to show that they are in fact their unique canonical extensions. The treatment of the nullary operator is trivial. For the binary operators, we will show that they are smooth and thus equal to the unique canonical extensions $\alpha_{\otimes}^\sigma, \alpha_{-\otimes}^\sigma$ and $\alpha_{\otimes-}^\sigma$ respectively, by Proposition 6. We start by proving the following claim which readily generalises to the n -ary case.

Claim: Let A, B be DLs and let $f : \mathcal{U}A \rightarrow \mathcal{U}B$. Assume that $\tilde{f} : \mathcal{U}A^\sigma \rightarrow \mathcal{U}B^\sigma$ is an extension of f that (anti-)preserves all binary joins or (anti-)preserves binary meets, then \tilde{f} is (σ, γ) -continuous and thus smooth.

Proof: Since \tilde{f} preserve all binary joins, its restriction f preserves binary joins, and is thus smooth. Moreover, we also know that the canonical extension $f^\sigma = f^\pi$ preserves all non-empty joins. If \tilde{f} preserves all non-empty joins, then in particular it preserves all up-directed ones, and f is consequently $(\gamma^\uparrow, \gamma^\uparrow)$ -continuous. Thus we need only show that it is $(\sigma, \gamma^\downarrow)$ -continuous too. In fact, we show the stronger statement that \tilde{f} is $(\sigma^\downarrow, \sigma^\downarrow)$ -continuous. To see this, note first that since \tilde{f} extends f and preserves non-empty joins we have for every $u \in O(A)$

$$\begin{aligned}
\tilde{f}(u) &= \tilde{f}(\bigvee \{a \in A \mid a \leq u\}) \\
&= \bigvee \{\tilde{f}(a) \mid A \ni a \leq u\} \\
&= \bigvee \{f(a) \mid A \ni a \leq u\} = f^\pi(u) = f^\sigma(u).
\end{aligned}$$

That is, \tilde{f} agrees with f^σ on open elements. Now we use the fact that since f^σ preserves non-empty joins, it is $(\sigma^\downarrow, \sigma^\downarrow)$ -continuous (see Proposition 8); that is, that for any open $v \in O(B)$, there exists $u \in O(A)$ such that $f^\sigma(u) = v$ and thus

$(f^\sigma)^{-1}(\downarrow v) = \downarrow u$. But since \tilde{f} and f^σ coincide on open elements, this also means that $\tilde{f}(u) = v$; that is, that $(\tilde{f})^{-1}(\downarrow v) = \downarrow u$; that is, \tilde{f} is $(\sigma^\downarrow, \sigma^\downarrow)$ -continuous.

The proof for the other preservation properties are very similar. Assume for example that \tilde{f} preserves non-empty meets, it preserve down-directed ones and is thus $(\gamma^\downarrow, \gamma^\downarrow)$ continuous. Moreover $f^\sigma = f^\pi$ preserves non-empty meets and \tilde{f} agrees with f^σ on all closed elements. Since f^σ preserve non-empty meets, it is $(\sigma^\uparrow, \sigma^\uparrow)$ -continuous, and thus so is \tilde{f} by definition of σ^\uparrow and the fact that \tilde{f} and f^σ agree on closed elements. The proof for the anti-preservation properties are similar, with $(\sigma^\downarrow, \sigma^\downarrow)$ and $(\sigma^\uparrow, \sigma^\downarrow)$ -continuity being shown for anti-join preservation and anti-meet preservation respectively.

This having been established, we can now return to our proof. Since $\mathcal{UPf}\alpha$ and $\mathcal{U}\zeta_A$ are inverse images, they preserve any meet and any join, and in particular down-directed meets and up-directed joins. They are therefore (γ, γ) -continuous. Note that this concept makes sense for maps between any DLs, not just canonical extensions. All that needs to be shown now is that $\delta_\otimes, \delta_{-\otimes}$ and $\delta_{\otimes-}$ have one of the preservation properties of proposition 8. We start with δ_\otimes : for any $u_i, v \in \mathcal{UPf}A, i \in I$ we have

$$\begin{aligned} \delta_\otimes(\bigvee_{i \in I} u_i, v) &= \{t \in T_{\text{RL}}\text{Pf}A \mid \exists(x, y) \in \pi_2(t) \exists i \in I, x \in u_i, y \in v\} \\ &= \bigcup_{i \in I} \{t \in T_{\text{RL}}\text{Pf}A \mid \exists(x, y) \in \pi_2(t) \ x \in u_i, y \in v\} \\ &= \bigvee_{i \in I} \delta_\otimes(u_i, v) \end{aligned}$$

and similarly for the second argument. Thus δ_\otimes preserves all non-empty joins, and by Proposition 8 it is therefore (σ^2, γ) -continuous. Since $\mathcal{UPf}\alpha$ and $\mathcal{U}\zeta_A$ are (γ, γ) -continuous, we get that $\mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_\otimes$ is (σ^2, γ) -continuous and thus

$$\alpha_\otimes^\sigma = \mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_\otimes$$

by Proposition 6.

We can similarly show that $\delta_{/}$ (resp. $\delta_{-\otimes}$) preserves all non-empty meets in its first (resp. second) argument and anti-preserves non-empty joins in its second (resp. first) argument. As an illustration,

$$\begin{aligned} \delta_{-\otimes}(\bigvee_{i \in I} u_i, v) &= \{t \in T_{\text{RL}}\text{Pf}A \mid \forall(x, y) \in \pi_3(t) \ x \in \bigvee_{i \in I} u_i \Rightarrow y \in v\} \\ &= \{t \in T_{\text{RL}}\text{Pf}A \mid \forall(x, y) \in \pi_3(t) \ x \notin \bigvee_{i \in I} u_i \text{ or } y \in v\} \\ &= \bigcap_{i \in I} \{t \in T_{\text{RL}}\text{Pf}A \mid \forall(x, y) \in \pi_3(t) \ x \notin u_i \text{ or } y \in v\} \\ &= \bigwedge_{i \in I} \delta_{-\otimes}(u_i, v) \end{aligned}$$

In consequence we also get that

$$\alpha_{-\otimes}^\sigma = \mathcal{UPf}\alpha \circ \mathcal{U}\zeta_A \circ \delta_{-\otimes}$$

and similarly for $\otimes-$, which concludes the proof. \square