


Peripheral nerve injury increases contribution of L-type calcium channels to synaptic transmission in spinal lamina II: Role of $\alpha 2\delta-1$ subunits

Molecular Pain
Volume 14: 1–12
© The Author(s) 2018
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1744806918765806
journals.sagepub.com/home/mpx


Sascha RA Alles^{1,2}, Esperanza Garcia^{1,2},
Sridhar Balasubramanian^{3,4}, Karen Jones^{1,2}, John R Tyson^{1,2},
Twinkle Joy^{3,4}, Terrance P Snutch^{1,2} and Peter A Smith^{3,4}

Abstract

Background: Following peripheral nerve chronic constriction injury, the accumulation of the $\alpha 2\delta-1$ auxiliary subunit of voltage-gated Ca^{2+} channels in primary afferent terminals contributes to the onset of neuropathic pain. Overexpression of $\alpha 2\delta-1$ in *Xenopus* oocytes increases the opening properties of $\text{Ca}_v1.2$ L-type channels and allows Ca^{2+} influx at physiological membrane potentials. We therefore posited that L-type channels play a role in neurotransmitter release in the superficial dorsal horn in the chronic constriction injury model of neuropathic pain.

Results: Whole-cell recording from lamina II neurons from rats, subject to sciatic chronic constriction injury, showed that the L-type Ca^{2+} channel blocker, nitrendipine (2 μM) reduced the frequency of spontaneous excitatory postsynaptic currents. Nitrendipine had little or no effect on spontaneous excitatory postsynaptic current frequency in neurons from sham-operated animals. To determine whether $\alpha 2\delta-1$ is involved in upregulating function of $\text{Ca}_v1.2$ L-type channels, we tested the effect of the $\alpha 2\delta-1$ ligand, gabapentin (100 μM) on currents recorded from HEK293F cells expressing $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-1$ channels and found a significant decrease in peak amplitude with no effect on control $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-3$ expressing cells. In PC-12 cells, gabapentin also significantly reduced the endogenous dihydropyridine-sensitive calcium current. In lamina II, gabapentin reduced spontaneous excitatory postsynaptic current frequency in neurons from animals subject to chronic constriction injury but not in those from sham-operated animals. Intraperitoneal injection of 5 mg/kg nitrendipine increased paw withdrawal threshold in animals subject to chronic constriction injury.

Conclusion: We suggest that L-type channels show an increased contribution to synaptic transmission in lamina II dorsal horn following peripheral nerve injury. The effect of gabapentin on $\text{Ca}_v1.2$ via $\alpha 2\delta-1$ may contribute to its anti-allodynic action.

Keywords

Allodynia, pharmacology, neuropathic pain, dihydropyridine, gabapentin, anti-allodynic, chronic constriction injury, central sensitization, calcium channel blocker, substantia gelatinosa

Date Received: 20 December 2017; revised 16 February 2018; accepted: 20 February 2018

Introduction

The auxiliary alpha-2-delta ($\alpha 2\delta$) subunits of voltage-gated Ca^{2+} channels control the trafficking, localization, and biophysical properties of the pore-forming α_1 -subunits.^{1–3} The $\alpha 2\delta-1$ subunit also serves as the binding partner of the gabapentinoid drugs, pregabalin (PGB), and gabapentin (GBP),^{4,5} which are first-line treatments in the management of neuropathic pain.⁶

In experimental animal models, allodynia and hyperalgesia can be induced by chronic constriction injury

¹Michael Smith Laboratories, University of British Columbia, Vancouver, BC, Canada

²Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver, BC, Canada

³Department of Pharmacology, University of Alberta, Edmonton, AB, Canada

⁴Neuroscience and Mental Health Institute, University of Alberta, Edmonton, AB, Canada

Corresponding Author:

Peter A Smith, Department of Pharmacology, 9.75 Medical Sciences Building, University of Alberta, Edmonton, AB, Canada T6G 2H7. Email pas3@ualberta.ca



(CCI) of the sciatic nerve.^{7,8} This leads to a decrease in paw withdrawal threshold (PWT) to tactile stimulation by von Frey filaments that correlates with increased excitability and altered expression of ion channels in dorsal root ganglia (DRG) neurons^{9–17} and with increased excitatory synaptic transmission in the superficial dorsal horn.^{3,18–21} This latter process of “central sensitization” is initiated in part by upregulation of expression of $\alpha 2\delta-1$ in primary afferent terminals^{22–27} and leads to increased spontaneous neurotransmitter release.²³

Recent experiments in *Xenopus* oocytes showed that overexpression of $\alpha 2\delta-1$ influenced the properties of the voltage sensors of co-expressed $\text{Ca}_v1.2$ L-type channels such that Ca^{2+} influx occurs at physiological membrane potentials.²⁸ Since sensory neurons express dihydropyridine sensitive, L-type channels,^{10,29} we wondered whether the increase in $\alpha 2\delta-1$ expression produced by nerve injury could lead to persistent Ca^{2+} influx in primary afferent terminals. This in turn may contribute to an increase in spontaneous transmitter release, which would be dependent on the activity of L-type Ca^{2+} channels. To test this possibility, we examined the effect of the L-type Ca^{2+} channel blocker, nitrendipine on spontaneous transmitter release in spinal lamina II of rats subject to sciatic CCI. Nitrendipine was chosen by virtue of its especially high affinity for its binding site in rat brain ($K_i = 55$ pM compared to nifedipine $K_i = 384$ pM).³⁰ In humans, nitrendipine is three times more potent than nifedipine in reducing peripheral vascular resistance, and arterial blood pressure, and in increasing leg blood flow.³¹

To determine whether augmentation of L-type channel function is contingent on $\alpha 2\delta-1$, we applied GBP to HEK293F cells expressing $\text{Ca}_v1.2/\alpha 2\delta-1$ and observed a reduction in peak amplitude of calcium current. GBP binds with high affinity to the $\alpha 2\delta-1$ and $\alpha 2\delta-2$ subunits but not to $\alpha 2\delta-3$ subunits³² and failed to affect calcium current in $\text{Ca}_v1.2/\alpha 2\delta-3$ expressing cells. We further confirmed these results on native L-type channels in PC-12 cells and observed that GBP reduced the peak amplitude of the dihydropyridine-sensitive current.

We also found that intraperitoneal injection of 5 mg/kg nitrendipine increased PWT in animals subject to CCI. Taken together, our findings suggest that L-type $\text{Ca}_v 1.2$ channels show an increased contribution to synaptic transmission in lamina II dorsal horn following peripheral nerve injury and that an effect of GBP on $\text{Ca}_v1.2$ via $\alpha 2\delta-1$ may contribute to its anti-allodynic action.

Methods

Nerve injury surgery and assessment of mechanical allodynia

All procedures were approved by the University of Alberta Animal Welfare Committee in accordance with

Canadian Council on Animal Care (CCAC) guidelines. Male Sprague–Dawley rats (19–23-day old) were subject to CCI of the sciatic nerve using polyethylene cuffs⁸ as described previously.¹⁸ Surgery was performed under isoflurane anesthesia (5% induction and 2% maintenance). For sham surgery, the sciatic nerve was exposed but not manipulated. Ten to 14 days after surgery, animals were assessed for the presence of allodynia using von Frey filaments as previously described.¹⁸ Animals exhibiting a PWT <6 g were assumed to be expressing allodynia, a sign of neuropathic pain.

Spinal cord slice preparation and electrophysiology

Rats were euthanized with an intraperitoneal injection of urethane (1.5 g/kg). Following cessation of respiration and loss of nociceptive reflexes (paw pinch with forceps), the spinal cord was removed by laminectomy, and 300 μm transverse spinal cord slices prepared on a vibratome as previously described.^{18,21} Whole-cell patch-clamp recordings were made under infrared differential-interference optics in neurons from slices from CCI or sham-operated animals. Recordings were made from lamina II (*substantia gelatinosa*) neurons ipsilateral to the sciatic injury.²¹ *Substantia gelatinosa* was identified by its translucent appearance in the slice. For recording, slices were superfused at room temperature ($\sim 22^\circ\text{C}$) with 95% O_2 –5% CO_2 saturated aCSF which contained (in mM): 127 NaCl, 2.5 KCl, 1.2 NaH_2PO_4 , 26 NaHCO_3 , 1.3 MgSO_4 , 2.5 CaCl_2 , 25 D-glucose, pH 7.4. Recording pipettes had resistances of 5–10 M Ω when filled with an internal solution containing (in mM) 130 potassium gluconate, 1 MgCl_2 , 2 CaCl_2 , 10 HEPES, 10 EGTA, 4 Mg-ATP, 0.3 Na-GTP, pH 7.2, 290–300 mOsm. Recordings were made using an NPI SEC-05LX amplifier in discontinuous, single-electrode voltage-clamp or current clamp mode. Data were only collected from neurons that exhibited clear overshooting action potentials >60 mV in amplitude. Spontaneous excitatory postsynaptic currents (sEPSC) were recorded at -70 mV. Mini Analysis Program (Synaptosoft, Decatur, GA, USA) was used to distinguish sEPSC from baseline noise. All detected events were then re-examined and visually accepted or rejected based on visual examination. Acceptable events had a sharp onset and exponential offset, a total duration of <50 ms, and an amplitude at least double the baseline noise.

EPSCs were evoked at 0.05 Hz using a teflon-coated nichrome stimulating electrode (7620, A-M Systems, Carlsborg, WA, USA) on the dorsal root or near the dorsal root entry zone to activate primary afferent fibers. The voltage-gated ion channel blocker QX-314 (5 mM) was included in the internal solution to prevent action potential discharge while recording evoked EPSPs

(eEPSCs) in the presence of 10 μM bicuculline and 1 μM strychnine. Stimulus intensity for eEPSCs was between 2 and 40 V and the stimulus duration was 100 or 400 μs . Measurements were made from six averaged responses.

Drugs and chemicals

All chemicals used in slice preparation and whole-cell electrophysiology were from Sigma (St. Louis, MO, USA), except GBP, which was from TCI America (Portland OR, USA) and nitrendipine, which was from Tocris Bioscience Minneapolis (MN, USA). For IP injections, 18 mg of nitrendipine was dissolved in 1 ml of dimethyl sulfoxide (DMSO). This was further diluted by the addition of 5 ml of deionized water and the solution made up to 20 ml by the addition of 14 ml saline (0.9% NaCl). This yielded a suspension of 0.9 mg/ml of nitrendipine. Animals received IP injections of 5 mg/kg nitrendipine. Control vehicle solution contained a mixture of DMSO, water, and saline in the ratio 1:5:14.

Cell culture, transfection, and electrophysiology

PC12 cells (ATCC CRL-1721) were grown on Bovine Collagen Type I-coated flasks in F-12K (Invitrogen 21127022) medium supplemented with 2.5% fetal bovine serum (FBS, GIBCO 12483-020) and 15% horse serum (Invitrogen 26050-088) at 37°C in a humidified atmosphere with 5% CO₂-95% air. Whole-cell electrophysiological recordings were performed on cells plated on Poly-L-Lysine-coated glass coverslips 48–54 h after seeding.

HEK293F (Invitrogen 11625-019) cells were maintained in Dulbecco's Modified Eagle's Medium (Invitrogen 12800-017) supplemented with 10% FBS and 1% non-essential amino acids (Invitrogen 11140). Cells were plated on Poly-L-Lysine-coated glass coverslips and transfected after 24 h using TurboFect transfection reagent (Fisher Cat R0531), following the procedure recommended by the manufacturer. Plasmids encoding rat Ca_v1.2 α_1 subunit and ancillary subunits β_4 , $\alpha_2\delta_1$, or $\alpha_2\delta_3$ were combined at equimolar ratio and co-transfected with enhanced green fluorescent protein (EGFP) as a reporter. A total of 1.1 μg or 1.6 μg cDNA per 35 mm dish was used for the transfection mix with $\alpha_2\delta_1$ or $\alpha_2\delta_3$, respectively. Current recordings were performed 18–24 h after transfection.

Whole-cell patch-clamp recordings from PC12 or HEK cells were obtained using a 200B Axopatch amplifier and data were acquired with a Digidata 1332A system (Molecular Devices, CA) controlled by pCLAMP 9 software. Data were low-pass filtered at 2 kHz using the Bessel filter of the amplifier and sampled at 10 kHz. Experiments were performed at room temperature (20–22°C). Borosilicate glass (Sutter BF150-86-10)

patch pipettes were made with a horizontal puller P97 (Sutter Instrument Co) and polished with a microforge (Narishige MF-900). Electrodes had a resistance of 2.8–3.5 M Ω when filled with the following internal solution (mM): 105 Cesium methanesulphonate (CH₃CsO₃S), 25 TEA-Cl, 10 HEPES, 11 EGTA, 1 CaCl₂, 5 Mg-ATP, 3 Tris₂-Phosphocreatine, 3 Na₂-Phosphocreatine, and 0.4 Tris-GTP (pH 7.2; 290 mOsm). The composition of the bathing external solution for recordings from cells expressing channels associated with $\alpha_2\delta_1$ ancillary subunit was (in mM) 88 CsCl, 40 TEA-Cl, 10 HEPES, 10 Glucose, 1 MgCl₂, and 5 BaCl₂. Osmolality was adjusted to 300 mOsm and pH at 7.4. Native macroscopic currents from PC12 cells and from recombinant Ca_v1.2 co-transfected with $\alpha_2\delta_3$ ancillary subunit were recorded with an extracellular solution containing (in mM) 95 CsCl, 30 TEA-Cl, 10 HEPES, 10 Glucose, 1 MgCl₂, and 20 BaCl₂ (300 mOsm; pH 7.4). S(-)-Bay K8644 (Tocris 1546) stock solution was prepared in DMSO and stored in aliquots at -20°C; 100 nM working solutions were made fresh in external recording solution (final concentration of vehicle was 0.025%). ω -Conotoxin GVIA and ω -Agatoxin VIA (Alomone Labs) were reconstituted in sterile PBS and stored at -20°C; appropriate volume was added directly into external solution to a final concentration of 1 μM and 200 nM, respectively. GBP (TCI Chemicals, Portland, OH, USA) 100 mM stock solution was prepared in nanopure H₂O and diluted into external solution to a final concentration of 100 μM . Perfusion rate was 1.2 mL/min. No corrections were made for liquid junction potentials.

To examine the effect of GBP on recombinant channels in HEK cells, barium currents were evoked by 120 ms repetitive step depolarizations from -100 mV to 0 mV ($\alpha_2\delta_1$) or from -100 mV to +10 mV ($\alpha_2\delta_3$) at 0.1 Hz. Current amplitudes were divided by the average value obtained during the perfusion with external control solution and the fraction of remaining (unblocked) current (I/I_{Control}) was plotted against time. The steady state value of unblocked current after 3 min of GBP perfusion is reported as fraction of inhibition ($I_{\text{GBP}}/I_{\text{Control}}$). Current-voltage relationship for channels expressing $\alpha_2\delta_1$ subunit was studied with 180 ms pulses from -60 mV to +30 mV at 5 mV increments ($V_h = -100$ mV). Fitting of peak current values obtained with increasing depolarizations was performed using the modified Boltzmann equation:

$$I_{\text{Ba}}/I_{\text{Ba,Max}} = [(V_m - E_r)] / (1 + \exp[(V_m - V_{50})/k])$$

where I_{Ba} is the peak current at test potential V_m , $I_{\text{Ba,Max}}$ is the maximum peak current observed, E_r is the

apparent reversal potential, V_{50} is the midpoint voltage for activation, and k is the slope factor.

The effect of GBP on neuronal L-type currents was studied by examining native barium currents from undifferentiated PC12 cells, evoked by 200 ms voltage ramps from -100 to $+100$ mV applied every 10 s. Current amplitude at the peak of the resultant IV curve was measured in the presence of ω -Conotoxin GVIA and ω -Agatoxin VIA as control value, and compared to the values obtained after the consecutive application of S(-)-Bay K8644 and GBP. Nitrendipine ($1 \mu\text{M}$) was applied to demonstrate that GBP affects the dihydropyridine-sensitive current component.

mRNA expression analysis using quantitative real-time PCR

PC12 cell pellets were homogenized in the presence of TRI-Reagent (Ambion) and total RNA isolated using the MagMAXTM-96 for Microarrays preparation kit (Life Technologies, AM1839). Total cDNA was synthesized from total RNA using a high capacity cDNA reverse transcription kit (Applied Biosystems, 4368814). Gene-specific qPCR reactions utilized the KAPA Probe Fast Universal qPCR Kit (KK4702) and TaqManTM probes from Life Technologies and Integrated DNA Technologies (IDT). Quantitative PCR reactions were performed on a Bio-Rad CFX Real-Time Systems using cycling conditions of 95°C for 3 min then 95°C for 15 s followed by 60°C for 45 s for 40 cycles. Specific TaqManTM probes used in this study: rCav1.1(Rn01490941_m1), rCav1.2 (Rn00709287_m1), rCav1.3(Rn00692157_m1), rCav2.1 (Rn00563825_m1), rCav2.2(Rn00595911_m1), rCav2.3 (Rn00494444_m1), rA2D1(Rn01442581_m1), rA2D2 (Rn00457825_m1), and rA2D3(Rn00598241_m1) from Life Technologies, and Glyceraldehyde 3-phosphate dehydrogenase (rGAPDH) (Rn.PT.56a.35727291) from IDT.

Statistical analysis

The Kolmogorov–Smirnov two-sample test (K-S test) was used to compare distributions of sEPSC amplitudes and inter-event intervals.³³ All other data were compared using the two-tailed t-test or by one-way analysis of variance (ANOVA) where appropriate. Significance was attributed if $p < 0.05$. Mini Analysis program (Synaptosoft, Decatur, GA, USA) was used to generate cumulative probability plots. All other graphing and statistical analyses were performed using Origin (v8.6 or 2018, Origin Lab., Northampton, MA, USA).

Results

Nitrendipine reduces the frequency of lamina II sEPSCs in animals subject to CCI

As already mentioned, peripheral nerve injury increases the excitability of the superficial dorsal horn.^{3,18–21} To assess the role of presynaptic L-type Ca^{2+} channels in this process, we examined the effect of nitrendipine ($2 \mu\text{M}$) on spontaneous excitatory postsynaptic currents (sEPSCs) in lamina II neurons of rats subject to sciatic CCI. Nitrendipine was highly effective in reducing frequency (increasing inter-event interval, IEI) of sEPSCs in rats subject to CCI (Figure 1(b); $p < 0.001$; K-S test: Figure 1(d); $p < 0.001$; t-test). By contrast in sham-operated rats, cumulative probability plots show that nitrendipine was without effect on IEI (Figure 1(a); $p > 0.15$; K-S test) of sEPSCs in lamina II neurons. There was also no significant effect of nitrendipine on average sEPSC IEI (Figure 1(c); $p > 0.05$; t-test).

Nitrendipine slightly decreased sEPSC amplitude in sham-operated rats according to a cumulative probability plot (Figure 1(e); $p < 0.05$; K-S test), but the averaged amplitude was unaffected (Figure 1(g); $p > 0.05$; t-test). Somewhat unexpectedly, nitrendipine produced a slight increase in sEPSC amplitude in neurons from animals subject to CCI and this was significant according to K-S statistics (Figure 1(f); $p < 0.02$) and from consideration of average event amplitude (Figure 1(h); $p < 0.05$; t-test).

Nitrendipine reduces the amplitude of lamina II eEPSCs in animals subject to CCI

In neurons from sham-operated animals, superfusion of nitrendipine ($2 \mu\text{M}$) caused a small reduction in the amplitude of evoked EPSCs (eEPSCs) by an average of $14.85 \pm 5.53\%$ ($n = 13$, $p < 0.05$, paired t-test). This reduction is in agreement with earlier reports on effect of L-type channel blocker on evoked EPSCs in the superficial dorsal horn neurons.^{34,35} By contrast, in rats subject to CCI, nitrendipine caused a much larger reduction of $40.23 \pm 6.77\%$ ($n = 12$, $p < 0.005$, paired t-test) in the amplitude of eEPSCs. Sample traces are illustrated in Figure 1(i) and (j). Figure 1(k) illustrates the time course of the effect of nitrendipine on eEPSCs in CCI and sham-operated rats. The time course of action of nitrendipine varied from cell to cell. In general, the effect took about 10–15 min to develop and the washout time was 30 min or longer. Although we cannot rule out the possibility that some of the recorded events were polysynaptic, the eEPSCs observed under the conditions of our experiments were of short duration and monophasic and thus may have been predominantly monosynaptic.




Figure 1. Effects of nitrendipine on sEPSC and eEPSCs recorded from lamina II neurons of CCI and sham-operated rats. Cumulative probability plots for effect of nitrendipine on inter-event interval (IEI) of sEPSCs from sham-operated animals (a) and those subject to CCI (b). (c and d) Same data replotted as mean IEI. Cumulative probability plots for effect of nitrendipine on amplitude of sEPSCs from sham-operated animals (e) and those subject to CCI (f). (g and h) same data replotted as mean IEI. For shams, IEI data for 1629 events and nitrendipine data for 1398 events in 24 neurons. For CCI, IEI data for 1853 events and nitrendipine data for 1379 events in 27 neurons. For shams, amplitude data for 1561 events and nitrendipine data for 1415 events in 24 neurons. For CCI, amplitude data for 1898 events and nitrendipine data for 1389 events in 27 neurons. Note that p values were derived from K-S test for cumulative probability graphs and paired t-test for the bar graphs. ** $p < 0.001$ and * $p < 0.05$ compared to pre-drug amplitude (paired t-test). In each case, recordings were obtained from only one neuron per slice. Sample recordings of eEPSCs following DRZ stimulation from sham-operated animals (i) and those subject to CCI (j), before, during, and after superfusion of 2 μ M nitrendipine. (k) Time course of effect of nitrendipine on eEPSC amplitude in 13 neurons from sham-operated animals and 12 neurons from animals subject to CCI. Error bars = SEM. ** $p < 0.001$ and * $p < 0.05$ compared to pre-drug amplitude (paired t-test). CCI: chronic constriction injury; eEPSC: evoked excitatory postsynaptic currents.

Effects of nitrendipine on mechanical allodynia

Following CCI, animals expressing a PWT of <6 g were defined as allodynic. PWT were measured 30 min after an IP injection of 5 mg/kg nitrendipine. Pharmacokinetic studies of nitrendipine in rats are scarce but literature suggests that plasma concentration peaks at 1.2 h following intravenous, oral, or intraduodenal administration.³⁶ The changes in PWT in response to a series of nitrendipine injections are shown in Figure 2(a). Injections given on day 1 increased PWT strongly in some animals and weakly in others. On day 2, two injections of 5 mg/kg

were administered 60 min apart and PWT addressed 30 min after the second injection. On day 3, animals received a single injection of nitrendipine and all five animals responded with an increase in PWT. The progressive improvement in effect may reflect drug accumulation following repeated injections. The elimination half time in rats is 57 h.³⁶ Data for each animal are shown in Figure 2(b). Vehicle injections failed to affect PWT of the ipsilateral limb (Figure 2(a)). The PWT for the paw contralateral to CCI was >15 g and therefore any non-specific effect of nitrendipine on could not be determined.




Figure 2. Effects of nitrendipine on PWT in animals subject to CCI. (a) Averaged data from five animals, arrows labeled N indicate times of IP injections of 5 mg/kg nitrendipine. Dashed line with gray symbols illustrates minimal effect of vehicle injections (two animals) (b) Data from each animal at the 36 h time point to illustrate the variation in nitrendipine effectiveness. PWT: paw withdrawal threshold.

Effect of $\alpha 2\delta-1$ on properties of exogenous L-type channel complexes in HEK cells

The above results suggest that L-type Ca^{2+} channels contribute to neurotransmitter release in nerve injured animals. Since Savalli et al. (2016) observed a potentiation of specifically $\text{Ca}_v1.2$ L-type channel current following $\alpha 2\delta-1$ overexpression in *Xenopus* oocytes, we tested whether GBP (100 μM) has a direct effect on $\text{Ca}_v1.2$ -mediated currents in the presence of $\alpha 2\delta-1$. We used HEK293 cells expressing $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-1$ and examined peak amplitude of calcium current before and after acute application of GBP (100 μM). Since GBP does not bind to the $\alpha 2\delta-3$ subunit,³¹ we performed experiments on HEK293 cells expressing $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-3$ as a control. GBP (100 μM) produced a significant reduction (Figure 3(a, i), $p < 0.05$, ANOVA) of $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-1$ peak calcium currents with no significant effect on the peak amplitude on $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-3$ currents (Figure 3(a, ii), $p > 0.05$, ANOVA). This relative effect is summarized in Figure 3(a, iii) showing the effect of GBP after a 3 min application. The average inhibition of peak calcium current by GBP in $\text{Ca}_v1.2/\beta 4/\alpha 2\delta-1$ cells was $\sim 25\%$ ($p < 0.05$, ANOVA) as shown by current traces in Figure 3(b, ii). There were no significant effects of GBP on voltage dependence of activation ($p > 0.05$, ANOVA) as shown in Figure 3(b, i) by comparing the values of fitted parameters corresponding to the continuous lines obtained by the Boltzmann function before ($V_{50} = -12.73 \pm 0.37$ mV, $k = -5.23 \pm 0.14$) and after ($V_{50} = -14.53 \pm 0.53$ mV, $k = -5.55 \pm 0.10$) GBP application.

Gabapentin reduces the native dihydropyridine-sensitive current in PC-12 cells

To further confirm that GBP inhibits L-type channels, we examined native L-type Ca^{2+} currents expressed in undifferentiated pheochromocytoma cells (PC-12 cells). PC-12 cells have long been studied as a model system

of neuronal physiology and biochemistry and the effects of pharmacological agents due to their similarity to sympathetic neurons upon differentiation.³⁷ Undifferentiated PC-12 cells mainly express N- and L-type Ca^{2+} channels although studies have also reported the presence of an ω -Agatoxin-VIA-sensitive (P/Q-type) component.³⁸ In order to isolate the L-type current in undifferentiated PC-12 cells, it was necessary to remove both N-type and P/Q-type currents with ω -Conotoxin GVIA (1 μM) and ω -Agatoxin VIA (200 nM; Figure 4 (b)). Further, in order to obtain a large enough L-type current from which to establish any effects of GBP, we also applied the L-type agonist S-(S)-BayK8644 (100 nM) (Figure 4(b)). The resulting whole current represents a dihydropyridine-sensitive inward current as it was blocked by nitrendipine (1 μM) (Figure 4(c)). GBP (100 μM) significantly inhibited the peak amplitude of this current by approximately 40% (Figure 4(c), $p < 0.05$, ANOVA).

In order to confirm the expression of selected Ca^{2+} channel complex proteins, we performed qRT-PCR on RNA isolated from PC-12 cells. Figure 4(c) confirms the endogenous expression of the $\text{Cav}1.2$ L-type channel together with the $\text{Cav}2.2$ (N-type) and $\text{Cav}2.1$ (P/Q-type) channels. Figure 4(d) demonstrates that PC-12 cells also express significant amounts of the $\alpha 2\delta-1$, $\alpha 2\delta-2$, and $\alpha 2\delta-3$ proteins in support of the data from HEK293F cells showing that GBP requires the $\alpha 2\delta-1$ subunit in order to elicit an inhibitory effect on L-type currents.

Gabapentin reduces sEPSC frequency in CCI but not sham-operated rats

To test whether the effectiveness of nitrendipine after CCI corresponds to increased expression of $\alpha 2\delta-1$ in primary afferent neurons as reported in Luo et al.,^{24,25} we next examined the effect of the $\alpha 2\delta-1$ ligand GBP at a clinically relevant concentration (100 μM).³⁹ GBP did not alter IEI in neurons from sham-operated animals




Figure 3. Gabapentin block of L-type $\text{Ca}_v1.2$ channel currents expressed in HEK293F cells. (a) Time course of inhibitory effect of $100 \mu\text{M}$ GBP on Ba^{2+} currents recorded from HEK cells expressing $\text{Ca}_v1.2$ channels in combination with either $\alpha_2\delta-1$ (i) or $\alpha_2\delta-3$ (ii) auxiliary subunits. The averaged fraction of peak current amplitude is plotted against time. Cells were perfused with GBP during the periods indicated by horizontal bars. Inserts correspond to representative macroscopic currents obtained before (black) and after (red) the application of GBP. (iii) The fraction of inhibition after 3 min of drug application is shown as a bar graph. Number of cells examined are indicated in brackets; the asterisk indicates a statistical significance $p < 0.05$. (b) Normalized current-voltage curves (i) obtained for the recombinant $\text{Ca}_v1.2$ channel in the presence (red symbols) and in the absence (black symbols) of $100 \mu\text{M}$ GBP. The values of fitted parameters corresponding to the continuous lines obtained by Boltzmann function fits are shown. Application of GBP produced an average current inhibition of $\sim 25\%$ with no significant changes in the voltage dependence parameters. Data are plotted as mean \pm SEM ($n = 5$ for each data point). (ii) Typical traces from a representative cell show the effect of GBP on the I-V relationship with no apparent modification of current kinetics.

(Figure 5(a) and (c); $p > 0.05$) but profoundly decreased it in neurons from animals subject to CCI (Figure 5(b) and (d); $p < 0.0001$). GBP produced a small but significant reduction in the sEPSC amplitude in neurons from sham-operated animals (Figure 5(e) and (g); $p < 0.001$), but there was no significant effect on neurons from animals subject to CCI (Figure 5(f) and (h)). A comparison

of the effects of nitrendipine and GBP on sEPSCs in CCI and sham-operated rats is shown in Table 1.

Discussion

We tested the hypothesis that the increased expression of $\alpha_2\delta-1$ at nerve terminals⁵ after nerve injury⁵ alters the




Figure 4. Gabapentin inhibits a dihydropyridine-sensitive inward current in undifferentiated PC12 cells. a. Current-voltage relation of barium currents were examined in the presence of 1 mM ω -Conotoxin GVIA and 200nM ω -Agatoxin VIA to eliminate the contribution of native non-L-type channels. Currents were elicited by voltage ramps from -100 to $+100$ mV at 1mV/ms, and cells were exposed to 100nM of the agonist (S)-(-)-BayK 8644 prior to the application of 100 μ M GBP. b. The bar graph shows the average of peak amplitude values ($n = 5$) obtained in cells perfused with external recording solution (Ba^{2+} 20mM), compared to that obtained after the consecutive addition of peptide toxins, BayK 8644 and GBP. Data are plotted as Mean \pm S.E.M. The asterisk indicates the change in current amplitude after perfusion with GBP is statistically significant at $p < 0.05$. c. Endogenous expression of ancillary $\alpha_2\delta$ subunits (Left) and HVA α_1 subunits (Right) in PC12 cells was quantified by RT-PCR. Expression levels were calculated relative to the “housekeeping gene” GAPDH (Glyceraldehyde 3-phosphate dehydrogenase). Mean values \pm S.D. of three samples.

properties of L-type $Ca_v1.2$ channels such that influx of Ca^{2+} occurs at physiological membrane potential.²⁷ This notion is supported by the observation that both GBP and nitrendipine produced a substantial increase in the IEI of sEPSCs in neurons from nerve injured animals (Figure 1(b) and (d) and Figure 5(b) and (d), Table 1), whereas in sham-operated animals, both nitrendipine

and GBP were without effect (Figure 1(a) and (c) and Figure 5(a) and (c)). This is also supported by a recent report that intrathecal administration of the L-type blocker nifedipine reversed tactile allodynia in mice engineered to overexpress $\alpha_2\delta-1$.⁴⁰ It is also consistent with our observation that nitrendipine exerted an antiallo-dynamic effect in nerve-injured animals (Figure 2).




Figure 5. Effects of GBP on sEPSCs recorded from lamina II neurons of CCI and sham-operated rats. Cumulative probability plots for effect of 100 μ M GBP on (a) inter-event interval (IEI) of sEPSCs in sham-operated animal and (b) in animals subject to CCI. (c and d) same data replotted as bar graphs. Cumulative probability plots for effect of 100 μ M GBP on (e) amplitude of sEPSCs in sham-operated animal and (f) in animals subject to CCI. (g and h) same data replotted as bar graphs. For sham-operated rats, control data for 515 events and GBP data for 625 events in 7 neurons from 5 rats. For CCI, control data for 886 events and GBP data for 721 events in 9 neurons from 8 rats. *p*-values were derived from K-S test for cumulative probability graphs and paired *t*-test for the bar graphs. ***p* < 0.001 and **p* < 0.05 compared to pre-drug amplitude (paired *t*-test). In each case, recordings were obtained from only one neuron per slice. sEPSC: spontaneous excitatory postsynaptic currents; CCI: chronic constriction injury.

Table 1. Comparison of effects of nitrendipine and gabapentin on sEPSCs recorded from CCI and sham-operated rats.

	NTR (2 μ M)	GBP (100 μ M)
CCI sEPSC frequency	↓ 62.23%	↓ 52.8%
CCI sEPSC amplitude	↑ 8.38%	Not significant
Sham sEPSC frequency	Not significant	Not significant
Sham sEPSC amplitude	Not significant	↓ 13.4%

Note: L-type channel blockade using 2 μ M NTR produces similar effects on sEPSCs as a clinically relevant concentration (100 μ M) of α 2 δ -1 ligand GBP suggesting that GBP may produce effects on sEPSCs by binding to α 2 δ -1 associated with L-type channels. The effects of GBP are exclusively on frequency in CCI rats suggesting that GBP acts presynaptically. GBP: gabapentin; CCI: chronic constriction injury; sEPSCs: spontaneous excitatory postsynaptic currents; NTR: nitrendipine.

Following CCI, nitrendipine acts predominantly presynaptically in lamina II as effects on sEPSC amplitude were small (Figures 1(f) and (h)), whereas those on sEPSC frequency were highly significant (Figure 1(b) and (d)). The observed slight increase in sEPSC amplitude in animals subject to CCI was unexpected (Figure 1(f) and (h)). One possibility is that there exists a weak blockade of postsynaptic dendritic K^+ channels by nitrendipine,^{41,42} and that a resulting increase in space constant may increase the amplitude of events recorded in the soma. It remains to be determined why this effect was observed in animals subject to CCI

(Figure 1(f) and (h)) but not from sham-operated animals (Figure 1(e) and (g)).

Given the ubiquitous distribution of L-type Ca^{2+} channels in the nervous system,⁴³ it is possible that some of the anti-allodynic actions of nitrendipine are exerted at thalamic or cortical levels or perhaps by actions on descending pain control mechanisms. We think this rather unlikely as no overt CNS effects are observed in patients receiving nitrendipine for the management of cardiovascular disorders.

Like nitrendipine, the effects of GBP were predominantly presynaptic as there was a large increase in IEI in neurons from animals subject to CCI (Figure 5(b) and (d)) but not in neurons from sham-operated animals (Figure 5(a) and (c)). This agrees with literature showing that α 2 δ -1 upregulation occurs in primary afferent neurons after CCI but not postsynaptically in the dorsal horn.^{24,25} There is no obvious explanation for the observation that GBP reduces sEPSC amplitude in neurons from sham-operated animals (Figure 5(e) and (g)) but not in those subject to CCI (Figure 5(f) and (h)).

Generally, N-type Ca^{2+} channels play a predominant role in neurotransmitter release from primary afferent terminals.^{34,35,44} Thus, our observation that nitrendipine increases the IEI of sEPSC and the amplitude of eEPSCs after CCI is intriguing as it suggests that L-type channels contribute to this process when nerves are injured and α 2 δ -1 is upregulated. This may be explicable in terms of

the findings of Savalli et al. (2016) who showed that $\alpha 2\delta-1$ upregulation alters voltage sensitivity of $\text{Ca}_v1.2$ channels. This may indicate that they are more likely to be activated by presynaptic action potentials and is supported by the classical observation that in DRG neurons N-type Ca^{2+} channels begin to activate around -20 mV and with depolarization to approximately -10 mV being required to open L-type channels.²⁸ Such activation would be more likely to participate in the release process as $\alpha 2\delta-1$ indiscriminately recruits voltage gated Ca^{2+} channels to release sites.⁴⁵

Using both the HEK293F expression system and PC-12 cells, we have shown that acute application of a clinically relevant concentration of GBP produces a robust inhibition of L-type currents. In HEK293F cells, this depression is dependent on the co-expression of $\alpha 2\delta-1$ (Figure 3(a), (i)) but is not observed when $\alpha 2\delta-3$ is expressed (Figure 3(a), (ii)) as these subunits do not bind GBP.³¹ In rats subject to peripheral nerve injury, we have noted that GBP reduces the amplitude and frequency of sEPSCs recorded in excitatory neurons⁴⁶ which receive input from C-fibers.⁴⁷ GBP is less effective at synapses onto inhibitory neurons which receive input from both non-nociceptive $\text{A}\beta$ -fibers⁴⁸ and unmyelinated fibers.⁴⁹ If the effects of GBP involve an action on $\text{Ca}_v1.2$ L-type channels as a result of inhibition of $\alpha 2\delta-1$ function, its lack of effect on input to inhibitory neurons and preferential effect on excitatory neurons may reflect the relative paucity of L-type Ca^{2+} channels on large, non-nociceptive DRG neurons compared to small, nociceptive DRG neurons.^{10,50} It should also be noted that long-term application of GBP is more effective in suppressing high-voltage activated I_{Ca} in small DRG neurons compared to large neurons,³⁹ suggesting that $\alpha 2\delta-1$ may be preferentially expressed in nociceptive neurons which, as mentioned above, predominate the excitatory synaptic input to excitatory neurons.

While cardiovascular actions of dihydropyridines may limit their use in the treatment of neuropathic pain, their use as adjunct agents has been suggested.⁵¹ There is also increasing interest in the possible use of broad spectrum dihydropyridine-related Ca^{2+} channel blockers such as M4 that inhibits $\text{Ca}_v1.2$ (L-type), $\text{Ca}_v2.2$ (N-type), and the $\text{Ca}_v3.2$ and 3.3 T-type channels.⁵²

Authors' contributions

SRAA designed aspects of the study, carried out experiments with GBP, and wrote first draft of manuscript. SRAA and EG designed and carried out experiments on PC12 and HEK cells and contributed to writing final manuscript. KJ performed transfections and cell culture of HEK and PC-12 cells. JRT performed qPCR experiments and analysis. SB designed, carried out, and interpreted experiments with nitrendipine in spinal cord. TJ designed, executed, and interpreted behavioral experiments with nitrendipine. TPS provided CIHR funding

and laboratory facilities, supervised the work of SRAA, EG, JRT, and KJ, and contributed to experimental design. PAS provided NSERC funding and laboratory facilities, supervised the work of SRAA, SB, and TJ, contributed to experimental design, edited and wrote final manuscript. All authors reviewed drafts of manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work is supported by grants from the Natural Sciences and Engineering Research Council (NSERC-Canada; PAS) and the Canadian Institutes of Health Research (CIHR, T.P.S.; #10677). TPS is also supported by the Canada Research Chair in Biotechnology and Genomics-Neurobiology.

References

1. Dolphin AC. Calcium channel auxiliary alpha2delta and beta subunits: trafficking and one step beyond. *Nat Rev Neurosci* 2012; 13: 542–555.
2. Dolphin AC. Voltage-gated calcium channels and their auxiliary subunits: physiology and pathophysiology and pharmacology. *J Physiol* 2016; 594: 5369–5390.
3. Alles SRA and Smith PA. The etiology and pharmacology of neuropathic pain. *Pharmacol Rev* 2018; 70: 315–347.
4. Gee NS, Brown JP, Dissanayake VU, Offord J, Thurlow R and Woodruff GN. The novel anticonvulsant drug, gabapentin (Neurontin), binds to the alpha2delta subunit of a calcium channel. *J Biol Chem* 1996; 271: 5768–5776.
5. Bauer CS, Nieto-Rostro M, Rahman W, Tran-Van-Minh A, Ferron L, Douglas L, Kadurin I, Sri Ranjan Y, Fernandez-Alacid L, Millar NS, Dickenson AH, Lujan R and Dolphin AC. The increased trafficking of the calcium channel subunit alpha2delta-1 to presynaptic terminals in neuropathic pain is inhibited by the alpha2delta ligand pregabalin. *J Neurosci* 2009; 29: 4076–4088.
6. Finnerup NB, Sindrup SH and Jensen TS. The evidence for pharmacological treatment of neuropathic pain. *Pain* 2010; 150: 573–581.
7. Kim KJ, Yoon YW and Chung JM. Comparison of three rodent models of neuropathic pain. *Exp Brain Res* 1997; 113: 200–206.
8. Mosconi T and Kruger L. Fixed-diameter polyethylene cuffs applied to the rat sciatic nerve induce a painful neuropathy: ultrastructural morphometric analysis of axonal alterations. *Pain* 1996; 64: 37–57.
9. Abdulla FA and Smith PA. Axotomy and autotomy-induced changes in the excitability of rat dorsal root ganglion neurons. *J Neurophysiol* 2001; 85: 630–643.
10. Abdulla FA and Smith PA. Axotomy- and autotomy-induced changes in Ca^{2+} and K^{+} channel currents of rat

- dorsal root ganglion neurons. *J Neurophysiol* 2001; 85: 644–658.
11. Abdulla FA and Smith PA. Changes in Na⁺ channel currents of rat dorsal root ganglion neurons following axotomy and axotomy-induced autotomy. *J Neurophysiol* 2002; 88: 2518–2529.
 12. Cummins TR and Waxman SG. Downregulation of tetrodotoxin resistant sodium currents and upregulation of rapidly repriming tetrodotoxin sensitive sodium current in small sensory neurons after nerve injury. *J Neurosci* 1997; 17: 3503–3514.
 13. Everill B, Cummins TR, Waxman SG and Kocsis JD. Sodium currents of large (Aβ-type) adult cutaneous afferent dorsal root ganglion neurons display rapid recovery from inactivation before and after axotomy. *Neuroscience* 2001; 106: 161–169.
 14. Sleeper AA, Cummins TR, Dib-Hajj SD, Hormuzdiar W, Tyrrell L, Waxman SG and Black JA. Changes in expression of two tetrodotoxin-resistant sodium channels and their currents in dorsal root ganglion neurons after sciatic nerve injury but not rhizotomy. *J Neurosci* 2000; 20: 7279–7289.
 15. Zamponi GW, Lewis RJ, Todorovic SM, Arneric SP and Snutch TP. Role of voltage-gated calcium channels in ascending pain pathways. *Brain Res Rev* 2009; 60: 84–89.
 16. Waxman SG and Zamponi GW. Regulating excitability of peripheral afferents: emerging ion channel targets. *Nat Neurosci* 2014; 17: 153–163.
 17. Wall PD and Devor M. Sensory afferent impulses result from dorsal root ganglia as well as from the periphery in normal and nerve-injured rats. *Pain* 1983; 17: 321–339.
 18. Balasubramanian S, Stemkowski PL, Stebbing MJ and Smith PA. Sciatic chronic constriction injury produces cell-type specific changes in the electrophysiological properties of rat *Substantia Gelatinosa* neurons. *J Neurophysiol* 2006; 96: 579–590.
 19. Woolf CJ. Evidence for a central component of post-injury pain hypersensitivity. *Nature* 1983; 306: 686–688.
 20. Sandkuhler J. Models and mechanisms of hyperalgesia and allodynia. *Physiol Rev* 2009; 89: 707–758.
 21. Chen Y, Balasubramanian S, Lai AY, Todd KG and Smith PA. Effects of sciatic nerve axotomy on excitatory synaptic transmission in rat *Substantia Gelatinosa*. *J Neurophysiol* 2009; 102: 3203–3215.
 22. Zhou C and Luo ZD. Electrophysiological characterization of spinal neuron sensitization by elevated calcium channel alpha-2-delta-1 subunit protein. *Eur J Pain* 2014; 18: 649–658.
 23. Zhou C and Luo ZD. Nerve injury-induced calcium channel alpha-2-delta-1 protein dysregulation leads to increased pre-synaptic excitatory input into deep dorsal horn neurons and neuropathic allodynia. *Eur J Pain* 2015; 19: 1267–1276.
 24. Luo ZD, Chaplan SR, Higuera ES, Sorkin LS, Stauderman KA, Williams ME and Yaksh TL. Upregulation of dorsal root ganglion (alpha)2(delta) calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats. *J Neurosci*. 2001; 21: 1868–1875.
 25. Luo ZD, Calcutt NA, Higuera ES, Valder CR, Song YH, Svensson CI and Myers RR. Injury type-specific calcium channel alpha 2 delta-1 subunit up-regulation in rat neuropathic pain models correlates with antiallodynic effects of gabapentin. *J Pharmacol Exp Ther* 2002; 303: 1199–1205.
 26. Boroujerdi A, Zeng J, Sharp K, Kim D, Steward O and Luo ZD. Calcium channel alpha-2-delta-1 protein upregulation in dorsal spinal cord mediates spinal cord injury-induced neuropathic pain states. *Pain* 2011; 152: 649–655.
 27. Li CY, Song YH, Higuera ES and Luo ZD. Spinal dorsal horn calcium channel $\alpha 2\delta$ -1 subunit upregulation contributes to peripheral nerve injury-induced tactile allodynia. *J Neurosci* 2004; 24: 8494–8499.
 28. Savalli N, Pantazis A, Sigg D, Weiss JN, Neely A and Olcese R. The alpha2delta-1 subunit remodels CaV1.2 voltage sensors and allows Ca²⁺ influx at physiological membrane potentials. *J Gen Physiol* 2016; 148: 147–159.
 29. Fox AP, Nowycky MC and Tsien RW. Kinetic and pharmacological properties distinguishing three types of calcium currents in chick sensory neurons. *J Physiol* 1987; 394: 149–172.
 30. Gould RJ, Murphy KM and Snyder SH. [³H]nitrendipine-labeled calcium channels discriminate inorganic calcium agonists and antagonists. *Proc Natl Acad Sci U S A* 1982; 79: 3656–3660.
 31. Eichelbaum Mikus M, Mast G, Fischer V, Kuhlmann C and Machleidt UC. Pharmacokinetics and pharmacodynamics of nitrendipine in healthy subjects and patients with kidney and liver disease. *J Cardiovasc Pharmacol* 1988; 12: S6–S10.
 32. Marais E, Klugbauer N and Hofmann F. Calcium channel alpha(2)delta subunits-structure and Gabapentin binding. *Mol Pharmacol* 2001; 59: 1243–1248.
 33. Prescott SA and de Koninck Y. Four cell types with distinctive membrane properties and morphologies in lamina I of the spinal dorsal horn of the adult rat. *J Physiol* 2002; 539: 817–836. [Database][Mismatch]
 34. Bao J, Li JJ and Perl ER. Differences in Ca²⁺ channels governing generation of miniature and evoked excitatory synaptic currents in spinal laminae I and II. *J Neurosci* 1998; 18: 8740–8750.
 35. Heinke B, Balzer E and Sandkuhler J. Pre- and postsynaptic contributions of voltage-dependent Ca²⁺ channels to nociceptive transmission in rat spinal lamina I neurons. *Eur J Neurosci* 2004; 19: 103–111.
 36. Krause HP, Ahr HJ, Beermann D, Siefert HM, Suwelack D and Weber H. The pharmacokinetics of nitrendipine. I. Absorption, plasma concentrations, and excretion after single administration of [¹⁴C]nitrendipine to rats and dogs. *Arzneimittelforschung* 1988; 38: 1593–1599.
 37. Shafer TJ and Atchison WD. Transmitter, ion channel and receptor properties of pheochromocytoma (PC12) cells: a model for neurotoxicological studies. *Neurotoxicology* 1991; 12: 473–492.
 38. Liu Felix HR, Gurnett CA, Waard MD, Witcher DR and Campbell KP. Expression and subunit interaction of voltage-dependent Ca²⁺ channels in PC12 cells. *J Neurosci* 1996; 16: 7557

39. Biggs JE, Boakye PA, Ganesan N, Stemkowski PL, Lantero A, Ballanyi K and Smith PA. Analysis of the long-term actions of gabapentin and pregabalin in dorsal root ganglia and substantia gelatinosa. *J Neurophysiol* 2014; 112: 2398–2412.
40. Chang E, Chen X, Kim M, Gong N, Bhatia S and Luo ZD. Differential effects of voltage-gated calcium channel blockers on calcium channel alpha-2-delta-1 subunit protein-mediated nociception. *Eur J Pain* 2015; 19: 639–648.
41. Hatano N, Ohya S, Muraki K, Giles W and Imaizumi Y. Dihydropyridine Ca²⁺ channel antagonists and agonists block Kv4.2, Kv4.3 and Kv1.4 K⁺ channels expressed in HEK293 cells. *Br J Pharmacol* 2003; 139: 533–544.
42. Fagni L, Bossu JL and Bockaert J. Inhibitory effects of dihydropyridines on macroscopic K⁺ currents and on the large-conductance Ca(2+)-activated K⁺ channel in cultured cerebellar granule cells. *Pflugers Arch* 1994; 429: 176–182.
43. Tanaka O, Sakagami H and Kondo H. Localization of MRNAs of voltage-dependent Ca(2+)-channels: four subtypes of alpha 1- and beta-subunits in developing and mature rat brain. *Brain Res Mol Brain Res* 1995; 30: 1–16.
44. Motin L and Adams DJ. Omega-conotoxin inhibition of excitatory synaptic transmission evoked by dorsal root stimulation in rat superficial dorsal horn. *Neuropharmacology* 2008; 55: 860–864.
45. Hoppa MB, Lana B, Margas W, Dolphin AC and Ryan TA. Alpha2delta expression sets presynaptic calcium channel abundance and release probability. *Nature* 2012; 486: 122–125.
46. Alles SRA, Bandet MV, Eppler K, Winship IR, Noh MC, Baker G, Ballanyi K and Smith PA. Acute anti-allodynic action of gabapentin in dorsal horn and primary somatosensory cortex: correlation of behavioural and physiological data. *Neuropharmacology* 2017; 113A: 576–590.
47. Peirs C and Seal RP. Neural circuits for pain: recent advances and current views. *Science* 2016; 354: 578–584.
48. Daniele CA and MacDermott AB. Low-threshold primary afferent drive onto gabaergic interneurons in the superficial dorsal horn of the mouse. *J Neurosci* 2009; 29: 686–695.
49. Zheng J, Lu Y and Perl ER. Inhibitory neurones of the spinal substantia gelatinosa mediate interaction of signals from primary afferents. *J Physiol* 2010; 588: 2065–2075.
50. Scroggs RS and Fox AP. Calcium current variation between acutely isolated adult rat dorsal root ganglion neurons of different size. *J Physiol* 1992; 445: 639–658.
51. Lipman AG. Analgesic drugs for neuropathic and sympathetically maintained pain. *Clin Geriatr Med* 1996; 12: 501–515.
52. Gadotti VM, Bladen C, Zhang FX, Chen L, Gunduz MG, Simsek R, Safak C and Zamponi GW. Analgesic effect of a broad-spectrum dihydropyridine inhibitor of voltage-gated calcium channels. *Pflugers Arch* 2015; 467: 2485–2493.