

**FHS PUBLIC ACCESS**

Author manuscript

*J Occup Environ Med.* Author manuscript; available in PMC 2017 September 01.

Published in final edited form as:

*J Occup Environ Med.* 2016 September ; 58(9): 940–945. doi:10.1097/JOM.0000000000000832.**Biomarkers of human cardiopulmonary response after short-term exposures to medical laser generated particulate matter from simulated procedures: a pilot study****Ramon Lopez, PhD<sup>1</sup>, Mark O. Farber, MD<sup>2</sup>, Vincent Wong, MD<sup>3</sup>, and Steven E. Lacey, PhD<sup>1</sup>**<sup>1</sup>Indiana University, Fairbanks School of Public Health, Department of Environmental Health Science<sup>2</sup>Indiana University Health, Pulmonology-Critical Care<sup>3</sup>Indiana University, Krannert Institute of Cardiology, Department of Medicine**Abstract**

**Objective**—We conducted an exposure chamber study in humans using a simulated clinical procedure lasing porcine tissue to demonstrate evidence of effects of exposure to laser generated particulate matter (LGPM).

**Methods**—We measured pre- and post-exposure changes in exhaled nitric oxide (eNO), spirometry, heart rate variability (HRV), and blood markers of inflammation in five volunteers.

**Results**—Change in pre- and post-exposure measurements of eNO and spirometry were unremarkable. Neutrophil and lymphocyte counts increased and fibrinogen levels decreased in four of the five subjects. Measures of HRV showed decreases in the standard deviation of normal between beat intervals and sequential five-minute intervals.

**Conclusion**—These data represent the first evidence of human physiologic response to LGPM exposure. Further exploration of coagulation effects and HRV are warranted.

**Keywords**

Biomarkers; medical laser; particulate matter; exposure

**Background**

In the United States, an estimated half million healthcare professionals were exposed to surgical smoke in 2008 (2), and the rapid development of new clinical laser technologies and their applications, as well as growth in the sale of medical lasers (3) promises continued growth of human exposure. Medical laser-generated aerosol is created from the heating of the target tissue leading to the vaporization, pyrolysis, and combustion of cellular material, and the release of steam, cell content, and combustion by-products (4, 5). Our preliminary

\*Corresponding author: Steven E. Lacey, PhD, CIH, CSP, Indiana University, Fairbanks School of Public Health, Department of Environmental Health Science, 1050 Wishard Blvd. (RG 6086), Indianapolis, IN 46202 USA, selacey@iu.edu.

Conflict of interest: The authors have no relationships/conditions/circumstances that present potential conflict of interest.

work measuring the concentration of gases and particulate matter present in the laser generated aerosol has directed us to focus specifically on the particulate matter fraction (6, 7), and we have demonstrated that operational parameters (e.g., power, beam diameter) impact size-specific laser generated particulate matter (LGPM) emission rates (8) which may be important in determining possible health implications and designing control strategies.

The current evidence-based understanding of LGPM exposure health effects has been limited to animal studies, which have demonstrated inflammatory responses in pulmonary tissue defined by interstitial pneumonia, bronchiolitis, and emphysema (9–12). One study noted a decreased response with increased air filtration, indicating a potential dose-response relationship (9). Long-term outcomes of human exposure associated with the chronic inhalation of the aerosol have not been studied. This pilot study measured for the first time the human physiological response to LGPM aerosol exposure by measuring exhaled nitric oxide (eNO), spirometry, heart rate variability (HRV), and blood biomarkers including white blood cell counts, fibrinogen, and platelets. A symptom survey was also used to document any noticeable effects the participants may have experienced.

### Measures of cardiopulmonary response to particulate matter exposures

Nitric oxide is produced by the endothelial and epithelial cells in lung tissue, and its concentration in exhaled breath has been demonstrated to increase with airway inflammation (13, 14). Real-time eNO analyzers have been developed, tested and approved for medical use (15, 16) and studies examining the repeatability of eNO measurements have shown low variability not impacted by season, food intake, body weight, or height (17). The utility of eNO for measuring particulate matter (PM)-associated airway inflammation has been demonstrated in studies of ambient air pollution in which increased eNO was measured after short trips along high-density roadways, after exercise activities in urban locations, and after normal activities on days with high ambient PM concentrations (16, 18–22).

Spirometry studies have measured forced expiratory volume in one second ( $FEV_1$ ) and forced vital capacity (FVC) and demonstrated an inverse relationship with PM concentration in susceptible populations, including elderly and asthmatics, after short-term ambient PM exposures such as walking in urban settings during heavy traffic (23–25).

While heart rate remains fairly constant in most normal individuals, the time between two successive beats can vary significantly (26). Time-domain measures of HRV have been established to measure response to environmental exposures, including the standard deviation of all normal RR intervals (SDNN, representing overall HRV); standard deviation of sequential five-minute intervals (SDANN, long-term changes in HRV); and root mean square of the successive differences (RMSSD, short term changes in HRV) (27). Heart rate variability is a predictor of cardiovascular mortality and morbidity after exposure to high levels of ambient PM (28–31); disruption of the autonomic nervous system, activation of pro-inflammatory pathways and accelerated atherosclerosis, are linked to HRV and the increased risk of cardiovascular events. Reductions in heart rate variability from PM exposure has been demonstrated in animals and humans, even when signs of hypoxia and respiratory distress are absent (32, 33).

It has been proposed that ultrafine particles may penetrate into the blood stream and cause a systemic inflammatory response that can be measured by biomarkers present in blood (34). The particles interact with platelets in the bloodstream, increasing coagulability (35). It has also been demonstrated that changes occur in plasma fibrinogen (36–39) and peripheral neutrophils (40–42) in animals and humans; both conditions are associated with coronary heart disease and myocardial infarction (43, 44) and are believed to be responses to oxidative stress and systemic inflammation caused by exposure to ambient PM (35, 39, 45, 46). Other studies have demonstrated increases in lymphocytes and eosinophils in animals with exposure to outdoor PM (47); the increases were noticed six hours after exposure, and peaked at 12 hours post-exposure before declining (47).

## Methods

We measured pre- and post-exposure changes in eNO, spirometry, HRV, and blood markers of systemic inflammation in five volunteer participants under controlled laboratory conditions. Study subjects were healthy men and women with no current pulmonary or cardiovascular illness or disease, self-described as non-smokers, sedentary, and between 35 and 55 years of age. We used an Ultra MD™ 60 Laser System (max power = 60 W,  $\lambda$ = 10,600 nm, pulsed) (Laser Engineering Inc., Franklin, Tennessee) in a simulated laser clinical procedure, and lased porcine skin in the hood of an exposure chamber in a controlled manner. Size-selective particle concentration was measured. The study design was reviewed and approved by the Indiana University Institutional Review Board, protocol #1505847764.

### Exposure chamber

The exposure chamber was a hybrid design of the emission chamber described in Lippert et al. (2014) (48) and an exposure chamber designed by Morawska et al. (2009) (Figure 1) (49). The chamber system is composed of a rectangular glass hood, connected via the transition section to an aluminum duct. An opening in the glass hood allows participants to comfortably position their heads in the system and maintain eye contact with the research team.

### Exposure concentrations

Two previously reported field studies that simulated medical laser procedures in hospital operating rooms documented concentrations of laser generated particulate matter ranging from 590–1,690  $\mu\text{g}/\text{m}^3$  (50, 51). We used 1,690  $\mu\text{g}/\text{m}^3$  as upper limit to establish the range of exposures in the chamber study, performing our evaluations at 0%, 50% and 100% of 1,690  $\mu\text{g}/\text{m}^3$ . Our earlier work demonstrated that our system is capable of reliably generating predictable mean concentrations inside the exposure chamber through manipulation of operational parameter settings (Power: 2 and 3 W; PRF: 5 Hz; pulse duration: 0.1 seconds; beam diameter: 1.0 mm). Real-time monitoring of LGPM concentration in the chamber was measured using a Aerotrak® 9306 (TSI® Inc. Shorewood, Minnesota), and a P-Trak® 8525 (TSI® Inc. Shorewood, Minnesota) as described in Lopez et al. (2015) (7). At the start of each day, we verified background concentration of particulate matter and air-flow rate in the exposure chamber. We recognize that gases and vapors are also present in laser generated aerosol, but our previous work has demonstrated these contaminants are transient and, when

present, have extremely low concentrations, so we have focused on the particulate matter fraction. (52).

### Human exposure events

Since cardiopulmonary response to LGPM exposure has never been measured systematically, it was unclear at what level of exposure a response would be noticed. We used an accelerated study design with two phases (phase 1 and phase 2), with a new set of study participants in each phase. Five participants (one male, four female) comprised the study group. All participants wore laser eye protection during the lasing procedure.

In phase 1, we performed a response range-finding experiment with two subjects. In their first session, each participant sat in the exposure chamber for 15-minutes, but no lasing was performed (zero exposure). Three days later, study subjects returned and were exposed to four sequential 15-minute sessions, with a 10-minute break between sessions; the first two sessions were at  $\sim 850 \mu\text{g}/\text{m}^3$  (50% exposure) and the latter two sessions were at  $\sim 1690 \mu\text{g}/\text{m}^3$  (100% exposure). The initial range-finding experiment in phase 1 was intended to scale up exposure concentrations and to monitor for any unanticipated acute adverse reaction as a safety measure. Exposure time of 15 minutes for each session was chosen because it is considered a typical duration for many clinical laser procedures (53), and it is typical for several procedures to be performed in a single day.

Similarly, in phase 2, three new subjects were first exposed to a control session (zero exposure), then three days later study subjects returned and were exposure to four sequential 15-minute sessions, with a 10-minute break between sessions, at  $\sim 1690 \mu\text{g}/\text{m}^3$  (100% exposure).

### Measuring a response

**Exhaled NO**—Participants provided pre- and post-exposure exhaled breath samples using a Niox Vero<sup>®</sup> eNO monitor (Aerocrine, Solna, Sweden) (17). Participants had 20 minutes to adjust to the laboratory environment before two pre-exposure eNO measurements were made. Post-exposure eNO measurements were made immediately after the exposure event, and then 20 minutes after the event concluded. In phase 2, additional post-exposure measurements were collected every 20 minutes for 1 hour and 20 minutes.

**Spirometry**—Pulmonary function testing was performed in accordance with the guidelines of the American Thoracic Society/European Respiratory Society Task Force: Standardization of Spirometry (54). A Spirodoc<sup>®</sup> Spirometer (MIR, Waukesha, WI) with logging capabilities was used to measure FVC and FEV<sub>1</sub>, and the study participants were coached by our team pulmonologist. Each participant completed the spirometry tests immediately before the exposure event(s), and again five minutes post-exposure.

**Heart rate variability**—Heart rate variability was measured using a Datrix VX3 Holter (Biomedical Systems, Brussels, Belgium) with seven lead attachments. Our team cardiologist prepped and connected the monitors and participants wore the devices for two

consecutive 24-hour monitoring sessions that represented pre- and post-exposure event. Post-exposure monitoring began at the start of the exposure event.

**Blood analysis**—Participants provided peripheral venous blood 20 minutes before the exposure event, and three hours post-exposure. Samples were collected by a certified phlebotomist and analyzed by the Indiana University Health Pathology Laboratory for white blood cells (total, monocytes, neutrophils, lymphocytes, eosinophils, and basophils), fibrinogen and platelets.

**Symptom survey**—A short survey was verbally administered to participants pre- and post-exposure, and in the evening after the exposure event. The questionnaire was based on a tool developed by the National Institute for Occupational Safety and Health to assess possible occupational exposure health outcomes from medical laser aerosol exposure (55). Participants were asked if they were experiencing symptoms including irritation, headache, dizziness, cough, and noticeable lung problems.

## Results

For all measures of response, no material changes were observed in the control (zero-exposure) sessions. Tabulated results present pre- and post-measures of response from actual exposure sessions.

### Exhaled nitric oxide

There was no notable change in eNO measurements for any participant (Table 1); change in post-exposure measures varied within  $\pm 4$  ppb, similar to measures observed in control exposure sessions. During phase 2, additional post-exposure eNO measures were made, but again no material change was observed.

### Spirometry

Spirometry results were unremarkable for all participants (Table 2). The data demonstrate no change from pre- to post-exposure for all five study participants.

### Heart rate variability

Three measures of heart rate variability were calculated for each 24-hour time interval: SDNN, SDANN, and RMSSD (27). All participants demonstrated a post-exposure decrease in SDNN and SDANN, and three of five participants demonstrated a post-exposure decrease in RMSSD (Table 3). Control session measurement showed greater change both above and below baseline measurements.

### Biomarkers in blood

There was a general increase in white blood cell counts from pre- to post-exposure in four of the five participants; three participants demonstrated increases between 20 to 25% (Table 4). Neutrophils and lymphocytes had the greatest post-exposure change. Changes in monocyte and eosinophil counts were unremarkable.

Post-exposure fibrinogen levels decreased between 6 and 10% in four out of five participants. One participant experienced a 22% post-exposure increase in fibrinogen during the control exposure.

Post-exposure platelet counts increased 5 to 12% in three out of five participants; the two other participants showed modest decreases.

### Symptom survey

One participant reported feeling dizzy and lightheaded for 3 to 5 minutes immediately post-exposure. No other participants reported any noticeable effects.

### Discussion

Exhaled nitric oxide is an effective measure of inflammatory response in asthmatic patients, but studies related to ambient air pollution have been inconclusive, with increases in eNO demonstrated in some studies, and no change in other studies. Most of these investigations involve longer exposure periods than our study (19–21, 56). Our investigation looked at short exposures of 15-minutes, with a total one hour of exposure, for a single day. We were uncertain if a change would occur instantaneously, or if there was lag time between exposure and outcome. During phase 2, participants provided exhaled breath samples every twenty minutes post-exposure for 80 minutes but no change was observed. Future exploration to determine if a response is dose- or time-dependent is warranted.

In our study, concentration of fibrinogen decreased and platelet count increased post-exposure for most subjects. In a previous chamber study, human subjects were exposed to airborne particle concentrations up to 200  $\mu\text{g}/\text{m}^3$  for two hours; measures of fibrinogen showed an increase 18 hours post-exposure, but measures immediately after exposure showed no change (36). A second study measured fibrinogen after exposure to welding fume, and noticed a significant decrease in fibrinogen concentration six and 24 hours post-exposure (57). In our study, it was unclear when a response began or when it peaked since we collected a single blood sample 3-hours post-exposure. The change in levels of fibrinogen and platelets may be explained by the coagulation cascade. Endothelial damage activates the increased synthesis of platelets and fibrinogen, and the conversion of fibrinogen to fibrin fibers, to reduce blood loss and stabilizes platelet plugs (58). Since this is a time-dependent process, when a post-exposure blood sample is collected may affect the observed fibrinogen and platelet count. In future studies, additional blood samples may elucidate time-dependent effects.

We noted post-exposure increases of neutrophils, lymphocytes, monocytes and total WBC, similar to increases seen in ambient PM studies as an inflammatory response from oxidative stress (28, 40, 45, 57, 59). In our study, one participant (#4) demonstrated a high pre-exposure WBC count and a post-exposure decrease in WBC count; in a follow-up conversation with the participant, he suggested he was developing a cold at that time, which may explain this specific result.

Other biomarkers in blood exist that may also be helpful in determining response to PM exposures. C-reactive protein is believed to be a marker of inflammation, and is associated with risk of myocardial infarctions (44), and in some studies has been found to be elevated with high PM exposures (60–62). Interleukin-6 is believed to be stimulated as an immune and inflammatory response, and also has been demonstrated to increase with high levels of PM (62, 63). Future studies should explore these additional potential indicators of response.

A high degree of heart rate variability is normal for healthy individuals. The observed decline in HRV measures, including SDNN and RMSSD, are similar to those found in other studies comparing changes in HRV to levels of PM in ambient air, or from activities at home that produce PM, and they generally noted decreases in measures of HRV with increasing PM exposure (64–67).

The purpose of our study was to determine a human exposure-response from LGPM exposure, and the efficacy of respiratory protection was outside the scope of this pilot study, so participants did not use respiratory protection during exposure events. It has been reported that healthcare professionals do use surgical masks during laser use, but these are not filtering respirators (68, 69)

## Conclusion

Our pilot study is the first attempt to measure a response to short-term LGPM exposure in humans. The clearest evidence of response was demonstrated by decreased HRV, increased WBC counts, decreased fibrinogen, and limited evidence of increased platelets. Under our experimental conditions, eNO, and spirometry did not prove to be effective measures of response. These lines of investigation were highly exploratory, and the limits of our study design mean we may have missed detecting a response that may be time- or dose-dependent. Further exploration of eNO and spirometry is warranted since if we determine a study strategy that demonstrates a response, these measures are relatively inexpensive and noninvasive. Other blood biomarkers should also be explored.

These results warrant further investigation in exploring human responses to workplace LGPM exposure as the implications for risk management of the health effects are critical for ensuring safe work environments. Improved understanding of human health implications will make more compelling risk communication and the need for improved control strategies.

## Acknowledgments

Funding source: This study was supported by a Project Development Team within the ICTSI NIH/NCRR Grant Number UL1TR001108.

Sources of support requiring acknowledgment: This study was supported by a Project Development Team within the ICTSI NIH/NCRR Grant Number UL1TR001108.

The authors would like to thank Sarah Eberhardt for her participation in the initial design of the chamber, and Jeries Smirat for his help in the laboratory.



## References


1. OSHA. Technical manual: laser hazards, section III. Vol. chapter 6. Occupational Safety and Health Administration; 1999.
2. OSHA. Administration OSAH. Safety and health topics: laser/electrosurgery plume. Washington, DC: Occupational Safety And Health Administration; 2008.
3. Overton, G.; Belforte, D.; Noguee, A.; Holton, C. Laser Marketplace 2015: Lasers surround us in the Year of Light. 2015.
4. Alp E, Bijl D, Bleichrodt RP, Hansson B, Voss A. Surgical smoke and infection control. *The Journal of Hospital Infection*. 2006; 62:1–5. [PubMed: 16002179]
5. Ulmer BC. The hazards of surgical smoke. *AORN Journal*. 2008; 87:721–738. [PubMed: 18461735]
6. Lopez R, Lacey SE, Jones RM. Application of a two-zone model to estimate medical laser generated particulate matter exposures. *Journal of Occupational and Environmental Hygiene*. 2014
7. Lopez R, Lacey SE, Lippert JF, Liu LC, Esmen NA, Conroy LM. Characterization of Size-Specific Particulate Matter Emission Rates for a Simulated Medical Laser Procedure-A Pilot Study. *Ann Occup Hyg*. 2015
8. Lopez R, Lacey SE, Lippert JF, Liu L, Esmen N, Conroy L. Characterization of size-specific particulate matter emission rates for a simulated medical laser procedure. *Annals of Occupational Hygiene*. 2014
9. Baggish MS, Baltoyannis P, Sze E. Protection of the rat lung from the harmful effects of laser smoke. *Lasers in Surgery and Medicine*. 1988; 8:248–253. [PubMed: 3393053]
10. Baggish MS, Elbakry M. The effects of laser smoke on the lungs of rats. *American Journal of Obstetrics and Gynecology*. 1987; 156:1260–1265. [PubMed: 3107392]
11. Freitag L, Chapman GA, Sielczak M, Ahmed A, Russin D. Laser smoke effect on the bronchial system. *Lasers in Surgery and Medicine*. 1987; 7:283–288. [PubMed: 3626753]
12. Wenig BL, Stenson KM, Wenig BM, Tracey D. Effects of plume produced by the Nd:YAG laser and electrocautery on the respiratory system. *Lasers in Surgery and Medicine*. 1993; 13:242–245. [PubMed: 8464311]
13. Guo FH, De Raeve HR, Rice TW, Stuehr DJ, Thunnissen FB, Erzurum SC. Continuous nitric oxide synthesis by inducible nitric oxide synthase in normal human airway epithelium in vivo. *Proceedings of the National Academy of Sciences*. 1995; 92:7809–7813.
14. Asano K, Chee CB, Gaston B, et al. Constitutive and inducible nitric oxide synthase gene expression, regulation, and activity in human lung epithelial cells. *Proceedings of the National Academy of Sciences USA*. 1994; 91:10089–10093.
15. Dweik RA, Boggs PB, Erzurum SC, et al. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FENO) for clinical applications. *American journal of respiratory and critical care medicine*. 2011; 184:602–615. [PubMed: 21885636]
16. Hatziaorou E, Tsanakas J. Assessment of airway inflammation with exhaled NO measurement. *Hippokratia*. 2007; 11:51–62. [PubMed: 19582178]
17. Kharitonov SA, Gonio F, Kelly C, Meah S, Barnes PJ. Reproducibility of exhaled nitric oxide measurements in healthy and asthmatic adults and children. *European Respiratory Journal*. 2003; 21:433–438. [PubMed: 12661997]
18. Steerenberg PA, Snelder JB, Fischer PH, Vos JG, van Loveren H, van Amsterdam JG. Increased exhaled nitric oxide on days with high outdoor air pollution is of endogenous origin. *European Respiratory Journal*. 1999; 13:334–337. [PubMed: 10065677]
19. Adar SD, Adamkiewicz G, Gold DR, Schwartz J, Coull BA, Suh H. Ambient and microenvironmental particles and exhaled nitric oxide before and after a group bus trip. *Environmental health perspectives*. 2007; 115:507–512. [PubMed: 17450216]
20. Barraza-Villarreal A, Sunyer J, Hernandez-Cadena L, et al. Air pollution, airway inflammation, and lung function in a cohort study of Mexico City schoolchildren. *Environmental health perspectives*. 2008; 116:832–838. [PubMed: 18560490]



21. Bos I, De Boever P, Vanparijs J, Pattyn N, Panis LI, Meeusen R. Subclinical effects of aerobic training in urban environment. *Medicine & Science in Sports & Exercise*. 2013; 45:439–447. [PubMed: 23073213]
22. Benor S, Alcalay Y, Domany KA, et al. Ultrafine particle content in exhaled breath condensate in airways of asthmatic children. *J Breath Res*. 2015; 9:026001. [PubMed: 25830607]
23. McCreanor J, Cullinan P, Nieuwenhuijsen MJ, et al. Respiratory effects of exposure to diesel traffic in persons with asthma. *N Engl J Med*. 2007; 357:2348–2358. [PubMed: 18057337]
24. Gent JF, Triche EW, Holford TR, et al. Association of low-level ozone and fine particles with respiratory symptoms in children with asthma. *JAMA*. 2003; 290:1859–1867. [PubMed: 14532314]
25. Hoek G, Dockery DW, Pope A, Neas L, Roemer W, Brunekreef B. Association between PM10 and decrements in peak expiratory flow rates in children: reanalysis of data from five panel studies. *Eur Respir J*. 1998; 11:1307–1311. [PubMed: 9657571]
26. Achten J, Jeukendrup AE. Heart rate monitoring: applications and limitations. *Sports medicine (Auckland, NZ)*. 2003; 33:517–538.
27. Crawford MH, Bernstein SJ, Deedwania PC, et al. ACC/AHA guidelines for ambulatory electrocardiography: executive summary and recommendations. A report of the American College of Cardiology/American Heart Association task force on practice guidelines (committee to revise the guidelines for ambulatory electrocardiography). *Circulation*. 1999; 100:886–893. [PubMed: 10458728]
28. Pope CA 3rd, Hansen ML, Long RW, et al. Ambient particulate air pollution, heart rate variability, and blood markers of inflammation in a panel of elderly subjects. *Environmental health perspectives*. 2004; 112:339–345. [PubMed: 14998750]
29. Simkhovich BZ, Kleinman MT, Kloner RA. Air pollution and cardiovascular injury epidemiology, toxicology, and mechanisms. *J Am Coll Cardiol*. 2008; 52:719–726. [PubMed: 18718418]
30. Tsuji H, Larson MG, Venditti FJ Jr, et al. Impact of reduced heart rate variability on risk for cardiac events. The Framingham Heart Study. *Circulation*. 1996; 94:2850–2855. [PubMed: 8941112]
31. Tsuji H, Venditti FJ Jr, Manders ES, et al. Reduced heart rate variability and mortality risk in an elderly cohort. The Framingham Heart Study. *Circulation*. 1994; 90:878–883. [PubMed: 8044959]
32. Watkinson WP, Campen MJ, Costa DL. Cardiac arrhythmia induction after exposure to residual oil fly ash particles in a rodent model of pulmonary hypertension. *Toxicol Sci*. 1998; 41:209–216. [PubMed: 9520357]
33. Pope, CAR; Dockery, DW.; Kanner, RE.; Villegas, GM.; Schwartz, J. Oxygen saturation, pulse rate, and particulate air pollution: A daily time-series panel study. *American journal of respiratory and critical care medicine*. 1999; 159:365–372. [PubMed: 9927345]
34. Seaton A, MacNee W, Donaldson K, Godden D. Particulate air pollution and acute health effects. *Lancet*. 1995; 345:176–178. [PubMed: 7741860]
35. Schwartz J. Air pollution and blood markers of cardiovascular risk. *Environmental health perspectives*. 2001; 109(Suppl 3):405–409. [PubMed: 11427390]
36. Ghio AJ, Kim C, Devlin RB. Concentrated ambient air particles induce mild pulmonary inflammation in healthy human volunteers. *American journal of respiratory and critical care medicine*. 2000; 162:981–988. [PubMed: 10988117]
37. Gardner SY, Lehmann JR, Costa DL. Oil fly ash-induced elevation of plasma fibrinogen levels in rats. *Toxicol Sci*. 2000; 56:175–180. [PubMed: 10869466]
38. Chen R, Zhao Z, Sun Q, et al. Size-fractionated particulate air pollution and circulating biomarkers of inflammation, coagulation, and vasoconstriction in a panel of young adults. *Epidemiology (Cambridge, Mass)*. 2015; 26:328–336.
39. Wang C, Chen R, Zhao Z, et al. Particulate air pollution and circulating biomarkers among type 2 diabetic mellitus patients: the roles of particle size and time windows of exposure. *Environmental research*. 2015; 140:112–118. [PubMed: 25863184]
40. Salvi S, Blomberg A, Rudell B, et al. Acute inflammatory responses in the airways and peripheral blood after short-term exposure to diesel exhaust in healthy human volunteers. *American journal of respiratory and critical care medicine*. 1999; 159:702–709. [PubMed: 10051240]

41. Gordon T, Nadziejko C, Schlesinger R, Chen LC. Pulmonary and cardiovascular effects of acute exposure to concentrated ambient particulate matter in rats. *Toxicol Lett.* 1998; 96–97:285–288.
42. Sussan TE, Ingole V, Kim JH, et al. Source of biomass cooking fuel determines pulmonary response to household air pollution. *Am J Respir Cell Mol Biol.* 2014; 50:538–548. [PubMed: 24102120]
43. Danesh J, Collins R, Appleby P, Peto R. Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA.* 1998; 279:1477–1482. [PubMed: 9600484]
44. Lind P, Hedblad B, Stavenow L, Janzon L, Eriksson KF, Lindgarde F. Influence of plasma fibrinogen levels on the incidence of myocardial infarction and death is modified by other inflammation-sensitive proteins: a long-term cohort study. *Arteriosclerosis, thrombosis, and vascular biology.* 2001; 21:452–458.
45. Karotki DG, Spilak M, Frederiksen M, et al. Indoor and Outdoor Exposure to Ultrafine, Fine and Microbiologically Derived Particulate Matter Related to Cardiovascular and Respiratory Effects in a Panel of Elderly Urban Citizens. *International Journal of Environmental Research and Public Health.* 2015; 12:1667–1686. [PubMed: 25648225]
46. Frampton MW. Does inhalation of ultrafine particles cause pulmonary vascular effects in humans? *Inhalation toxicology.* 2007; 19(Suppl 1):75–79. [PubMed: 17886054]
47. Walters DM, Breyse PN, Wills-Karp M. Ambient urban Baltimore particulate-induced airway hyperresponsiveness and inflammation in mice. *American journal of respiratory and critical care medicine.* 2001; 164:1438–1443. [PubMed: 11704592]
48. Lippert JF, Lacey SE, Lopez R, et al. A pilot study to determine medical laser generated air contaminant emission rates for a simulated surgical procedure. *Journal of Occupational and Environmental Hygiene.* 2014:11.
49. Morawska L, Johnson GR, Ristovski ZD, et al. Size distribution and sites of origin of droplets expelled from the human respiratory tract during expiratory activities. *Journal of Aerosol Science.* 2009; 40:256–269.
50. Albrecht H, Washce W, Muller GJ. Assessment of the risk potential of pyrolysis products in plume produced during laser treatment under OR conditions. *Proceedings of SPIE. The International Society for Optics and Photonics.* 2005:455.
51. Tanpowpong K, Koytong W. Suspended particulate matter in an office and laser smoke particles in an operating room. *Journal of the Medical Association of Thailand = Chotmaihet thangphaet.* 2002; 85:53–57. [PubMed: 12075720]
52. Lippert JF, Lacey SE, Jones RM. Modeled occupational exposures to gas-phase medical laser generated air contaminants. *Journal of Occupational and Environmental Hygiene.* 2014
53. Sroka R, Janda P, Killian T, Vaz F. Comparison of long term results after Ho:YAG and diode laser treatment of hyperplastic inferior nasal turbinates. *Lasers in Surgery and Medicine.* 2007:324. [PubMed: 17304563]
54. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *European Respiratory Journal.* 2005; 26:319–338. [PubMed: 16055882]
55. Moss, EC.; Bryant, C.; Stewart, J.; Wen-Zong, W.; Fleegeer, ABJG. Health NifOS. HETA 88-101-2008. Washington D.C: 1990.
56. Jacobs L, Nawrot TS, de Geus B, et al. Subclinical responses in healthy cyclists briefly exposed to traffic-related air pollution: an intervention study. *Environmental health : a global access science source.* 2010; 9:64. [PubMed: 20973949]
57. Kim JY, Chen JC, Boyce PD, Christiani DC. Exposure to welding fumes is associated with acute systemic inflammatory responses. *Occup Environ Med.* 2005; 62:157–163. [PubMed: 15723880]
58. Laurens N, Koolwijk P, de Maat MP. Fibrin structure and wound healing. *Journal of thrombosis and haemostasis : JTH.* 2006; 4:932–939. [PubMed: 16689737]
59. Rubel C, Fernandez GC, Dran G, Bompadre MB, Isturiz MA, Palermo MS. Fibrinogen promotes neutrophil activation and delays apoptosis. *J Immunol.* 2001; 166:2002–2010. [PubMed: 11160249]

60. Michikawa T, Okamura T, Nitta H, et al. Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. *Environmental pollution (Barking, Essex : 1987)*. 2016; 213:460–467.
61. Dabass A, Talbott EO, Venkat A, et al. Association of exposure to particulate matter (PM2.5) air pollution and biomarkers of cardiovascular disease risk in adult NHANES participants (2001–2008). *International journal of hygiene and environmental health*. 2016; 219:301–310. [PubMed: 26725170]
62. Westberg H, Elihn K, Andersson E, et al. Inflammatory markers and exposure to airborne particles among workers in a Swedish pulp and paper mill. *International archives of occupational and environmental health*. 2016
63. Burchiel SW, Lauer FT, MacKenzie D, et al. Changes in HPBMC markers of immune function following controlled short-term inhalation exposures of humans to hardwood smoke. *Inhalation toxicology*. 2016; 28:61–70. [PubMed: 26895307]
64. Huang YL, Chen HW, Han BC, et al. Personal exposure to household particulate matter, household activities and heart rate variability among housewives. *PloS one*. 2014; 9:e89969. [PubMed: 24594880]
65. Gold DR, Litonjua A, Schwartz J, et al. Ambient pollution and heart rate variability. *Circulation*. 2000; 101:1267–1273. [PubMed: 10725286]
66. Liao D, Creason J, Shy C, Williams R, Watts R, Zweidinger R. Daily variation of particulate air pollution and poor cardiac autonomic control in the elderly. *Environmental health perspectives*. 1999; 107:521–525. [PubMed: 10378998]
67. Mordukhovich I, Coull B, Kloog I, Koutrakis P, Vokonas P, Schwartz J. Exposure to sub-chronic and long-term particulate air pollution and heart rate variability in an elderly cohort: the Normative Aging Study. *Environmental health : a global access science source*. 2015; 14:87. [PubMed: 26546332]
68. Commission TJ. Commission TJ. *Implementing Hospital Respiratory Protection Programs: Strategies from the Field*. Oakbrook Terrace, IL: The Joint Commission; 2014.
69. Steege AL, Boiano JM, Sweeney MH. NIOSH health and safety practices survey of healthcare workers: training and awareness of employer safety procedures. *American journal of industrial medicine*. 2014; 57:640–652. [PubMed: 24549581]



**Figure 1.** Glass exposure chamber for lasing of tissue and exposure to study participants, air is pulled through the exposure chamber and exhausted to a fume cabinet.

Pre- & post-exposure levels of exhaled NO in PPB

**Table 1**

Participant ID	Pre	Post	20 min Post	40 min Post	60 min Post	80 min Post
<b>Phase 1</b>						
1	19	17	16	-	-	-
2	29.5	32	31	-	-	-
<b>Phase 2</b>						
3	21	21	20	20	21	20
4	34.5	29	29	32	28	31
5	36	33	33	28	29	38

\* Post-exposure measurements after the first 20 minutes were not made during phase 1

**Table 2**

Pre- & post-exposure levels of spirometry in liters

	Participant ID	FEV <sub>1</sub>		FVC	
		Pre	Post	Pre	Post
<b>Phase 1</b>	1	2.93	2.86	3.4	3.3
	2	2.18	2.07	2.84	2.82
<b>Phase 2</b>	3	2.15	2.54	2.92	2.56
	4	4.35	4.39	4.49	4.47
	5	4.43	4.42	5.21	5.0

FEV<sub>1</sub>, forced expiratory volume in 1 second; FVC, forced vital capacity.

**Table 3**

Pre- & post- exposure levels of HRV measures

Participant ID	SDNN		SDANN		RMSSD		
	Pre	Post	Pre	Post	Pre	Post	
<b>Phase 1</b>	1	154	134	140	120	33	28
	2	139	135	120	119	61	42
<b>Phase 2</b>	3	110	97	99	79	27	35
	4	116	101	106	85	35	44
	5	110	92	87	68	35	33

SDNN, standard deviation of all normal RR intervals; SDANN standard deviation of sequential five-minute intervals; RMSSD, root mean square of the successive difference.



Pre- & post-exposure levels of fibrinogen and white blood cell counts on days with an exposure

**Table 4**

Participant ID	Fibrinogen (mg/dl)		Total WBC (#/µl)		Neutrophils (#/µl)		Lymphocytes (#/µl)		Platelets (# K/ µl)*	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
<b>Phase 1</b>	1	295	263	5300	3300	4300	1600	2000	244	262
	2	333	311	5600	3600	4400	1400	1700	215	227
<b>Phase 2</b>	3	302	342	6700	4400	4500	1200	2000	201	226
	4	414	378	9700	8300	5800	2700	2300	279	257
	5	440	406	5900	7300	3100	4200	2100	2200	254

\* Platelet count in [# \* 1000]