

1 Running head: Race/ethnicity and retention in TBI research

2

3 Race/Ethnicity and Retention in Traumatic Brain Injury Outcomes Research:

4 A Traumatic Brain Injury Model Systems National Database Study

5

6 Angelle M. Sander, Ph.D., Anthony H. Lequerica, Ph.D., Jessica M. Ketchum, Ph.D.,

7 Flora M. Hammond, Ph.D., Kelli Williams Gary, Ph.D., Monique R. Pappadis, Ph.D.,

8 Elizabeth R. Felix, Ph.D., Douglas Johnson-Greene, Ph.D., Tamara Bushnik, Ph.D.

9 From the Departments of Physical Medicine and Rehabilitation & Psychiatry, Baylor College of

10 Medicine/Harris Health System and Brain Injury Research Center, TIRR Memorial Hermann,

11 Houston, TX (Sander); Kessler Foundation and Rutgers, New Jersey Medical School, West

12 Orange, NJ (Lequerica); Craig Hospital, Englewood, CO (Ketchum); Department of Physical

13 Medicine and Rehabilitation, Indiana University School of Medicine, Indianapolis, IN

14 (Hammond); Departments of Physical Medicine and Rehabilitation/Neuropsychology and

15 Occupational Therapy, Virginia Commonwealth University, Richmond, VA (Gary); Division of

16 Rehabilitation Sciences, University of Texas Medical Branch, Galveston, TX and Brain Injury

17 Research Center, TIRR Memorial Hermann, Houston, TX (Pappadis); University of Miami

18 Miller School of Medicine, Miami, FL (Felix and Johnson-Greene); Rusk Rehabilitation, New

19 York University Langone School of Medicine, New York, NY (Bushnik)

This is the preprint of the article published in final edited form as:

Sander, A., Lequerica, A.H., Ketchum, J.M., Hammond, F.M., et al. (2018). Race/Ethnicity and Retention in Traumatic Brain Injury Outcomes Research: A Traumatic Brain Injury Model Systems National Database Study. *Journal of Head Trauma Rehabilitation*. Preprint.

20 The contents of this publication were developed under grants from the National Institute on
21 Disability, Independent Living, and Rehabilitation Research (NIDILRR grant numbers
22 90DPTB0016, 90DP0028, 90DP0060, and 90RT5007 (TIRR), 90DP0032 (Kessler), 90DP0034
23 and 90DP0084 (Craig), 90DP0036 (Indiana), 90DP0046 (Miami), 90DP0047 (New York), and
24 90DP0033 (Virginia)); the Agency for Healthcare Research and Quality (AHRQ grant number
25 R24HS022134); UTMB Claude D. Pepper Older Americans Independence Center funded by the
26 National Institute on Aging (NIA grant number 5P30-AG024832); the UTMB Institute for
27 Translational Sciences supported by a National Institutes of Health (NIH) Clinical and
28 Translational Science Award (UL1RR029876). NIDILRR is a Center within the Administration
29 for Community Living (ACL), Department of Health and Human Services (HHS). The contents
30 of this publication do not necessarily represent the policy or official views of NIDILRR, ACL,
31 AHRQ, NIA, NIH, or HHS, and you should not assume endorsement by the Federal
32 Government.

33

34 We certify that no party having a direct interest in the results of the research supporting this
35 article has or will confer a benefit on us or on any organization with which we are associated.

36 We also certify that all financial and material support for this research and work are clearly
37 identified in the title page of the manuscript.

38 Correspondence to Angelle M. Sander, Ph.D., TIRR Memorial Hermann Research Center, 1333
39 Moursund St., Houston, TX 77030, Email: asander@bcm.edu; Phone: 713-797-7161; Fax: 713-
40 797-7150. Reprints will not be available from the author.

41 Word count manuscript: 3,546

42 Word count abstract: 171

43 **ABSTRACT**

44 **Objective:** To investigate the contribution of race/ethnicity to retention in traumatic brain injury
45 (TBI) research at 1 to 2 years post-injury. **Setting:** Community. **Participants:** 5548 Whites,
46 1347 Blacks, and 790 Hispanics enrolled in the Traumatic Brain Injury Model Systems National
47 Database with dates of injury between October 1, 2002 and March 31, 2013. **Design:**
48 Retrospective database analysis. **Main Measure:** Retention, defined as completion of at least
49 one question on the follow-up interview by the person with TBI or a proxy. **Results:** Retention
50 rates 1-2 years post-TBI were significantly lower for Hispanic (85.2%) than for White (91.8%)
51 or Black participants (90.5%) and depended significantly on history of problem drug or alcohol
52 use. Other variables associated with low retention included older age, lower education, violent
53 cause of injury, and discharge to an institution versus private residence. **Conclusions:** The
54 findings emphasize the importance of investigating retention rates separately for Blacks and
55 Hispanics rather than combining them or grouping either with other races or ethnicities. The
56 results also suggest the need for implementing procedures to increase retention of Hispanics in
57 longitudinal TBI research.

58 **Keywords:** *Traumatic brain injury, Cultural Competency, Follow-Up Studies*

59

60 **LIST OF ABBREVIATIONS:**

61 IRB – Institutional Review Board

62 NIDILRR – National Institute on Disability, Independent Living and Rehabilitation Research

63 PTA – posttraumatic amnesia

64 TBI – Traumatic Brain Injury

65 TBIMS – Traumatic Brain Injury Model Systems

66

67

68

69

70

71

72

73

74

75

76

77

78 **INTRODUCTION**

79 Cognitive,¹ emotional,² and physical³ impairments resulting from traumatic brain injury
80 (TBI) contribute to decreased independent living,⁴ employment,⁵ and participation in leisure
81 activities.⁶ Approximately 3.2 million United States residents are estimated to have long-term or
82 life-long disability resulting from TBI.⁷ Unfortunately, minorities are disproportionately
83 represented among those who sustain TBI⁸ and those with poor outcomes.⁹ Blacks and Hispanics
84 with TBI have been shown to have poorer outcomes than Whites in overall functioning,¹⁰
85 functional independence,¹¹ independence in home activities,¹² employment outcomes,^{13,14} and
86 satisfaction with participation.¹⁵ Minorities have also been shown to utilize rehabilitation
87 services less than Whites in both civilian¹⁶ and military¹⁷ samples. These findings emphasize the
88 need for inclusion of minorities in longitudinal research and clinical trials targeting TBI, as their
89 exclusion can yield a biased view of outcomes.

90 Recruitment and retention of minorities is challenging for health research as a whole.
91 National Institutes of Health investigators are less likely to meet recruitment goals for minorities
92 compared to Whites.¹⁸ Minimal empirical evidence exists to support specific retention
93 strategies.¹⁹ Greater loss of minorities to follow-up is a common problem in research on TBI
94 outcomes, posing a threat to internal and external validity.²⁰

95 The relationship between race/ethnicity and loss to follow-up in TBI research has been
96 investigated in prior studies. Corrigan and colleagues²¹ studied predictors of loss to follow-up in
97 three longitudinal samples, including the Colorado TBI registry, five TBI Model System
98 (TBIMS) centers, and a single brain injury rehabilitation unit. Minorities with TBI were less
99 likely to be followed at 1 year compared to Whites in two of the three samples investigated.
100 Other variables that predicted loss to follow-up included violent injury, elevated blood alcohol

101 level at hospital admission, lower FIMTM motor score at rehabilitation admission, non-private
102 health insurance, and discharge to an institution. Krellman and colleagues²² studied predictors of
103 longitudinal follow-up patterns in the TBIMS National Database. Each participant had the
104 opportunity to complete follow-up at four time points- 1, 2, 5, and 10 years post-injury. Findings
105 were that non-responders (did not complete any follow-ups, but did not formally withdraw) and
106 wave responders (completed some follow-ups and skipped others) were more likely to be
107 minorities; however, Whites were also more likely to be in one of these groups if they were
108 missing data on pre-injury education. Missing data on pre-injury education, acute care payer, or
109 pre-injury employment status was also associated with non-responding and wave responding.
110 Recently, Jourdan and colleagues²³ studied a sample of 504 adults with severe TBI in Paris.
111 While they did not include race/ethnicity as a variable, they found other associations with loss to
112 follow-up that can inform covariate analyses. Specifically, loss to follow-up at 1 year post-injury
113 was associated with pre-injury unemployment and violent mechanism of injury. Pre-injury
114 unemployment and alcohol abuse were predictive of loss to follow-up at 4 years post-injury.

115 Research findings to date indicate that race/ethnicity likely contributes to retention in
116 longitudinal TBI research, with the pattern being lower retention of minorities; however, the
117 extant research is limited by methodological issues. First, prior studies have either combined
118 Blacks with Hispanics or grouped Hispanics with other minorities and compared them to Blacks
119 and Whites.^{21,22} The importance of investigating retention of Hispanics as a separate group is
120 justified by the fact that Hispanics currently represent the largest racial/ethnic minority group in
121 the United States, comprising 17% of the total population.²⁴ Persons of Hispanic ethnicity make
122 up approximately 10% of current enrollees in the TBIMS National Database.²⁵ Given their
123 substantial numbers and their likelihood of having poor outcomes compared to Whites,

124 investigation of retention of Hispanics as a separate group is warranted. Additionally, prior
125 studies have not investigated the potential interaction of race/ethnicity with other variables that
126 may impact retention in longitudinal TBI studies. For example, prior research has shown that
127 minorities with TBI are more likely to be unemployed at the time of injury and to be injured via
128 violence.^{26,27} As these variables have also been shown to predict loss to follow-up, they may
129 interact with race/ethnicity to impact retention.

130 The aims of the current study are: (1) to investigate retention in the TBIMS database for
131 Whites, Blacks, and Hispanics; (2) to investigate the contribution of being White, Black, or
132 Hispanic to retention at 1-or 2-years post-injury, after controlling for other variables that may
133 impact retention; and (3) to investigate the interaction of race/ethnicity with other variables that
134 may impact retention.

135 **METHODS**

136 **Participants**

137 Participants included were those enrolled in the National Database of the National
138 Institute for Disability, Independent Living, and Rehabilitation Research (NIDILRR) TBIMS
139 program. The TBIMS National Database includes individuals with newly acquired TBI who
140 receive comprehensive inpatient rehabilitation services at one of the NIDILRR-funded centers in
141 the US. Twenty-two centers contributed to the dataset for this analysis, with 7685 individuals
142 with dates of injury between October 1, 2002 and March 31, 2013. The start and end dates were
143 selected based on the availability of key variables (variables are periodically added and deleted
144 from the National Database) and to ensure all subjects had become eligible for 2-year follow-up.

145 Criteria for inclusion in the TBIMS National Database include: age \geq 16 at time of injury;
146 medically documented complicated mild, moderate, or severe TBI (emergency department

147 Glasgow Coma Scale score ≤ 12 or duration of posttraumatic amnesia > 24 hours or loss of
148 consciousness > 30 minutes or evidence of intracranial trauma on neuroimaging); admission to a
149 TBIMS acute-care hospital within 72 hours of injury; completion of inpatient rehabilitation
150 within the TBIMS; and informed consent obtained. During the interval covered by this study,
151 race/ethnicity was coded as a mixed variable rather than two separate variables in the TBIMS
152 National Database. Race/ethnicity was coded as White, Black, Asian/Pacific Islander, Native
153 American, Hispanic Origin, Other, or Unknown. Only participants coded as White, Black, or
154 Hispanic origin were included in the current analysis because the numbers in the other categories
155 were too low to provide a meaningful comparison. As shown in Figure 1, 348 people were
156 excluded for race/ethnicity other than White, Black, or Hispanic.

157 **Procedure**

158 IRB approval was obtained at all participating TBIMS institutions. Medical and injury
159 information was abstracted from participants' medical records according to TBIMS National
160 Database standardized procedures.⁴⁰ Demographic information was obtained by trained research
161 personnel who interviewed the individual with TBI or a proxy.

162 Follow-up interviews were conducted in person, via phone, or through the mail at 1 (± 2
163 months) and 2 years (± 3 months) post-injury. Sample derivation is shown in the flowchart in
164 Figure 1. In the TBIMS National Database, follow-up status is coded as followed, lost, refused,
165 withdrew, expired, incarcerated, or follow-up not attempted due to a center losing TBIMS
166 funding. Persons eligible for 1 or 2 year follow-up were excluded from the sample if they had
167 expired prior to 1 year follow-up, were incarcerated at both follow-ups, or if no attempt was
168 made to contact them at either follow-up due to loss of funding. Participants were considered to
169 be retained if the interview status variable was coded as "followed" at either year 1 or year 2.

170 Participants were considered to be not retained if interview status was coded as lost, refused, or
171 withdrew at both year 1 and year 2.

172 Standard follow-up procedures used by all centers included: 1) attempting contact as soon
173 as the follow-up window opened; 2) making at least 12 phone contact attempts during various
174 times of day and night and days of the week (including weekends) using the most reliable phone
175 numbers available; 3) sending a letter to the participant and any known contacts at their last
176 known mailing addresses; 4) using phone directory assistance in the last known city of residence,
177 internet searches, fee-based location services, and medical records to identify updated phone
178 numbers, addresses, or other contact information; 5) conducting a search for potential death
179 information; and 6) conducting a search of an inmate database to determine if the participant was
180 incarcerated.

181 **Measures**

182 Race/ethnicity category was preferentially determined by asking the person with injury or
183 their caregiver, but medical record information was used if information could not be obtained in
184 the preferred way. The outcome variable, retention, was a dichotomous variable defined as
185 completion of at least one question on either the 1 or 2 year follow-up interview by the person
186 with TBI or a proxy.

187 Covariates:

188 Injury severity was measured by the duration of post-traumatic amnesia (PTA) – a period
189 marked by confusion and inability to form new memories after TBI that is predictive of global
190 outcomes after moderate-severe TBI.²⁸ Duration of PTA was calculated as the number of days
191 between the TBI and the first of two occasions within a 72-hour period in which the participant
192 was fully oriented, as defined by a score ≥ 76 on the Galveston Orientation and Amnesia Test,²⁹

193 a score over 25 on the Orientation Log,³⁰ or documentation of two days with consistent
194 orientation within a three day period in the acute medical record with no intervening days at less
195 than full orientation. For the 1442 individuals discharged before emerging from PTA, missing
196 values were imputed by using total length of stay (acute plus rehabilitation)+ 1 day.³¹ Injury
197 severity categories, based on the Mississippi PTA Intervals, were: moderate (0 to 14 days);
198 moderate-severe (15 to 28 days); severe (29 to 70 days); and extremely severe (>70 days).³²

199 The FIM™ is an 18-item rating scale of functional independence.³³ Rasch analysis has
200 indicated that items can be divided into a motor factor ranging from 13 to 91 and a cognitive
201 factor ranging from 5 to 35.^{34,35} Each item is rated on a scale of 1 (total assistance required) to 7
202 (complete independence), and higher scores indicate greater independence. FIM™ has good
203 internal consistency (Cronbach's α between .86 and .97) and has been shown to be sensitive to
204 changes in functional ability from admission to discharge and follow-up.^{36, 37}

205 History of problem substance use was determined by questions adapted from the Centers
206 for Disease Control and Prevention's Risk Factor Surveillance System.³⁸ These questions pertain
207 to frequency of alcohol consumption and average quantity consumed per occasion. Using
208 established criteria,^{38,39} participants were classified as having a history of problem substance use
209 if they endorsed more than 7 drinks per week for women, more than 14 drinks per week for men,
210 or had consumed more than 5 or more drinks on one occasion in the month prior to injury or had
211 used illicit drugs in the year before injury.

212 Residence at the time of rehabilitation discharge was categorized as private or non-
213 private (nursing home, adult home, correctional institution, hotel/motel, homeless, hospital,
214 subacute care, other). Job stability was defined as the number of weeks worked in the year prior
215 to injury. Cause of injury was classified as violent (gunshot wound; assault with blunt

216 instrument; stabbing; impalement; explosions) or non-violent (vehicular; sports-related; falls;
217 auto-pedestrian; hit by falling or flying object). Sex and pre-injury marital status, education, and
218 incarcerations were categorized as shown in Table 1.

219 **Data Analysis**

220 The demographic and injury characteristics of the sample were summarized separately
221 for each of the three race/ethnicity groups using means and standard deviations for continuous
222 variables and frequency counts and percentages for categorical variables. These characteristics
223 were compared between the race/ethnicity groups using chi-square tests and ANOVA models.

224 The probability of being retained was initially modeled as a function of race/ethnicity
225 using logistic regression unadjusted for other patient characteristics. Multivariate logistic
226 regression was then used to model the relationship between race/ethnicity and retention status
227 controlling for 12 patient characteristics that may impact retention in longitudinal studies (age,⁴¹
228 gender,⁴² marital status,⁴² education,⁴¹ residence at rehabilitation discharge,²¹ pre-injury
229 incarceration, problem substance use,^{21,23} violent cause of injury,^{21,23} PTA, FIM™ scores at
230 rehabilitation discharge,²¹ and job stability for the year prior to injury²³). The assumption of
231 linearity in the logit was assessed for all continuous variables and was found to be adequate.
232 Significant interactions between race/ethnicity and patient characteristics were also examined
233 and included in the final adjusted model if significant. All statistical analyses were conducted
234 using SAS v.9.4 with a significance level of 0.05. Significant interactions were investigated
235 using a Bonferroni correction for the level of significance, as shown in Table 3.

236 **RESULTS**

237 **Description of the Sample**

238 The demographic and injury characteristics of the sample are summarized by
239 race/ethnicity in Table 1. Overall, the sample was primarily White and single, with at least a high
240 school education and moderate to severe TBI. A substantial number had a history of pre-injury
241 problem substance use and most were discharged from rehabilitation to a private residence. The
242 race/ethnicity groups showed significant differences in all patient characteristics (all p 's \leq
243 0.0004) except for pre-injury problem substance use ($p = 0.9906$). Compared to Whites, Blacks
244 and Hispanics were less likely to be female, less likely to have a high school education, more
245 likely to have been incarcerated prior to injury, and more likely to have violent cause of injury.
246 Blacks were less likely to be married, and Hispanics were more likely to have less than an 8th
247 grade education.

248 **Unadjusted Relationship between Race/Ethnicity and Retention Status**

249 As shown in Table 2, there was a significant difference in retention rates among the three
250 race/ethnicity groups (chi-square = 32.5, $df = 2$, p -value < 0.0001). Retention rates were 91.8%
251 for Whites, 90.5% for Blacks, and 85.2% for Hispanics. The unadjusted odds of being retained
252 were 1.9 times greater for Whites as compared to Hispanics (p -value < 0.0001) and 1.7 times
253 greater for Blacks as compared to Hispanics (p -value = 0.0002). There was not a significant
254 difference in unadjusted retention rates between Whites and Blacks (OR = 1.18, p -value =
255 0.1239).

256 **Adjusted Relationship between Race/Ethnicity and Retention Status**

257 There was a significant interaction effect between race/ethnicity and pre-injury problem
258 substance use ($p = 0.0330$) on retention rates. Table 3 shows the effects of race/ethnicity on
259 retention status for those with and without a pre-injury history of problem substance use, as well

260 as the effect of a pre-injury history of problem substance use on retention status for each
261 race/ethnicity group. Odds ratios with a p -value less than a Bonferroni adjusted significance level
262 of $\alpha = 0.05/9 = 0.0056$ were considered significant. For subjects without a history of pre-injury
263 problem substance use, the odds of being retained were 2.09 times greater for Whites as
264 compared to Hispanics and 2.45 times greater for Blacks as compared to Hispanics; the odds of
265 retention did not differ between Whites and Blacks without a history of substance problem use.
266 For subjects with a history of pre-injury problem substance use, the odds of being retained did
267 not differ among the race/ethnicity groups. The odds of being retained were 1.28 greater for
268 Whites without a history of problem use as compared to those with a history and 1.81 greater for
269 Blacks without a history of problem substance use as compared to Blacks with a history. For
270 Hispanics, the odds of retention did not differ between those with and without a history of
271 problem substance use. The relationship between retention and race/ethnicity was not found to
272 depend significantly on any of the other examined covariates.

273 **Adjusted Relationship between Covariates and Retention Status**

274 There was a significant relationship between retention status and age ($p = 0.0011$),
275 education ($p < 0.0001$), residence ($p = 0.0019$), and violent cause of injury ($p = 0.0006$). As
276 shown in Table 4, the odds of being retained were 0.99 times lower for each year increase in age
277 at injury, 1.44 times greater for those discharged to a private versus non-private residence, and
278 1.57 times greater for those with injuries due to non-violent causes. Furthermore, increases in
279 levels of pre-injury education were associated with increases in the odds of retention. No other
280 variables were associated with retention status.

281 **DISCUSSION**

282 The findings are consistent with prior studies that have shown lower retention of
283 minorities in TBI outcomes research;^{21,22} however, the results are unique in showing that
284 Hispanics are less likely to be retained than Whites or Blacks, with retention rates being similar
285 for Whites and Blacks. The findings emphasize the importance of investigating Blacks and
286 Hispanics separately, rather than combining them or grouping either with other races or
287 ethnicities, when investigating retention in longitudinal rehabilitation research. This would
288 increase the probability of study samples accurately reflecting the broader population, as
289 Hispanics are currently the largest minority group in the United States. The results justify efforts
290 to facilitate Hispanics' participation in research through targeted retention strategies. A unique
291 finding is that pre-injury problem substance use interacts with race/ethnicity. Hispanics did not
292 differ from Whites or Blacks in the group with pre-injury problem substance use. Problem
293 substance use history was associated with a slight decrease in retention rates for Whites and a
294 more substantial decrease for Blacks, while no decrease was noted for Hispanics. It is possible
295 that sociocultural factors associated with Hispanic race/ethnicity impact retention in research to
296 the extent that problem substance abuse does not have any additive predictive value. Such a
297 hypothesis could be investigated in future research.

298 Findings are consistent with previous research that showed a lower retention rate for
299 persons with TBI injured by violent means^{21,23} and those discharged to an institution versus a
300 private residence.²¹ In addition, older persons and those with lower education were less likely to
301 be retained. These variables were predictive of lower retention regardless of race/ethnicity and
302 can be used to target retention strategies.

303 Factors influencing retention of study participants may be participant-specific or study-
304 specific. Public health studies report a lesser likelihood of study retention for males and those

305 with multiple comorbidities,⁴³ persons with low income,⁴⁴ and immigrants.⁴⁵ Our results showed
306 no relationship between sex and retention, but the other factors were not included in our study
307 and may have impacted retention, particularly for our Hispanic participants. Relocation to their
308 country of origin is common among Hispanic research participants at some centers included in
309 this analysis and may have influenced retention. Study-specific factors that might have
310 influenced lower retention of Hispanic participants include cultural and linguistic barriers
311 between research staff and participants, inexperience of data collectors with the Hispanic
312 population, and few bilingual data collectors. In recent years, the TBIMS national data and
313 statistical center has implemented procedures to increase cultural competence of data collectors
314 and investigators, including training in cultural sensitivity. However, this may not substitute for
315 in-person contact with a bilingual research staff member and/or person of similar race/ethnicity.

316 Retaining participants from minority groups in rehabilitation research has been
317 recognized as challenging.^{20,46} Creative recruitment/retention strategies that focus on cultural
318 factors, language preferences, and community resources are needed to maximize retention. To
319 enhance retention of U.S. born and non-U.S. born Hispanic participants, acknowledging cultural
320 values of *familismo* (importance of family), *personalismo* (building rapport or personal
321 connection), *confianza* (being trustworthy), and *respeto* (being respectful) are key to conducting
322 culturally competent research.⁴⁴ Employing research staff from the same cultural and linguistic
323 background as participants can increase rapport, reduce mistrust, and increase comfort with
324 discussing sensitive information. These strategies have been shown to increase Hispanics’
325 satisfaction with and motivation to participate in psychological research.⁴⁷ Community
326 partnerships can also be effective for recruiting and retaining minority groups. Hispanic research
327 participants referred by community agencies/activities have been shown to have greater

328 engagement and study completion.⁴⁸ The researcher's connection with community-based
329 organizations familiar to Hispanic participants fosters trust and motivates consistency in
330 research involvement through social networking.⁴⁸

331 Making research participation convenient and less burdensome can increase retention.
332 Transportation can be a major problem for persons with TBI,⁴⁹ and this problem can be
333 exacerbated for newer immigrants and persons with low income. Compensating participants for
334 the cost of transportation and parking may increase engagement and retention. In addition,
335 offering follow-up outside normal work hours could facilitate participation by those who work in
336 industries with irregular work hours.

337 **Study Limitations**

338 This study assessed the impact of race/ethnicity on study retention among individuals
339 who received inpatient rehabilitation following primarily moderate-to-severe TBI and were
340 enrolled in the TBIMS National Database. Findings may differ among individuals with mild
341 TBI, veterans with TBI, and those with moderate-to-severe TBI who received acute care but not
342 inpatient rehabilitation. Analyses were also limited to variables available in the TBIMS database
343 during the study period. Retention was defined as being followed at a specific time point (1 or 2
344 years post-injury). This study was also limited by using a combined race/ethnicity variable, not
345 allowing for distinctions between White Hispanics and Black Hispanics. Race and ethnicity are
346 coded separately in other federally funded databases. The TBIMS has recently changed its
347 coding to reflect this, although not for the period covered by current analyses. We also
348 acknowledge that there is a plethora of environmental and sociopolitical factors that are
349 associated with race/ethnicity and that may impact retention in longitudinal rehabilitation

350 research. These factors were not quantified in this retrospective database study, but are important
351 to consider for future prospective studies.

352 **Conclusions**

353 Lower retention of Hispanic participants in TBI research can bias outcomes and threaten
354 external validity. Researchers should implement strategies to improve retention of Hispanic
355 participants in TBI research. Other variables, including primary language spoken, acculturation,
356 citizenship or visa status, country of residence at time of injury, and proximity of residence to
357 rehabilitation hospital, may contribute to retention and should be investigated in future studies.
358 Future research should examine whether longitudinal patterns of retention differ for Hispanics
359 compared to Blacks and Whites.

360

361 References

362 1. Benedictus MR, Spikman JM, Naalt JVD. Cognitive and behavioral impairment in
363 traumatic brain injury related to outcome and return to work. *Arch Phys Med Rehabil*
364 2010;91(9):1436-1441.
365

366 2. Arciniegas DB, Wortzel HS. Emotional and behavioral dyscontrol after traumatic brain
367 injury. *Psych Clinics North America* 2014;37(1):31-53.
368

369 3. Masel BF, Dewitt DS. Traumatic brain injury: A disease process, not an event. *J*
370 *Neurotrauma* 2010;27(8):1529-1540.
371

372 4. Andelic N, Sigurdardottir S, Schanke A-K, Sandvik L, Sveen U, Roe C. Disability,
373 physical health and mental health 1 year after traumatic brain injury. *Disability and*
374 *Rehabil* 2010;32:1122-1131.
375

376 5. Grauwmeijer E, Heijenbrok-Kal MH, Haitisma IK, Ribbers GM. A prospective study on
377 employment outcome 3 years after moderate to severe traumatic brain injury. *Arch Phys*
378 *Med Rehabil* 2012;93:993-999.
379

380 6. Bier N, Dutil E, Couture M. Factors affecting leisure participation after a traumatic brain
381 injury: An exploratory study. *J Head Trauma Rehabil* 2009;24:187-194.
382

383 7. Corrigan JD, Selassie AW, Orman JA. The epidemiology of traumatic brain injury. *J*
384 *Head Trauma Rehabil* 2010;25:72-80.
385

386 8. Coronado VG, McGuire LC, Sarmiento K, et al. Trends in traumatic brain injury in the
387 U.S. and the public health response: 1995-2009. *J Safety Res* 2012;43:299-307.
388

389 9. Gary KW, Arango-Lasprilla J, Stevens LF. Do racial/ethnic differences exist in post-
390 injury outcomes after TBI? A comprehensive review of the literature. *Brain Inj*
391 2009;23:775-789.
392

393 10. Shafi S, Marquez de la Plata C, Diaz-Arrastia R, Shipman K, Carlile M, Frankel H, Parks
394 J, Gentilello LM. Racial disparities in long-term functional outcome after traumatic brain
395 injury. *J Trauma* 2007;63:1263-1268.
396

397 11. Arango-Lasprilla JC, Rosenthal M, Deluca J, Komaroff E, Sherer M, Cifu D, Hanks R.
398 Traumatic brain injury and functional outcomes: Does minority status matter? *Brain Inj*
399 2007;21:701-708.
400

401 12. Sander AM, Pappadis MR, Davis LC, Clark AN, Evans G, Struchen MA, Mazzei DK.
402 Relationship of race/ethnicity and income to community integration following traumatic
403 brain injury: Investigation in a non-rehabilitation trauma sample. *NeuroRehabilitation*
404 2009;24:15-27.
405


- 406 13. Arango-Lasprilla JC, Ketchum JM, Lewis AN, Krch D, Gary KW, Dodd BA. Racial and
407 ethnic disparities in employment outcomes for persons with traumatic brain injury: A
408 longitudinal investigation 1-5 years after injury. *PM&R* 2001;3:1083-1091.
409
- 410 14. Jamison L, Kolakowsky-Hayner SA, Wright J. Preliminary investigation of longitudinal
411 sociodemographic, injury and psychosocial characteristics in a group of non-English
412 speaking Latinos with brain injury. *Brain Injury* 2012;26:805-813.
413
- 414 15. Mascialino G, Hirshson C, Egan M, Cantor J, Ashman T, Tsaousides T, Spielman L.
415 Objective and subjective assessment of long-term community integration in minority
416 groups following traumatic brain injury. *NeuroRehabilitation* 2009;24:29-36.
417
- 418 16. Asemota AO, George BP, Cumpsty-Fowler CJ, Haider AH, Schneider EB. Race and
419 insurance disparities in discharge to rehabilitation for patients with traumatic brain injury.
420 *J Neurotrauma* 2013;30:2057-2065.
421
- 422 17. Dismuke CE, Gebregziabher M, Yeager D, Egede LE. Racial/ethnic differences in VA
423 services utilization as a partial pathway to mortality differences among veterans
424 diagnosed with TBI. *Glob Health Sci* 2016;8:260-272.
425
- 426 18. Ejiogu N, Norbeck JH, Mason MA, Cromwell BC, Zonderman AB, Evans MK.
427 Recruitment and retention strategies for minority or poor clinical research participants:
428 Lessons from the Healthy Aging in Neighborhoods of Diversity Across the Life Span
429 Study. *The Gerontologist* 2011;51(Supplement 1).
430
- 431 19. Carroll JK, Yancey AK, Spring B, et al. What are successful recruitment and retention
432 strategies for underserved populations? Examining physical activity interventions in
433 primary care and community settings. *Translational Behavioral Medicine* 2011;1:234-
434 251.
435
- 436 20. Bell KR, Hammond F, Hart T, Bickett AK, Temkin NR, Dikmen S. Participant
437 recruitment and retention in rehabilitation research. *Am J Phys Med Rehabil*
438 2008;87(4):330-8.
439
- 440 21. Corrigan JD, Harrison-Felix C, Bogner J, Dijkers M, Terrill MS, Whiteneck G.
441 Systematic bias in traumatic brain injury outcome studies because of loss to follow-up.
442 *Arch Phys Med Rehabil* 2003;84(2):153-60.
443
- 444 22. Krellman JW, Kolakowsky-Hayner SA, Spielman L, Dijkers M, Hammond FM, Bogner
445 J, Hart T, Cantor JB, Tsaousides T. *Arch Phys Med Rehabil* 2014;95:633-641.
446
- 447 23. Jourdan C, Bayen E, Bahrami S, Ghout I, Darnoux E, Azerad S, Ruet A, Vallat-Azouvi
448 C, Weis J, Aegerter P, Mateo J, Vigue B, Tazarourte K, Pradat-Diehl P, Azouvi P. Loss
449 to follow-up and social background in an inception cohort of patients with severe
450 traumatic brain injury: Results from the PariS-TBI study. *J Head Trauma Rehabil*
451 2016;31:E42-E48.

- 452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
24. Annual Estimates of the Resident Population by Sex, Age, Race, and Hispanic Origin for the United States and States: April 1, 2010 to July 1, 2015 Source: U.S. Census Bureau, Population Division Release Date: June 2016.
 25. Traumatic Brain Injury Model Systems National Data and Statistical Center. (2016). *National database: 2015 profile of people within the Traumatic Brain Injury Model Systems* (Grant No. H133A110006 and 90DP0012). Englewood, CO: Craig Hospital.
 26. Asemota AO, George BP, Cumpsty-Fowler CJ, Haider AH, Schneider EB. Race and insurance disparities in discharge to rehabilitation for patients with traumatic brain injury. *J Neurotrauma* 2013;30:2057-2065.
 27. Ketchum JM, Getachew AM, Krch D, Banos JH, Kolakowsky-Hayner SA, Lequerica A, Jamison L, Arango-Lasprilla JC. Early predictors of employment outcomes 1 years post traumatic brain injury in a population of Hispanic individuals. *Neurorehabilitation* 2012;30:13-22.
 28. Walker WC, Ketchum JM, Marwitz JH, et al. A multicenter study on the clinical utility of post-traumatic amnesia duration in predicting global outcome after moderate-severe traumatic brain injury. *J Neurol Neurosurg Psychiatry* 2010;81:87e9.
 29. Levin HS, O'Donnell VM, Grossman RG. The Galveston Orientation and Amnesia Test. A practical scale to assess cognition after head injury. *J Nerv Ment Dis.* 1979;167:675–684.
 30. Jackson WT, Novack TA, Dowler RN. Effective serial measurement of cognitive orientation in rehabilitation: the Orientation Log. *Arch Phys Med Rehabil.* 1998;79:718–720.
 31. Nakase-Richardson R, Yablon SA, Sherer M. Prospective comparison of acute confusion severity with duration of post-traumatic amnesia in predicting employment outcome after traumatic brain injury. *J Neurol Neurosurg Psychiatry* 2007;78:872-876.
 32. Nakase-Richardson R, Sepehri A, Sherer M, Yablon SA, Evans C, Mani T. Classification schema of posttraumatic amnesia duration-based injury severity relative to 1-year outcome: analysis of individuals with moderate and severe traumatic brain injury. *Arch Phys Med Rehabil* 2009;90:17-19.
 33. Keith, R.A., Granger, C.V., Hamilton, B.B., and Sherwin, F.S., The functional independence measure: a new tool for rehabilitation. *Adv Clin Rehabil*, 1987. 1: p. 6-18.
 34. Hall KM, Hamilton BB, Gordon WA, Zasler ND. Characteristics and comparisons of functional assessment indices: Disability Rating Scale, Functional Independence Measure, and Functional Assessment Measure. *J Head Trauma Rehabil* 1993;8(2):60-74.

- 498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
35. Linacre JM, Heinemann AW, Wright BD, Granger CV, Hamilton BB. The structure and stability of the Functional Independence Measure. *Arch Phys Rehabil* 1994;75:127-132.
 36. Dodds, T.A., Martin, D.P., Stolov, W.C., and Deyo, R.A., A validation of the functional independence measurement and its performance among rehabilitation inpatients. *Arch Phys Med Rehabil* 1993. 74(5): p. 531-6.
 37. Ottenbacher, K.J., Hsu, Y., Granger, C.V., and Fiedler, R.C., The reliability of the functional independence measure: a quantitative review. *Arch Phys Med Rehabil* 1996. 77(12): p. 1226-32.
 38. Centers on Disease Control and Prevention. Behavioral Risk Factor Surveillance System User's Guide. Report. Atlanta, GA:US Department of Health and Human Services, Centers for Disease Control and Prevention, 1998.
 39. Substance Abuse and Mental Health Services Administration, Office of Applied Studies. National Household Survey on Drug Abuse: Population Estimates, 1997. Rockville, MD: US Department of Health and Human Services, 1998.
 40. Traumatic Brain Injury Model Systems National Database Syllabus. Traumatic Brain Injury Model Systems National Data and Statistical Center; 2017.
[URL:http://www.tbindsc.org](http://www.tbindsc.org).
 41. Bharna S, Tinker A, Ashcroft R, Askham J. The retention of older people in longitudinal studies: A review of the literature. *Quality in Ageing and Older Adults* 2008;9:27-35.
 42. Radler BT, Ryff CD. Who participates? Longitudinal retention in the MIDUS National Study of Health and Well-Being. *J Aging Health* 2010;22:307-331.
 43. Eakin EG, Bull SS, Riley K, Reeves MM, Gutierrez S, McLaughlin P. Recruitment and retention of Latinos in a primary care-based physical activity and diet trial: The Resources for Health study. *Health Educ Res* 2007;22(3):361-71.
 44. Ojeda L, Flores LY, Meza RR, Morales A. Culturally competent qualitative research with Latino immigrants. *Hisp J Behav Sci* 2011; 33(2): 184-203.
 45. Martinez Jr CR, McClure HH, Eddy JM, Ruth B, Hyers MJ. Recruitment and retention of Latino immigrant families in prevention research. *Prev Sci* 2012;13(1):15-26.
 46. Paskett ED, Reeves KW, McLaughlin JM, Katz ML, McAlearney AS, Ruffin MT, Halbert CH, Merete C, Davis F, Gehlert S. Recruitment of minority and underserved populations in the United States: the Centers for Population Health and Health Disparities experience. *Contemp Clin Trials* 2008;29:847-861.

- 543 47. Haack LM, Gerdes AC, Lawton KE. Conducting research with Latino families:
544 Examination of strategies to improve recruitment, retention, and satisfaction with an at-
545 risk and underserved population. *J Child Fam Stud* 2014;23:410-421.
546
- 547 48. Sankaré I C, Bross R, Brown AF, del Pino HE, Jones LF, Morris D, Porter C, Lucas-
548 Wright A, Forge N, Norris KC, Kahn KL. Strategies to build trust and recruit African
549 American and Latino community residents for health research: a cohort study. *Clin Trans*
550 *Sci* 2015;8: 412-420.
551
- 552 49. Fleming J, Nalder E, Alves-Stein S, Cornwell P. The effect of environmental barriers on
553 community integration for individuals with moderate to severe traumatic brain injury. *J*
554 *Head Trauma Rehabil* 2014;29:125-135.
555
- 556

Figure 1



*Traumatic Brain Injury Model Systems centers are funded for 5-year periods and then must compete for another funding cycle. This sometimes results in existing centers losing funding for one or more cycles, and thus participants in the national database may not be followed due to lack of funds.

Table 1: Summary of Sample Characteristics by Race/Ethnicity

	White (N = 5548)	Black (N = 1347)	Hispanic (N = 790)	Total Count	p-value
Sex					< 0.0001
Male	3991 (71.9%)	1049 (77.9%)	613 (77.6%)	5653	
Female	1557 (28.1%)	297 (22.1%)	177 (22.4%)	2031	
Pre-Injury Marital Status					< 0.0001
Married	2063 (37.2%)	267 (19.9%)	217 (27.5%)	2547	
Not Married	3485 (62.8%)	1078 (80.1%)	571 (72.5%)	5134	
Pre-Injury Education					< 0.0001
≤ 8 th Grade	186 (3.4%)	84 (6.3%)	196 (25.0%)	466	
9 th – 11 th Grade	844 (15.3%)	366 (27.4%)	199 (25.4%)	1409	
12 th Grade (HS/GED)	2074 (37.7%)	523 (39.1%)	216 (27.6%)	2813	
> 12 th Grade	2411 (43.6%)	366 (27.3%)	172 (22.1%)	2950	
Residence at Discharge*					0.0004
Private Residence	4549 (82.1%)	1098 (81.8%)	691 (87.7%)	6338	
Non-Private Residence	989 (17.9%)	244 (18.2%)	97 (12.3%)	1330	
Pre-Injury Penal Incarcerations					< 0.0001
No	5081 (92.9%)	1080 (81.0%)	691 (89.6%)	6852	
Yes	390 (7.1%)	254 (19.0%)	80 (10.4%)	724	
Pre-Injury Substance Problem Use					0.9906
No	3061 (57.8%)	753 (57.9%)	426 (57.6%)	4240	
Yes	2236 (42.2%)	548 (42.1%)	314 (42.4%)	3098	
PTA Group					< 0.0001
Moderate	1852 (34.24%)	384 (29.20%)	207 (27.79%)	2443	
Moderate/Severe	1200 (22.19%)	274 (20.84%)	150 (20.13%)	1624	
Severe	1247 (23.05%)	321 (24.41%)	183 (24.56%)	1751	
Extremely Severe	1110 (20.52%)	336 (25.55%)	205 (27.52%)	1651	
Cause of Injury					< 0.0001
Violent	336 (6.1%)	318 (23.6%)	136 (17.3%)	790	
Not Violent	5204 (93.9%)	1029 (76.4%)	651 (82.7%)	6884	
	Mean (SD)	Mean (SD)	Mean (SD)		
Age	42.82(19.96)	38.58(17.03)	35.86(17.56)		<0.0001
FIM TM Motor at Discharge	66.44(18.24)	63.22(17.86)	66.02 (17.53)		<0.0001
FIM TM Cognitive at Discharge	24.00(6.72)	22.80(6.63)	23.40(6.82)		<.00001
Job Stability**	29.87 (24.15)	23.41(24.08)	30.76(23.77)		<0.0001

* Non-private residence=nursing home, adult home, correctional institution, hotel/motel, homeless, hospital, subacute care, or other

**number of weeks worked in the year prior to injury, modeled as a continuous variable

Table 2: Differences in Retention Rates Between Whites, Blacks, and Hispanics

	Retained		Total
	No	Yes	
White	455 (8.2%)	5093 (91.8%)	5548
Black	128 (9.5%)	1219 (90.5%)	1347
Hispanic	117 (14.8%)	673 (85.2%)	790
Total	700 (9.1%)	6985 (90.9%)	7075

Table 3: Odds Ratios Comparing Race/Ethnicity and Pre-Injury Problem Substance Use Groups from Adjusted[§] Model

History of Pre-Injury Problem Substance Use	Race/Ethnicity	OR [†]	95% CI	<i>p</i> -value
No	White vs. Black	0.853	(0.607, 1.200)	0.3608
	White vs. Hispanic	2.091	(1.489, 2.936)	< 0.0001 ‡
	Black vs. Hispanic	2.451	(1.607, 3.739)	< 0.0001 ‡
Yes	White vs. Black	1.212	(0.883, 1.666)	0.2347
	White vs. Hispanic	1.341	(0.904, 1.989)	0.1453
	Black vs. Hispanic	1.106	(0.710, 1.723)	0.6567
No vs. Yes	White	1.275	(1.012, 1.606)	0.0388
No vs. Yes	Black	1.812	(1.211, 2.712)	0.0038 ‡
No vs. Yes	Hispanic	0.818	(0.518, 1.290)	0.3866

[§]Model variables include race/ethnicity, age, gender, marital status, education, residence at rehabilitation discharge, pre-injury incarceration, problem substance use, violent cause of injury, PTA, discharge FIM, pre-injury job stability, and race/ethnic by problem substance use

[†] Odds ratios represent the odds of being retained versus not retained for one sub group versus another subgroup

[‡] Significant at a Bonferroni corrected level of significance $\alpha = 0.05/9 = 0.0056$

Table 4: Odds Ratios Comparing Covariate Subgroups from Adjusted[§] Model

Variable	Comparison	OR [†]	95% CI	<i>p</i> -value
Age	1 year increase	0.991	(0.986, 0.997)	0.0011
Sex	Female vs. Male	1.193	(0.963, 1.478)	0.1070
Marital Status	Married vs. Not Married	1.232	(0.999, 1.519)	0.0512
Education	> 12 th Grade vs. ≤ 8 th Grade	2.832	(2.042, 3.927)	< 0.0001
	9 th – 11 th Grade vs. ≤ 8 th Grade	1.638	(1.177, 2.279)	0.0034
	12 th Grade (HS/GED) vs. ≤ 8 th Grade	1.739	(1.279, 2.365)	0.0004
	> 12 th Grade vs. 9 th – 11 th Grade	1.730	(1.329, 2.250)	< 0.0001
	12 th Grade (HS/GED) vs. 9 th – 11 th Grade	1.062	(0.838, 1.347)	0.6195
	> 12 th Grade vs. 12 th Grade (HS/GED)	1.628	(1.309, 2.026)	< 0.0001
Residence	Private vs. Not Private	1.443	(1.145, 1.818)	0.0019
Incarcerated	No vs. Yes	1.013	(0.767, 1.338)	0.9275
PTA Group	Moderate vs. Moderate/Severe	0.855	(0.670, 1.091)	0.2082
	Moderate vs. Severe	0.813	(0.634, 1.042)	0.1019
	Moderate vs. Extremely Severe	0.878	(0.649, 1.189)	0.4001
	Moderate/Severe vs. Severe	0.951	(0.727, 1.243)	0.7112
	Moderate/Severe vs. Extremely Severe	1.027	(0.753, 1.402)	0.8664
	Severe vs. Extremely Severe	1.081	(0.807, 1.447)	0.6038
FIM Motor	1 unit increase	0.998	(0.992, 1.005)	0.6195
FIM Cognitive	1 unit increase	0.991	(0.973, 1.009)	0.3050
Job Stability	1 week increase	1.003	(0.999, 1.007)	0.1170
Violent Injury	No vs. Yes	1.565	(1.212, 2.021)	0.0006

[§]Model variables include race/ethnicity, age, gender, marital status, education, residence at rehabilitation discharge, pre-injury incarceration, problem substance use, violent cause of injury, PTA, discharge FIM, pre-injury job stability, and race/ethnic by problem substance use

[†] Odds ratios represent the odds of being retained versus not retained for one subgroup versus another subgroup