

# On existence of an $x$ -integral for a semi-discrete chain of hyperbolic type

K Zheltukhin<sup>1</sup> and N Zheltukhina<sup>2</sup>

<sup>1</sup> Department of Mathematics, Middle East Technical University, Ankara, Turkey

<sup>2</sup> Department of Mathematics, Bilkent University, Ankara, Turkey

E-mail: zheltukh@metu.edu.tr

**Abstract.** A class of semi-discrete chains of the form  $t_{1x} = f(x, t, t_1, t_x)$  is considered. For the given chains easily verifiable conditions for existence of  $x$ -integral of minimal order 4 are obtained.

## 1. Introduction

In the present paper we consider the integrable differential-difference chains of hyperbolic type

$$t_{1x} = f(x, t, t_1, t_x), \quad (1)$$

where the function  $t(n, x)$  depends on discrete variable  $n$  and continuous variable  $x$ . We use the following notations  $t_x = \frac{\partial}{\partial x}t$  and  $t_1 = t(n+1, x)$ . It is also convenient to denote  $t_{[k]} = \frac{\partial^k}{\partial x^k}t$ ,  $k \in \mathbb{N}$  and  $t_m = t(n+m, x)$ ,  $m \in \mathbb{Z}$ .

The integrability of the chain (1) is understood as Darboux integrability that is existence of so called  $x$ - and  $n$ -integrals [1, 4]. Let us give the necessary definitions.

**Definition 1** Function  $F(x, t, t_1, \dots, t_k)$  is called an  $x$ -integral of the equation (1) if

$$D_x F(x, t, t_1, \dots, t_k) = 0$$

for all solutions of (1). The operator  $D_x$  is the total derivative with respect to  $x$ .

**Definition 2** Function  $G(x, t, t_x, \dots, t_{[m]})$  is called an  $n$ -integral of the equation (1) if

$$DG(x, t, t_x, \dots, t_{[m]}) = G(x, t, t_x, \dots, t_{[m]})$$

for all solutions of (1). The operator  $D$  is a shift operator.

To show the existence of  $x$ - and  $n$ -integrals we can use the notion of characteristic ring. The notion of characteristic ring was introduced by Shabat to study hyperbolic systems of exponential type (see [11]). This approach turns out to be very convenient to study and classify the integrable equations of hyperbolic type (see [12] and references there in).

For difference and differential-difference chains the notion of characteristic ring was developed by Habibullin (see [3]-[8]). In particular, in [4] the following theorem was proved



**Theorem 3** (see [4]). *A chain (1) admits a non-trivial  $x$ -integral if and only if its characteristic  $x$ -ring is of finite dimension.*

*A chain (1) admits a non-trivial  $n$ -integral if and only if its characteristic  $n$ -ring is of finite dimension.*

For known examples of integrable chains the dimension of the characteristic ring is small. The differential-difference chains with three dimensional characteristic  $x$ -ring were considered in [6]. We consider chains with four dimensional characteristic  $x$ -ring, such chains admit  $x$ -integral of minimal order four. That is we obtain necessary and sufficient conditions for a chain to have a four dimensional characteristic  $x$ -ring. This conditions can be easily checked by direct calculations.

Note that if a chain (1) admits a nontrivial  $x$ -integral  $F(x, t, t_1, \dots, t_k)$  and a non trivial  $n$ -integral  $G(x, t, t_x, \dots, t_{[m]})$  its solutions satisfy two ordinary equations

$$F(x, t, t_1, \dots, t_k) = a(n),$$

$$G(x, t, t_x, \dots, t_{[m]}) = b(x)$$

for some functions  $a(n)$  and  $b(x)$ . This allows to solve (1) (see [9]).

The paper is organized as follows. In Section 2 we derive necessary and sufficient conditions on function  $f(x, t, t_1, t_x)$  so that the chain (1) has four dimensional characteristic ring and in Section 3 we consider some applications of the derived conditions.

## 2. Chains admitting four dimensional $x$ -algebra.

Suppose  $F$  is an  $x$ -integral of the chain (1) then its positive shifts and negative shifts  $D^k F$ ,  $k \in \mathbb{Z}$ , are also  $x$ -integrals. So, looking for an  $x$ -integral it is convenient to assume that it depends on positive and negative shifts of  $t$ .

To express  $x$  derivatives of negative shifts we can apply  $D^{-1}$  to the chain (1) and obtain

$$t_x = f(x, t_{-1}, t, t_x).$$

Solving the above equation for  $t_{-1x}$  we get

$$t_{-1x} = g(x, t_{-1}, t, t_x).$$

Let  $F(x, t, t_1, t_{-1}, \dots)$  be an  $x$ -integral of the chain (1). Then on solutions of (1) we have

$$D_x F = \frac{\partial F}{\partial x} + t_x \frac{\partial F}{\partial t} + t_{1x} \frac{\partial F}{\partial t_1} + t_{-1x} \frac{\partial F}{\partial t_{-1}} + t_{2x} \frac{\partial F}{\partial t_2} + t_{-2x} \frac{\partial F}{\partial t_{-2}} + \dots = 0$$

or

$$D_x F = \frac{\partial F}{\partial x} + t_x \frac{\partial F}{\partial t} + f \frac{\partial F}{\partial t_1} + g \frac{\partial F}{\partial t_{-1}} + Df \frac{\partial F}{\partial t_2} + D^{-1}g \frac{\partial F}{\partial t_{-2}} + \dots = 0.$$

Define a vector field

$$K = \frac{\partial}{\partial x} + t_x \frac{\partial}{\partial t} + f \frac{\partial}{\partial t_1} + g \frac{\partial}{\partial t_{-1}} + Df \frac{\partial}{\partial t_2} + D^{-1}g \frac{\partial}{\partial t_{-2}} + \dots, \quad (2)$$

then

$$D_x F = K F.$$

Note that  $F$  does not depend on  $t_x$  but the coefficients of  $K$  do depend on  $t_x$ . So we introduce a vector field

$$X = \frac{\partial}{\partial t_x} \quad (3)$$

The vector fields  $K$  and  $X$  generate the characteristic  $x$ -ring  $L_x$ .

Let us introduce some other vector fields from  $L_x$ .

$$C_1 = [X, K] \quad \text{and} \quad C_n = [X, C_{n-1}] \quad n = 2, 3, \dots \quad (4)$$

and

$$Z_1 = [K, C_1] \quad \text{and} \quad Z_n = [K, Z_{n-1}] \quad n = 2, 3, \dots \quad (5)$$

Thus

$$C_1 = \frac{\partial}{\partial t} + f_{tx} \frac{\partial}{\partial t_1} + g_{tx} \frac{\partial}{\partial t_{-1}} + \dots$$

$$C_2 = f_{tx} \frac{\partial}{\partial t_1} + g_{tx} \frac{\partial}{\partial t_{-1}} + \dots$$

$$Z_1 = (f_{t_{xx}} + t_x f_{t_{xt}} + f f_{t_{xt_1}} - f_t - f_{tx} f_{t_1}) \frac{\partial}{\partial t_1} + (g_{t_{xx}} + t_x g_{t_{xt}} + g g_{t_{xt_1}} - g_t - g_{tx} g_{t_1}) \frac{\partial}{\partial t_{-1}} + \dots$$

and so on.

It is easy to see that if  $f_{t_{xx}} \neq 0$  then the vector fields  $X, K, C_1$  and  $C_2$  are linearly independent and must form a basis of  $L_x$  provided  $\dim L_x = 4$ . By Lemma 3.6 in [6], if  $f_{t_{xx}} = 0$  and  $(f_{t_{xx}} + t_x f_{t_{xt}} + f f_{t_{xt_1}} - f_t - f_{tx} f_{t_1}) = 0$  then  $\dim L_x = 3$ . So in the case  $f_{t_{xx}} = 0$  we may assume  $(f_{t_{xx}} + t_x f_{t_{xt}} + f f_{t_{xt_1}} - f_t - f_{tx} f_{t_1}) \neq 0$ . Then the vector fields  $X, K, C_1$  and  $Z_1$  are linearly independent and must form a basis of  $L_x$  provided  $\dim L_x = 4$ . We consider this two cases separately.

In the rest of the paper we assume that the characteristic ring  $L_x$  is four dimensional.

**Remark 4** *It is convenient to check equalities between vector fields using the automorphism  $D(\ )D^{-1}$ . Direct calculations show that*

$$DXD^{-1} = \frac{1}{f_x} X,$$

$$DKD^{-1} = K - \frac{f_x + t_x f_t + f f_{t_1}}{f_{tx}} X.$$

*The images of other vector fields under this automorphism can be obtained by commuting  $DXD^{-1}$  and  $DKD^{-1}$ .*

2.1.  $f(x, t, t_1, t_x)$  is non linear with respect to  $t_x$ .

Let  $f(x, t, t_1, t_x)$  be non linear with respect to  $t_x$ ,  $f_{t_{xx}} \neq 0$ . Then the vector fields  $X, K, C_1$  and  $C_2$  form a basis of  $L_x$ . For the algebra  $L_x$  to be spanned by  $X, K, C_1$  and  $C_2$  it is enough that  $C_3$  and  $Z_1$  are linear combinations of  $X, K, C_1$  and  $C_2$ . From the form of the vector fields it follows that we must have

$$C_3 = \lambda C_2 \quad \text{and} \quad Z_1 = \mu C_2$$

for some functions  $\mu$  and  $\lambda$ . The conditions for the above equalities to hold are given by the following theorem.

**Theorem 5** *The chain (1) with  $f_{t_{xx}} \neq 0$  has characteristic ring  $L_x$  of dimension four if and only if the following conditions hold*

$$D \left( \frac{f_{t_x t_x t_x}}{f_{t_x t_x}} \right) = \frac{f_{t_x t_x t_x} f_{t_x} - 3 f_{t_x t_x}^2}{f_{t_x t_x} f_{t_x}^2}. \quad (6)$$

$$D \left( \frac{f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1} - f_t - f_{t_x} f_{t_1}}{f_{t_x t_x}} \right) = \frac{f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1} - f_t - f_{t_x} f_{t_1}}{f_{t_x t_x}} f_{t_x} - (f_x + t_x f_t + f_{t_1}). \tag{7}$$

The characteristic ring is generated by the vector fields  $X, K, C_1, C_2$ .

**Proof.** By Remark 4 we have

$$DC_2 D^{-1} = \frac{1}{f_{t_x}^2} C_2 - \frac{f_{t_x t_x}}{f_{t_x}^3} C_1 + \frac{f_{t_x t_x} f_t}{f_{t_x}^4} X$$

$$DC_3 D^{-1} = \frac{1}{f_{t_x}^3} C_2 - \frac{3f_{t_x t_x}}{f_{t_x}^4} C_2 - \frac{f_{t_x t_x t_x} f_{t_x} - 3f_{t_x t_x}^2}{f_{t_x}^5} C_1 + f_t \frac{f_{t_x t_x t_x} f_{t_x} - 3f_{t_x t_x}^2}{f_{t_x}^6} X$$

$$DZ_1 D^{-1} = \frac{1}{f_{t_x}} Z_1 - \left( \frac{m f_{t_x} + p}{f_{t_x}^2} \right) \left( C_1 - \frac{f_t}{f_{t_x}} X \right),$$

where  $p = \frac{f_x + t_x f_t + f f_{t_1}}{f_{t_x}}$  and  $m = \frac{-(f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1}) + f_t + f_{t_x} f_{t_1}}{f_{t_x}}$ .

The equality  $C_3 = \lambda C_2$  implies that

$$DC_3 D^{-1} = (D\lambda) DC_2 D^{-1}. \tag{8}$$

Substituting expressions for  $DC_2 D^{-1}$  and  $DC_3 D^{-1}$  into (8) and comparing coefficients of  $C_1, C_2$  and  $X$  we obtain that  $\lambda$  satisfies

$$\lambda = f_{t_x} (D\lambda) + \frac{3f_{t_x t_x}}{f_{t_x}}$$

$$(D\lambda) = \frac{f_{t_x t_x t_x} f_{t_x} - 3f_{t_x t_x}^2}{f_{t_x t_x} f_{t_x}^2}.$$

We can find  $\lambda$  and  $D\lambda$  independently and condition that  $D\lambda$  is a shift of  $\lambda$  leads to (6). The equality  $Z_1 = \mu C_2$  implies that

$$DZ_1 D^{-1} = (D\mu) DC_2 D^{-1}. \tag{9}$$

Substituting expressions for  $DC_2 D^{-1}$  and  $DC_3 D^{-1}$  into (9) and comparing coefficients of  $C_1, C_2$  and  $X$  we obtain that  $\mu$  satisfies

$$\mu - \frac{f_x + t_x f_t + f f_{t_1}}{f_{t_x}} = \frac{(D\mu)}{f_{t_x}}$$

and

$$-(f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1} - f_t - f_{t_x} f_{t_1}) + \frac{f_x + t_x f_t + f f_{t_1}}{f_{t_x}} f_{t_x t_x} = -\frac{f_{t_x t_x} (D\mu)}{f_{t_x}}$$

We can find  $\mu$  and  $D\mu$  independently and condition that  $D\mu$  is a shift of  $\mu$  leads to (7).  $\square$

**Remark 6** Let  $\dim L_x = 4$  and  $f_{t_x} \neq 0$ . Then the characteristic ring  $L_x$  have the following multiplication table

	$X$	$K$	$C_1$	$C_2$
$X$	0	$C_1$	$C_2$	$\mu C_2$
$K$	$-C_1$	0	$\lambda C_2$	$\rho C_2$
$C_1$	$-C_2$	$-\lambda C_2$	0	$\eta C_2$
$C_2$	$-\mu C_2$	$-\rho C_2$	$-\eta C_2$	0

where  $\rho = \lambda\mu + X(\lambda)$  and  $\eta = X(\rho) - K(\mu)$ .

**Example 7** Consider the following chain

$$t_{1x} = \frac{tt_x - \sqrt{t_x^2 - M^2}(t_1 + t)}{t_1}$$

introduced by Habibullin and Zheltukhina [10]. We can easily check that the function

$$f(t, t_1, t_x) = \frac{tt_x - \sqrt{t_x^2 - M^2}(t_1 + t)}{t_1}$$

satisfies the conditions of Theorem 5. Hence the corresponding  $x$ -algebra is four dimensional. The chain has the following  $x$ -integral

$$F = \frac{(t_1^2 - t^2)(t_1^2 - t_2^2)}{t_1^2}.$$

2.2.  $f(x, t, t_1, t_x)$  is linear with respect to  $t_x$ .

Let  $f(x, t, t_1, t_x)$  be linear with respect to  $t_x$ ,  $f_{t_x t_x} = 0$ . Then vector fields  $X, K, C_1$  and  $Z_1$  form a basis of  $L_x$ . The condition  $f_{t_x t_x} = 0$  also implies that the vector field  $C_2 = 0$ , see [6]. For the algebra  $L_x$  to be spanned by  $X, K, C_1$  and  $Z$  it is enough that  $Z_2$  is a linear combination of  $X, K, C_1$  and  $Z_1$ . From the form of the vector fields it follows that we must have

$$Z_2 = \alpha Z_1$$

for some function  $\alpha$ . The conditions for the above equality to hold given by the following theorem.

**Theorem 8** The chain (1) with  $f_{t_x t_x} = 0$  has the characteristic ring  $L_x$  of dimension four if and only if the following condition hold

$$D \left( \frac{K(m)}{m} - m + \frac{f_t}{f_{t_x}} \right) = \frac{K(m)}{m} + m - f_{t_1}. \tag{10}$$

where  $m = \frac{-(f_{xt_x} + t_x f_{t_x t} + f f_{t_x t_1}) + f_t + f_{t_x} f_{t_1}}{f_{t_x}}$ . The characteristic ring is generated by the vector fields  $X, K, C_1, Z_1$ .

**Proof.** By Remark 4 we have

$$DZ_1 D^{-1} = \frac{1}{f_{t_x}} Z_1 - \left( \frac{m f_{t_x} + p}{f_{t_x}^2} \right) \left( C_1 - \frac{f_t}{f_{t_x}} X \right),$$

and

$$DZ_2 D^{-1} = \left( K \left( \frac{1}{f_{t_x}} \right) + \frac{\alpha + m}{f_{t_x}} \right) Z_1 + \left( K \left( \frac{m}{f_{t_x}} \right) + \frac{m f_t}{f_{t_x}^2} - p X \left( \frac{m}{f_{t_x}} \right) \right) \left( C_1 - \frac{f_t}{f_{t_x}} X \right)$$

The equality  $Z_2 = \alpha Z_1$  implies that

$$DZ_2D^{-1} = (D\alpha)DZ_1D^{-1}. \tag{11}$$

Substituting expressions for  $DZ_1D^{-1}$  and  $DZ_2D^{-1}$  into (11) and comparing coefficients of  $C_1$ ,  $Z_1$  and  $X$  we obtain that  $\alpha$  and  $D(\alpha)$  satisfy

$$K \left( \frac{1}{f_{t_x}} \right) + \frac{m}{f_{t_x}} + \frac{\alpha}{f_{t_x}} = \frac{D(\alpha)}{f_{t_x}}$$

$$K \left( \frac{m}{f_{t_x}} \right) + \frac{mf_t}{f_{t_x}^2} = \frac{mD(\alpha)}{f_{t_x}}$$

We can find  $\alpha$  and  $D(\alpha)$  independently and condition that  $D(\alpha)$  is a shift of  $\alpha$  leads to (10).  $\square$

**Remark 9** Let  $\dim L_x = 4$  and  $f_{t_{xx}} = 0$ . Then the characteristic ring  $L_x$  have the following multiplication table

	$X$	$K$	$C_1$	$Z_1$
$X$	0	$C_1$	0	0
$K$	$-C_1$	0	$Z_1$	$\alpha Z_1$
$C_1$	0	$-Z_1$	0	$X(\alpha)Z_1$
$Z_1$	0	$-\alpha Z_1$	$-X(\alpha)Z_1$	0

**Example 10** Consider the following chain

$$t_{1x} = t_x + e^{\frac{t+t_1}{2}}$$

introduced by Dodd and Bullough [2]. We can easily check that the function

$$f(t, t_1, t_x) = t_x + e^{\frac{t+t_1}{2}}$$

satisfies the conditions of Theorem 8. Hence the corresponding  $x$ -algebra is four dimensional. The chain has the following  $x$ -integral

$$F = e^{\frac{t_1-t}{2}} + e^{\frac{t_1-t_2}{2}}$$

### 3. Applications

The conditions derived in the previous section can be used to determine some restrictions on the form of the function  $f(x, t, t_1, t_x)$  in (1).

**Lemma 11** Let the chain (1) have four dimensional characteristic  $x$ -ring. Then

$$f = M(x, t, t_x)A(x, t, t_1) + t_x B(x, t, t_1) + C(x, t, t_1), \tag{12}$$

where  $M$ ,  $A$ ,  $B$  and  $C$  are some functions.

**Proof.** Let  $f_{t_x t_x} \neq 0$  (if  $f_{t_x t_x} = 0$  then  $f$  obviously has the above form). Since characteristic  $x$ -ring has dimension four the condition (6) holds. It is easy to see that (6) implies that  $\frac{f_{t_x t_x t_x}}{f_{t_x t_x}}$  does not depend on  $t_1$ . Hence

$$X(\ln |f_{t_x t_x}|) = M_1(x, t, t_x) \quad \text{and} \quad \ln |f_{t_x t_x}| = M_2(x, t, t_x) + A_1(x, t, t_1).$$

The last equality implies (12).  $\square$

We can also put some restrictions on the shifts of the function  $f(x, t, t_1, t_x)$  in (1).

**Lemma 12** *Let the chain (1) have four dimensional characteristic x-ring and  $f_{t_x t_x} \neq 0$ . Then*

$$Df = -H_1(x, t, t_1, t_2)t_x + H_2(x, t, t_1, t_2)f + H_3(x, t, t_1, t_2), \quad (13)$$

where  $H_1$ ,  $H_2$  and  $H_3$  are some functions.

**Proof.** Note that the shift operator  $D$  and the vector field  $X$  satisfy

$$DX = \frac{1}{f_{t_x}}XD. \quad (14)$$

The condition (6) can be written as

$$DX(\ln |f_{t_x t_x}|) = \frac{1}{f_{t_x}}X(\ln |f_{t_x t_x}| - \ln |f_{t_x}|^3)$$

Using (14) we get

$$\frac{1}{f_{t_x}}XD(\ln |f_{t_x t_x}|) = \frac{1}{f_{t_x}}X(\ln |f_{t_x t_x}| - \ln |f_{t_x}|^3)$$

which implies that

$$X\left(\ln \left|f_{t_x}^3 \frac{Df_{t_x t_x}}{f_{t_x t_x}}\right|\right) = 0 \quad \text{or} \quad X\left(f_{t_x}^3 \frac{Df_{t_x t_x}}{f_{t_x t_x}}\right) = 0.$$

Thus  $Df_{t_x t_x} = H_1(x, t, t_1, t_2)\frac{f_{t_x t_x}}{f_{t_x}^3}$ . Since  $Df_{t_x t_x} = DX(f_{t_x})$  and  $\frac{f_{t_x t_x}}{f_{t_x}^3} = -\frac{1}{f_{t_x}}X\left(\frac{1}{f_{t_x}}\right)$  we can rewrite previous equality using (14) as

$$X\left(Df_{t_x} + H_1(x, t, t_1, t_2)\frac{1}{f_{t_x}}\right) = 0$$

which implies

$$Df_{t_x} = -H_1(x, t, t_1, t_2)\frac{1}{f_{t_x}} + H_2(x, t, t_1, t_2).$$

Writing

$$DX(f) = -H_1(x, t, t_1, t_2)\frac{1}{f_{t_x}} + H_2(x, t, t_1, t_2)\frac{f_{t_x}}{f_{t_x}}$$

and applying (14) as before we get

$$X(Df + H_1(x, t, t_1, t_2)t_x - H_2(x, t, t_1, t_2)f) = 0.$$

The last equality gives (13).  $\square$

Note that the equality (13) can be written as

$$t_{2x} = H_2(x, t, t_1, t_2)t_{1x} - H_1(x, t, t_1, t_2)t_x + H_3(x, t, t_1, t_2).$$

## References

- [1] Darboux G 1915 *Leçons sur la théorie générale des surfaces et les applications géométriques du calcul infinitesimal* **2** (Paris: Gautier Villas)
- [2] R K Dodd and R K Bullough 1976 *Proc. R. Soc. London Ser. A* **351** 499
- [3] Habibullin I T 2005 *SIGMA Symmetry Integrability Geom.: Methods Appl.* **1**
- [4] Habibullin I T and Pekcan A 2007 *Theoret. and Math. Phys.* **151** 781790
- [5] Habibullin I 2007 *Characteristic algebras of discrete equations. Difference equations, special functions and orthogonal polynomials* (Hackensack NJ World Sci. Publ.) 249-257
- [6] Habibullin I Zheltukhina N and Pekcan A 2008 *Turkish J. Math.* **32** 277292
- [7] Habibullin I Zheltukhina N and Pekcan A 2008 *J. Math. Phys.* **49** 102702
- [8] Habibullin I Zheltukhina N and Pekcan A 2009 *J. Math. Phys.* **50** 102710
- [9] Habibullin I Zheltukhina N and Sakieva A 2010 *J. Phys. A* **43** 434017
- [10] Habibullin I and Zheltukhina N 2014 Discretization of Liouville type nonautonomous equations *Preprint* nlin.SI:1402.3692v1
- [11] Shabat A B and Yamilov R I 1981 Exponential systems of type I and Cartan matrices *Preprint* BBAS USSR Ufa
- [12] Zhiber A B, Murtazina R D, Habibullin I T and Shabat A B 2012 *Ufa Math. J.* **4** 17-85