

1 **TITLE**  
2 NATURAL VITAMIN B12 AND FUCOSE SUPPLEMENTATION OF GREEN  
3 SMOOTHIES WITH EDIBLE ALGAE AND RELATED QUALITY CHANGES  
4 DURING THEIR SHELF LIFE

5 **RUNNING TITLE**  
6 Natural vitamin B12 and fucose supplementation of green smoothies

7  
8 **AUTHORS' NAMES**  
9 Noelia Castillejo<sup>a</sup>, Ginés Benito Martínez-Hernández<sup>a,b</sup>, Valentina Goffi<sup>a</sup>, Perla A.  
10 Gómez<sup>b</sup>, Encarna Aguayo<sup>a,b</sup>, Francisco Artés<sup>a,b</sup> and Francisco Artés-Hernández<sup>a,b\*</sup>

11  
12 <sup>a</sup> Postharvest and Refrigeration Group, Department of Food Engineering, Universidad  
13 Politécnica de Cartagena, Paseo Alfonso XIII, 48, 30203, Cartagena, Murcia, Spain.

14 <sup>b</sup> Institute of Plant Biotechnology, Universidad Politécnica de Cartagena, Campus  
15 Muralla del Mar s/n, 30202, Cartagena, Murcia, Spain.

16  
17 \* To whom correspondence should be addressed: Tel: +34-968-325509; Fax:  
18 +34-968-325433. E-mail: [fr.artes-hdez@upct.es](mailto:fr.artes-hdez@upct.es) Web site: [www.upct.es/gpostref](http://www.upct.es/gpostref)

19  
20 **ABSTRACT**  
21 BACKGROUND: Some algae are an excellent source of vitamin B12, of special  
22 interest for vegetarian/vegan consumers, and fucose to supplement fruit and vegetables  
23 beverages like smoothies. Nevertheless, the algae supplementation of smoothies may  
24 lead to possible quality changes during smoothie shelf life that need to be studied.  
25 Accordingly, the quality changes of fresh green smoothies supplemented (2.2%) with 9

26 edible algae (sea lettuce, kombu, wakame, thongweed, dulse, Irish moss, nori, spirulina  
27 and chlorella) were studied throughout 24 days at 5°C.

28 **RESULTS:** The initial vitamin C content (238.7–326.0 mg kg<sup>-1</sup> fw) of a 200 g-portion  
29 of any of the smoothies ensured a full coverage of its recommended daily intake, being  
30 still covered a 50–60% of the recommended intake after 7 days. Chlorella and  
31 spirulina-smoothies showed the highest vitamin B12 content (33.3 and 15.3 µg kg<sup>-1</sup> fw,  
32 respectively) while brown algae showed fucose contents of 141.1–571.3 mg kg<sup>-1</sup> fw.  
33 Such vitamin B12 and fucose contents were highly maintained during smoothies'  
34 shelf-lives.

35 **CONCLUSION:** The spirulina supplementation of a 200 g-smoothie portion ensured a  
36 full coverage of the recommended vitamin B12 intakes with lower vitamin C  
37 degradation during a shelf-life of 17 days. Furthermore, thongweed and kombu are also  
38 considered as excellent fucose sources with the same shelf-lives.

39

40 **Keywords:** Seaweed, beverages, health-promoting compounds, fucoidans, phenols,  
41 antioxidants.

42

43

## INTRODUCTION

44 Fruit and vegetables represent a rich source of phytochemicals with health-promoting  
45 properties related to preventative effects on cardiovascular diseases, cancers,  
46 hypertension and other chronic conditions such as diabetes and obesity.<sup>1</sup> White grapes,  
47 broccoli and cucumber have high contents of such phytochemicals such as phenolic  
48 compounds, vitamin C and other antioxidant compounds, among others.<sup>2-4</sup> However,  
49 fruit and vegetables consumption is below the recommended daily intake.<sup>5</sup> Beverages,  
50 and more recently smoothies, represent an excellent and convenient alternative to

51 promote the daily consumption of fruit and vegetables.<sup>6, 7</sup> Smoothies are non-alcoholic  
52 beverages prepared from fresh or frozen fruit and/or vegetables, which are blended and  
53 usually mixed with crushed ice to be immediately consumed. Often, some smoothies  
54 may include other components like yogurt, milk, ice-cream, lemonade or tea<sup>8</sup>.

55 The current consumer searches for innovative food products with new tastes, which also  
56 cover the nutritional needs together with additional health-promoting properties.  
57 'Fortification' or 'enrichment' is the 'addition of one or more essential nutrients to a  
58 food whether or not it is normally contained in it, for the purpose of preventing or  
59 correcting a demonstrated deficiency of one or more nutrients in the population or  
60 specific population  
61 groups'.<sup>9</sup> Nevertheless, the actual consumer looks for food products with natural  
62 ingredients. Accordingly, fortified products with natural ingredients are attracting much  
63 attention. Vitamins B12 and C cannot be synthesized by humans so they must be  
64 ingested with food. Usual dietary sources of vitamin B12 are animal food products, but  
65 not plant food products, being such fact of crucial interest for some populations groups  
66 such as vegetarians/vegans. Some edible algae have been reported to shown large  
67 amounts of vitamin B12.<sup>10, 11</sup> High contents of phenolic compounds can be also found in  
68 marine algae, being phlorotannins the main phenolic group, which provide a wide range  
69 of potential biological activities (antioxidant, anticancer, antibacterial, anti-allergic,  
70 anti-diabetes, anti-aging, anti-inflammatory and anti-HIV activities)<sup>12, 13</sup>. Brown  
71 algae are also rich sources of fucoidans, L-fucose sulphated polysaccharides, which  
72 have several health-promoting properties such as anticancer, antioxidant, antiviral and  
73 antioxidant, among others, as recently reviewed.<sup>12, 14</sup> Algae have been traditionally used  
74 for culinary purposes in Asian countries although their consumption has recently spread  
75 to Western countries as bioactive ingredients included in functional foods. Algae are

76 commonly classified into three groups based on their pigmentation: brown  
77 (*Phaeophyceae*), red (*Rhodophyceae*) and green (*Chlorophyceae*) algae. Furthermore,  
78 such scenario also promotes the creation of edible algae industries in other countries  
79 different from Asian area which quality may be excellent, and even higher for some  
80 purposes, compared to those imported dried seaweeds from East Asia <sup>15</sup>.

81 The natural vitamin B12 fortification of fruit/vegetable smoothies with algae may have  
82 a high relevance in the food industry to supply to the consumer food products with  
83 natural ingredients, which covers their nutritional needs. Furthermore, such natural  
84 fortification may lead to extra health-promoting properties derived from the high  
85 phenolics and fucose contents, among other compounds, of such marine plants.  
86 However, there are no previous reports of possible side effects of algae fortification on  
87 the quality of fruit/vegetables smoothies. Accordingly, the aim of the present work was  
88 to study the main quality changes and bioactive contents of several fresh  
89 fruit/vegetables smoothies formulated with 9 different edible algae during 24 days of  
90 storage at 5°C.

91

92

## MATERIALS AND METHODS

### 93 **Plant material and smoothie preparation**

94 Fresh white grapes and cucumbers were purchased at a local supermarket and  
95 kalia-hybrid broccoli (Bimi<sup>®</sup>) was obtained from a local producer (Campo de  
96 Lorca–Juan Marín S.L.; Lorca, Murcia, Spain) in June. Plant material was transported  
97 within 1 h to the Pilot Plant at the Universidad Politécnica de Cartagena, where it was  
98 stored at 4°C and 90–95% relative humidity (RH) until next day.

99 The 9 edible algae used were sea lettuce, kombu, wakame, thongweed, dulse, Irish  
100 moss, nori, chlorella and spirulina, which are described in Table 1. They were

101 purchased from Porto–Muiños (La Coruña, Galicia, Spain). Algae were supplied as  
102 ground dried powder (200 g) in plastic bottles. Since all samples had different particle  
103 sizes, they were grinded with a mill (IKA, A 11 Basic, Berlin, Germany) using liquid  
104 nitrogen to fine powder with a measured (Scirocco 2000, Malvern Instruments;  
105 Malvern, Worcestershire, UK) average particle size of 300  $\mu\text{m}$ .

106 Preparation of smoothies was accomplished in a disinfected cold room at 8°C. Plant  
107 material was carefully inspected, selecting those free from defects and with similar  
108 visual appearance. Subsequently, plant material was sanitized with 75 mg L<sup>-1</sup> NaClO  
109 during 2 min and then rinsed with cold tap water for 1 min. Then, cucumbers were  
110 peeled, grape berries detached from the cluster and broccoli was cut with total length of  
111 approximately 15 cm with a sharp knife. Nine different smoothies containing the  
112 different algae were prepared. The vegetables, fruit and alga proportions for preparation  
113 of smoothies were: 56.5% white grapes, 15.5% broccoli, 25.8% cucumber and 2.2%  
114 alga. A smoothie without alga was prepared as control (CTRL) containing: 57.8%  
115 grapes, 15.8% broccoli and 26.4% cucumber. The smoothie composition was selected  
116 among several formulations according to sensory pre–evaluations conducted by a  
117 sensory panel focussing on the maximum broccoli quantity in order to maximize the  
118 bioactive contents of the smoothie. Smoothies were prepared in a food processor (Robot  
119 Cook®, Robot Coupe; Vincennes, Île-de-France, France) and immediately cooled to  
120 4°C with an ice–water bath. Immediately after smoothie preparation, approximately 80  
121 g of each smoothie were filled (Infantino Squeeze station, Infantino; San Diego,  
122 California, USA) under aseptic conditions into a sterile squeeze polyvinyl chloride  
123 pouch (9 cm×13 cm; 118 mL; Infantino; San Diego, California, USA). Samples were  
124 stored in darkness at 4°C being conducted sampling times up to 24 days. Three  
125 replicates per treatment, storage temperature and sampling day were prepared. Samples

126 of each treatment were taken on each sampling day to be analysed storing also samples  
127 for bioactive compounds at -80°C until further analyses.

128

### 129 **Microbial analysis**

130 Psychrophilic, and yeast and moulds (Y+M) growth was determined using standard  
131 enumeration methods according to Castillejo *et al.*<sup>6</sup>. All microbial counts were reported  
132 as log colony forming units per gram of smoothie (log CFU g<sup>-1</sup>). Each of the three  
133 replicates was analysed in duplicate. *Salmonella* spp., *Listeria monocytogenes* and  
134 generic *Escherichia coli* were monitored meeting the obtained results the food safety  
135 European legislation for these products.<sup>16</sup>

136

### 137 **Physiochemical analyses**

138 The total soluble solids content (SSC), pH, titratable acidity (TA) and colour of  
139 smoothies were determined as previously described.<sup>6</sup> The SSC of the smoothie was  
140 determined by a digital hand-held refractometer (Atago N1; Tokyo, Kanto, Japan) at  
141 20°C and expressed as % (g sugar equivalents 100 g<sup>-1</sup>). A pH-meter (Basic20, Crison;  
142 Alella, Cataluña, Spain) was used to determine the pH. TA was determined by titration  
143 of 5 mL of smoothie plus 35 mL of distilled water with 0.1 M NaOH to pH 8.1 (T50,  
144 Metter Toledo; Milan, Lombardia, Italy) and expressed as % (g tartaric acid 100 mL<sup>-1</sup>).  
145 Colour was determined using a colorimeter (Chroma Meter CR-300, Minolta; Tokyo,  
146 Kanto, Japan) calibrated with a white reference plate (light source C), 2° observer and  
147 8-mm viewing aperture. Samples were introduced in a special glass tube mounted on a  
148 device connected to the colorimeter. Three colour readings were taken turning the tube  
149 every caption and all three measurements were automatically averaged by the device  
150 and recorded. Measurements were recorded using the standard tristimulus parameters

151 ( $L^*$ ,  $a^*$ ,  $b^*$ ) of the CIE Lab system. Total colour differences ( $\Delta E$ ) throughout storage  
152 compared to their respective initial values were calculated according to equations  
153 previously described.<sup>17</sup>

154

### 155 **Sensory evaluation**

156 Sensory analyses were performed according to international standards.<sup>18</sup> Tests were  
157 conducted in a standard room<sup>19</sup> equipped with 10 individual taste booths. Smoothie  
158 samples (about 30 mL) were served at room temperature in transparent plastic glasses  
159 coded with three random digit numbers. Still mineral water was used as palate cleanser.  
160 The panel consisted of 12 assessors (6 women/6 men, aged 22–70 years) screened for  
161 sensory ability (visual appearance, colour, aroma, flavour and texture). A 5–point scale  
162 of damage incidence and severity was scored for off–colours, off–flavours, off–odours,  
163 lumpiness and phase separation (5: none; 4: slight; 3: moderate, limit of usability (LU);  
164 2: severe; 1: extreme). Visual appearance, aroma, flavour, texture and overall quality  
165 were assessed using a 5–point hedonic scale of acceptability (5: excellent; 4: good; 3:  
166 fair, LU; 2: poor; 1: extremely bad).

167

### 168 **Vitamin C**

169 The ascorbic (AA) and dehydroascorbic (DHA) acids were measured as previously  
170 described.<sup>20, 21</sup> Briefly, 5 g ground frozen (-80°C) sample was placed into a 25–mL  
171 Falcon tube and 10 mL of cold (4°C) buffer (0.1 M citric acid, 0.05% EDTA, 4 mM  
172 sodium fluoride and 5% MeOH) were added. The mixture was homogenised  
173 (UltraTurrax T25 basic, IKA; Berlin, Germany) for 10 s, filtered (four–layer  
174 cheesecloth) and the pH was adjusted (6N NaOH) to 2.35–2.40. Subsequently, 750  $\mu$ L  
175 filtered (0.45– $\mu$ m polytetrafluoroethylene (PTFE) membrane filters) purified extract

176 (Sep–Pak cartridges C18, Waters; Dublin, Leinster, Ireland) was derivatised with 250  
177  $\mu\text{L}$  of 7.7 M 1,2–phenylenediamine for 37 min in darkness at room temperature.  
178 Immediately after derivatisation, 20  $\mu\text{L}$  were injected in a Gemini NX (250 mm $\times$ 4.6  
179 mm, 5  $\mu\text{m}$ ) C18 column (Phenomenex; Torrance, California, USA), using an HPLC  
180 (Series 1100 Agilent Technologies; Waldbronn, Baden-Württemberg, Germany)  
181 equipped with a G1322A degasser, G1311A quaternary pump, G1313A autosampler,  
182 G1316A column heater and G1315B photodiode array detector. AA and DHA were  
183 quantified using commercial standards (Sigma; St Louis, Missouri, USA). Calibration  
184 curves were made with at least six data points for each standard. AA and DHA were  
185 expressed as  $\text{mg kg}^{-1}$  fresh weight (fw). Each sample was analysed in duplicate.

186

#### 187 **Total phenolic content**

188 Frozen samples of 1 g were placed in glass bottles and 4 mL of methanol was added.  
189 The extraction was carried out in an orbital shaker (Stuart; Staffordshire, West  
190 Midlands, UK) for 1 h at 200 rpm in darkness inside a polystyrene (PS) box with an ice  
191 bed. The extracts were transferred in eppendorf tubes and centrifuged at 15,000 $\times g$  for  
192 10 min at 4°C. The supernatant was used as total phenolic content (TPC) and total  
193 antioxidant capacity (TAC) extracts.<sup>22, 23</sup> The TPC was determined as previously  
194 described based on, but with modifications proposed by. Briefly, 19  $\mu\text{L}$  of TPC extract  
195 was placed on a flat–bottom PS 96–well plate (Greiner Bio–One; Frickenhausen,  
196 Baden-Württemberg, Germany) and 29  $\mu\text{L}$  of 1 N Folin–Ciocalteu reagent was added.  
197 The latter mixture was incubated for 3 min in darkness at room temperature. Then, 192  
198  $\mu\text{L}$  of a solution containing  $\text{Na}_2\text{CO}_3$  (0.4%) and NaOH (2%) was added. After 1 h of  
199 incubation at room temperature in darkness, the absorbance was measured at 750 nm  
200 using a Multiscan plate reader (Tecan Infinite M200; Männedorf, Meilen,



201 Switzerland). The TPC was expressed as mg gallic acid equivalents (GAE) kg<sup>-1</sup> fw.  
202 Each sample was analysed in duplicate.

203

#### 204 **Total antioxidant capacity**

205 The extracts were analysed for TAC using the same instruments and methodology as  
206 previously described<sup>8</sup> using three different methods: free radical scavenging capacity  
207 with 2,2-diphenyl-1-picrylhydrazil (DPPH),<sup>24</sup> ferric reducing antioxidant power  
208 (FRAP)<sup>25</sup> and 2,20-azino-bis (3-ethylbenzothiazoline-6-sulphonicacid) (ABTS).<sup>26</sup>  
209 Results were expressed as mg Trolox equivalent antioxidant capacity kg<sup>-1</sup> fw. Each  
210 sample was analysed in duplicate.

211

#### 212 **Vitamin B12**

213 Vitamin B12 was determined according to a commercial microbiological kit for vitamin  
214 B12 (VitaFast, r-biopharm; Berlin, Germany). Briefly, 1 g of smoothie was mixed with  
215 40 mL of distilled water, vortex and incubated at 95°C for 30 min. After cooling down  
216 at room temperature, the solution was centrifuged at 32,000×g for 15 min at 15°C and  
217 filtered through 0.45 µm PTFE membrane filters. Subsequently, 150 µL of vitamin B12  
218 assay medium (available from the kit) was disposed on the wells of the microtiter plate  
219 (pre-coated with *Lactobacillus delbrueckii* subsp. *Lactis* (leichmannii)) supplied by the  
220 vitamin B12-kit. Then, 150 µL of the vitamin B12 extract was added and the microtiter  
221 plate was incubated at 37°C in the dark for 46 h. Finally, the absorbance was measured  
222 at 620 nm using the Multiscan plate reader. Vitamin B12 was quantified using the  
223 vitamin B12 standard supplied by the vitamin B12-kit. The vitamin B12 was expressed  
224 as µg kg<sup>-1</sup> fw. Each of the samples was analysed in duplicate.

225

## 226 **Furoidans/Fucose**

227 Fucose (*L*-fucose) was determined using a commercial kit (*L*-fucose, Megazyme;  
228 Bray, Leinster, Ireland). Briefly, 2.5 g of smoothie was mixed with 2.5 mL 1.3 M HCl,  
229 vortex and incubated at 100°C for 1 h. After cooling down at room temperature, 2.5 mL  
230 of 1.3 M NaOH were added, vortex and filtered through 0.45 µm PTFE membrane  
231 filters. Subsequently, 200 µL of water, 20 µL of fucose extract, 40 µL of buffer  
232 (supplied by the fucose kit) and 10 µL of NADP<sup>+</sup> solution (supplied by the kit) were  
233 placed on a flat-bottom PS 96-well plate. After 4 min of incubation at room  
234 temperature, 2 µL of *L*-fucose dehydrogenase suspension (supplied by the kit) was  
235 added and it was incubated at 37°C for 1 h. Finally, the absorbance was measured at 340  
236 nm using the Multiscan plate reader. Fucose was quantified using the *L*-fucose standard  
237 supplied by the kit. The fucose content was expressed as g kg<sup>-1</sup> fw. Each of the samples  
238 was analysed in duplicate.

239

## 240 **Statistical Analysis**

241 The experiment was a two-factor (smoothie type×storage time) design subjected to  
242 analysis of variance (ANOVA) using Statgraphics Plus software (vs. 5.1, Statpoint  
243 Technologies Inc.; Warrenton, Virginia, USA). Statistical significance was assessed at  
244 the level  $p=0.05$ , and Tukey's multiple range test was used to separate means.

245

## 246 **RESULTS AND DISCUSSION**

### 247 **Physicochemical quality**

248 The physicochemical quality of smoothies can be evaluated based on SSC, pH, TA  
249 and colour being closely related to sensory quality.<sup>6</sup> Table 2 represents the effect of  
250 algae supplementation on the physicochemical quality of smoothies throughout

251 storage. CTRL smoothie samples showed an initial high SSC of 12.4% being owed to  
252 the high content of grapes in the smoothie. A similar SSC has been also reported in  
253 other fruit-containing smoothies differing from other vegetables smoothies without  
254 fruit.<sup>8, 27</sup> The SSC of the smoothie was not significantly ( $p < 0.05$ ) changed after algae  
255 supplementation. Particularly, smoothies supplemented with brown and red  
256 macroalgae showed higher SSC ( $p < 0.05$ ) than those with green algae (hereinafter  
257 including both macro and microalgae). Such finding may be explained by the higher  
258 content of SSC, mainly sugars, of brown and red algae regarding green algae.<sup>28, 29</sup> The  
259 CTRL smoothie showed an initial pH of 4.24 allowing such acidic medium a moderate  
260 shelf life of the beverage under refrigeration conditions without the need of thermal  
261 treatments<sup>30</sup> which may reduce the sensory and nutritional/bioactive quality of the  
262 smoothie. Algae supplementation of smoothies led to a light pH increase up to  
263 4.32–4.77 owed to the high mineral contents of algae, which may achieve up to 40%  
264 of total weight.<sup>31</sup> The CTRL smoothie showed an initial TA of 0.30% that was slightly  
265 reduced after algae supplementation according to previous slight pH increment.

266 In general, SSC of samples did not highly change throughout storage ( $< 0.6$  SSC units),  
267 with a particular general SSC decrease on day 3, except for sea lettuce and CTRL  
268 smoothies, being newly upregulated on days 7–10. The latter slight SSC decrease may  
269 be owed to a sugar consumption by microorganisms, which initiated to growth after  
270 such initial adaptation period to the smoothie medium according to psychrophiles data  
271 (shown later). Furthermore, no high pH and TA changes were observed ( $< 0.3$  pH and  
272  $< 0.28$  TA units) after 24 days at 5°C, showing sea lettuce and CTRL smoothies the  
273 lowest pH/TA variations. Similarly, no pH changes were either observed in a fresh  
274 (unheated) green vegetable puree after 43 days at 4°C.<sup>32</sup> Nevertheless, smoothies  
275 supplemented with the microalgae spirulina and chlorella particularly showed TA

276 increments of 0.35 and 0.23 units in the last 7 days of storage although such  
277 acidification was not negatively scored by the sensory panel even showing spirulina  
278 smoothie the best flavour scores after 24 days of storage (see sensory data).  
279 The addition of algae to the green smoothie induced a decrease of luminosity ( $L^*$ ) and  
280 yellowness ( $b^*$ ), and an increase of redness ( $a^*$ ). The microalgae spirulina and  
281 chlorella showed the highest colour changes, as expected due to their intense green  
282 colour, with  $\Delta E$  of 23.0 and 20.8 on processing day, respectively (Table 2).  
283 Nevertheless, such colour changes did not negatively affect to the consumer  
284 acceptance of algae-supplemented smoothies since the sensory panel highly scored (>  
285 4) general appearance and colour of all samples on processing day (see sensory data).  
286 On the other side, algae-smoothies showed lower colour changes ( $\Delta E=6.8-9.8$ ) than  
287 CTRL smoothie ( $\Delta E=11.9$ ) after 24 days, showing spirulina-smoothie the lowest  
288 colour differences. Therefore, the colour degradation of the smoothie due to enzymatic  
289 activity, as previously observed,<sup>8</sup> was reduced with the algae supplementation,  
290 probably owed to enzymatic-inhibiting compounds from such marine plants.  
291 Accordingly, the physicochemical quality of smoothies was not highly affected after  
292 algae supplementation even showing lower colour changes compared to  
293 non-supplemented smoothie.

294

### 295 **Microbiological analysis**

296 The smoothie preparation included several unit operations such as peeling, cutting,  
297 blending and the addition of bioactive ingredients like algae, which may highly increase  
298 the microbial growth during refrigerated storage, limiting its shelf life and  
299 compromising its food safety. Consequently, microbial quality should be determined in  
300 such products in order to monitor spoilage microorganisms and pathogens.

301 Psychrophilic and Y+M loads were monitored throughout storage of smoothies at this  
302 low storage temperature and the adaptation of Y+M to grow under such acidic  
303 beverages (Table 3). The initial psychrophilic and Y+M loads of the green smoothie  
304 (3.9 and 2.9 log CFU g<sup>-1</sup>, respectively) were not highly altered with the addition of  
305 algae to the smoothies, reporting increments lower than 0.3 and 0.6 log units,  
306 respectively, on processing day.

307 A general microbial reduction was observed in the psychrophilic and Y+M growth  
308 during storage of smoothies showing loads of 2.8–3.8 and 2.3–2.9 log CFU g<sup>-1</sup>,  
309 respectively, after 7 days at 5°C. Nevertheless, microbial loads of all smoothies were  
310 increased after day 7. Particularly, psychrophilic growth was higher in algae–smoothies  
311 compared to CTRL samples, showing brown algae–smoothies loads of 7 log CFU g<sup>-1</sup>  
312 after 17 days at 5 °C while such levels were only exceeded in red–algae after 21 days at  
313 °C. Furthermore, CTRL and sea lettuce–smoothies showed the lowest psychrophilic  
314 loads after 24 days at 5°C, with 4.9 and 5.3 log CFU g<sup>-1</sup>, respectively. *Brassica* species,  
315 i.e. broccoli, have high glucosinolates contents,<sup>2</sup> which after plant cell disruption, i.e.  
316 smoothie preparation, come in contact with plant myrosinase that is previously located  
317 in separate cell compartments. The activity of myrosinase transforms glucosinolates to  
318 unstable intermediate compounds, which rearranges mainly to isothiocyanates under  
319 acidic conditions and presence of mineral ions, among other factors, instead of other  
320 breakdown products.<sup>33, 34</sup> High antimicrobial properties have been reported by  
321 sulforaphane, the isothiocyanate resulting from the glucosinolate glucoraphanin, one of  
322 the main glucosinolates of broccoli.<sup>6</sup> Therefore, the higher psychrophilic growth in all  
323 algae–smoothies may be owed to the higher pH and mineral contents regarding the  
324 CTRL smoothie without algae supplementation. Nevertheless, the lower psychrophilic  
325 growth in sea–lettuce smoothie may be explained by the lower mineral contents from

326 this alga compared to the remaining algae (data not shown). However, such  
327 antimicrobial effect throughout storage was not observed for Y+M of CTRL smoothie,  
328 which showed the highest Y+M load, together with kombu-smoothie (low SSC and  
329 TA), of 4.9 log CFU g<sup>-1</sup> after 24 days at 5°C. Meanwhile, the remaining samples  
330 showed Y+M loads that ranged among 3.3 to 3.6 log CFU g<sup>-1</sup>. The latter behaviour may  
331 be explained by the high adaptation of Y+M to grow under acidic conditions.  
332 Furthermore, the lower Y+M growth of most of algae-smoothies may be owed to the  
333 early known fungistatic properties of marine algae.<sup>35</sup>

334 *Salmonella* spp., *L. monocytogenes* and generic *E. coli* were monitored throughout  
335 storage, meeting the obtained results the food safety European legislation for these  
336 products.<sup>16</sup>

337 Conclusively, algae-smoothies could be stored up to 17–21 days at 5°C showing  
338 psychrophilic loads close to 7 log units, while Y+M levels were highly inhibited  
339 (1.3–1.7 lower log units) after 24 days compared to the CTRL smoothie without algae  
340 supplementation.

341

### 342 **Sensory analysis**

343 As expected, the used algae concentration within smoothies led to a mild marine taste  
344 detected by the panellists (Figure 1). Irish moss and chlorella addition led to the lowest  
345 overall quality scores on processing day, mainly due to a stronger marine odour/flavour  
346 of these algae, showing their smoothies scores of 2.3/3.0 and 2.4/3.1, respectively. As  
347 depicted in material and methods section, a general 2.2% algae content was used for all  
348 smoothie formulations in order to avoid quality differences owed to different algae  
349 contents. Nevertheless, the Irish moss and chlorella contents are recommended to be  
350 reduced from the 2.2% tested in order to achieve a higher consumer acceptance. The

351 sensory quality of the remaining algae–smoothies was highly scored (>4) showing  
352 spirulina and kombu the highest overall quality scores with 4.5–4.6. No  
353 off–flavours/odours/colours were detected among all the smoothies on processing day,  
354 showing a pleasant texture without a remarkable lumpiness justified by the appropriated  
355 blending program used with the used semi–industrial food processor.

356 Algae–smoothies still showed overall quality scores over the limit of acceptability (3)  
357 after 14 days at 5°C showing Irish moss, chlorella and wakame the lowest scores of  
358 3.1–3.2 mainly owed to low flavour and aroma scores (Figure 1). Nevertheless, overall  
359 quality of algae–smoothies was below the limit of acceptability after 17 days at 5°C,  
360 except sea lettuce–smoothie. The latter low overall quality scores were mainly owed to  
361 the low aroma and flavour scores with remarkable off–flavours, mainly for chlorella  
362 and brown macroalgae, and increased lumpiness, which reached a score of 2.7 for  
363 wakame–smoothie. No high phase separation was observed for the smoothies with  
364 scores of 4–4.5 and 3–3.5 after 17 and 24 days at 5°C, respectively. The overall quality  
365 of sea lettuce–smoothie was scored with 3.0 after 24 days at 5°C (Figure 1) with similar  
366 scores to the CTRL smoothie (data not shown). The latter finding is explained by the  
367 milder marine taste of sea lettuce compared to the remaining algae.

368 Conclusively, the shelf life of algae–smoothies could be established in 17 days at 5°C  
369 based on sensory and microbiological quality. Particularly, the shelf life of the  
370 lettuce–smoothie was even extended up to 24 days at °C due to its previously discussed  
371 low psychrophilic load throughout the storage period.

372

### 373 **Vitamin C**

374 Ascorbic acid is stable when dry but in solutions it readily oxidises to the intermediate  
375 compound monodehydroascorbate (MDHA) through the activity of the enzyme

376 ascorbate oxidase. Subsequently, MDHA may be converted to DHA that can be reduced  
377 newly to AA or hydrolysed to 2,3-diketogulonic acid (DKG).<sup>36</sup> DHA also exhibits  
378 antioxidant properties in addition to antiscorbutic activity equivalent to that of AA,  
379 contrary to the non-bioactive compound DKG.<sup>37</sup> Therefore, vitamin C content of fruit  
380 and vegetables has been proposed as the sum of AA and DHA.<sup>38</sup> The CTRL smoothie  
381 showed an initial vitamin C content of 326.0 mg kg<sup>-1</sup> fw (Table 4). Similar total vitamin  
382 C contents have been reported in other fruit/vegetables fresh smoothies, also containing  
383 broccoli and grapes.<sup>6, 27</sup> Nevertheless, no AA was detected in the smoothie samples,  
384 contrary to previous data on vegetables smoothies.<sup>6</sup> The latter finding may be explained  
385 by a high AA degradation by ascorbate oxidase due to the cucumber included in our  
386 smoothie, being this vegetable, and *Cucurbitaceae* family in general, among the most  
387 abundant sources of this enzyme.<sup>39</sup> The role of metal ions, such as those contained in  
388 algae, in the oxidation of AA has been widely known for more than 95 years<sup>40</sup>  
389 explaining the mild vitamin C reduction (up to 27%) observed after algae  
390 supplementation of smoothies. Nevertheless, a 200 g-portion of the algae-smoothie  
391 with the lowest ( $p<0.05$ ) initial vitamin C content (Irish moss) still ensured the  
392 recommended daily intake (RDI) of vitamin C.<sup>41</sup>

393 The vitamin C content of smoothies decreased throughout storage, showing levels  
394 50–60% lower after 7 days at 5°C. Latter finding may be explained since DHA is itself  
395 very unstable in aqueous solution (half-life of 6 min at 37°C) and undergoes  
396 irreversible hydrolytic ring cleavage to the non-bioactive DKG.<sup>42</sup> Particularly,  
397 chlorella-smoothie showed the highest vitamin C reduction of 70% probably owed to  
398 the high content in this alga of iron, one of the main metal ions which highly induce  
399 vitamin C degradation.<sup>43</sup> A vitamin C degradation of approximately 90%, for all  
400 smoothies without high differences among them, was observed on day 14, regarding



401 their respective initial levels, being such low levels maintained until the last day of  
402 storage. Likewise, high vitamin C degradation has been previously observed in fresh  
403 fruit/vegetables smoothies stored under similar low storage temperature.<sup>6, 44, 45</sup>

404 Conclusively, all smoothies covered the vitamin C recommended daily intake by the  
405 WHO while a 200 g-portion stored for 7 days at 5°C still ensured the 50–60% of the  
406 recommended daily intake of this vitamin.

407

#### 408 **Vitamin B12**

409 Smoothies supplemented with all macroalgae, except kombu and thongweed  
410 (undetected levels), showed similar ( $p<0.05$ ) initial vitamin B12 contents of  
411 approximately  $1 \mu\text{g kg}^{-1}$  fw ( $0.4 \mu\text{g kg}^{-1}$  fw for Irish moss) (Figure 2). Nevertheless,  
412 chlorella and spirulina-smoothies showed initial vitamin B12 levels of 33.3 and  $15.3 \mu\text{g}$   
413  $\text{kg}^{-1}$  fw, respectively. Accordingly, chlorella and spirulina-smoothies portions of just  
414 70 and 160 g would cover the recommended vitamin B12 daily intake.<sup>41</sup> Spirulina and  
415 chlorella are algae with high vitamin B12 content, as previously reported.<sup>10, 46</sup>

416 Vitamin B12 contents of previous smoothies did not change ( $p<0.05$ ) throughout  
417 storage. As observed, the supplementation of the smoothie with all algae (except kombu  
418 and thongweed) may be considered as a natural tool to fortify fruit/vegetable beverages  
419 with vitamin B12, being of special interest for some populations groups such as  
420 vegetarians/vegans, elderly, individuals with disorders of malnutrition, etc. Vitamin B12  
421 belongs to the corrinoid group and is usually restricted to cyanocobalamin although  
422 microbiological analytical method, hereby used and approved by the Association of  
423 Analytical Communities,<sup>47</sup> may also detect other corrinoids non-bioavailable for  
424 humans known as pseudo-vitamin B12.<sup>48-51</sup> Accordingly, active vitamin B12  
425 coenzymes comprised about 60 % of total vitamin B12 in nori and chlorella

426 supplements.<sup>52</sup> Accordingly, the total vitamin B12 contained in 250g-portion of  
427 chlorella and spirulina-smoothies stored for 24 days at 5 °C represents 475 and 245% of  
428 the recommended vitamin B12 daily intake which would lead to a full coverage of the  
429 needed biologically-active B12 levels.

430

#### 431 **Fucose**

432 Brown algae are natural sources of fucoidans, or fucans, which are naturally occurring  
433 L-fucose sulphated polysaccharides. Several health-promoting properties (anticancer,  
434 antioxidant, antiviral and antioxidant, among others) have been linked to fucoidans as  
435 previously reviewed.<sup>12</sup> The fucoidans composition is complex and still unclear being  
436 due to its high heterogeneity, which is influenced by the alga specie, part of the plant or  
437 even the extraction method used.<sup>53</sup> In this sense, each fucoidan extracted from a  
438 different specie with a specific method will be unique regarding to structure and  
439 composition, leading to differences related to biological activities. Therefore, fucoidans  
440 were indirectly studied in this work by their conversion to the fucose monomer by  
441 depolymerisation and desulphation by strong acid and high temperature. The fucose  
442 data regarding to analysed brown macroalgae-smoothies, being not detected in the  
443 remaining smoothies, showed contents of 571.3, 455.7 and 141.1 mg kg<sup>-1</sup> fw for  
444 thongweed, kombu and wakame, respectively. Such contents were not changed ( $p<0.05$ )  
445 after 24 days at 5°C (data not shown).

446

#### 447 **Total phenolic content and antioxidant capacity**

448 Polyphenols from terrestrial plants are derived from gallic and ellagic acid, whereas the  
449 algal polyphenols are derived from polymerized phloroglucinol units.<sup>12</sup> The TPC were  
450 expressed as gallic acid equivalents due to the higher fruit and vegetables contents in the

451 smoothie compared to algae. The CTRL smoothie showed an initial TPC of 280.2 mg  
452 gallic acid equivalent  $\text{kg}^{-1}$  fw (Table 5). Such high TPC may be owed to the high grapes  
453 content together with broccoli which are fruit and vegetable with high phenolic  
454 contents.<sup>3, 4, 54</sup> The initial TPC content of the CTRL smoothie was increased ( $p < 0.05$ ) by  
455 69–70% after supplementation with kombu and dulse algae. Brown algae have shown  
456 higher phenolic content than red and green algae being phlorotannins the major phenolic  
457 compounds.<sup>12</sup> Furthermore, thongweed algae has shown the higher TPC compared to  
458 the other brown algae.<sup>55</sup> The alga addition and the plant cell wounding implied during  
459 smoothie preparation may generate different stresses conditions in the smoothie, which  
460 may lead to the generation of free radicals. Consequently, phenols from high source  
461 pools like thongweed may be highly used to prevent such oxidative stresses.  
462 Accordingly, thongweed–smoothie showed the lowest TPC and TAC values among  
463 brown and red algae–smoothies on processing day (Table 5).

464 A general TPC decrease of 20–50% after 3 days was observed in the smoothies,  
465 probably owed to the use of such phenolic compounds to counterbalance the stress  
466 generated during smoothie preparation (Table 5). According to such data, a TAC  
467 increase was observed on day 3 by the three TAC methods (Table 5). Subsequently, a  
468 general TPC increase from day 3 to day 7 was observed with increments ranging from  
469 60–140 and 10–30% in macro and microalgae–smoothies, respectively. Such  
470 increments may be due to a phenolic biosynthesis to counterbalance the stress during  
471 processing. Therefore, similar phenolic biosynthesis has been observed in smoothies  
472 during cold storage correlated with the activation of phenylalanine ammonia-lyase  
473 (PAL) which is considered the key enzyme in the phenylpropanoid pathway.<sup>8</sup> Higher  
474 TPC increments were observed in thongweed and Irish moss–smoothies of 400 and  
475 340%, respectively, from day 3 to day 7 regarding the remaining smoothies. That

476 finding may be explained by the low TPC of latter two smoothies on day 3, which could  
477 generate a higher PAL activation sign due to such low contents of those needed  
478 antioxidants. No remarkable TPC and TAC changes were observed from day 7 to day  
479 21 being a new general TPC decrease/TAC increase observed from day 21 to day 24.  
480 The latter second antioxidants biosynthesis may be explained by the stress generated  
481 during storage of smoothies under such low storage temperatures.

482

483

### CONCLUSIONS

484 Main quality changes of green vegetables smoothies supplemented with 9 of the most  
485 consumed/known edible algae were determined during refrigerated shelf life. Generally,  
486 the shelf life of algae–smoothies, based on microbiological and sensory quality, was  
487 established in 17 days at 5°C. Sea lettuce showed the longest shelf life (24 days) although  
488 their bioactive contents were lower than the rest of algae–smoothies. Among them, the  
489 brown algae thongweed, kombu and wakame–smoothies showed high fucose contents  
490 reporting wakame also high vitamin B12 contents. The smoothies with the microalgae  
491 chlorella and spirulina showed the highest vitamin B12 contents although the  
492 chlorella–smoothie was scored with low sensory quality and the highest vitamin C  
493 degradation during storage. Accordingly, a reduction of chlorella concentration in the  
494 smoothie formulation should be further studied for supplying a high vitamin B12  
495 contents. Therefore, fortification of smoothies with spirulina ensured a full coverage of  
496 the recommended vitamin B12 intakes with lower vitamin C degradation regarding  
497 chlorella during 17 days at 5°C. Among macroalgae–smoothies, thongweed and kombu  
498 are also considered as excellent fucose sources.

499

500 **ACKNOWLEDGEMENTS**

501 The authors are grateful to Spanish Ministry of Economy and Competitiveness, Project  
502 AGL2013-48830-C2-1-R and FEDER for financial support. We would also like to  
503 thank SAKATA SEED IBERICA S.L. for providing Bimi<sup>®</sup> samples.

504 **REFERENCES**

- 505 1. Dillard CJ and German JB, Phytochemicals: nutraceuticals and human health. *J Sci Food*  
506 *Agric* **80**:1744-1756 (2000).
- 507 2. Rosa EAS, Heaney RK, Portas CAM and Fenwick GR, Changes in glucosinolate  
508 concentrations in *Brassica* crops (*B oleracea* and *B napus*) throughout growing seasons. *J Sci*  
509 *Food Agric* **71**:237-244 (1996).
- 510 3. Martínez-Hernández GB, Gómez PA, Artés F and Artés-Hernández F, Nutritional quality  
511 changes throughout shelf-life of fresh-cut kailan-hybrid and 'Parthenon' broccoli as affected by  
512 temperature and atmosphere composition. *Food Sci Technol Int* **21**:14-23 (2015).
- 513 4. Artés-Hernández F, Artés F and Tomás-Barberán FA, Quality and enhancement of  
514 bioactive phenolics in cv. Napoleon table grapes exposed to different postharvest gaseous  
515 treatments. *J Agric Food Chem* **51**:5290-5295 (2003).
- 516 5. WHO/FAO, Population nutrient intake goals for preventing diet-related chronic  
517 diseases, in *Diet, nutrition and the prevention of chronic diseases*. World Health Organization  
518 and Food and Agriculture Organization of the United Nations, Geneva (2003).
- 519 6. Castillejo N, Martínez-Hernández GB, Gómez PA, Artés F and Artés-Hernández F, Red  
520 fresh vegetables smoothies with extended shelf life as an innovative source of health-  
521 promoting compounds. *J Agric Food Chem* **53**:1-12 (2016).
- 522 7. Hurtado A, Guàrdia MD, Picouet P, Jofré A, Ros JM and Bañón S, Stabilisation of red  
523 fruit-based smoothies by high-pressure processing. Part II: effects on sensory quality and  
524 selected nutrients. *J Sci Food Agric* **97**:777-783 (2017).
- 525 8. Rodríguez-Verástegui LL, Martínez-Hernández GB, Castillejo N, Gómez PA, Artés F and  
526 Artés-Hernández F, Bioactive compounds and enzymatic activity of red vegetable smoothies  
527 during storage. *Food Bioproc Tech* **9**:137-146 (2015).
- 528 9. FAO/WHO, General principles for the addition of essential nutrients to foods CAC/GL  
529 09-1987 (amended 1989, 1991), Ed, Rome, Italy (2015).
- 530 10. Watanabe F, Takenaka S, Kittaka-Katsura H, Ebara S and Miyamoto E, Characterization  
531 and bioavailability of vitamin B12-compounds from edible algae. *J Nutr Sci Vitaminol* **48**:325-  
532 331 (2002).
- 533 11. Taboada C, Millán R and Míguez I, Composition, nutritional aspects and effect on  
534 serum parameters of marine algae *Ulva rigida*. *J Sci Food Agric* **90**:445-449 (2010).
- 535 12. Holdt SL and Kraan S, Bioactive compounds in seaweed: functional food applications  
536 and legislation. *J Appl Phycol* **23**:543-597 (2011).
- 537 13. Machu L, Misurcova L, Vavra Ambrozova J, Orsavova J, Mlcek J, Sochor J and Jurikova T,  
538 Phenolic content and antioxidant capacity in algal food products. *Molecules* **20**:1118 (2015).
- 539 14. Black WAP, The seasonal variation in the combined L-fucose content of the common  
540 British Laminariaceae and fucaceae. *J Sci Food Agric* **5**:445-448 (1954).
- 541 15. EC, Production of edible seaweed species in Danish waters: the beginning of a new  
542 profession, Ed by FARNET (2012).
- 543 16. Regulation\_EC, Commission regulation on microbiological criteria for foodstuffs.  
544 *Official Journal of the European Union* **32** (2007).
- 545 17. Rico D, Martín-Diana AB, Frías JM, Henahan GTM and Barry-Ryan C, Effect of ozone  
546 and calcium lactate treatments on browning and texture properties of fresh-cut lettuce. *J Sci*  
547 *Food Agric* **86**:2179-2188 (2006).
- 548 18. ASTM, *Physical requirements guidelines for sensory evaluation laboratories*. American  
549 Society for Testing Materials, Philadelphia, USA (1986).
- 550 19. ISO\_8586:2012, *Sensory analysis—General guidance for selection, training and*  
551 *monitoring of assessors*. ISO, Geneva, Switzerland (2012).
- 552 20. Zapata S and Dufour J-P, Ascorbic, dehydroascorbic and isoascorbic acid simultaneous  
553 determinations by reverse phase ion interaction HPLC. *J Food Sci* **57**:506-511 (1992).

- 554 21. Martínez-Hernández GB, Artés-Hernández F, Colares-Souza F, Gómez PA, García-  
555 Gómez P and Artés F, Innovative cooking techniques for improving the overall quality of a  
556 kailan-hybrid broccoli. *Food Bioproc Tech* **6**:2135-2149 (2013).
- 557 22. Martínez-Hernández GB, Gómez PA, Pradas I, Artés F and Artés-Hernández F,  
558 Moderate UV-C pretreatment as a quality enhancement tool in fresh-cut Bimi® broccoli.  
559 *Postharvest Biol Technol* **62**:327-337 (2011).
- 560 23. Singleton VL, Orthofer R and Lamuela-Raventós RM, Analysis of total phenols and  
561 other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods*  
562 *Enzymol* **299**:152-178 (1999).
- 563 24. Brand-Williams W, Cuvelier ME and Berset C, Use of a free radical method to evaluate  
564 antioxidant activity. *Lebenson Wiss Technol* **28**:25-30 (1995).
- 565 25. Benzie IF and Strain JJ, Ferric reducing/antioxidant power assay: Direct measure of  
566 total antioxidant activity of biological fluids and modified version for simultaneous  
567 measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol*  
568 **299**:15-27 (1999).
- 569 26. Cano A, Hernández-Ruiz J, García-Cánovas F, Acosta M and Arnao MB, An end-point  
570 method for estimation of the total antioxidant activity in plant material. *Phytochem Anal*  
571 **9**:196-202 (1998).
- 572 27. Di Cagno R, Minervini G, Rizzello CG, De Angelis M and Gobbetti M, Effect of lactic acid  
573 fermentation on antioxidant, texture, color and sensory properties of red and green  
574 smoothies. *Food Microbiol* **28**:1062-1071 (2011).
- 575 28. Gómez-Ordóñez E, Jiménez-Escrig A and Rupérez P, Dietary fibre and physicochemical  
576 properties of several edible seaweeds from the northwestern Spanish coast. *Food Res Int*  
577 **43**:2289-2294 (2010).
- 578 29. Carvalho AFU, Portela MCC, Sousa MB, Martins FS, Rocha FC, Farias DF and Feitosa  
579 JPA, Physiological and physico-chemical characterization of dietary fibre from the green  
580 seaweed *Ulva fasciata* Delile. *Braz J Biol* **69**:969-977 (2009).
- 581 30. Castillejo N, Martínez-Hernández GB, Monaco K, Gómez PA, Aguayo E, Artés F and  
582 Artés-Hernández F, Preservation of bioactive compounds of a green vegetable smoothie using  
583 short time-high temperature mild thermal treatment. *Food Sci Technol Int* **23**:46-60 (2017).
- 584 31. Ortega-Calvo JJ, Mazuelos C, Hermosin B and Saiz-Jimenez C, Chemical composition of  
585 Spirulina and eukaryotic algae food products marketed in Spain. *J Appl Phycol* **5**:425-435  
586 (1993).
- 587 32. Wang R, Xu Q, Yao J, Zhang Y, Liao X, Hu X, Wu J and Zhang Y, Post-effects of high  
588 hydrostatic pressure on green color retention and related properties of spinach puree during  
589 storage. *Innov Food Sci Emerg Technol* **17**:63-71 (2013).
- 590 33. Hanschen FS, Klopsch R, Oliviero T, Schreiner M, Verkerk R and Dekker M, Optimizing  
591 isothiocyanate formation during enzymatic glucosinolate breakdown by adjusting pH value,  
592 temperature and dilution in *Brassica* vegetables and *Arabidopsis thaliana*. *Scientific Reports*  
593 **7**:40807 (2017).
- 594 34. Uda Y, Kurata T and Arakawa N, Effects of pH and Ferrous Ion on the Degradation of  
595 Glucosinolates by Myrosinase. *Agric Biol Chem* **50**:2735-2740 (1986).
- 596 35. Welch AM, Preliminary survey of fungistatic properties of marine algae. *J Bacteriol*  
597 **83**:97-99 (1962).
- 598 36. Deutsch JC, Spontaneous hydrolysis and dehydration of dehydroascorbic acid in  
599 aqueous solution. *Anal Biochem* **260**:223-229 (1998).
- 600 37. Munyaka AW, Makule EE, Oey I, Van Loey A and Hendrickx M, Thermal stability of L-  
601 ascorbic acid and ascorbic acid oxidase in broccoli (*Brassica oleracea* var. *italica*). *J Food Sci*  
602 **75**:C336-340 (2010).
- 603 38. Lee SK and Kader AA, Preharvest and postharvest factors influencing vitamin C content  
604 of horticultural crops. *Postharvest Biol Technol* **20**:207-220 (2000).

- 605 39. Batth R, Singh K, Kumari S and Mustafiz A, Transcript profiling reveals the presence of  
606 abiotic stress and developmental stage specific ascorbate oxidase genes in plants. *Front Plant*  
607 *Sci* **8**:198 (2017).
- 608 40. Hess AF and Unger LJ, The destruction of the antiscorbutic vitamin in milk by the  
609 catalytic action of minute amounts of copper. *Proc Soc Exp Biol Med* **19**:119-120 (1921).
- 610 41. FAO/WHO, *Vitamin and mineral requirements in human nutrition*. World Health  
611 Organization and Food and Agriculture Organization of the United Nations, Bangkok (2004).
- 612 42. Rose RC, Solubility properties of reduced and oxidized ascorbate as determinants of  
613 membrane permeation. *Biochim Biophys Acta* **924**:254-256 (1987).
- 614 43. Bradshaw MP, Barril C, Clark AC, Prenzler PD and Scollary GR, Ascorbic acid: A review  
615 of its chemistry and reactivity in relation to a wine environment. *Crit Rev Food Sci Nutr* **51**:479-  
616 498 (2011).
- 617 44. Andrés V, Villanueva MJ and Tenorio MD, The effect of high-pressure processing on  
618 colour, bioactive compounds, and antioxidant activity in smoothies during refrigerated  
619 storage. *Food Chem* **192**:328-335 (2016).
- 620 45. Keenan DF, Brunton NP, Gormley TR, Butler F, Tiwari BK and Patras A, Effect of thermal  
621 and high hydrostatic pressure processing on antioxidant activity and colour of fruit smoothies.  
622 *Innov Food Sci Emerg Technol* **11**:551-556 (2010).
- 623 46. Watanabe F-, Yabuta Y, Bito T and Teng F, Vitamin B(12)-containing plant food sources  
624 for vegetarians. *Nutrients* **6**:1861-1873 (2014).
- 625 47. AOAC, VitaFast® B12 microbiological microtiter test for the determination of vitamin  
626 B12. **101002** (2014).
- 627 48. Herbert V and Drivas G, Spirulina and vitamin B 12. *Jama* **248**:3096-3097 (1982).
- 628 49. van den Berg H, Daqnelie PC and van Staveren WA, Vitamin B12 and seaweed. *Lancet*  
629 **1**:242-243 (1988).
- 630 50. Dagnelie PC, van Staveren WA and van den Berg H, Vitamin B-12 from algae appears  
631 not to be bioavailable. *Am J Clin Nutr* **53**:695-697 (1991).
- 632 51. Watanabe F, Katsura H, Takenaka S, Fujita T, Abe K, Tamura Y, Nakatsuka T and Nakano  
633 Y, Pseudovitamin B(12) is the predominant cobamide of an algal health food, spirulina tablets.  
634 *J Agric Food Chem* **47**:4736-4741 (1999).
- 635 52. Takenaka S, Sugiyama S, Ebara S, Miyamoto E, Abe K, Tamura Y, Watanabe F, Tsuyama  
636 S and Nakano Y, Feeding dried purple laver (nori) to vitamin B12-deficient rats significantly  
637 improves vitamin B12 status. *Br J Nutr* **85**:699-703 (2001).
- 638 53. Eluvakkal T, Sivakumar SR and Arunkumar K, Fucoïdan in some Indian brown seaweeds  
639 found along the coast gulf of Mannar. *Int J Botany* **6**:176-181 (2010).
- 640 54. Zhao L, Peralta-Videa JR, Rico CM, Hernandez-Viezcas JA, Sun Y, Niu G, Servin A, Nunez  
641 JE, Duarte-Gardea M and Gardea-Torresdey JL, CeO(2) and ZnO nanoparticles change the  
642 nutritional qualities of cucumber (*Cucumis sativus*). *J Agric Food Chem* **62**:2752-2759 (2014).
- 643 55. Jiménez-Escrig A, Gómez-Ordóñez E and Rupérez P, Brown and red seaweeds as  
644 potential sources of antioxidant nutraceuticals. *J Appl Phycol* **24**:1123-1132 (2012).

645



646 **TABLE AND FIGURE CAPTIONS**

647

648 **Table 1.** Classification and details of the nine edible marine algae studied.

649

650 **Table 2.** Total soluble solids content (SSC, %), pH, titratable acidity (TA, expressed in  
651 %: g tartaric acid 100 g<sup>-1</sup>) and total colour differences ( $\Delta E$ ) of fresh fruit/vegetable  
652 smoothies with or without algae fortification stored at 5 °C (n=5±SD). Different capital  
653 letters denote significant differences ( $P \leq 0.05$ ) among smoothies for the same sampling  
654 day. Different lowercase letters denote significant differences ( $P \leq 0.05$ ) among sampling  
655 days for the same smoothie.

656

657 **Table 3.** Psychrophilic, and yeast and moulds counts (log CFU g<sup>-1</sup>) of fresh  
658 fruit/vegetable smoothies with or without algae fortification stored at 5 °C (n=5±SD).  
659 Different capital letters denote significant differences ( $P \leq 0.05$ ) among smoothies for  
660 the same sampling day. Different lowercase letters denote significant differences  
661 ( $P \leq 0.05$ ) among sampling days for the same smoothie.

662

663 **Table 4.** Vitamin C (mg kg<sup>-1</sup>) of fresh fruit/vegetable smoothies with or without algae  
664 fortification stored at 5 °C (n=5±SD). Different capital letters denote significant  
665 differences ( $P \leq 0.05$ ) among smoothies for the same sampling day. Different lowercase  
666 letters denote significant differences ( $P \leq 0.05$ ) among sampling days for the same  
667 smoothie.

668

669 **Table 5.** Total phenolic content (TPC, mg gallic acid equivalent kg<sup>-1</sup>) and total  
670 antioxidant capacity (three methods; mg Trolox equivalent kg<sup>-1</sup>) of fresh fruit/vegetable

671 smoothies with or without algae fortification stored at 5 °C (n=5±SD). Different capital  
672 letters denote significant differences ( $P \leq 0.05$ ) among smoothies for the same sampling  
673 day. Different lowercase letters denote significant differences ( $P \leq 0.05$ ) among sampling  
674 days for the same smoothie.

675

676 **Figure 1.** Sensory attributes of fresh fruit/vegetable smoothies with or without algae  
677 fortification on processing day and after 14, 17 and 1 days at 5 °C (n=5±SD).

678

679 **Figure 2.** Vitamin B12 ( $\mu\text{g kg}^{-1}$ ) of fresh fruit/vegetable smoothies with algae  
680 fortification on processing day and after 24 days at 5 °C (n=5±SD). Different capital  
681 letters denote significant differences ( $P \leq 0.05$ ) among smoothies for the same sampling  
682 day. Different lowercase letters denote significant differences ( $P \leq 0.05$ ) among sampling  
683 days for the same smoothie.