



UNIVERSITI  
TEKNOLOGI  
MARA

Institut  
Pengajian  
Siswazah

# THE DOCTORAL RESEARCH ABSTRACTS

TWELFTH  
ISSUE

Volume: 12, Issue 12

October 2017

IGS Biannual Publication



**Name :** SALWA MOHAMMED RAWEH ABDULLAH AL-FAQEER

**Title :** PHYTOCHEMICAL AND ANTIOXIDANT STUDIES OF MALYSIAN MEDICINAL PLANTS SYZYGIUM POLYANTHUM AND OCTOMELES SUMATRANA

**Supervisor :** PROF. DR. JEAN FREDERIC WEBER @ FAIZAL WEBER (MS)  
 PROF. DR. AISHAH ADAM (CS)  
 DR. MIZATON HAZIZUL HASAN (CS)  
 DR. HUMERA NAZ (CS)

The leaves of *S. polyanthum* (Myrtaceae) and barks of *O. sumatrana* (Datiaceae) were investigated for their chemical constituents, antioxidant and cytoprotective activities. Their aqueous extracts were first subjected to acidic hydrolysis and the organic layers were dissolved in water and partitioned using hexane, ethyl acetate (EtOAc) and n-butanol (BuOH). Six compounds (betulinic acid, ellagic acid, kaempferol, myricetin, quercetin, and  $\beta$ -sitosterol) were isolated and identified from the EtOAc and BuOH extracts of *S. polyanthum* and four compounds (quercetin, kaempferol, rutin, bryonolic acid) were purified from the n-butanol extract of *O. sumatrana* by means of MPLC and HPLC. The structures of the above compounds were determined by comparing their NMR and LCMS-TOF data with reported values. The structure of bryonolic acid was further confirmed by X-ray crystallography. Eleven essential oil components ( $\alpha$ -caryophyllene,  $\beta$ -caryophyllene, caryophyllene oxide, 1,8-cineole,  $\beta$ -elemene, eugenol, eugenol acetate, isoeugenol,  $\alpha$ -pinene,  $\beta$ -pinene, terpinen-4-ol) were identified from *S. polyanthum* and four fatty acid derivatives (linoleic acid, methyl linoleate, myristic acid, palmitic acid) and three steroids [5,6-dihydroergosterol, ergosta-5,8(14)-dien-3 $\beta$ -ol, ergosta-5-en-3 $\beta$ -ol] were determined from *O. sumatrana* by GC-MS analysis of their hexane extracts. The hexane, EtOAc and BuOH extracts were subjected to DPPH, FRAP and cytoprotective activities. The EtOAc and BuOH extracts of both plants showed potent DPPH activity with the EC<sub>50</sub> values of 159.12  $\pm$  0.11  $\mu$ g/mL and 186.40  $\pm$  0.58  $\mu$ g/mL in *S. polyanthum* and 125.3  $\pm$  0.17  $\mu$ g/mL and 136.4  $\pm$  0.17  $\mu$ g/mL in *O. sumatrana*, respectively. It was found that bryonolic acid (EC<sub>50</sub> = 26.7  $\pm$  0.74) only marginally quenched DPPH radical

but ellagic acid, myricetin, quercetin, rutin and kaempferol (92.4  $\pm$  3.82, 74.1  $\pm$  1.29, 76.04  $\pm$  2.63, 76.8  $\pm$  1.11 and 71.22  $\pm$  1.09  $\mu$ M), respectively) showed strong DPPH radical scavenging activity. Then, the isolated compounds from *S. polyanthum* and *O. sumatrana* (myricetin, ellagic acid, betulinic acid,  $\beta$ -sitosterol, rutin, quercetin, kaempferol and bryonolic acid) were tested for their cytotoxic effects towards three types of cells including normal human embryonic liver (WRL-68), normal green monkey kidney (Vero) and human hepatocarcinoma (HepG2) cell lines. The cells were treated with different concentrations of the compounds and the results showed that the compounds from *S. polyanthum* and *O. sumatrana* were non-toxic towards normal cells. However, betulinic acid and bryonolic acid had high cytotoxicity towards HepG2 cells. Next, the cytoprotective effects of the isolated compounds against hydrogen peroxide-induced WRL-68 and Vero cells were investigated. Quercetin, kaempferol, myricetin, ellagic acid, betulinic acid,  $\beta$ -sitosterol and bryonolic acid showed significant protective effects compared to control against oxidative stress-induced WRL-68 and Vero cells. Furthermore, betulinic acid and bryonolic acid showed higher protective effect compared to ellagic acid, kaempferol, myricetin and quercetin and the activities of the antioxidative enzymes such as superoxide dismutase (SOD) and catalase (CAT) were enhanced in a dose-dependent manner. In conclusion, this study demonstrated that most compounds from *S. polyanthum* and *O. sumatrana* were cytoprotective against oxidative stress induced by H<sub>2</sub>O<sub>2</sub> with betulinic acid and bryonolic acid having the highest potential to be developed to be used as anticancer candidates and alternative medicine.