

- **12408 | Fracture of the proximal third of the femur and BMI in elderly women**

Rodrigues, Cinara, Faculdade de Ciências da Nutrição e Alimentação, Portugal

Correia, Flora, Faculdade de Ciências da Nutrição e Alimentação, Portugal

Oliveira, Bruno, Faculdade de Ciências da Nutrição e Alimentação, Portugal

Background: Increasing life expectancy has provided a demographic aging, which results in an increased prevalence of chronic diseases, as well as falls and bone fractures in elderly. Regarding the fractures of the proximal third of the femur, it's known that they are the most common traumatic injuries among elderly, with high morbidity and mortality. The relationship between Body Mass Index (BMI) and the prognosis of these episodes remains unclear. The aim of this study was to evaluate this possible association.

Methodology: The study group consisted of 40 women aged between 65 and 99 years old, hospitalized in Hospital São João' Orthopedic and Traumatology Department due to a fracture of the proximal third of the femur. The control group was composed of 36 institutionalized elderly women, aged between 65 and 94 years old, without any history of this type of fracture. We carried out anthropometric measurements (weight and height), body composition assessment by bioelectrical impedance analysis, sociodemographic and clinical data collection and cognitive and functional assessment.

Results: The control group mean BMI was 28,3kg/m² (sd=6,0) and the study group mean BMI was 25.5kg/m² (sd=4,9), which is significantly lower (p=0,029). We also noted that the fracture risk is lower in single or divorced women (p=0,020), smaller (p=0,016), with higher BMI (p=0,012) and without hypertension (p=0,016). By analysing the ROC curve, we can see that a BMI below 22,6kg/m² is predictor of fracture risk.

Conclusion: We found that in elderly women, a higher BMI protects them from the fracture of the proximal third of the femur, therefore we should not promote a weight loss corresponding to a BMI below 22,6kg/m², which is near the lower limit of the range proposed by Lipschitz, in 1994.

Keywords: Aging; falls; fracture of the proximal third of the femur; BMI.