



Core skin DC signatures control immune tolerance to skin cancer and limit anti-tumor immunity

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Nirschl, Christopher, Yong Liu, Kavita Sarin, Mayte Suarez-Farinas, David Chau, Peter Sage, Arlene Sharpe, and Niroshana Anandasabapathy. 2015. "Core skin DC signatures control immune tolerance to skin cancer and limit anti-tumor immunity." <i>Journal for Immunotherapy of Cancer</i> 3 (Suppl 2): P205. doi:10.1186/2051-1426-3-S2-P205. http://dx.doi.org/10.1186/2051-1426-3-S2-P205 .
Published Version	doi:10.1186/2051-1426-3-S2-P205
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:23845199
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

POSTER PRESENTATION

Open Access

Core skin DC signatures control immune tolerance to skin cancer and limit anti-tumor immunity

Christopher Nirschl¹, Yong Liu¹, Kavita Sarin², Mayte Suarez-Farinas³, David Chau¹, Peter Sage⁴, Arlene Sharpe⁴, Niroshana Anandasabapathy^{1*}

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Background

Dendritic cells (DC) are gatekeepers of immunity, critical to both an initiate immune response upon infection and promote tolerance to self-antigens.

Methods

We have just established that DC subsets in the skin that constitutively migrate into LNs can individually and collectively temper an on-going immune response. Tissue DC originating in skin share a unique transcriptome signature geared towards immune dampening when compared to their lymphoid counterparts in both mouse and human.

Results

Here we demonstrate expression of unique core skin DC transcripts are closely associated with increased clinical aggressiveness of BCC in humans and the stratify stage 4 melanoma outcomes. In mice, loss of signature genes in DC, which include but are not limited to PD-L1/PD-L2, lead to enhanced antigen-specific immunity, decreased tumor growth, and improved anti-tumor vaccine priming to melanoma skin cancer by distinguishable molecular control of T cell effector function and clonal proliferation.

Conclusions

These data suggest core skin DC signatures regulate the immune-epithelial interface.

Authors' details

¹Brigham and Women's Hospital, Department of Dermatology, Harvard Medical School, Boston, MA, USA. ²Stanford University, Stanford, CA, USA. ³Mount Sinai school of Medicine, New York, NY, USA. ⁴Harvard Medical School, Boston, MA, USA.

Published: 4 November 2015

¹Brigham and Women's Hospital, Department of Dermatology, Harvard Medical School, Boston, MA, USA

Full list of author information is available at the end of the article

doi:10.1186/2051-1426-3-S2-P205

Cite this article as: Nirschl *et al.*: Core skin DC signatures control immune tolerance to skin cancer and limit anti-tumor immunity. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P205.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

