



Broad targeting of angiogenesis for cancer prevention and therapy

The Harvard community has made this
article openly available. [Please share](#) how
this access benefits you. Your story matters

| | |
|-------------------|---|
| Citation | Wang, Z., C. Dabrosin, X. Yin, M. M. Fuster, A. Arreola, W. K. Rathmell, D. Generali, et al. 2015. "Broad targeting of angiogenesis for cancer prevention and therapy." <i>Seminars in Cancer Biology</i> 35 (Suppl): S224-S243. doi:10.1016/j.semcancer.2015.01.001. http://dx.doi.org/10.1016/j.semcancer.2015.01.001 . |
| Published Version | doi:10.1016/j.semcancer.2015.01.001 |
| Citable link | http://nrs.harvard.edu/urn-3:HUL.InstRepos:25658434 |
| Terms of Use | This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA |



Review

Broad targeting of angiogenesis for cancer prevention and therapy



Zongwei Wang^{a,**}, Charlotta Dabrosin^{b,c}, Xin Yin^d, Mark M. Fuster^d, Alexandra Arreola^e, W. Kimryn Rathmell^e, Daniele Generali^f, Ganji P. Nagaraju^g, Bassel El-Rayes^g, Domenico Ribatti^{h,i}, Yi Charlie Chen^j, Kanya Honoki^k, Hiromasa Fujii^k, Alexandros G. Georgakilas^l, Somaira Nowsheen^m, Amedeo Amedeiⁿ, Elena Niccolaiⁿ, Amr Amin^{o,p}, S. Salman Ashraf^q, Bill Helferich^r, Xujuan Yang^r, Gunjan Guha^s, Dipita Bhakta^s, Maria Rosa Ciriolo^t, Katia Aquilano^t, Sophie Chen^u, Dorota Halicka^v, Sulma I. Mohammed^w, Asfar S. Azmi^x, Alan Bilsland^y, W. Nicol Keith^y, Lasse D. Jensen^{z,A,*}

^a Department of Urology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

^b Department of Oncology, Linköping University, Linköping, Sweden

^c Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden

^d Medicine and Research Services, Veterans Affairs San Diego Healthcare System & University of California, San Diego, San Diego, CA, USA

^e Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA

^f Molecular Therapy and Pharmacogenomics Unit, AO Isituti Ospitalieri di Cremona, Cremona, Italy

^g Department of Hematology and Medical Oncology, Emory University, Atlanta, GA, USA

^h Department of Basic Medical Sciences, Neurosciences and Sensory Organs, University of Bari Medical School, Bari, Italy

ⁱ National Cancer Institute Giovanni Paolo II, Bari, Italy

^j Department of Biology, Alderson Broaddus University, Philippi, WV, USA

^k Department of Orthopedic Surgery, Arthroplasty and Regenerative Medicine, Nara Medical University, Nara, Japan

^l Physics Department, School of Applied Mathematics and Physical Sciences, National Technical University of Athens, Athens, Greece

^m Mayo Graduate School, Mayo Clinic College of Medicine, Rochester, MN, USA

ⁿ Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy

^o Department of Biology, College of Science, United Arab Emirate University, United Arab Emirates

^p Faculty of Science, Cairo University, Cairo, Egypt

^q Department of Chemistry, College of Science, United Arab Emirate University, United Arab Emirates

^r University of Illinois at Urbana Champaign, Urbana, IL, USA

^s School of Chemical and Bio Technology, SASTRA University, Thanjavur, India

^t Department of Biology, University of Rome "Tor Vergata", Rome, Italy

^u Ovarian and Prostate Cancer Research Trust Laboratory, Guilford, Surrey, UK

^v New York Medical College, New York City, NY, USA

^w Department of Comparative Pathobiology, Purdue University Center for Cancer Research, West Lafayette, IN, USA

^x School of Medicine, Wayne State University, Detroit, MI, USA

^y Institute of Cancer Sciences, University of Glasgow, Glasgow, UK

^z Department of Medical, and Health Sciences, Linköping University, Linköping, Sweden

^A Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden

ARTICLE INFO

Article history:

Available online 16 January 2015

Keywords:

Angiogenesis
Cancer
Phytochemicals
Treatment
Anti-angiogenic

ABSTRACT

Deregulation of angiogenesis – the growth of new blood vessels from an existing vasculature – is a main driving force in many severe human diseases including cancer. As such, tumor angiogenesis is important for delivering oxygen and nutrients to growing tumors, and therefore considered an essential pathologic feature of cancer, while also playing a key role in enabling other aspects of tumor pathology such as metabolic deregulation and tumor dissemination/metastasis. Recently, inhibition of tumor angiogenesis has become a clinical anti-cancer strategy in line with chemotherapy, radiotherapy and surgery, which underscore the critical importance of the angiogenic switch during early tumor development. Unfortunately the clinically approved anti-angiogenic drugs in use today are only effective in a subset of the

* Corresponding author at: Linköping University, Department of Medical and Health Sciences, Farmakologen, Ingång 68, Pl. 08, SE-581 85 Linköping, Sweden.

Tel.: +46 101034004.

** Corresponding author at: Massachusetts General Hospital, Harvard Medical School, Department of Urology, 55 Fruit Street, Warren Building 324, Boston, MA 02114, USA.

Tel.: +1 617 643 1956.

E-mail addresses: zwang0@partners.org (Z. Wang), lasse.jensen@liu.se (L.D. Jensen).

patients, and many who initially respond develop resistance over time. Also, some of the anti-angiogenic drugs are toxic and it would be of great importance to identify alternative compounds, which could overcome these drawbacks and limitations of the currently available therapy. Finding “the most important target” may, however, prove a very challenging approach as the tumor environment is highly diverse, consisting of many different cell types, all of which may contribute to tumor angiogenesis. Furthermore, the tumor cells themselves are genetically unstable, leading to a progressive increase in the number of different angiogenic factors produced as the cancer progresses to advanced stages. As an alternative approach to targeted therapy, options to broadly interfere with angiogenic signals by a mixture of non-toxic natural compound with pleiotropic actions were viewed by this team as an opportunity to develop a complementary anti-angiogenesis treatment option. As a part of the “Halifax Project” within the “Getting to know cancer” framework, we have here, based on a thorough review of the literature, identified 10 important aspects of tumor angiogenesis and the pathological tumor vasculature which would be well suited as targets for anti-angiogenic therapy: (1) endothelial cell migration/tip cell formation, (2) structural abnormalities of tumor vessels, (3) hypoxia, (4) lymphangiogenesis, (5) elevated interstitial fluid pressure, (6) poor perfusion, (7) disrupted circadian rhythms, (8) tumor promoting inflammation, (9) tumor promoting fibroblasts and (10) tumor cell metabolism/acidosis. Following this analysis, we scrutinized the available literature on broadly acting anti-angiogenic natural products, with a focus on finding qualitative information on phytochemicals which could inhibit these targets and came up with 10 prototypical phytochemical compounds: (1) oleanolic acid, (2) tripterine, (3) silibinin, (4) curcumin, (5) epigallocatechin-gallate, (6) kaempferol, (7) melatonin, (8) enterolactone, (9) withaferin A and (10) resveratrol. We suggest that these plant-derived compounds could be combined to constitute a broader acting and more effective inhibitory cocktail at doses that would not be likely to cause excessive toxicity. All the targets and phytochemical approaches were further cross-validated against their effects on other essential tumorigenic pathways (based on the “hallmarks” of cancer) in order to discover possible synergies or potentially harmful interactions, and were found to generally also have positive involvement in/effects on these other aspects of tumor biology. The aim is that this discussion could lead to the selection of combinations of such anti-angiogenic compounds which could be used in potent anti-tumor cocktails, for enhanced therapeutic efficacy, reduced toxicity and circumvention of single-agent anti-angiogenic resistance, as well as for possible use in primary or secondary cancer prevention strategies.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction to tumor angiogenesis

Vessel formation in both health and disease occur through either vasculogenesis – i.e. the recruitment of bone marrow-derived endothelial progenitor cells to form new vessels, angiogenesis – i.e. the sprouting and growth of new vessels from an existing vasculature or intussusception – i.e. the division or splitting of a blood vessel into two or more new vessels [1]. The most common pathway for neo-vessel growth in malignancy is angiogenesis (reviewed in [2]) and the process is therefore called tumor angiogenesis.

In 1971, Judah Folkman first advanced the hypothesis that tumor growth depends on angiogenesis [3]. According to this hypothesis, endothelial cells may be switched from a resting state to a rapid growth phase by a diffusible chemical signal emanating from the tumor cells. The switch depends on increased production of one or more positive regulators of angiogenesis, such as vascular endothelial growth factor (VEGF), fibroblast growth factor-2 (FGF-2), interleukin-8 (IL-8), placental growth factor (PlGF), transforming growth factor-beta (TGFbeta), platelet derived growth factor (PDGF), angiopoietins (Angs) and others (reviewed in [4]). These can be exported from tumor cells, mobilized from the extracellular matrix, or released from host cells recruited to the tumor. The switch may also involve down-regulation of endogenous inhibitors of angiogenesis such as endostatin, angiostatin or thrombospondin (reviewed in [5]) and has thus been regarded as the result of tipping the net balance between positive and negative regulators. Mature microRNAs (miRNAs) can furthermore regulate the levels of pro- or anti-angiogenic gene expression at the post-transcriptional level (reviewed in [6]).

Angiogenic signals lead to the preferential differentiation of certain endothelial cells into so-called tip cells, which start to migrate and exist at the leading front of the growing vessels. A number of factors including VEGF receptor (VEGFR)-3 (for lymphatic endothelial cells), VEGFR-1 and -2 (for blood endothelial cells), PDGF-B, and the Notch ligand delta-like ligand (DLL)-4 have been

shown to contribute to the endothelial tip cell phenotype [7,8]. In healthy angiogenesis during development for example, the number of tip-cells are limited leading to an orderly and organized expansion of the vasculature. Endothelial cells located behind the tip cell, so-called stalk cells, express other factors such as VEGFR-1 and Notch-1 and -4 which are important for inducing a quiescent state of these cells [9,10], maturation of the vascular wall, lumen formation and to support perfusion. However, in pathological angiogenesis including tumor angiogenesis this process is usually disrupted by either excess production of pro-angiogenic signals, lack of angiogenesis inhibitors, path-finding signals or maturation factors, thus leading to excessive tip-cell formation and migration of endothelial cells [11,12], which do not assume a quiescent phenotype associated with a healthy vasculature.

1.1. Structural and dysfunctional features of tumor blood vessels

As a result of the imbalance of angiogenic activators and inhibitors, tumor blood vessels display many structural and functional abnormalities including unusual leakiness (reviewed in [13]), potential for rapid growth and remodeling [14], high tortuosity and sinusoidal appearance (reviewed in [13]), poor coverage by vascular supportive cells including pericytes and smooth muscle cells [15], lack of arterial or venous identity leading to chaotic blood flow, poor functionality and perfusion [16], incorporation of tumor stem-like cells to endothelial cells which contribute to the tumor vasculature – a process known as vascular mimicry [17]. These phenotypes, which can be considered “hallmarks of the tumor vasculature”, mediate the dissemination of tumor cells in the bloodstream and maintain the pathological characteristics of the tumor microenvironment.

Tumor vessel density is furthermore very heterogeneous: the highest values are found in the invading tumor edge, where the density is 4–10 times greater than inside the tumor and

the arrangement of vessels in the center of a tumor is much more chaotic than at its edges (reviewed in [18]). Importantly, mechanical stress generated by proliferating tumor cells also compress vessels in tumors, with some vessels being oversized, whereas others are more immature and smaller. These structural abnormalities result in disturbed blood flow, hypoxia, hyperpermeability, and elevated interstitial pressure in many solid tumors (reviewed in [19]), responsible, in turn, for impaired delivery of anti-cancer drugs as well as oxygen, the former being critical for the success of chemotherapy and the latter for radiation therapy.

1.2. Tumor hypoxia – an emerging target

There is a complex interrelationship between tumor hypoxia and tumor angiogenesis. The production of several angiogenic cytokines and growth factors is regulated by hypoxia, but as mentioned, tumor angiogenesis also further elevates tumor hypoxia. This vicious circle is critical for driving many of the most pathogenic features of cancer including poor treatment outcome and progression to severe and metastatic disease.

Much of the dynamic regulation of this process involves transcriptionally mediated changes that promote the enhanced production of ligands and receptors, signaling aspects of both mitogenic growth and directing the organization of an endothelial network. While hypoxia and hypoxia signaling play major roles, they are only two means by which angiogenesis can be triggered. Other processes involving proteomic signaling, including those effects directly attributed to the tumor biology [20], can also be activated in support of the angiogenic response to various tissue conditions.

Hypoxia in tumors develops in the form of chronic hypoxia, resulting from long diffusion distances between (perfused) tumor vessels, and/or of acute hypoxia, resulting from a transient collapse of tumor vessels. One essential pathway activated in this series of events is the activation of hypoxia inducible factors (HIFs), heterodimeric transcription factors composed from alpha and beta subunits, which can be rapidly stabilized to fluidly adapt to and overcome the effects of a hypoxic environment. There are three HIFalpha subunits, HIF1alpha, HIF2alpha, HIF3alpha, which are regulated in an oxygen dependent manner. HIFalpha subunits are hydroxylated by prolyl hydroxylases (PHDs) [21] allowing HIFalpha subunits to be recognized by the von Hippel–Lindau (VHL) ubiquitin-ligase complex [22]. VHL poly-ubiquitinates HIFalpha subunits, leading to their subsequent proteasome-mediated degradation. Under low oxygen conditions PHDs have reduced activity, allowing for HIFalpha subunits to escape VHL-mediated degradation. HIFalpha subunits accumulate in the cytoplasm where they bind HIFbeta to form a heterodimer that subsequently translocates to the nucleus to activate transcription of target genes, including genes important for various processes such as metabolism (glucose transporter (GLUT)-1, hexokinase (HK)-1), cell growth (cyclin (CCN)-D1 [23]) and also angiogenesis, such as erythropoietin, VEGF and PDGF [24] (summarized in Fig. 1). In some cancers, mutations of the machinery (such as VHL or other components) regulating HIF stability can result in oxygen-independent constitutive stabilization of the HIFalpha factors, and as a result these tumors are notoriously highly vascularized [25].

Additional factors besides HIF-mediated VEGF transcriptional activation have also been identified as promoting VEGF expression under hypoxic conditions. Environmental stress as a result of low oxygen and proper nutrient deprivation, such as glucose deprivation, are capable of inducing VEGF mRNA stabilization resulting in increased levels of the secreted ligand and angiogenic growth [26]. Hypoxic stress has also been described as inducing changes in miRNAs which can further influence the host microenvironment with effects on angiogenesis [27].

Targeting hypoxia signaling is a promising approach to provide more options for intervention in this critical pathway of tumor/host biology. Additionally, HIFalpha factors can be regulated by mammalian target of rapamycin (mTOR) family translational signals (Fig. 1), which provides a rich alternate source of targeting (reviewed in [28]), and mTOR drugs are already in the market or emerging.

1.3. Tumor lymphangiogenesis and lymphatic metastasis

Metastatic spread of tumor cells, via either blood or lymphatic vascular systems, accounts for the majority of morbidity and mortality in cancer patients. The presence of tumor cells within sentinel lymph nodes (LN) that accept afferent lymphatic vessels draining lymph from the primary tumor often indicates initial metastasis that precedes (or predicts) distant metastasis to other organs. It is also one of the most important markers for predicting patient prognosis and deciding on therapeutic options [29,30].

Lymphangiogenesis, is often enhanced in malignant tumors, and associated with positive LN metastasis as well as poor survival of cancer patients [31–35]. Tumor-associated lymphangiogenesis may occur either at the immediate tumor periphery (peri-tumoral lymphatics) or within the tumor mass (intra-tumoral lymphatics), the former having been demonstrated to be functionally responsible for tumor cell dissemination [36,37]. Tumor lymphangiogenesis is – as tumor blood- (or hem)angiogenesis – also regulated by a balance of pro- and anti-lymphangiogenesis factors. The most frequently studied tumor lymphangiogenic factors are members of the VEGF family, most predominantly VEGF-C and -D, through interactions with in particular VEGFR-3, with some additional evidence for VEGF-A interacting with VEGFR-2 (reviewed in [38]). These factors were found to increase LN metastasis and their expression correlated with poor prognosis in both animal models and human cancers [39–44]. Other important lymphangiogenic factors include FGF-2 and the key receptor FGF receptor (FGFR)-1 [45–47], hepatocyte growth factor (HGF) and the cognate receptor c-met [48], insulin-like growth factors (IGF)-1, -2 and IGF receptor (IGFR) [49,50], EphrinB-2 and Eph receptor tyrosine kinase [51], Ang-1, -2 and Tie2 [52], PDGF-BB and PDGF receptor (PDGFR)alpha and -beta [53], growth hormone and the growth hormone receptor [54], among others.

Tumor cells not only stimulate lymphangiogenesis within or around the primary tumor site, but also have the capability to induce neo-lymphangiogenesis in the LN itself, so as to prepare a “pro-metastatic niche” for the spread of tumor cells [55–57]. Lymphangiogenesis at the sentinel LN appears to occur beforehand and is further enhanced upon the arrival of metastatic cancer cells [55,56], suggesting that the LN (possibly conditioned by certain tumor effectors, such as VEGF-A or VEGF-C) helps to provide a favorable environment for tumor metastasis (reviewed in [58]).

As lymphangiogenesis is associated with increased LN metastases (reviewed in [59]), blocking the process (or possibly inducing lymphatic endothelial apoptosis/regression) may serve as a favorable strategy to prevent lymph node metastasis. However, even if the strategy may result in fewer lymph-borne metastases over time, there's still the possibility that other biophysical functions are affected by the alteration in lymphatic flow within and surrounding the tumor. For example, a reduction in lymphatic drainage from the tumor may result in increased interstitial fluid pressure (IFP) within the tumor (reviewed in [60]). This in turn may increase tumor necrosis, hypoxia, and progression, while (at least temporarily) reducing the ability to deliver chemotherapy or other agents via the compressed tumor vasculature.

Adding to the complexity of the regulation and functions of tumor lymphatics, a variety of factors have been found to play key roles in the separation of lymphatic vasculature from blood




Fig. 1. Molecular mechanisms behind HIF regulation and responses in cells. The cellular oxygen sensing response is tightly regulated by a family of prolyl hydroxylases (PHD) which under normal oxygen conditions (normoxia; blue arrows) are responsible for hydroxylating proline residues on hypoxia inducible factor (HIF) alpha subunits. These hydroxylated residues are recognized by a pVHL-E3 ubiquitin ligase complex, whereby HIFalpha subunits are marked for polyubiquitination and subsequent proteosomal degradation. When oxygen levels are low (hypoxia; red arrow) PHDs cannot hydroxylate HIFalphas thereby allowing them to escape pVHL-mediated degradation. HIFalpha subunits accumulate and bind to their heterodimeric partner, HIFbeta, translocate into the nucleus and activate a cascade of hypoxic signaling pathways. Alternatively, c-Src is also capable of activating HIFs by indirectly inhibiting PHD activity via the NADPH oxidase/Rac pathway. mTOR can also promote stabilization and HIF transcriptional activity. Critical points for therapeutic intervention include the use of c-Src and mTOR inhibitors to prevent HIFalpha accumulation and activation.

vessels during development. While these pathways may also contribute to neolymphatic outgrowth in tumors [61], the application of inhibitory strategies to block the relevant pathways/effectors (which include podoplanin, spleen tyrosine kinase (SYK)/SH2-domain containing leukocyte protein of 76 kDa (SLP-76), Rac1/Ras homology gene family member (Rho), and sprout-related EVH1-domain containing (Spred)-1, -2 molecules) and examine the effects on tumor lymphatic investment as well as tumor vascular progression/remodeling has not been examined till date. Whether this might disrupt (or complement the inhibition of) tumor lymphangiogenesis while maintaining the delivery of chemotherapy to tumors during the relevant phase of treatment in the wider cancer treatment program remains to be examined.

1.4. Disrupted circadian rhythms in cancer

Social and occupational jetlag is a consequence of the disruption of our internal time-keeping system known as the circadian clock. Social jetlag arise in people who are often rotating between day and night shifts – often seen in healthcare workers, but a feature that is becoming increasingly prevalent in people employed in other types of jobs as well [62]. Occupational jetlag arises from traveling across several time zones and thus exposing oneself to a prolonged, unnatural day/light period, which is often experienced by airline pilots, cabin personnel and globally acting businessmen. It is becoming increasingly clear that such disruptions in the circadian rhythm are associated with higher risk of various diseases, most prominently sleep, metabolic, cardiovascular disorders and cancer [63,64]. In a large epidemiological study following more than 100,000 American nurses over 10 years, it was found that nurses who worked rotating day and night shifts more than five times per month were at significantly increased risk of various types of cancer [65,66].

Recently several research groups have found that the circadian rhythm is intimately involved in regulation of angiogenesis both during development [64,67] as well as in disease [68–70]. As such, circadian transcription factors were found to directly regulate VEGF levels and were responsible for the elevated night-time spikes in VEGF which are very important for physiological, developmental angiogenesis [64,71]. Due to the wide-range of cancers that have been associated with disrupted circadian rhythms, and the profound role of angiogenesis in the development of malignancy, it is tempting to speculate that circadian disruption may be an important player in pathological tumor angiogenesis.

2. Angiogenesis enables essential tumorigenic pathways

During tumor progression the amount and complexity of deregulated pathways which are essential for full blown malignancy, increase. Whereas pathological deregulation of cell cycle control in (often epithelial) cells is the first step toward tumor development, it is becoming increasingly clear that most of the essential tumorigenic pathways that lead to cancer are dependent on pathological deregulation of non-malignant host cells, and in particular angiogenesis and tumor vascular functions. As such, the tumor vasculature enables pathological tumor metabolism, genetic instability, inflammation, microenvironmental disruption and tumor cell invasion/metastasis.

Tumors are often hypoxic in spite of high vascularization, due to the poor structure and functionality of tumor blood vessels [11,12]. Intratumoral hypoxia is, somewhat paradoxically, a main cause of high reactive oxygen species (ROS) formation within the tumor cells (reviewed in [72]), and also coupled to pathological tumor cell metabolism and acidosis (reviewed in [73,74]). Thus, improving the quality of the tumor vasculature has been considered a way to improve perfusion, reduce the pathological leakiness of the tumor

vessels and reduce tumor hypoxia [16,19], which would also result in a more stable tumor genome.

Tumor angiogenesis and pathological activation of the endothelium, tumor vessel leakiness and hypoxia-induced apoptosis/necrosis in the tumor core, are resulting in a massive recruitment and activation of inflammatory cells, such as lymphocytes, neutrophils, macrophages and mast cells. These cells communicate by means of a complex network of intercellular signaling pathways mediated by surface adhesion molecules, cytokines and their receptors. These infiltrating immune cells, generate an environment abundant in growth and angiogenic factors and are implicated in enhancing cancer growth and subsequent resistance to therapy [4,75]. The inflammatory cytokines interleukin (IL)-1 α and IL-1 β as well as a wide panel of other signaling molecules produced by infiltrated inflammatory cells including VEGF and matrix-metalloproteinases (MMPs) may contribute to angiogenesis, tumor proliferation, and local invasion of cancer [76,77].

The deregulated tumor vasculature not only affects the recruitment and activation of inflammatory cells, also cancer-associated fibroblasts (CAFs), myofibroblasts and a number of other cell types, which contribute to tumor progression and resistance to treatment [14,78], is activated by endothelial cell- or hypoxia-derived factors such as PDGF-BB. On the other hand, CAFs are also rich sources of tumor angiogenic growth factors and cytokines, and thereby play an active role in sustaining tumor angiogenesis and providing resistance to anti-angiogenic therapy.

Tumors may disseminate both by local invasion as well as via blood or lymph vessels (hematogenous or lymphatic dissemination). Clinically the latter two processes are the most problematic in most tumor types as they lead to multifocal metastases. Also in tumors that mostly disseminate locally (i.e. neurological cancers for example), this dissemination often occurs via the vasculature as tumor cells coopt the blood vessels and invade the tissue by crawling along the endothelium [79]. Thus in all cases blood and lymphatic vessels in or around the tumor are prerequisite for tumor invasion and metastasis. Metastasis is further enabled due to the poor structural integrity of the tumor blood vessels and pathological angiogenesis-associated tumor hypoxia. Recently, anti-angiogenic therapy has been reported to cause an increased metastatic phenotype, possibly via elevated tumor hypoxia and hypoxia-induced epithelial to mesenchymal transition (EMT) [80–82]. Anti-angiogenic therapy may however also increase the metastatic potential of tumor cells through adaptive resistance pathways not associated with hypoxia [83], indicating that anti-angiogenic therapy-induced changes in the tumor phenotype may lead to a more aggressive disease through a number of different mechanisms. It is therefore not clear if targeting the tumor vessels would be beneficial or detrimental from a tumor metastasis point of view. However, it may be possible to merely reduce tumor vascularization, improve the structure of the tumor blood vessels and perfusion in the tumor and thus reduce the pathological characteristics of the vessels, for example via sub-maximal dosing of anti-angiogenic drugs [19]. There are still not much clinical data supporting a beneficial role for promoting formation of less pathological vessels in tumors, and it is not known how to best achieve this in patients.

3. Targets for anti-angiogenic therapy

The complexity in the angiogenic system provides many targets for therapeutic intervention. On the other hand redundancy in the angiogenic pathways raises the possibility of resistance to selective therapeutic agents (reviewed in [84,85]). Examples of agents that target circulating angiogenic factors include monoclonal antibodies

targeted against VEGF (bevacizumab) [86] or fusion proteins that trap angiogenic factors (aflibercept or AMG386) [87]. Agents that target synthesis of angiogenic factors include inhibitors of mTOR, cyclo-oxygenase (COX) or heat shock protein 90 (HSP90) [88,89]. These groups of agents in addition to inhibiting the synthesis of angiogenic factors can inhibit several other aspects of cancer biology such as growth, resistance to apoptosis or metastasis. Agents that target the angiogenic receptors are mainly tyrosine kinase inhibitors (sorafenib, sunitinib, pazopanib, regorafenib or axitinib) with multiple targets [90–92]. These agents are currently being used in the treatment of several malignant diseases ranging from breast, lung, gastric, colorectal, hepatocellular, glioblastoma, and neuroendocrine tumors.

Similarly, there are several agents in clinical trials aimed at blocking lymphangiogenesis/metastasis, mostly via neutralizing VEGF-A, -C or -D-induced receptor activation. For example, a major approach involves application of a variety of tyrosine kinase inhibitors such as Ki23057, used to block gastric cancer spread in mice through blockade of VEGFR3 autophosphorylation [93]. Other agents in clinical testing include PTK787/ZK222584 (Phase III – colorectal cancer); BAY43-9006 (Phase II, multiple carcinomas); CEP7055 (Phase I – various malignancies); or JNJ-26483327 (Phase I for multiple advanced solid tumors). These, among other anti-lymphangiogenic agents under study are reviewed in Ref. [59].

It is worthwhile mentioning that hormone- and chemotherapy could also have anti-angiogenic activity, particularly metronomic therapy which refers to the frequent, even daily, administration of chemotherapy (e.g. cyclophosphamide, methotrexate or capecitabine) in doses below the maximum tolerated dose, for long periods of time, with no prolonged drug-free breaks [94,95]. Other chemotherapeutics, routinely used in clinic, may also have anti-angiogenic activity *in vitro* or *in vivo* [96] as: (1) paclitaxel [97], doxorubicin and thalidomide [98] which seems to be mediated via inhibition of VEGF and bFGF [99]; (2) celecoxib, which may cause a time-dependent reduction in circulating angiogenic markers; (3) bisphosphonates may have anti-angiogenic effects [100] via reduction of VEGF and PDGF serum levels [101]; (4) PI3K inhibitors (including rapamycin analogues as temsirolimus (CCI-779) and everolimus (RAD001)) decrease tumor angiogenesis [102–104] via the inhibition of HIF-1 α caused by the blockade of mTOR activity.

Trials that have combined monoclonal antibodies and tyrosine kinase inhibitors have given rise to an increase in the side effects profile. A more rational approach would be to consider combinations of agents that block production of angiogenic factors with such that target angiogenic factors or receptors. The rationale behind such combinations include the fact that anti-angiogenic agents can improve the delivery of cytotoxic agents to the tumor site, may alter hypoxia in the tumor and sensitize it to chemotherapy or may impede the ability of the tumor to recover from cytotoxic effects of chemotherapeutic agents [105]. As tumors express more than one angiogenic cytokine and the fact that during tumor progression the palette of tumor-derived angiogenic factors grows more and more complex, any single inhibitor would not be sufficient for achieving sustained anti-tumor responses [74]. We hypothesize that simultaneously hitting multiple important aspects of tumor angiogenesis, each outlined in detail in the sections above, with a cocktail of compounds might create a more effective treatment. This could particularly be the case for indications such as cancer prevention in high risk settings, or maintenance therapies.

To facilitate the use of plant-derived compounds in cancer treatment, we have selected 10 key mechanisms that lead to pathological growth and functions of the tumor vasculature, such as EC migration/tip cell formation, phenotypic changes in the tumor microenvironment or pathogenic activation of stromal cells

(macrophages or fibroblasts). These pathways (top row in Table 1) were selected as candidate targets for anti-angiogenic natural compound-based therapy development.

4. Therapeutic potential of plant-derived compounds

During the last decades, phytochemicals have gained significant recognition for their potential therapeutic uses in cancer [106–108]. Extensive research has revealed enormous potential and exciting pharmacological properties of plant-based medicinal compounds, and demonstrated synergistic effects in combination with other agents to inhibit tumor angiogenesis, although the use of phytochemicals alone is still a limited option for cancer treatment. Some phytochemicals used in cancer therapies demonstrate relatively low side-effects, and some even limit the side-effects of chemotherapeutics or anti-angiogenic drugs. Fruits, vegetables, cereals, pulses, legumes, herbs, spices and medicinal plants – such as *Artemisia annua* (Chinese wormwood), *Viscum album* (European mistletoe), *Curcuma longa* (turmeric), *Scutellaria baicalensis* (Chinese skullcap), *Vitis vinifera* (grape seed extract), *Magnolia officinalis* (Chinese magnolia tree), *Camellia sinensis* (green tea), *Ginkgo biloba* (ginkgo), *Poria cocos* (tuckahoe), *Zingiber officinalis* (ginger), *Panax ginseng* (ginseng), *Rabdosia rubescens hora* (rabdosia), and Chinese destagnation herbs – are all considered to be good sources of phytochemicals exhibiting anti-cancer, and in particular anti-angiogenesis activities. The active ingredients in these plants are sometimes extracted and given in doses higher than what can be achieved from consuming the plants of which they are derived in order to give stronger therapeutic effect.

Many medicinal herbs and purified phytochemicals have recently been evaluated for anti-lymphangiogenic and anti-angiogenic properties in cancer (reviewed in [109–111]). The potential mechanisms underlying their anti-lymphangiogenic features involve (1) the control on cell proliferation, tube formation and cell cycle progression of lymphatic endothelial cells, as exhibited by multiple compounds fractionated from Korean and Japanese *Saussureae radix*, *Psoraleae semen* and *Aurantii fructus immaturus* [112]; (2) the inhibition of COX-2 expression and IL-1beta production and the subsequent reduction in VEGF-C-induced VEGFR-3 phosphorylation, as observed with wogonin [113]; and (3) the down-regulation of VEGFR3 and small GTPases, as well as the inhibition of VEGFR3-mediated extracellular-signal regulated kinase (ERK)-1 and -2 phosphorylation by cryptotanshinone [114]. The mechanisms behind their anti-angiogenic effects include: (1) inhibition of MMPs, (2) prevention of capillary sprout formation, endothelial cell proliferation and migration, and (3) modulation of angiogenic ligand/receptor mediated signaling pathways.

MMP inhibition blocks the degradation of endothelial basement membrane proteins, a process that otherwise leads to increased permeability and poor structure and stability of the tumor vasculature as well as release of sequestered angiogenic factors. However, caution is warranted regarding the effects on MMP/tissue inhibitor of metalloproteinases (TIMP) system as MMP inhibitors in clinical trials have failed and even induced worse survival rates compared to placebo treated patients [115]. In addition, several experimental studies have shown potent anti-tumorigenic activities of many MMPs including MMP-9 [116–121]. In the case of TIMPs both tumor protective and tumor enhancing properties have been reported [122–125]. For example, in breast cancer patients, high tumor and serum levels of TIMP-1 have been associated with decreased response to chemotherapy and consequently reduced survival [126–128].

Targeting capillary sprout formation and directional endothelial cell migration may depend on diverse inhibitors of the Notch

Table 1
Effects of the selected targets for anti-angiogenic cancer therapy on other cancer “hallmarks”.

| | Other cancer hallmarks | | | | | | | | | |
|---|---|--|------------------|-----------------------------|---|--------------------------|---|---|---|--|
| | Angiogenesis targets | | | | | Other cancer hallmarks | | | | |
| | (Inhibit) endothelial cell migration/tip cell formation | (Reduce) structural abnormalities of tumor vessels | (Reduce) hypoxia | (Inhibit) lymphangiogenesis | (Reduce) elevated interstitial fluid pressure | (Reverse) poor perfusion | (Normalize) disrupted circadian rhythms | (Suppress) tumor promoting inflammation | (De-activate) tumor promoting fibroblasts | (Normalize) tumor cell metabolism/acidosis |
| Genetic instability | 0 | 0 | + [239] | 0 | 0 | 0 | + [240] | 0 | 0 | + [241] |
| Sustained proliferating signaling | 0 | 0 | + [242] | +/- [243,244] | + [245,246] | 0 | + [247,248] | 0 | + [249,250] | 0 |
| Tumor-promoting inflammation | + [251] | + [252] | + [253,254] | 0 | + [255] | + [252,256] | + [257] | NA | + [258] | + [259] |
| Evasion of anti-growth signaling | + [260] | 0 | +/- [261,262] | 0 | + [246] | +/- [263–265] | +/- [266,267] | + [268] | + [269,270] | + [271,272] |
| Resistance to apoptosis | 0 | + [273] | + [274] | 0 | + [275] | + [276] | + [277] | + [278] | + [279] | + [280] |
| Replicative immortality | 0 | 0 | + [281–283] | 0 | 0 | 0 | +/- [284,285] | 0 | 0 | 0 |
| Dysregulated metabolism | + [286] | + [287,288] | + [289–291] | 0 | + [292] | + [293,294] | + [295] | 0 | + [296] | + [297] |
| Immune system evasion | + [298] | + [299,300] | + [301,302] | + [303] | + [304,305] | + [306] | + [307] | + [308,309] | + [310,311] | + [312,313] |
| Invasion and metastasis | + [287] | + [15,314] | + [315] | + [316] | + [317–319] | + [320] | + [321–323] | + [324] | + [325] | + [325,326] |
| Interactions in the tumor micro-environment | + [260] | + [327] | + [328] | + [329] | + [330] | + [327] | + [331] | + [332] | + [333,334] | + [335] |

Our 10 identified targets of anti-angiogenesis therapy are presented in the top row. 10 other cancer “hallmarks” are listed in the column to the left. Positive interactions (i.e. if the anti-angiogenesis target could also be a target for the indicated “hallmark”) are denoted “+”, controversial interactions (i.e. if the anti-angiogenesis target could both promote and inhibit the indicated “hallmark”) are denoted “+/-” and no interaction (i.e. if we have not been able to find any interaction between the anti-angiogenesis target and the indicated “hallmark”) is denoted “0”.

signaling pathway, for which commercial drugs are in development. There are also phytochemicals which exhibit such effects.

Finally, angiogenic signaling inhibitors include the modulation of mitogen-activated protein kinase (MAPK) and Akt signaling, inhibition of activator protein (AP)-1 activation, and the down-regulation of VEGF, transforming growth factor (TGF)-beta, MMP-9, as well as the upregulation of TIMP-1 which lead to reduction of tumor cell invasion and blood vessel growth [129–132].

4.1. Tea polyphenols

Tea is one of the most highly consumed beverages in the world and is rich in compounds exhibiting multiple health benefits (reviewed in [108]). The pharmacological action of tea is mainly attributed to large quantities of polyphenolic compounds known as catechins, which include epicatechin (EC), epigallocatechin (EGC), epicatechin-3-gallate (ECG), and epigallocatechin-3-gallate (EGCG) particularly in green tea and thearubigin and theaflavins in black tea. Tea polyphenols have been shown to act on angiogenesis through different signaling pathways. For instance, EGCG has been found to directly inhibit capillary endothelial cell proliferation at low concentrations, which illustrates the importance of this molecule as an important tumor angiogenesis inhibitor [133]. EGCG also inhibits VEGF production in MDA-MB231 breast cancer cells and human umbilical vein endothelial cells which correlate with the inhibition of protein kinase C, c-fos and c-jun RNA transcripts, suggesting that AP-1 responsive regions present in the human VEGF promoter may be involved [134]. In neuroblastoma, fibrosarcoma, glioblastoma, prostate cancer, and human gastric cancer cells, EGCG inhibited MMP-2 and MMP-9 while inducing the activity of their inhibitors TIMP-1 and TIMP-2 [135–138]. In human breast cancer cells, EGCG treatment reduced MMP-2 activity and the expression of focal adhesion kinase (FAK), membrane type-1-MMP (MT1-MMP), nuclear factor (NF)-kappaB, VEGF, and the adhesion of cells to the extracellular matrix (ECM) [139]. Similar signaling pathways have also been demonstrated in animal studies. EGCG also targets urokinase plasminogen activator (u-PA), leading to a down-regulation of VEGF production in tumor cells and subsequent repression of AP-1, NF-kappaB and signal transducer and activator of transcription (STAT)-1 transcription factor pathway [140,141]. EGCG furthermore inhibits aryl hydrocarbon receptor (AhR)-AhR-mediated transcription by binding to HSP90. Furthermore, a range of semi-synthetic and synthetic derivatives are more potent than EGCG in a luciferase refolding assay for HSP90 activity [106]. In transgenic adenocarcinoma of the mouse prostate (TRAMP) mice, green tea polyphenol infusion resulted in marked inhibition of effectors of angiogenesis and metastasis, notably VEGF, uPA, MMP-2, and MMP-9 [142]. In a dimethylaminoazobenzene (DAB) induced hepatoma model, administration of black tea polyphenols not only reduced the incidence of invasion, but also inhibited tumor hypoxia and angiogenesis [143]. In addition, EGCG targets tissue plasminogen activator (t-PA), which is one of the critical proteases that enable tumors to metastasize [144]. Several studies based on cell culture and animal models further demonstrated the cancer preventive function of tea, which is associated not only with the inhibition of VEGF, NF-kappaB, c-fos and cyclin D1 promoter activity, but also the decrease of Bcl-XL and the stabilization of p53 [145,146].

4.2. Curcumin

Curcumin is a polyphenol isolated from *Curcuma longa*. As a yellow dye, curcumin has been used widely for centuries not only in cooking, but also in traditional therapies for various diseases, especially inflammatory diseases. Curcumin and tetrahydro-curcumin, one of its metabolites have been extensively investigated as

anti-inflammatory and anti-cancer molecules [147,148]. Curcumin inhibits the expression of epidermal growth factor receptor (EGFR), VEGFR-1, VEGFR-2 and VEGFR-3, and the kinase activity of Src and FAK, which are responsible for the induction of angiogenic genes as well as endothelial cell polarity and migration [149]. Curcumin also reduces the MMP-2 and MMP-9 expression, along with the suppression of growth and invasion potential of tumor cells in culture and xenograft experiments [150]. Oral administration of curcumin in nude mice xenografted with hepatocarcinoma cells led to significant lowering of tumor neocapillary density. The expression of angiogenic biomarkers COX-2 and serum levels of VEGF were significantly reduced in the curcumin-treated group [151]. This inhibition of both VEGF and VEGFR in different cancers, has fueled the interest in this compound for further investigations as a potential anti-angiogenesis agent [152].

4.3. Resveratrol

Resveratrol (3,4',5-trihydroxy-trans-stilbene), a dietary polyphenol derived from grapes, berries, peanuts and other plant sources has been shown to affect several steps in the tumorigenic process including tumor angiogenesis [153–155]. Resveratrol inhibits capillary endothelial cell growth and new blood vessel growth in animals [130]. It also prevents diethylnitrosamine-initiated and phenobarbital-promoted hepatocarcinogenesis in rats through interrupting cell proliferation, inducing apoptosis [155] and impeding angiogenesis by suppressing VEGF expression through down-regulation of HIF-1alpha [156]. In in vitro studies, resveratrol was reported to inhibit cell proliferation of human ovarian cancer cells and human osteosarcoma cells by attenuating HIF-1alpha [157,158]. Through abrogation of VEGF-mediated tyrosine phosphorylation of vascular endothelial cadherins and beta-catenin, resveratrol further prevents cytokine-induced vascular leakage and tumor metastasis [130]. The underlying molecular mechanisms include: blocking VEGF- and FGF-receptor-mediated MAPK activation, inhibiting Akt- and MAPK-driven HIF-1alpha basal expression and its induction by IGF-1, stimulating the proteasomal degradation of HIF-1alpha, inhibiting phosphatidylinositol (PI)-3K/Akt and Ras/mitogen/extracellular signal-regulated kinase (MEK)/ERK pathways, and activation of forkhead box (FOX)O transcription factors [130,157–159].

Furthermore, the synthetic stilbene derivatives of resveratrol have stronger inhibitory effect on angiogenesis than resveratrol, as measured by cell proliferation and tube formation in bovine aorta endothelial cells [160]. For instance, the stilbene derivative rhapontigenin inhibits angiogenesis-induction by prostate cancer cells through HIF-1alpha degradation [161]. Likewise, resveratrol and a series of natural or synthetic stilbenes inhibit the growth of colon cancer-xenografts in mice through attenuation of VEGF and pro-MMP-9 [162]. While pre-clinical studies have been done broadly and for many different indications, resveratrol is yet to be clinically assessed for safety and prevention of cancer as well as in cancer therapeutic regimens.

4.4. Flavonoids

In the family of polyphenols, flavonoids have been found to suppress tumorigenesis via anti-angiogenesis, anti-oxidant as well as anti-proliferation effects on tumor- as well as tumor-associated stromal cells including endothelial cells [163,164]. Flavonoids, including flavones (apigenin, luteolin), flavanols (quercetin, kaempferol), flavanones (hesperetin, naringenin), anthocyanins (cyanidin, delphinidin) and isoflavones (genistein, daidzein) function as scavengers of free radicals and thus inhibit ROS formation and hypoxia-signaling cascades, modulate COX-2, and inhibit EGFR, IGF-1 and NF-kappaB signaling pathways [165].

For instance, kaempferol (3,5,7-trihydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one) has been reported to reduce ovarian cancer risk [166,167]. Kaempferol exhibits anti-inflammatory effects through inhibition of IL-4 [168] and COX-2 expression by suppressing Src kinase [169], and further down-regulates the NF-kappaB pathway [170]. Flavonoids furthermore inhibit angiogenesis through multiple mechanisms, including interaction with the COX-2 and lipoxygenase-5 enzymes, EGFR and the human epidermal growth factor receptor (HER)-2 intracellular signaling pathway [171].

Silymarin is the standardized extract of the seeds of *Silibium marianum* (milk thistle). Silibinin, the major active constituent was initially developed as a hepatoprotective product. Recently, however, it has been reported that silibinin inhibit MMP-2 expression and suppresses capillary formation of human umbilical cord venous endothelial cells (HUVECs) on matrigel [172–174]. Other naturally occurring flavonoids have been showing anti-angiogenesis and anti-oxidant effects via interference with diverse signaling pathways. For example, myricetin has been shown to inhibit inhibitor of kappaB kinase (IKK) kinase activity and prevent degradation of I-kappaBalpha and I-kappaBbeta in tumor necrosis factor (TNF)-activated endothelial cells [175]. Sulforaphane has been demonstrated to inhibit VEGFR2 at the transcriptional level [176].

Studies also showed that licochalcone A (LicA), a major constituent of *Glycyrrhiza inflata*, significantly inhibits proliferation, migration, and tube formation of HUVECs as well as microvessel growth from rat aortic rings. Furthermore, LicA significantly inhibits the growth of CT-26 colon cancer implants in BALB/c mice, with fewer CD31- and Ki-67-positive cells but more apoptotic cells [177]. Isoliquiritigenin, another flavonoid found in *G. inflata* inhibits cell migration and invasion of human prostate cancer cells (DU145 and LNCaP) mediated by decreased c-Jun N-terminal kinase (JNK)/AP-1 signaling and reduced production of proangiogenic factors [178]. Taken together, these findings provide evidence that flavonoids inhibit angiogenesis in vitro and in vivo, via antioxidant, anti-inflammatory and anti-angiogenic signaling pathways.

4.5. Terpenoids

Terpenoids are the most diverse constituents in many plant species. They form a group of natural substances which includes steroids and sterols, exhibiting anti-inflammatory and anti-carcinogenic properties [20,179–182].

The bioactive terpenoid, tripterine, also known as celastrol, a quinine methide triterpenoid is the most abundant bioactive compound derived from the root of *Trypterigium wilfordii*. Tripterine modulates the expression of proteins with angiogenic and metastatic activities (MMP-9, intercellular adhesion molecule (ICAM)-1, VEGF) and those involved in cell survival (inhibitor of apoptosis protein (IAP)-1, X-linked (X)-IAP, B-cell lymphoma (Bcl)-2, Bcl-xL, fllice inhibitory protein (cFLIP), survivin) or cell proliferation (cyclin D1, COX-2) in tumor cells [183,184]. These findings provide evidence for its potential anti-angiogenic and anti-tumor activities.

Escin is a pentacyclic triterpenoid which is isolated from the seeds of *Aesculus hippocastanum* (horse chestnuts) [185]. Escin sodium has been shown to inhibit endothelial cell migration and motility. This anti-angiogenic activity was mediated partly by inhibiting ERK and p38 MAPK pathways, which are involved in cell proliferation, motility and apoptosis [186].

Withaferin A is a major steroidal lactone constituent of the medicinal plant *Withania somnifera*, consumed as a dietary supplement around the world and used in the treatment of tumors and inflammation in several Asian countries [187]. Withaferin A exerts potent anti-angiogenic activity in vivo even at 500-times lower dose than that exerting direct anti-tumor activity [188,189].

Similar trends toward a more potent anti-angiogenic rather than anti-tumorigenic effects are also observed for other phytochemicals (such as tubulysin A as discussed below) suggesting that the vasculature may be a good target for treatments with phytochemicals.

In addition, carotenoids have anti-cancer activity in breast cancer animal models. The carotenoid group includes alpha-carotene, beta-carotene, lycopene, lutein, astaxanthin, cryptoxanthin and zeaxanthin [190,191]. The anti-oxidant action is one of the presumed mechanisms for the anti-angiogenic effects of the carotenoids.

4.6. Phytoestrogens

Phytoestrogens are plant compounds which are structurally similar to estrogen. Thereby, these compounds may have beneficial effects in prevention of steroid hormone-dependent cancers such as breast and prostate cancer. There are two major classes of phytoestrogens, isoflavones, found in soy at high concentrations, and lignans, found in flaxseed in high amounts. Lignans are ubiquitous compounds in plant material and are present in seeds, whole grains, vegetables, and berries [192–194]. Genistein (GEN), the representative of isoflavone, and enterolactone (ENL) is the main active and most potent metabolite of lignin, both have agonistic and antagonistic actions in cancer, depending on their dose and the tumor-type. In Eastern societies, the diet contains large amounts of phytoestrogens, and accordingly, the incidence of cancers such as breast cancer and prostate cancer is low [195,196]. In the diet, phytoestrogens are consumed in combination and various phytoestrogens may have different effects on tumor biology. GEN has been shown to reduce the angiogenic and metastatic potential of various cancer cell types in vitro [197]. However, the pro-estrogenic effects of GEN in breast cancer [198] suggest that caution with this particular phytoestrogen in hormone-dependent cancers is warranted. In addition, as the major circulating lignan in the human body, the majority of dietary lignans are converted into ENL. ENL inhibits cancer growth in vivo by enhancing apoptosis, inhibiting angiogenesis, and reducing inflammation without severe side-effects [199–201].

5. Prophylaxis, pharmacodynamics and safety issues

As outlined above, there is convincing evidence in support of beneficial effects of phytochemicals in cancer-related pathways, particularly with regard to anti-angiogenesis. Furthermore, many natural molecules exhibit potent anti-angiogenic activity similar to the synthetically produced drugs currently in clinical use. For example curcumin, EGCG, finasteride and barrigtozenol demonstrated comparable effects on VEGF receptor inactivation as pazopanib, the reference drug [202]. 10 μM curcumin significantly inhibited VEGF-induced HUVEC migration to a higher degree (52%) than the same concentration of a synthetic anti-angiogenic agent – the selective PDE4 inhibitor rolipram (41%) [203]. Furthermore, the EC50 dose-responses are usually lower for anti-angiogenic effects vs. cytotoxic effects on cancer cells both for synthetic and natural products. For example tubulysin A, a natural compound of myxobacterial origin, which inhibits tubulin polymerization has an EC50 for endothelial cell proliferation of 1.2 nM and up to 3.0 nM for cancer cell proliferation [204,205]. The synthetic derivatives of tubulysin A, AU816 and JB337, exhibit similar differences; EC50 of AU816 for endothelial cell proliferation is 4.4 nM, whereas inhibition of cancer cell proliferation is 10 nM and for JB337 the EC50 levels are 14 nM compared to 100 nM for endothelial and cancer cell proliferation, respectively [204,206].

Since most phytochemicals have poor water solubility and low bioavailability, the use of nano-carriers allows the preparation of these compounds in solid or liquid formulations with highly improved pharmacodynamics properties. Such encapsulation, however, render these natural compounds semi-synthetic, but the active ingredients in such preparations are still all-natural. Often, delivery mechanisms based on nano-carriers may in addition to improved pharmacokinetics also lead to improve targeting of the active compounds to the appropriate – i.e. tumor – site. For example in the case of anti-lymphatic drugs, nano-carriers, which are not able to cross the endothelium by themselves, but extravasate in tumors due to the high leakiness of tumor blood vessels, accumulate in draining intra-tumoral lymphatics and thus serve as a means to target the drugs to the leaky tumor environment. Nanoparticles made of biodegradable polymers have been utilized recently for the delivery of anti-cancer phytochemicals [207,208]. Siddiqui et al. [209] reported that nano-encapsulated EGCG retains its biological effectiveness, with over 10-fold dose advantage compared to nonencapsulated EGCG for exerting its pro-apoptotic, and anti-angiogenic effects. Encapsulation of kaempferol with nanoparticles significantly reduces cancer cell viability, as does coating it onto poly(DL-lactic acid-co-glycolic acid) (PLGA) nanoparticles [210]. Furthermore, nano-encapsulated curcumin has increased in vivo tumor anti-angiogenic activity relative to its non-encapsulated form [211]. Solid lipid nano-encapsulation of berberine also increased inhibition of cancer cell proliferation, cell cycle arrest, and apoptosis [212]. Nano-encapsulation therefore seems like a promising strategy to overcome pharmacodynamics issues associated with many phytochemicals.

The bioavailability, efficacy and stability of phytochemicals may also be increased by synthetic derivation – a procedure in which functional groups are added to or removed from the natural phytochemical, leading to improved pharmacokinetics of the resulting semi-synthetic phytochemical drug, while retaining or even improving its biological effects. Such an approach has proven particularly useful in rendering oleanolic acid attractive for therapeutic applications. Oleanolic acid is an oleanane-type triterpenoid that is widely found in dietary and medicinal plants [213]. Studies demonstrated that oleanolic acid has anti-angiogenic effects on bovine aortic endothelial cells and in the chick embryo chorioallantoic membrane assay [214]. However, 2-cyano-3,12-dioxoolean-1,9-dien-28-oic acid (CDDO), the synthetic derivative of oleanolic acid, its C-28 methyl ester (CDDO-Me) and C-28 imidazole (CDDO-Im), are all shown to be potent inhibitors of angiogenesis in the Matrigel sponge assay and in tumors established from immortalized Kaposi's sarcoma cells. These compounds prevent endothelial cell tubulogenesis on Matrigel and inhibit the VEGF-induced ERK-1 and -2 pathway in HUVECs [214,215]. It has been suggested that COX-2 overexpression may lead to increased tumor angiogenesis [216,217]. Therefore, inhibition of the COX-2 pathway might prevent carcinogenesis and angiogenesis via anti-inflammatory signaling. For example, CDDO-Me has been shown as an effective inhibitor of COX-2 and tumor growth in breast cancer models in mice.

These studies underscore the importance of phytochemicals and semi-synthetic derivatives or formulations of phytochemicals, as a novel anti-tumor treatment option in the future. Moreover, their anti-angiogenic properties could help to optimize the effectiveness of existing chemotherapeutic drugs. The use of phytochemicals as adjuvants in combination with anti-angiogenic drugs is promising. For example, tumor hypoxia can be a deleterious result of anti-angiogenic cancer treatment leading to changes in cellular metabolism [73,74,218], elevated intracellular ROS-levels [72], de-differentiation of tumor cells and increased metastatic propensity [218,219]. These hypoxia-induced pathways can be treated by various phytochemical antioxidants including EGCG,

melatonin, resveratrol and silibinin [143,157,220–223]. While this could provide a strong argument for synergy by combining such agents, it was traditionally believed that hypoxia should be the very goal of anti-angiogenic treatment and an important inducer of tumor cell apoptosis [224,225]. Therefore addition of such antioxidants may reduce the hypoxia-induced cell death, as it does for non-malignant cells [226,227], potentially leading to a poorer outcome from the anti-angiogenic treatment. Also in the interaction between tumor inflammation and angiogenesis the combination of anti-inflammatory phytochemicals with anti-angiogenic drugs may provide a potential for synergy as inflammatory cells are considered a source of alternative angiogenic factors and thus important for evasive resistance [4]. However, some inflammatory cells are thought to be anti-tumorigenic and important for tumor clearance [228,229], and inhibiting them could lead to antagonistic interactions in such combinations. Thus each combination of phytochemicals and other treatment options should be thoroughly tested in pre-clinical and clinical trials before adopted by medical practitioners.

Tumor angiogenesis is not only required for late-stage progressive disease, but is crucial even for initial growth of the tumor to clinically detectable masses. Therefore, the inhibition of angiogenesis has been proposed as an important preventive strategy [230]. This idea, also known as angioprevention, is highly efficacious in preventing tumor growth in animal models [156,231]. Angioprevention should be achieved using non-toxic agents such as phytochemicals, as they are often consumed on a daily basis. Ideally, we should however also aim to achieve anti-angiogenic treatment efficacy using phytochemical agents at sufficiently low doses to keep toxic side-effects are kept to a minimum. Thus safety concerns should weigh in heavily when deciding on the most appropriate drugs for anti-angiogenic treatment in cancer. Here, we intend to identify 10 suitable, prototypical anti-angiogenic therapeutic approaches based on phytochemicals that have proven to be effective against our previously defined targets. We aimed at finding compounds that are free from intellectual property constraints, readily available and cheap such that everyone, all over the world, could use them as an alternative or complementary treatment option. A modern approach to chemotherapy may be to use lower doses of combinations of chemicals, which work in complementary ways. Thus, while low-doses of a single compound may not achieve the full anti-angiogenic function, a mixture of compounds, or extracts from medical plants, may be able to specifically synergize in preventing angiogenesis while avoiding significant side-effects. Such an approach, combining phytochemicals in tea and soy extracts has been successfully applied to the anti-angiogenic treatment of prostate and breast cancer in animal models [47,49]. Therefore we aimed at selecting phytochemicals from as many different classes of natural compounds listed in Section 4 as possible.

In line with this rationale, we have identified 10 natural compounds as potential therapeutic approaches (top row of Table 2) against the previously identified targets (top row of Table 1). These compounds are safer than the synthetic anti-angiogenesis inhibitors in clinical use today [232,233]—and may even be considered as prophylactic (angioprevention) agents. These molecules were selected because they exhibit pleiotropic functions, i.e. disrupting both inflammation and vascularization during tumor onset and progression and while some of them share molecular targets, combined they will inhibit a broad spectrum of pro-angiogenic signaling pathways lowering the risk of adaptive resistance. As tumors often progressively turn on pro-angiogenic pathways during their development to more advanced stages (reviewed in [234]), which is one possible reason why many patient do not respond to single-agent therapies such as anti-VEGF drugs, the cocktail of 10 phytochemical-based therapeutic approaches mentioned here

Table 2
Effects of phytochemical approaches in anti-angiogenic therapy as effective also against other cancer “hallmarks”.

| Other cancer hallmarks | Phytochemical approach | | | | | | | | | |
|---|------------------------|-------------|-------------|-----------------|----------------------------------|-----------------|-------------|---------------|--------------|---------------|
| | Oleanolic acid | Tripterine | Silibinin | Curcumin | Epigallo-catechin-gallate (EGCG) | Kaempferol | Melatonin | Enterolactone | Withaferin A | Resveratrol |
| Genetic instability | + [336] | 0 | + [337] | + [338] | +/- [339,340] | 0 | + [341] | 0 | 0 | + [342] |
| Sustained proliferative signaling | 0 | 0 | +/- [343] | 0 | + [245,246] | 0 | + [344,345] | + [346] | + [249,250] | + [347,348] |
| Tumor-promoting inflammation | + [20] | 0 | + [349] | + [350,351] | + [352,353] | + [169] | 0 | + [354] | 0 | + [355–357] |
| Evasion of anti-growth signaling | + [341,358,359] | + [360–362] | + [363–366] | + [367–370] | + [371–374] | + [375–378] | + [379] | + [380] | + [381–383] | + [384–387] |
| Resistance to apoptosis | + [388,389] | + | + [390] | + [391] | + [374,392,393] | + [394] | + [395] | + [396] | + [397] | + [398] |
| Replicative immortality | + [399,400] | 0 | + [401–403] | + [401,404] | + [405–407] | 0 | + [408] | 0 | 0 | + [409] |
| Dysregulated metabolism | + [358,410] | 0 | + [411,412] | + [391,392,413] | + [414–416] | + [375,417,418] | + [419–421] | 0 | + [422–424] | + [425–427] |
| Immune system evasion | + [428,429] | 0 | + [430] | + [431] | + [432,433] | 0 | + [434] | 0 | + [435] | +/- [436–438] |
| Invasion and metastasis | + [439,440] | + | + [442] | + [443] | + [444–446] | + [447,448] | + [449,450] | + [451] | + [452] | +/- [453–455] |
| Interactions in the tumor micro-environment | + [456] | + [457] | + [458] | + [459] | + [433,460] | + [461,462] | + [331,463] | + [199] | + [464] | + [465] |

Our 20 identified therapeutic approaches for anti-angiogenesis therapy are presented in the top row. 10 other cancer “hallmarks” are listed in the column to the left. Positive interactions (i.e. if the compound could also exhibit therapeutic potential against the indicated “hallmark”) are denoted “+”, controversial interactions (i.e. if the compound could both promote and inhibit the indicated “hallmark”) are denoted “+/-” and no interaction (i.e. if we have not been able to find any therapeutic activity of the compound against the indicated “hallmark”) is denoted “0”.

would be a more broad-acting treatment option, not least for such patients that do not respond to commercially available anti-VEGF drugs.

6. Validation of targets and approaches

Anti-angiogenic agents have demonstrated activity in a broad spectrum of malignancies. The questions of finding the best drug, or combination of drugs, for each patient and how to best evaluate the response to treatment are still not adequately answered. An alternative to current efforts on targeted therapy development could be to inhibit an array of pathological targets of tumor blood vessels such as the 10 targets identified here (Table 1, top row). We envision that such a broad approach, simultaneously hitting multiple or all of these targets of the tumor vasculature at the same time, would produce an improved therapeutic outcome compared to single-target therapy. The current strategy we endorse for further investigation considers a combination of phytochemicals, identified to act on more than one or a few of these targets (Table 2, top row). We believe that this approach circumvents the numerous avenues of emergent resistance, and has the potential to be an active therapeutic in a broader range of tumors. We encourage researchers and clinicians alike to test formulations based on this natural compound cocktail-principle and in this way further establish the paradigm of phytochemicals as anti-angiogenic treatment options in cancer.

Given the extensive cross-talk between mechanisms regulating angiogenesis and other aspects of cell biology as well as the broad effects of phytochemicals, we believe that it is important to be able to anticipate synergies that might be achieved through acting not only on aspects of tumor angiogenesis but also on other essential aspects of tumor biology such as the “hallmarks” of cancer. Accordingly, the identified targets for anti-angiogenic therapy and the therapeutic phytochemical-based approaches were cross-validated against their effects on other “hallmark” processes, by undertaking a thorough, qualitative background literature research. An independent team of scientists consisting of specialists in various “hallmark” areas specifically sought to determine the relevance of our identified targets and the nominated phytochemical approaches across these other important areas of cancer biology. The result of this elaborate cross-validation effort is shown in Tables 1 and 2, for the targets and suggested phytochemical-based approaches respectively. We found that many of our identified targets and approaches were also relevant for other aspects of cancer biology. In such cases they were noted as having “complementary” effects (denoted with a + in Tables 1 and 2). In instances where reports on relevant actions in other aspects of cancer biology were mixed (i.e. reports showing both pro- and anti-potential), the effects are considered “controversial” and denoted +/- in the tables. Finally, in instances where no literature was found to support the relevance of a target or approach in other aspects of cancer biology, we consider that there is “no known relationship” and denote this with a 0 in the tables. We did not find any instances in which our targets or approaches acted to promote other pathological aspects of tumor biology. As an example, reducing tumor hypoxia would also lead to reduced tumor-promoting inflammation and therefore we mark this with a + in the cell intersecting the “(reduce) hypoxia” column with the “tumor promoting inflammation” row in Table 1. Conversely, our suggested approach resveratrol was found to both inhibit and induce metastasis and we therefore denote this as +/- in the cell intersecting the “resveratrol” column with the “Invasion and metastasis” row in Table 2.

As it can be determined from closely inspecting these tables, most of our targets and approaches are found to also inhibit other essential aspects or hallmarks of tumor pathology. As such, we

believe that the use of the suggested phytochemical approaches – or similarly acting derivatives or analogues – would be effective as alternative anti-angiogenic agents while also inhibiting multiple other aspects of tumor pathology at the same time. Especially oleanolic acid, silibinin, curcumin, EGCG, melatonin and resveratrol, have strong synergistic actions, i.e. anti-tumorigenic effects in almost all other hallmark areas in addition to being effective anti-tumor-angiogenic agents. It would be highly important to experimentally test the effects of such a cocktail of phytochemicals in pre-clinical and clinical studies in the future.

7. Future directions

The VEGF pathway holds the key in regulating angiogenesis and targeted therapies against this pathway have been established in the clinic. However, after initial positive results several studies have failed to show increased survival rates for many groups of patients treated with anti-VEGF drugs. It is clear that multiple factors contribute to enabling the angiogenic switch in tumors. However, till date an overwhelming focus has been placed on blocking the VEGF-VEGFR2 pathway. Using small synthetic molecules with a broader target profile, it has been possible to achieve better anti-angiogenic effects in tumors compared to using specific antibody based therapeutics. However, the former are more toxic, still not sufficiently effective and therefore other strategies need to be considered.

It is becoming increasingly clear that plant-derived medicinal compounds may in many cases be at least as effective in blocking angiogenesis as the currently used synthetic drugs, while exhibiting only a fraction of the negative side-effects. Some of these compounds may, however, be more effective in treatment of some tumors, such as the potential for using phytoestrogens in treatment of ER-positive breast cancer or that of melatonin in treating cancer in people with disrupted circadian rhythms. Identification of suitable biomarkers for selection of patients to different types of therapeutic regimens as well as response to treatment has however so far not been achieved but remain a very important avenue for future research.

The desired optimal effect of anti-angiogenic treatment is another issue, which requires further study. Is complete vascular regression what we aim for? Such an outcome would be expected to result in extensive tumor hypoxia, radically reduced perfusion and thus reduced delivery of not only oxygen and nutrients but also drugs to the tumor tissue. Some studies argue that such a result of anti-angiogenic treatment may in fact lead to a more invasive tumor phenotype and worse prognosis compared to the non-anti-angiogenesis treated group. Perhaps it would be a better option to merely reduce the vascular density, improve the structure and function of tumor blood vessels such that they lose their pathogenic characteristics and more closely resemble blood vessels in healthy tissues. Such a phenotype would be expected, however, to lead to improved perfusion, oxygenation and delivery of nutrients to the tumor, thereby potentially increasing tumor growth rate. On the other hand, this would improve the effects of co-administered chemotherapy, which would also be delivered more effectively to the tumor, and which would more effectively kill the tumor cells if these were well nourished and actively proliferating. Perhaps it should even be considered to withdraw anti-angiogenic treatment if it is found to result in dramatically reduced perfusion and oxygenation of the tumor, at least if such treatment is used in combination with chemo- or radiotherapy. The phytochemical approaches to inhibit tumor angiogenesis identified in [Table 2](#) can, however, in addition to pruning non-perfused and pathological blood vessels in the tumor, also reduce the detrimental effects of chemotherapy, leading to improved tolerance and therefore improved health and potentially extended survival when used

in combination with classical cytostatic therapies. It should therefore be considered whether such compounds could be beneficial for tumor patients receiving chemotherapy, even if they are also receiving anti-VEGF drugs as the ones in clinical use today do not reduce the side-effects associated with chemotherapy.

While many natural compounds have general beneficial effects, it is clear that some compounds are more effective in cancer treatment than others. For example, a recent meta-analysis of randomized placebo-controlled trials testing the effect of dietary supplementation of multi-vitamins and multi-minerals, failed to show any effect on cancer mortality suggesting that this is not a route to prioritize in future efforts [235]. Vitamin E and high doses of beta-carotene have been shown to increase prostate and lung cancer risk suggesting that caution is warranted with these powerful biological components [236,237]. Recently it has become evident that the function of healthy tissues also relies on the activity of angiogenic factors. For example, currently used anti-VEGF drugs cause a marked reduction in vascular density, with a subsequent decline in the function of healthy organs such as the thymus and other endocrine glands, intestine and other parts of the digestive tract, gonads and the kidney [238]. As such, the effects that we are seeking from the treatment – potent and broad-acting anti-angiogenic activity – may end up leading to side-effects. Possible ways to overcome this issue is by targeted delivery such as via nanoparticle carriers as discussed above, or by titrating the amount of phytochemicals given such that it is effective in reducing/preventing pathological angiogenesis but not high enough to impair healthy vascular functions. Such an approach could logically be considered together with the combination of for example the 10 phytochemicals identified here in one cocktail where each compound is selected to complement the other hopefully leading to synergistic anti-angiogenic effects while maintaining negligible toxicity. This concept, as well as the identification of which tumor types are more susceptible to such treatment and identification of readouts for efficacy and toxicity, are important areas for future research.

There are, however, also potential drawbacks associated with the use of phytochemicals, which needs to be resolved before effective formulations of such compounds can be achieved. Due to poor absorption of many phytochemicals in their natural form, it may be necessary to chemically modify some of the compounds suggested here, or use them in semi-synthetic formulations such as nanoparticle encapsulations, which would lower their “natural” status but could promote their therapeutic and pharmacokinetic profiles. How best to take advantage of such derivations or modifications is still an open question.

8. Final conclusion

The complex and multi-factorial nature of tumor angiogenesis, especially in advanced tumors, necessitate the use of therapeutic compounds which act broadly in order to reduce problems with developing resistance. Cancer is a chronic disorder, which in cases where no curative treatment exists, has to be treated medically for a long period of time. Therefore treatment options with very low if any toxicity are preferred. Cancer is furthermore a growing medical issue also in developing countries and in societies in which the high costs associated with cutting edge treatment at modern medical institutions is prohibitive to the majority of the patients. Therefore low-cost treatment options have great potential. Phytochemicals constitute a class of broadly acting and very cheap natural drugs with very low toxicity especially if given at relatively low doses in combination. The list of 10 prototypical phytochemical approaches to inhibit 10 important pro-angiogenic targets in cancer presented here, constitute a framework for further studies on how to mix a

potentially effective, non-toxic cocktail of natural compounds as a complement or alternative to other types of anti-cancer drugs available today. We hope that this review could serve as a primer for such investigations, which hopefully could lead to cheaper, safer and more effective anti-cancer treatment in the future.

Conflict of interest statement

The authors declare no conflict of interest.

Author contributions

Z.W., C.D., X.Y., M.F., A.A., W.K.R., D.G., G.R.N., B.E.R., D.R., Y.C.C. and L.D.J. wrote the manuscript, K.H., H.F., A.G.G., S.H., A.A., E.N., A.A., S.S.A., B.H., X.Y., G.G., D.B., M.R.C., K.A., S.C., D.H., S.I.M., A.A., A.B. and N.K. provided validation of the prioritized targets and approaches listed in Table 1, the result of which is shown in Tables 1 and 2.

Acknowledgments

We would like to thank Mr. Leroy Lowe for his vision and initiative in the GettingToKnowCancer organization. Dr. Jensen was funded by the Swedish Society for Medical Research, the Goesta Fraenkel Foundation, Åke Wiberg's Foundation, Ollie och Elof Ericsson's Foundation, the Karolinska Institute and Linköping University. Drs Keith and Bilsland were funded by the University of Glasgow, Beatson Oncology Center Fund and Cancer Research UK grant C301/A14762. Dr. Arreola was funded by NIH NRSA grant F31CA154080. Dr. Georgakilas was funded by EU grant MC-CIG-303514, the National Strategic Reference Framework grant MIS-379346 and COST action CM1201. Dr. Amedei was funded by the University of Florence and the Italian Ministry of Research. Dr. Amin was partially funded by the Terry Fox Foundation, 2013 and the UAEU Program for Advanced Research 2013. Dr. Yi Charlie Chen was funded by the West Virginia Experimental Program to Stimulate Competitive Research and NIH grants 5P20RR016477 and 8P20GM104434. Dr. Dabrosin was funded by the Swedish Research Council and the Swedish Research Society. Dr. Generali was funded by the EU Seventh Framework Program grant 278570. Dr. Halicka was funded by the NIH. Dr. Niccolai was funded by the Italian Ministry of Research and the University of Italy. Dr. Honoki was funded by the Japanese Ministry of Science grant 24590493. Dr. Rathmell was funded by Galaxo Smith Kline for an investigator-initiated clinical trial. Dr. Ciriolo was funded by the AIRC. Dr. Fuster was funded by the NIH/NHLBI grant R01HL107652 and an Eisai Healthcare Investigator grant. Dr. Sophie Chen was funded by the Ovarian and Prostate Cancer Research Trust. Dr. Yin was funded by NIH/NHLBI grant T32HL098062.

References

- Ribatti D, Djonov V. Intussusceptive microvascular growth in tumors. *Cancer Lett* 2012;316:126–31.
- Cao Y, Arbiser J, D'Amato RJ, D'Amore PA, Ingber DE, Kerbel R, et al. Forty-year journey of angiogenesis translational research. *Sci Transl Med* 2011;3:114rv113.
- Folkman J. Tumor angiogenesis: therapeutic implications. *N Engl J Med* 1971;285:1182–6.
- Bergers G, Hanahan D. Modes of resistance to anti-angiogenic therapy. *Nat Rev Cancer* 2008;8:592–603.
- Ribatti D. Endogenous inhibitors of angiogenesis: a historical review. *Leuk Res* 2009;33:638–44.
- Landskroner-Eiger S, Moneke I, Sessa WC. miRNAs as modulators of angiogenesis. *Cold Spring Harb Perspect Med* 2013;3:a006643.
- Hellstrom M, Phng LK, Hofmann JJ, Wallgard E, Coultas L, Lindblom P, et al. Dll4 signalling through Notch1 regulates formation of tip cells during angiogenesis. *Nature* 2007;445:776–80.
- Siekmann AF, Lawson ND. Notch signalling limits angiogenic cell behaviour in developing zebrafish arteries. *Nature* 2007;445:781–4.
- Jakobsson L, Franco CA, Bentley K, Collins RT, Ponsioen B, Aspalter IM, et al. Endothelial cells dynamically compete for the tip cell position during angiogenic sprouting. *Nat Cell Biol* 2010;12:943–53.
- Krueger J, Liu D, Scholz K, Zimmer A, Shi Y, Klein C, et al. Flt1 acts as a negative regulator of tip cell formation and branching morphogenesis in the zebrafish embryo. *Development* 2011;138:2111–20.
- Ridgway J, Zhang G, Wu Y, Stawicki S, Liang WC, Chantery Y, et al. Inhibition of Dll4 signalling inhibits tumour growth by deregulating angiogenesis. *Nature* 2006;444:1083–7.
- Noguera-Troise I, Daly C, Papadopoulos NJ, Coetsee S, Boland P, Gale NW, et al. Blockade of Dll4 inhibits tumour growth by promoting non-productive angiogenesis. *Nature* 2006;444:1032–7.
- Nagy JA, Dvorak AM, Dvorak HF. Vascular hyperpermeability, angiogenesis, and stroma generation. *Cold Spring Harb Perspect Med* 2012;2:a006544.
- Xue Y, Lim S, Yang Y, Wang Z, Jensen LD, Hedlund EM, et al. PDGF-BB modulates hematopoiesis and tumor angiogenesis by inducing erythropoietin production in stromal cells. *Nat Med* 2012;18:100–10.
- Hosaka K, Yang Y, Seki T, Nakamura M, Andersson P, Rouhi P, et al. Tumour PDGF-BB expression levels determine dual effects of anti-PDGF drugs on vascular remodelling and metastasis. *Nat Commun* 2013;4:2129.
- Hedlund EM, Yang X, Zhang Y, Yang Y, Shibuya M, Zhong W, et al. Tumor cell-derived placental growth factor sensitizes antiangiogenic and antitumor effects of anti-VEGF drugs. *Proc Natl Acad Sci U S A* 2013;110:654–9.
- van der Schaft DW, Seftor RE, Seftor EA, Hess AR, Gruman LM, Kirschmann DA, et al. Effects of angiogenesis inhibitors on vascular network formation by human endothelial and melanoma cells. *J Natl Cancer Inst* 2004;96:1473–7.
- Nagy JA, Dvorak HF. Heterogeneity of the tumor vasculature: the need for new tumor blood vessel type-specific targets. *Clin Exp Metastasis* 2012;29:657–62.
- Jain RK. Normalization of tumor vasculature: an emerging concept in antiangiogenic therapy. *Science* 2005;307:58–62.
- Shanmugam MK, Dai X, Kumar AP, Tan BK, Sethi G, Bishayee A. Oleonic acid and its synthetic derivatives for the prevention and therapy of cancer: preclinical and clinical evidence. *Cancer Lett* 2014;346:206–16.
- Bruick RK, McKnight SL. A conserved family of prolyl-4-hydroxylases that modify HIF. *Sci Signal* 2001;294:1337.
- Maxwell P, Wiesener M, Chang G, Clifford S, Vaux E, Cockman M, et al. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 1999;399:271–5.
- Raval RR, Lau KW, Tran MGB, Sowter HM, Mandriota SJ, Li JL, et al. Contrasting properties of hypoxia-inducible factor 1 (HIF-1) and HIF-2 in von Hippel–Lindau-associated renal cell carcinoma. *Mol Cell Biol* 2005;25:5675.
- Hu CJ, Wang LY, Chodosh LA, Keith B, Simon MC. Differential roles of hypoxia-inducible factor 1 α (HIF-1 α) and HIF-2 α in hypoxic gene regulation. *Mol Cell Biol* 2003;23:9361.
- Iliopoulos O, Levy AP, Jiang C, Kaelin WG, Goldberg MA. Negative regulation of hypoxia-inducible genes by the von Hippel–Lindau protein. *Proc Natl Acad Sci U S A* 1996;93:10595–9.
- Tosetti F, Ferrari N, De Flora S, Albini A. Angioprevention: angiogenesis is a common and key target for cancer chemopreventive agents. *FASEB J* 2002;16:2–14.
- Ivan M, Harris AL, Martelli F, Kulshreshtha R. Hypoxia response and microRNAs: no longer two separate worlds. *J Cell Mol Med* 2008;12:1426–31.
- Kaelin WG. The von Hippel–Lindau tumor suppressor gene and kidney cancer. *Clin Cancer Res* 2004;10:6290S–5S.
- Leong SP, Cady B, Jablons DM, Garcia-Aguilar J, Reintgen D, Jakob J, et al. Clinical patterns of metastasis. *Cancer Metastasis Rev* 2006;25:221–32.
- Sleeman JP. The lymph node as a bridgehead in the metastatic dissemination of tumors. *Recent Results Cancer Res* 2000;157:55–81.
- Bono P, Wasenius VM, Heikkilä P, Lundin J, Jackson DG, Joensuu H. High LYVE-1-positive lymphatic vessel numbers are associated with poor outcome in breast cancer. *Clin Cancer Res* 2004;10:7144–9.
- Nakamura Y, Yasuoka H, Tsujimoto M, Imabun S, Nakahara M, Nakao K, et al. Lymph vessel density correlates with nodal status, VEGF-C expression, and prognosis in breast cancer. *Breast Cancer Res Treat* 2005;91:125–32.
- Gombos Z, Xu X, Chu CS, Zhang PJ, Acs G. Peritumoral lymphatic vessel density and vascular endothelial growth factor C expression in early-stage squamous cell carcinoma of the uterine cervix. *Clin Cancer Res* 2005;11:8364–71.
- Chen W, Shen W, Chen M, Cai G, Liu X. Study on the relationship between lymphatic vessel density and distal intratumoral spread of rectal cancer. *Eur Surg Res* 2007;39:332–9.
- Renyi-Vamos F, Tovari J, Fillinger J, Timar J, Paku S, Kenessey I, et al. Lymphangiogenesis correlates with lymph node metastasis, prognosis, and angiogenic phenotype in human non-small cell lung cancer. *Clin Cancer Res* 2005;11:7344–53.
- Padera TP, Kadambi A, di Tomaso E, Carreira CM, Brown EB, Boucher Y, et al. Lymphatic metastasis in the absence of functional intratumor lymphatics. *Science* 2002;296:1883–6.
- Wang XL, Fang JP, Tang RY, Chen XM. Different significance between intratumoral and peritumoral lymphatic vessel density in gastric cancer: a retrospective study of 123 cases. *BMC Cancer* 2010;10:299.
- Ravindranath MH, Muthugounder S, Presser N, Viswanathan S. Anticancer therapeutic potential of soy isoflavone, genistein. *Adv Exp Med Biol* 2004;546:121–65.
- Mattila MM, Ruohola JK, Karpanen T, Jackson DG, Alitalo K, Harkonen PL. VEGF-C induced lymphangiogenesis is associated with lymph

- node metastasis in orthotopic MCF-7 tumors. *Int J Cancer* 2002;98:946–51.
- [40] Karpanen T, Egeblad M, Karkkainen MJ, Kubo H, Yla-Herttuala S, Jaattela M, et al. Vascular endothelial growth factor C promotes tumor lymphangiogenesis and intralymphatic tumor growth. *Cancer Res* 2001;61:1786–90.
- [41] Ochi N, Matsuo Y, Sawai H, Yasuda A, Takahashi H, Sato M, et al. Vascular endothelial growth factor-C secreted by pancreatic cancer cell line promotes lymphatic endothelial cell migration in an in vitro model of tumor lymphangiogenesis. *Pancreas* 2007;34:444–51.
- [42] Skobe M, Hawighorst T, Jackson DG, Prevo R, Janes L, Velasco P, et al. Induction of tumor lymphangiogenesis by VEGF-C promotes breast cancer metastasis. *Nat Med* 2001;7:192–8.
- [43] Mumprecht V, Detmar M. Lymphangiogenesis and cancer metastasis. *J Cell Mol Med* 2009;13:1405–16.
- [44] Ran S, Volk L, Hall K, Flister MJ. Lymphangiogenesis and lymphatic metastasis in breast cancer. *Pathophysiology* 2010;17:229–51.
- [45] Chang LK, Garcia-Cardena G, Farnese F, Fannon M, Chen EJ, Butterfield C, et al. Dose-dependent response of FGF-2 for lymphangiogenesis. *Proc Natl Acad Sci U S A* 2004;101:11658–63.
- [46] Kubo H, Cao R, Brakenhielm E, Makinen T, Cao Y, Alitalo K. Blockade of vascular endothelial growth factor receptor-3 signaling inhibits fibroblast growth factor-2-induced lymphangiogenesis in mouse cornea. *Proc Natl Acad Sci U S A* 2002;99:8868–73.
- [47] Zhou JR, Yu L, Zhong Y, Blackburn GL. Soy phytochemicals and tea bioactive components synergistically inhibit androgen-sensitive human prostate tumors in mice. *J Nutr* 2003;133:516–21.
- [48] Kajiya K, Hirakawa S, Ma B, Drinnenberg I, Detmar M. Hepatocyte growth factor promotes lymphatic vessel formation and function. *EMBO J* 2005;24:2885–95.
- [49] Zhou JR, Yu L, Mai Z, Blackburn GL. Combined inhibition of estrogen-dependent human breast carcinoma by soy and tea bioactive components in mice. *Int J Cancer* 2004;108:8–14.
- [50] Li H, Adachi Y, Yamamoto H, Min Y, Ohashi H, Li M, et al. Insulin-like growth factor-I receptor blockade reduces tumor angiogenesis and enhances the effects of bevacizumab for a human gastric cancer cell line, MKN45. *Cancer* 2011;117:3135–47.
- [51] Makinen T, Adams RH, Bailey J, Lu Q, Ziemiecki A, Alitalo K, et al. PDZ interaction site in ephrinB2 is required for the remodeling of lymphatic vasculature. *Genes Dev* 2005;19:397–410.
- [52] Helfrich I, Edler L, Sucker A, Thomas M, Christian S, Schadendorf D, et al. Angiopoietin-2 levels are associated with disease progression in metastatic malignant melanoma. *Clin Cancer Res* 2009;15:1384–92.
- [53] Cao R, Bjorndahl MA, Religa P, Clasper S, Garvin S, Galter D, et al. PDGF-BB induces intratumoral lymphangiogenesis and promotes lymphatic metastasis. *Cancer Cell* 2004;6:333–45.
- [54] Banziger-Tobler NE, Halin C, Kajiya K, Detmar M. Growth hormone promotes lymphangiogenesis. *Am J Pathol* 2008;173:586–97.
- [55] Hirakawa S, Kodama S, Kunstfeld R, Kajiya K, Brown LF, Detmar M. VEGF-A induces tumor and sentinel lymph node lymphangiogenesis and promotes lymphatic metastasis. *J Exp Med* 2005;201:1089–99.
- [56] Hirakawa S, Brown LF, Kodama S, Paavonen K, Alitalo K, Detmar M. VEGF-C-induced lymphangiogenesis in sentinel lymph nodes promotes tumor metastasis to distant sites. *Blood* 2007;109:1010–7.
- [57] Ishii H, Chikamatsu K, Sakakura K, Miyata M, Furuya N, Masuyama K. Primary tumor induces sentinel lymph node lymphangiogenesis in oral squamous cell carcinoma. *Oral Oncol* 2010;46:373–8.
- [58] Christiansen A, Detmar M. Lymphangiogenesis and cancer. *Genes Cancer* 2011;2:1146–58.
- [59] Alitalo K. The lymphatic vasculature in disease. *Nat Med* 2011;17:1371–80.
- [60] Fukumura D, Jain RK. Tumor microenvironment abnormalities: causes, consequences, and strategies to normalize. *J Cell Biochem* 2007;101:937–49.
- [61] Duong T, Koopman P, Francois M. Tumor lymphangiogenesis as a potential therapeutic target. *J Oncol* 2012;204946.
- [62] Roenneberg T. Chronobiology: the human sleep project. *Nature* 2013;498:427–8.
- [63] Jensen LD, Cao YH. Clock controls angiogenesis. *Cell Cycle* 2013;12:405–8.
- [64] Jensen LD, Cao Z, Nakamura M, Yang Y, Brautigam L, Andersson P, et al. Opposing effects of circadian clock genes *bmal1* and *period2* in regulation of VEGF-dependent angiogenesis in developing zebrafish. *Cell Rep* 2012;2:231–41.
- [65] Schernhammer ES, Laden F, Speizer FE, Willett WC, Hunter DJ, Kawachi I, et al. Night-shift work and risk of colorectal cancer in the nurses' health study. *J Natl Cancer Inst* 2003;95:825–8.
- [66] Schernhammer ES, Laden F, Speizer FE, Willett WC, Hunter DJ, Kawachi I, et al. Rotating night shifts and risk of breast cancer in women participating in the nurses' health study. *J Natl Cancer Inst* 2001;93:1563–8.
- [67] Rao S, Chun C, Fan J, Kofron JM, Yang MB, Hegde RS, et al. A direct and melanopsin-dependent fetal light response regulates mouse eye development. *Nature* 2013;494:243–6.
- [68] Busik JV, Tikhonenko M, Bhatwadekar A, Opreanu M, Yakubova N, Caballero S, et al. Diabetic retinopathy is associated with bone marrow neuropathy and a depressed peripheral clock. *J Exp Med* 2009;206:2897–906.
- [69] Anea CB, Cheng B, Sharma S, Kumar S, Caldwell RW, Yao L, et al. Increased superoxide and endothelial NO synthase uncoupling in blood vessels of *Bmal1*-knockout mice. *Circ Res* 2012;111:1157–65.
- [70] Wang CY, Wen MS, Wang HW, Hsieh IC, Li Y, Liu PY, et al. Increased vascular senescence and impaired endothelial progenitor cell function mediated by mutation of circadian gene *Per2*. *Circulation* 2008;118:2166–73.
- [71] Koyanagi S, Kuramoto Y, Nakagawa H, Aramaki H, Ohdo S, Soeda S, et al. A molecular mechanism regulating circadian expression of vascular endothelial growth factor in tumor cells. *Cancer Res* 2003;63:7277–83.
- [72] Guzy RD, Schumacker PT. Oxygen sensing by mitochondria at complex III: the paradox of increased reactive oxygen species during hypoxia. *Exp Physiol* 2006;91:807–19.
- [73] Denko NC. Hypoxia, HIF1 and glucose metabolism in the solid tumour. *Nat Rev Cancer* 2008;8:705–13.
- [74] Gillies RJ, Verduzco D, Gatenby RA. Evolutionary dynamics of carcinogenesis and why targeted therapy does not work. *Nat Rev Cancer* 2012;12:487–93.
- [75] Ferrara N. Role of myeloid cells in vascular endothelial growth factor-independent tumor angiogenesis. *Curr Opin Hematol* 2010;17:219–24.
- [76] Apte RN, Krelin Y, Song X, Dotan S, Recih E, Elkabets M, et al. Effects of micro-environment- and malignant cell-derived interleukin-1 in carcinogenesis, tumour invasiveness and tumour–host interactions. *Eur J Cancer* 2006;42:751–9.
- [77] Nicolini A, Carpi A, Rossi G. Cytokines in breast cancer. *Cytokine Growth Factor Rev* 2006;17:325–37.
- [78] Crawford Y, Kasman I, Yu L, Zhong C, Wu X, Modrusan Z, et al. PDGF-C mediates the angiogenic and tumorigenic properties of fibroblasts associated with tumors refractory to anti-VEGF treatment. *Cancer Cell* 2009;15:21–34.
- [79] Ricci-Vitiani L, Pallini R, Biffoni M, Todaro M, Invernici G, Cenci T, et al. Tumour vascularization via endothelial differentiation of glioblastoma stem-like cells. *Nature* 2010;468:824–8.
- [80] Maione F, Capano S, Regano D, Zentilin L, Giacca M, Casanovas O, et al. Semaphorin 3A overcomes cancer hypoxia and metastatic dissemination induced by antiangiogenic treatment in mice. *J Clin Invest* 2012;122:1832–48.
- [81] Paez-Ribes M, Allen E, Hudock J, Takeda T, Okuyama H, Vinals F, et al. Antiangiogenic therapy elicits malignant progression of tumors to increased local invasion and distant metastasis. *Cancer Cell* 2009;15:220–31.
- [82] Ebos JM, Lee CR, Cruz-Munoz W, Bjarnason GA, Christensen JG, Kerbel RS. Accelerated metastasis after short-term treatment with a potent inhibitor of tumor angiogenesis. *Cancer Cell* 2009;15:232–9.
- [83] Lu KV, Chang JP, Parachoniak CA, Pandika MM, Aghi MK, Meyronet D, et al. VEGF inhibits tumor cell invasion and mesenchymal transition through a MET/VEGFR2 complex. *Cancer Cell* 2012;22:21–35.
- [84] Jensen LD, Cao R, Cao Y. In vivo angiogenesis and lymphangiogenesis models. *Curr Mol Med* 2009;9:982–91.
- [85] Jensen LD, Rouhi P, Cao Z, Lanne T, Wahlberg E, Cao Y. Zebrafish models to study hypoxia-induced pathological angiogenesis in malignant and nonmalignant diseases. *Birth Defects Res C Embryo Today* 2011;93:182–93.
- [86] Willett CG, Boucher Y, di Tomaso E, Duda DG, Munn LL, Tong RT, et al. Direct evidence that the VEGF-specific antibody bevacizumab has antivascular effects in human rectal cancer. *Nat Med* 2004;10:145–7.
- [87] Lockhart AC, Rothenberg ML, Dupont J, Cooper W, Chevalier P, Sternas L, et al. Phase I study of intravenous vascular endothelial growth factor trap, aflibercept, in patients with advanced solid tumors. *J Clin Oncol* 2010;28:207–14.
- [88] Zhong H, Chiles K, Feldser D, Laughner E, Hanrahan C, Georgescu MM, et al. Modulation of hypoxia-inducible factor 1 α expression by the epidermal growth factor/phosphatidylinositol 3-kinase/PTE/AKT/FRAP pathway in human prostate cancer cells: implications for tumor angiogenesis and therapeutics. *Cancer Res* 2000;60:1541–5.
- [89] Mabbjeesh NJ, Post DE, Willard MT, Kaur B, Van Meir EG, Simons JW, et al. Geldanamycin induces degradation of hypoxia-inducible factor 1 α protein via the proteasome pathway in prostate cancer cells. *Cancer Res* 2002;62:2478–82.
- [90] Gotink KJ, Verheul HM. Anti-angiogenic tyrosine kinase inhibitors: what is their mechanism of action? *Angiogenesis* 2010;13:1–14.
- [91] Mihaly Z, Sztupinszki Z, Surowiak P, Gyorffy B. A comprehensive overview of targeted therapy in metastatic renal cell carcinoma. *Curr Cancer Drug Targets* 2012;12:857–72.
- [92] Grothey A, Van Cutsem E, Sobrero A, Siena S, Falcone A, Ychou M, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013;381:303–12.
- [93] Yashiro M, Shinto O, Nakamura K, Tendo M, Matsuoka T, Matsuzaki T, et al. Effects of VEGFR-3 phosphorylation inhibitor on lymph node metastasis in an orthotopic diffuse-type gastric carcinoma model. *Br J Cancer* 2009;101:1100–6.
- [94] Penel N, Adenis A, Bocci G. Cyclophosphamide-based metronomic chemotherapy: after 10 years of experience, where do we stand and where are we going? *Crit Rev Oncol Hematol* 2012;82:40–50.
- [95] Kerbel RS. Reappraising antiangiogenic therapy for breast cancer. *Breast* 2011;20(Suppl. 3):S56–60.
- [96] Miller KD, Sweeney CJ, Sledge Jr GW. Redefining the target: chemotherapeutics as antiangiogenics. *J Clin Oncol* 2001;19:1195–206.
- [97] Klauber N, Parangi S, Flynn E, Hamel E, D'Amato RJ. Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyestradiol and taxol. *Cancer Res* 1997;57:81–6.
- [98] Browder T, Butterfield CE, Kraling BM, Shi B, Marshall B, O'Reilly MS, et al. Antiangiogenic scheduling of chemotherapy improves efficacy against experimental drug-resistant cancer. *Cancer Res* 2000;60:1878–86.

- [99] Singhal S, Mehta J, Desikan R, Ayers D, Roberson P, Eddlemon P, et al. Antitumor activity of thalidomide in refractory multiple myeloma. *N Engl J Med* 1999;341:1565–71.
- [100] Wood J, Bonjean K, Ruetz S, Bellahcene A, Devy L, Foidart JM, et al. Novel antiangiogenic effects of the bisphosphonate compound zoledronic acid. *J Pharmacol Exp Ther* 2002;302:1055–61.
- [101] Santini D, Vincenzi B, Dicuonzo G, Avvisati G, Massacesi C, Battistoni F, et al. Zoledronic acid induces significant and long-lasting modifications of circulating angiogenic factors in cancer patients. *Clin Cancer Res* 2003;9:2893–7.
- [102] Land SC, Tee AR. Hypoxia-inducible factor 1alpha is regulated by the mammalian target of rapamycin (mTOR) via an mTOR signaling motif. *J Biol Chem* 2007;282:20534–43.
- [103] Hudson CC, Liu M, Chiang GG, Otterness DM, Loomis DC, Kaper F, et al. Regulation of hypoxia-inducible factor 1alpha expression and function by the mammalian target of rapamycin. *Mol Cell Biol* 2002;22:7004–14.
- [104] Guba M, von Breitenbuch P, Steinbauer M, Koehl G, Flegel S, Hornung M, et al. Rapamycin inhibits primary and metastatic tumor growth by antiangiogenesis: involvement of vascular endothelial growth factor. *Nat Med* 2002;8:128–35.
- [105] Zhang D, Hedlund EM, Lim S, Chen F, Zhang Y, Sun B, et al. Antiangiogenic agents significantly improve survival in tumor-bearing mice by increasing tolerance to chemotherapy-induced toxicity. *Proc Natl Acad Sci U S A* 2011;108:4117–22.
- [106] Singh BN, Singh BR, Sarma BK, Singh HB. Potential chemoprevention of N-nitrosodimethylamine-induced hepatocarcinogenesis by polyphenolics from *Acacia nilotica* bark. *Chem Biol Interact* 2009;181:20–8.
- [107] Singh BN, Singh HB, Singh A, Naqvi AH, Singh BR. Dietary phytochemicals alter epigenetic events and signaling pathways for inhibition of metastasis cascade: phyto blockers of metastasis cascade. *Cancer Metastasis Rev* 2014;33(1):41–85.
- [108] Singh BN, Shankar S, Srivastava RK. Green tea catechin, epigallocatechin-3-gallate (EGCG): mechanisms, perspectives and clinical applications. *Biochem Pharmacol* 2011;82:1807–21.
- [109] Gullett NP, Ruhul Amin AR, Bayraktar S, Pezzuto JM, Shin DM, Khuri FR, et al. Cancer prevention with natural compounds. *Semin Oncol* 2010;37:258–81.
- [110] Singh M, Singh P, Shukla Y. New strategies in cancer chemoprevention by phytochemicals. *Front Biosci (Elite Ed)* 2012;4:426–52.
- [111] Bishayee A. Editorial: current advances in cancer prevention and treatment by natural products. *Curr Pharm Biotechnol* 2012;13:115–6.
- [112] Jeong D, Watari K, Shirouzu T, Ono M, Koizumi K, Saiki I, et al. Studies on lymphangiogenesis inhibitors from Korean and Japanese crude drugs. *Biol Pharm Bull* 2013;36:152–7.
- [113] Kimura Y, Sumiyoshi M. Anti-tumor and anti-metastatic actions of wogonin isolated from *Scutellaria baicalensis* roots through anti-lymphangiogenesis. *Phytomedicine* 2013;20:328–36.
- [114] Luo Y, Chen W, Zhou H, Liu L, Shen T, Alexander JS, et al. Cryptotanshinone inhibits lymphatic endothelial cell tube formation by suppressing VEGFR-3/ERK and small GTPase pathways. *Cancer Prev Res (Phila)* 2011;4:2083–91.
- [115] Coussens LM, Fingleton B, Matrisian LM. Cancer therapy – matrix metalloproteinase inhibitors and cancer: trials and tribulations. *Science* 2002;295:2387–92.
- [116] Bendrik C, Robertson J, Gauldie J, Dabrosin C. Gene transfer of matrix metalloproteinase-9 induces tumor regression of breast cancer in vivo. *Cancer Res* 2008;68:3405–12.
- [117] Hamano Y, Zeisberg M, Sugimoto H, Lively JC, Maeshima Y, Yang CQ, et al. Physiological levels of tumstatin, a fragment of collagen IV alpha 3 chain, are generated by MMP-9 proteolysis and suppress angiogenesis via alpha V beta 3 integrin. *Cancer Cell* 2003;3:589–601.
- [118] Nilsson UW, Dabrosin C. Estradiol and tamoxifen regulate endostatin generation via matrix metalloproteinase activity in breast cancer in vivo. *Cancer Res* 2006;66:4789–94.
- [119] Pozzi A, LeVine WF, Gardner HA. Low plasma levels of matrix metalloproteinase 9 permit increased tumor angiogenesis. *Oncogene* 2002;21:272–81.
- [120] Decock J, Thirkettle S, Wagstaff L, Edwards DR. Matrix metalloproteinases: protective roles in cancer. *J Cell Mol Med* 2011;15:1254–65.
- [121] Leifler KS, Svensson S, Abrahamsson A, Bendrik C, Robertson J, Gauldie J, et al. Inflammation induced by MMP-9 enhances tumor regression of experimental breast cancer. *J Immunol* 2013;190:4420–30.
- [122] Partridge JJ, Madsen MA, Ardi VC, Papagiannakopoulos T, Kupriyanova TA, Quigley JP, et al. Functional analysis of matrix metalloproteinases and tissue inhibitors of metalloproteinases differentially expressed by variants of human HT-1080 fibrosarcoma exhibiting high and low levels of intravasation and metastasis. *J Biol Chem* 2007;282:35964–77.
- [123] Koptitz C, Gerg M, Bandapalli OR, Ister D, Pennington CJ, Hauser S, et al. Tissue inhibitor of metalloproteinases-1 promotes liver metastasis by induction of hepatocyte growth factor signaling. *Cancer Res* 2007;67:8615–23.
- [124] Lipton A, Ali SM, Leitzel K, Demers L, Evans DB, Hamer P, et al. Elevated plasma tissue inhibitor of metalloproteinase-1 level predicts decreased response and survival in metastatic breast cancer. *Cancer* 2007;109:1933–9.
- [125] Miyagi M, Aoyagi K, Kato S, Shirouzu K. The TIMP-1 gene transferred through adenovirus mediation shows a suppressive effect on peritoneal metastases from gastric cancer. *Int J Clin Oncol* 2007;12:17–24.
- [126] Klintman M, Wurtz SO, Christensen IJ, Hertel PB, Ferno M, Malmberg M, et al. Association between tumor tissue TIMP-1 levels and objective response to first-line chemotherapy in metastatic breast cancer. *Breast Cancer Res Treat* 2010;121:365–71.
- [127] Schroh AS, Christensen IJ, Pedersen AN, Jensen V, Mouridsen H, Murphy G, et al. Tumor tissue concentrations of the proteinase inhibitors tissue inhibitor of metalloproteinases-1 (TIMP-1) and plasminogen activator inhibitor type 1 (PAI-1) are complementary in determining prognosis in primary breast cancer. *Mol Cell Proteomics* 2003;2:164–72.
- [128] Schroh AS, Look MP, Meijer-van Gelder ME, Foekens JA, Brunner N. Tumor tissue levels of Tissue Inhibitor of Metalloproteinases-1 (TIMP-1) and outcome following adjuvant chemotherapy in premenopausal lymph node-positive breast cancer patients: a retrospective study. *BMC Cancer* 2009;9.
- [129] Stanley G, Harvey K, Slivova V, Jiang J, Sliva D. Ganoderma lucidum suppresses angiogenesis through the inhibition of secretion of VEGF and TGF-beta1 from prostate cancer cells. *Biochem Biophys Res Commun* 2005;330:46–52.
- [130] Brakenhielm E, Cao R, Cao Y. Suppression of angiogenesis, tumor growth, and wound healing by resveratrol, a natural compound in red wine and grapes. *FASEB J* 2001;15:1798–800.
- [131] Huang YT, Hwang JJ, Lee PP, Ke FC, Huang JH, Huang CJ, et al. Effects of luteolin and quercetin, inhibitors of tyrosine kinase, on cell growth and metastasis-associated properties in A431 cells overexpressing epidermal growth factor receptor. *Br J Pharmacol* 1999;128:999–1010.
- [132] Tosetti F, Benelli R, Albini A. The angiogenic switch in solid tumors: clinical implications. *Suppl Tumori* 2002;1:S9–11.
- [133] Cao Y, Cao R. Angiogenesis inhibited by drinking tea. *Nature* 1999;398:381.
- [134] Sartippour MR, Shao ZM, Heber D, Beatty P, Zhang L, Liu C, et al. Green tea inhibits vascular endothelial growth factor (VEGF) induction in human breast cancer cells. *J Nutr* 2002;132:2307–11.
- [135] Maeda-Yamamoto M, Kawahara H, Tahara N, Tsuji K, Hara Y, Isemura M. Effects of tea polyphenols on the invasion and matrix metalloproteinases activities of human fibrosarcoma HT1080 cells. *J Agric Food Chem* 1999;47:2350–4.
- [136] Ramos S. Effects of dietary flavonoids on apoptotic pathways related to cancer chemoprevention. *J Nutr Biochem* 2007;18:427–42.
- [137] Yang H, Sun DK, Chen D, Cui QC, Gu YY, Jiang T, et al. Antitumor activity of novel fluoro-substituted (–)-epigallocatechin-3-gallate analogs. *Cancer Lett* 2010;292:48–53.
- [138] Slivova V, Zaloga G, DeMichele SJ, Mukerji P, Huang YS, Siddiqui R, et al. Green tea polyphenols modulate secretion of urokinase plasminogen activator (uPA) and inhibit invasive behavior of breast cancer cells. *Nutr Cancer* 2005;52:66–73.
- [139] Sen T, Moulik S, Dutta A, Choudhury PR, Banerji A, Das S, et al. Multifunctional effect of epigallocatechin-3-gallate (EGCG) in downregulation of gelatinase-A (MMP-2) in human breast cancer cell line MCF-7. *Life Sci* 2009;84:194–204.
- [140] Lin JK, Liang YC, Lin-Shiau SY. Cancer chemoprevention by tea polyphenols through mitotic signal transduction blockade. *Biochem Pharmacol* 1999;58:911–5.
- [141] Jung YD, Kim MS, Shin BA, Chay KO, Ahn BW, Liu W, et al. EGCG, a major component of green tea, inhibits tumour growth by inhibiting VEGF induction in human colon carcinoma cells. *Br J Cancer* 2001;84:844–50.
- [142] Khan N, Adhami VM, Mukhtar H. Review: green tea polyphenols in chemoprevention of prostate cancer: preclinical and clinical studies. *Nutr Cancer* 2009;61:836–41.
- [143] Yang CS, Wang X, Lu G, Picinich SC. Cancer prevention by tea: animal studies, molecular mechanisms and human relevance. *Nat Rev Cancer* 2009;9:429–39.
- [144] Jankun J, Selman SH, Swiercz R, Skrzypczak-Jankun E. Why drinking green tea could prevent cancer. *Nature* 1997;387:561.
- [145] Hastak K, Gupta S, Ahmad N, Agarwal MK, Agarwal ML, Mukhtar H. Role of p53 and NF-kappaB in epigallocatechin-3-gallate-induced apoptosis of LNCaP cells. *Oncogene* 2003;22:4851–9.
- [146] Masuda M, Suzui M, Lim JT, Deguchi A, Soh JW, Weinstein IB. Epigallocatechin-3-gallate decreases VEGF production in head and neck and breast carcinoma cells by inhibiting EGFR-related pathways of signal transduction. *J Exp Ther Oncol* 2002;2:350–9.
- [147] Naksuriya O, Okonogi S, Schifferers RM, Hennink WE. Curcumin nanoformulations: a review of pharmaceutical properties and preclinical studies and clinical data related to cancer treatment. *Biomaterials* 2014;35:3365–83.
- [148] Aggarwal BB, Gupta SC, Sung B. Curcumin: an orally bioavailable blocker of TNF and other pro-inflammatory biomarkers. *Br J Pharmacol* 2013;169:1672–92.
- [149] Arbisser JL, Klauber N, Rohan R, van Leeuwen R, Huang MT, Fisher C, et al. Curcumin is an in vivo inhibitor of angiogenesis. *Mol Med* 1998;4:376–83.
- [150] Hong JH, Ahn KS, Bae E, Jeon SS, Choi HY. The effects of curcumin on the invasiveness of prostate cancer in vitro and in vivo. *Prostate Cancer Prostatic Dis* 2006;9:147–52.
- [151] Yoosungnoen P, Wirachwong P, Bhattarakosol P, Niimi H, Patumraj S. Effects of curcumin on tumor angiogenesis and biomarkers, COX-2 and VEGF, in hepatocellular carcinoma cell-implanted nude mice. *Clin Hemorheol Microcirc* 2006;34:109–15.
- [152] Hasima N, Aggarwal BB. Cancer-linked targets modulated by curcumin. *Int J Biochem Mol Biol* 2012;3:328–51.
- [153] Shukla Y, Singh R. Resveratrol and cellular mechanisms of cancer prevention. *Ann N Y Acad Sci* 2011;1215:1–8.
- [154] Garvin S, Ollinger K, Dabrosin C. Resveratrol induces apoptosis and inhibits angiogenesis in human breast cancer xenografts in vivo. *Cancer Lett* 2006;231:113–22.
- [155] Bishayee A. Cancer prevention and treatment with resveratrol: from rodent studies to clinical trials. *Cancer Prev Res (Phila)* 2009;2:409–18.

- [156] Bishayee A, Petit DM, Samtani K. Angioprevention is implicated in resveratrol chemoprevention of experimental hepatocarcinogenesis. *J Carcinog Mutagen* 2010;1, <http://dx.doi.org/10.4172/2157-2518.1000102>.
- [157] Cao Z, Fang J, Xia C, Shi X, Jiang BH. *trans*-3,4,5'-Trihydroxystilbene inhibits hypoxia-inducible factor 1 α and vascular endothelial growth factor expression in human ovarian cancer cells. *Clin Cancer Res* 2004;10:5253–63.
- [158] Liu Z, Li Y, Yang R. Effects of resveratrol on vascular endothelial growth factor expression in osteosarcoma cells and cell proliferation. *Oncol Lett* 2012;4:837–9.
- [159] Srivastava RK, Unterman TG, Shankar S. FOXO transcription factors and VEGF neutralizing antibody enhance antiangiogenic effects of resveratrol. *Mol Cell Biochem* 2010;337:201–12.
- [160] Marti-Centelles R, Cejudo-Marin R, Falomir E, Murga J, Carda M, Marco JA. Inhibition of VEGF expression in cancer cells and endothelial cell differentiation by synthetic stilbene derivatives. *Bioorg Med Chem* 2013;21:3010–5.
- [161] Jung DB, Lee HJ, Jeong SJ, Lee EO, Kim YC, Ahn KS, et al. Rhapontigenin inhibited hypoxia inducible factor 1 α accumulation and angiogenesis in hypoxic PC-3 prostate cancer cells. *Biol Pharm Bull* 2011;34:850–5.
- [162] Kimura Y, Sumiyoshi M, Baba K. Antitumor activities of synthetic and natural stilbenes through antiangiogenic action. *Cancer Sci* 2008;99:2083–96.
- [163] Fotsis T, Pepper MS, Aktas E, Breit S, Rasku S, Adlercreutz H, et al. Flavonoids, dietary-derived inhibitors of cell proliferation and in vitro angiogenesis. *Cancer Res* 1997;57:2916–21.
- [164] Fotsis T, Pepper M, Adlercreutz H, Hase T, Montesano R, Schweigerer L, Genistein, a dietary ingested isoflavonoid, inhibits cell proliferation and in vitro angiogenesis. *J Nutr* 1995;125:790S–7S.
- [165] van Meeteren ME, Hendriks JJ, Dijkstra CD, van Tol EA. Dietary compounds prevent oxidative damage and nitric oxide production by cells involved in demyelinating disease. *Biochem Pharmacol* 2004;67:967–75.
- [166] Chen AY, Chen YC. A review of the dietary flavonoid, kaempferol on human health and cancer chemoprevention. *Food Chem* 2013;138:2099–107.
- [167] Duthie G, Crozier A. Plant-derived phenolic antioxidants. *Curr Opin Clin Nutr Metab Care* 2000;3:447–51.
- [168] Cortes JR, Perez GM, Rivas MD, Zamorano J. Kaempferol inhibits IL-4-induced STAT6 activation by specifically targeting JAK3. *J Immunol* 2007;179:3881–7.
- [169] Lee KM, Lee KW, Jung SK, Lee EJ, Heo YS, Bode AM, et al. Kaempferol inhibits UVB-induced COX-2 expression by suppressing Src kinase activity. *Biochem Pharmacol* 2010;80:2042–9.
- [170] Garcia-Mediavilla V, Crespo I, Collado PS, Esteller A, Sanchez-Campos S, Tunon MJ, et al. The anti-inflammatory flavones quercetin and kaempferol cause inhibition of inducible nitric oxide synthase, cyclooxygenase-2 and reactive C-protein, and down-regulation of the nuclear factor kappaB pathway in Chang Liver cells. *Eur J Pharmacol* 2007;557:221–9.
- [171] Suh Y, Afaq F, Johnson JJ, Mukhtar H. A plant flavonoid fisetin induces apoptosis in colon cancer cells by inhibition of COX2 and Wnt/EGFR/NF-kappaB-signaling pathways. *Carcinogenesis* 2009;30:300–7.
- [172] Singh RP, Sharma G, Dhanalakshmi S, Agarwal C, Agarwal R. Suppression of advanced human prostate tumor growth in athymic mice by silibinin feeding is associated with reduced cell proliferation, increased apoptosis, and inhibition of angiogenesis. *Cancer Epidemiol Biomarkers Prev* 2003;12:933–9.
- [173] Singh RP, Dhanalakshmi S, Agarwal C, Agarwal R. Silibinin strongly inhibits growth and survival of human endothelial cells via cell cycle arrest and down-regulation of survivin, Akt and NF-kappaB: implications for angioprevention and antiangiogenic therapy. *Oncogene* 2005;24:1188–202.
- [174] Jiang C, Agarwal R, Lu J. Anti-angiogenic potential of a cancer chemopreventive flavonoid antioxidant, silymarin: inhibition of key attributes of vascular endothelial cells and angiogenic cytokine secretion by cancer epithelial cells. *Biochem Biophys Res Commun* 2000;276:371–8.
- [175] Tsai SH, Liang YC, Lin-Shiau SY, Lin JK. Suppression of TNF α -mediated NF κ B activity by myricetin and other flavonoids through downregulating the activity of IKK in ECV304 cells. *J Cell Biochem* 1999;74:606–15.
- [176] Bertl E, Bartsch H, Gerhauser C. Inhibition of angiogenesis and endothelial cell functions are novel sulforaphane-mediated mechanisms in chemoprevention. *Mol Cancer Ther* 2006;5:575–85.
- [177] Kim YH, Shin EK, Kim DH, Lee HH, Park JH, Kim JK. Antiangiogenic effect of licochalcone A. *Biochem Pharmacol* 2010;80:1152–9.
- [178] Kwon GT, Cho HJ, Chung WY, Park KK, Moon A, Park JH. Isoliquiritigenin inhibits migration and invasion of prostate cancer cells: possible mediation by decreased JNK/AP-1 signaling. *J Nutr Biochem* 2009;20:663–76.
- [179] Hasmeda M, Kweifio-Okai G, Maccrides T, Polya GM. Selective inhibition of eukaryote protein kinases by anti-inflammatory triterpenoids. *Planta Med* 1999;65:14–8.
- [180] Beveridge TH, Li TS, Drover JC. Phytosterol content in American ginseng seed oil. *J Agric Food Chem* 2002;50:744–50.
- [181] Thoppil RJ, Bishayee A. Triterpenoids as potential chemopreventive and therapeutic agents in liver cancer. *World J Hepatol* 2011;3:228–49.
- [182] Bishayee A, Ahmed S, Brankov N, Perloff M. Triterpenoids as potential agents for the chemoprevention and therapy of breast cancer. *Front Biosci (Landmark Ed)* 2011;16:980–96.
- [183] Sethi G, Ahn KS, Pandey MK, Aggarwal BB. Celastrol, a novel triterpene, potentiates TNF-induced apoptosis and suppresses invasion of tumor cells by inhibiting NF-kappaB-regulated gene products and TAK1-mediated NF-kappaB activation. *Blood* 2007;109:2727–35.
- [184] Huang Y, Zhou Y, Fan Y, Zhou D. Celastrol inhibits the growth of human glioma xenografts in nude mice through suppressing VEGFR expression. *Cancer Lett* 2008;264:101–6.
- [185] Matsuda H, Li Y, Murakami T, Ninomiya K, Yamahara J, Yoshikawa M. Effects of escins Ia, Ib, IIa, and IIb from horse chestnut, the seeds of *Aesculus hippocastanum* L., on acute inflammation in animals. *Biol Pharm Bull* 1997;20:1092–5.
- [186] Wang XH, Xu B, Liu JT, Cui JR. Effect of beta-escin sodium on endothelial cells proliferation, migration and apoptosis. *Vasc Pharmacol* 2008;49:158–65.
- [187] Yang H, Shi G, Dou QP. The tumor proteasome is a primary target for the natural anticancer compound Withaferin A isolated from "Indian winter cherry". *Mol Pharmacol* 2007;71:426–37.
- [188] Mohan R, Hammers HJ, Bargagna-Mohan P, Zhan XH, Herbstritt CJ, Ruiz A, et al. Withaferin A is a potent inhibitor of angiogenesis. *Angiogenesis* 2004;7:115–22.
- [189] Jayaprakasam B, Zhang Y, Seeram NP, Nair MG. Growth inhibition of human tumor cell lines by withanolides from *Withania somnifera* leaves. *Life Sci* 2003;74:125–32.
- [190] Park JS, Chew BP, Wong TS. Dietary lutein from marigold extract inhibits mammary tumor development in BALB/c mice. *J Nutr* 1998;128:1650–6.
- [191] Krinsky NI, Johnson EJ. Carotenoid actions and their relation to health and disease. *Mol Aspects Med* 2005;26:459–516.
- [192] Milder IE, Arts IC, van de Putte B, Venema DP, Hollman PC. Lignan contents of Dutch plant foods: a database including lariciresinol, pinoresinol, secoisolariciresinol and matairesinol. *Br J Nutr* 2005;93:393–402.
- [193] Thompson LU, Boucher BA, Liu Z, Cotterchio M, Kreiger N. Phytoestrogen content of foods consumed in Canada, including isoflavones, lignans, and coumestran. *Nutr Cancer* 2006;54:184–201.
- [194] Mazur W, Fotsis T, Wahala K, Ojala S, Salakka A, Adlercreutz H. Isotope dilution gas chromatographic-mass spectrometric method for the determination of isoflavonoids, coumestrol, and lignans in food samples. *Anal Biochem* 1996;233:169–80.
- [195] Touillaud MS, Thiebaut AC, Fournier A, Niravong M, Boutron-Ruault MC, Clavel-Chapelon F. Dietary lignan intake and postmenopausal breast cancer risk by estrogen and progesterone receptor status. *J Natl Cancer Inst* 2007;99:475–86.
- [196] Trock BJ, Hilakivi-Clarke L, Clarke R. Meta-analysis of soy intake and breast cancer risk. *J Natl Cancer Inst* 2006;98:459–71.
- [197] Banerjee S, Li Y, Wang Z, Sarkar FH. Multi-targeted therapy of cancer by genistein. *Cancer Lett* 2008;269:226–42.
- [198] Helferich WG, Andrade JE, Hoagland MS. Phytoestrogens and breast cancer: a complex story. *Inflammopharmacology* 2008;16:219–26.
- [199] Lindahl G, Saarinen N, Abrahamsson A, Dabrosin C. Tamoxifen, flaxseed, and the lignan enterolactone increase stroma- and cancer cell-derived IL-1Ra and decrease tumor angiogenesis in estrogen-dependent breast cancer. *Cancer Res* 2011;71:51–60.
- [200] Bergman Jungstrom M, Thompson LU, Dabrosin C. Flaxseed and its lignans inhibit estradiol-induced growth, angiogenesis, and secretion of vascular endothelial growth factor in human breast cancer xenografts in vivo. *Clin Cancer Res* 2007;13:1061–7.
- [201] Saarinen NM, Abrahamsson A, Dabrosin C. Estrogen-induced angiogenic factors derived from stromal and cancer cells are differently regulated by enterolactone and genistein in human breast cancer in vivo. *Int J Cancer* 2010;127:737–45.
- [202] Chatterjee S, Bhattacharjee B. Use of natural molecules as anti-angiogenic inhibitors for vascular endothelial growth factor receptor. *Bioinformation* 2012;8:1249–54.
- [203] Abusnina A, Keravis T, Zhou Q, Justiniano H, Lobstein A, Luginier C. Tumour growth inhibition and anti-angiogenic effects using curcumin correspond to combined PDE2 and PDE4 inhibition. *Thromb Haemost* 2015;113:319–28.
- [204] Rath S, Liebl J, Furst R, Ullrich A, Burkhart JL, Kazmaier U, et al. Anti-angiogenic effects of the tubulysin precursor pretubulysin and of simplified pretubulysin derivatives. *Br J Pharmacol* 2012;167:1048–61.
- [205] Schlupe T, Gunawan P, Ma L, Jensen GS, Düringer J, Hinton S, et al. Polymeric tubulysin-peptide nanoparticles with potent antitumor activity. *Clin Cancer Res* 2009;15:181–9.
- [206] Kubisch R, von Gamm M, Braig S, Ullrich A, Burkhart JL, Colling L, et al. Simplified pretubulysin derivatives and their biological effects on cancer cells. *J Nat Prod* 2014;77:536–42.
- [207] Peer D, Karp JM, Hong S, Farokhzad OC, Margalit R, Langer R. Nanocarriers as an emerging platform for cancer therapy. *Nat Nanotechnol* 2007;2:751–60.
- [208] Gref R, Minamitake Y, Peracchia MT, Trubetskoy V, Torchilin V, Langer R. Biodegradable long-circulating polymeric nanospheres. *Science* 1994;263:1600–3.
- [209] Siddiqui IA, Adhmi VM, Bharali DJ, Hafeez BB, Asim M, Khwaja SI, et al. Introducing nanochemoprevention as a novel approach for cancer control: proof of principle with green tea polyphenol epigallocatechin-3-gallate. *Cancer Res* 2009;69:1712–6.
- [210] Luo H, Jiang B, Li B, Li Z, Jiang BH, Chen YC. Kaempferol nanoparticles achieve strong and selective inhibition of ovarian cancer cell viability. *Int J Nanomed* 2012;7:3951–9.
- [211] Gou M, Men K, Shi H, Xiang M, Zhang J, Song J, et al. Curcumin-loaded biodegradable polymeric micelles for colon cancer therapy in vitro and in vivo. *Nanoscale* 2011;3:1558–67.
- [212] Wang L, Li H, Wang S, Liu R, Wu Z, Wang C, et al. Enhancing the antitumor activity of berberine hydrochloride by solid lipid nanoparticle encapsulation. *AAPS PharmSciTech* 2014;15:834–44.
- [213] Parikh NR, Mandal A, Bhatia D, Siveen KS, Sethi G, Bishayee A. Oleanane triterpenoids in the prevention and therapy of breast cancer: current evidence and future perspectives. *Phytochem Rev* 2014;13:793–810.

- [214] Sohn KH, Lee HY, Chung HY, Young HS, Yi SY, Kim KW. Anti-angiogenic activity of triterpene acids. *Cancer Lett* 1995;94:213–8.
- [215] Sogno I, Vannini N, Lorusso G, Cammarota R, Noonan DM, Generoso L, et al. Anti-angiogenic activity of a novel class of chemopreventive compounds: oleanic acid terpenoids. *Recent Results Cancer Res* 2009;181:209–12.
- [216] Masferrer JL, Leahy KM, Koki AT, Zweifel BS, Settle SL, Woerner BM, et al. Antiangiogenic and antitumor activities of cyclooxygenase-2 inhibitors. *Cancer Res* 2000;60:1306–11.
- [217] Parrett M, Harris R, Joarder F, Ross M, Clausen K, Robertson F. Cyclooxygenase-2 gene expression in human breast cancer. *Int J Oncol* 1997;10:503–7.
- [218] Voss MJ, Niggemann B, Zanker KS, Entschladen F. Tumour reactions to hypoxia. *Curr Mol Med* 2010;10:381–6.
- [219] Lofstedt T, Jogi A, Sigvardsson M, Gradin K, Poellinger L, Pahlman S, et al. Induction of ID2 expression by hypoxia-inducible factor-1: a role in dedifferentiation of hypoxic neuroblastoma cells. *J Biol Chem* 2004;279:39223–31.
- [220] Garcia-Maceira P, Mateo J. Silibinin inhibits hypoxia-inducible factor-1 α and mTOR/p70S6K/4E-BP1 signalling pathway in human cervical and hepatoma cancer cells: implications for anticancer therapy. *Oncogene* 2009;28:313–24.
- [221] Otolara BB, Madrid JA, Alvarez N, Vicente V, Rol MA. Effects of exogenous melatonin and circadian synchronization on tumor progression in melanoma-bearing C57BL6 mice. *J Pineal Res* 2008;44:307–15.
- [222] Trapp V, Parmakhtiar B, Papazian V, Willmott L, Fruehauf JP. Anti-angiogenic effects of resveratrol mediated by decreased VEGF and increased TSP1 expression in melanoma-endothelial cell co-culture. *Angiogenesis* 2010;13:305–15.
- [223] Park SY, Jeong KJ, Lee J, Yoon DS, Choi WS, Kim YK, et al. Hypoxia enhances LPA-induced HIF-1 α and VEGF expression: their inhibition by resveratrol. *Cancer Lett* 2007;258:63–9.
- [224] Li JL, Sainson RC, Shi W, Leek R, Harrington LS, Preusser M, et al. Delta-like 4 notch ligand regulates tumor angiogenesis, improves tumor vascular function, and promotes tumor growth in vivo. *Cancer Res* 2007;67:11244–53.
- [225] Lickliter JD, Francesconi AB, Smith G, Burge M, Coulthard A, Rose S, et al. Phase I trial of CYT997, a novel cytotoxic and vascular-disrupting agent. *Br J Cancer* 2010;103:597–606.
- [226] Lekli I, Ray D, Mukherjee S, Gurusamy N, Ahsan MK, Juhasz B, et al. Co-ordinated autophagy with resveratrol and gamma-tocotrienol confers synergistic cardioprotection. *J Cell Mol Med* 2010;14:2506–18.
- [227] Kuang X, Yao Y, Du JR, Liu YX, Wang CY, Qian ZM. Neuroprotective role of Z-ligustilide against forebrain ischemic injury in ICR mice. *Brain Res* 2006;1102:145–53.
- [228] Forssell J, Oberg A, Henriksson ML, Stenling R, Jung A, Palmqvist R. High macrophage infiltration along the tumor front correlates with improved survival in colon cancer. *Clin Cancer Res* 2007;13:1472–9.
- [229] Garaude J, Kent A, van Rooijen N, Blander JM. Simultaneous targeting of toll and nod-like receptors induces effective tumor-specific immune responses. *Sci Transl Med* 2012;4:120ra116.
- [230] Tosetti F, Noonan DM, Albini A. Metabolic regulation and redox activity as mechanisms for angioprevention by dietary phytochemicals. *Int J Cancer* 2009;125:1997–2003.
- [231] Deep G, Gangar SC, Rajamanickam S, Raina K, Gu M, Agarwal C, et al. Angiopreventive efficacy of pure flavonolignans from milk thistle extract against prostate cancer: targeting VEGF–VEGFR signaling. *PLoS ONE* 2012;7:e34630.
- [232] Surh YJ. Cancer chemoprevention with dietary phytochemicals. *Nat Rev Cancer* 2003;3:768–80.
- [233] Thomasset SC, Berry DP, Garcea G, Marczylo T, Steward WP, Gescher AJ. Dietary polyphenolic phytochemicals – promising cancer chemopreventive agents in humans? A review of their clinical properties. *Int J Cancer* 2007;120:451–8.
- [234] Bisacchi D, Benelli R, Vanzetto C, Ferrari N, Tosetti F, Albini A. Anti-angiogenesis and angioprevention: mechanisms, problems and perspectives. *Cancer Detect Prev* 2003;27:229–38.
- [235] Macpherson H, Pipingas A, Pase MP. Multivitamin-multimineral supplementation and mortality: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2013;97:437–44.
- [236] Klein EA, Thompson Jr IM, Tangen CM, Crowley JJ, Lucia MS, Goodman PJ, et al. Vitamin E and the risk of prostate cancer: the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA* 2011;306:1549–56.
- [237] Sommer A, Vyas KS. A global clinical view on vitamin A and carotenoids. *Am J Clin Nutr* 2012;96:1204S–6S.
- [238] Yang Y, Zhang Y, Cao Z, Ji H, Yang X, Iwamoto H, et al. Anti-VEGF- and anti-VEGF receptor-induced vascular alteration in mouse healthy tissues. *Proc Natl Acad Sci U S A* 2013;110:12018–23.
- [239] Kumareswaran R, Ludkovski O, Meng A, Sykes J, Pintlilie M, Bristow RG. Chronic hypoxia compromises repair of DNA double-strand breaks to drive genetic instability. *J Cell Sci* 2012;125:189–99.
- [240] Gaddameedhi S, Reardon JT, Ye R, Ozturk N, Sancar A. Effect of circadian clock mutations on DNA damage response in mammalian cells. *Cell Cycle* 2012;11:3481–91.
- [241] Dai C, Sun F, Zhu C, Hu X. Tumor environmental factors glucose deprivation and lactic acidosis induce mitotic chromosomal instability – an implication in aneuploid human tumors. *PLoS ONE* 2013;8:e63054.
- [242] Xiang L, Gilkes DM, Chaturvedi P, Luo W, Hu H, Takano N, et al. Ganetespib blocks HIF-1 activity and inhibits tumor growth, vascularization, stem cell maintenance, invasion, and metastasis in orthotopic mouse models of triple-negative breast cancer. *J Mol Med (Berl)* 2014;92:151–64.
- [243] Leelahavanichkul K, Amornphimoltham P, Molinolo AA, Basile JR, Koon-tongkaew S, Gutkind JS. A role for p38 MAPK in head and neck cancer cell growth and tumor-induced angiogenesis and lymphangiogenesis. *Mol Oncol* 2014;8:105–18.
- [244] Thiele W, Rothley M, Teller N, Jung N, Bulat B, Plaumann D, et al. Delphinidin is a novel inhibitor of lymphangiogenesis but promotes mammary tumor growth and metastasis formation in syngeneic experimental rats. *Carcinogenesis* 2013;34:2804–13.
- [245] Yu T, Liu K, Wu Y, Fan J, Chen J, Li C, et al. High interstitial fluid pressure promotes tumor cell proliferation and invasion in oral squamous cell carcinoma. *Int J Mol Med* 2013;32:1093–100.
- [246] Hofmann M, Guschel M, Bernd A, Bereiter-Hahn J, Kaufmann R, Tandl C, et al. Lowering of tumor interstitial fluid pressure reduces tumor cell proliferation in a xenograft tumor model. *Neoplasia* 2006;8:89–95.
- [247] Rana S, Munawar M, Shahid A, Malik M, Ullah H, Fatima W, et al. Deregulated expression of circadian clock and clock-controlled cell cycle genes in chronic lymphocytic leukemia. *Mol Biol Rep* 2014;41:95–103.
- [248] Sotak M, Polidarova L, Ergang P, Sumova A, Pacha J. An association between clock genes and clock-controlled cell cycle genes in murine colorectal tumors. *Int J Cancer* 2013;132:1032–41.
- [249] Orimo A, Gupta PB, Sgroi DC, Arenzana-Seisdedos F, Delaunay T, Naeem R, et al. Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth and angiogenesis through elevated SDF-1/CXCL12 secretion. *Cell* 2005;121:335–48.
- [250] Jia CC, Wang TT, Liu W, Fu BS, Hua X, Wang GY, et al. Cancer-associated fibroblasts from hepatocellular carcinoma promote malignant cell proliferation by HGF secretion. *PLoS ONE* 2013;8:e63243.
- [251] Wang YH, Dong YY, Wang WM, Xie XY, Wang ZM, Chen RX, et al. Vascular endothelial cells facilitated HCC invasion and metastasis through the Akt and NF- κ B pathways induced by paracrine cytokines. *J Exp Clin Cancer Res* 2013;32:51.
- [252] Shi S, Chen L, Huang G. Antiangiogenic therapy improves the antitumor effect of adoptive cell immunotherapy by normalizing tumor vasculature. *Med Oncol* 2013;30:698.
- [253] Casazza A, Laoui D, Wenes M, Rizzolio S, Bassani N, Mambretti M, et al. Impeding macrophage entry into hypoxic tumor areas by Sema3A/Nrp1 signaling blockade inhibits angiogenesis and restores antitumor immunity. *Cancer Cell* 2013;24:695–709.
- [254] Tripathi C, Tewari BN, Kanchan RK, Baghel KS, Nautiyal N, Shrivastava R, et al. Macrophages are recruited to hypoxic tumor areas and acquire a pro-angiogenic M2-polarized phenotype via hypoxic cancer cell derived cytokines Oncostatin M and Eotaxin. *Oncotarget* 2014;5:5350–68.
- [255] Tomei AA, Siegert S, Britschgi MR, Luther SA, Swartz MA. Fluid flow regulates stromal cell organization and CCL21 expression in a tissue-engineered lymph node microenvironment. *J Immunol* 2009;183:4273–83.
- [256] Mamlouk S, Wielockx B. Hypoxia-inducible factors as key regulators of tumor inflammation. *Int J Cancer* 2013;132:2721–9.
- [257] Lee JH, Sancar A. Regulation of apoptosis by the circadian clock through NF- κ B signaling. *Proc Natl Acad Sci U S A* 2011;108:12036–41.
- [258] Erez N, Truitt M, Olson P, Arron ST, Hanahan D. Cancer-associated fibroblasts are activated in incipient neoplasia to orchestrate tumor-promoting inflammation in an NF- κ B-dependent manner. *Cancer Cell* 2010;17:135–47.
- [259] Ohashi T, Akazawa T, Aoki M, Kuze B, Mizuta K, Ito Y, et al. Dichloroacetate improves immune dysfunction caused by tumor-secreted lactic acid and increases antitumor immunoreactivity. *Int J Cancer* 2013;133:1107–18.
- [260] Ghajar CM, Peinado H, Mori H, Matei IR, Evason KJ, Brazier H, et al. The perivascular niche regulates breast tumor dormancy. *Nat Cell Biol* 2013;15:807–17.
- [261] Gardner LB, Li Q, Park MS, Flanagan WM, Semenza GL, Dang CV. Hypoxia inhibits G1/S transition through regulation of p27 expression. *J Biol Chem* 2001;276:7919–26.
- [262] Goda N, Ryan HE, Khadivi B, McNulty W, Rickert RC, Johnson RS. Hypoxia-inducible factor 1 α is essential for cell cycle arrest during hypoxia. *Mol Cell Biol* 2003;23:359–69.
- [263] Yamanda S, Ebihara S, Asada M, Okazaki T, Niu K, Ebihara T, et al. Role of ephrinB2 in nonproductive angiogenesis induced by Delta-like 4 blockade. *Blood* 2009;113:3631–9.
- [264] Bussink J, Kaanders JH, Rijken PF, Martindale CA, van der Kogel AJ. Multiparameter analysis of vasculature, perfusion and proliferation in human tumour xenografts. *Br J Cancer* 1998;77:57–64.
- [265] Rupertus K, Dahlem C, Menger MD, Schilling MK, Kollmar O. Rapamycin inhibits hepatectomy-induced stimulation of metastatic tumor growth by reduction of angiogenesis, microvascular blood perfusion, and tumor cell proliferation. *Ann Surg Oncol* 2009;16:2629–37.
- [266] Li XM, Delaunay F, Dulong S, Claustrat B, Zampera S, Fujii Y, et al. Cancer inhibition through circadian reprogramming of tumor transcriptome with meal timing. *Cancer Res* 2010;70:3351–60.
- [267] Yang WS, Stockwell BR. Inhibition of casein kinase 1-epsilon induces cancer-cell-selective, PERIOD2-dependent growth arrest. *Genome Biol* 2008;9:R92.
- [268] Lin H, Yan J, Wang Z, Hua F, Yu J, Sun W, et al. Loss of immunity-supported senescence enhances susceptibility to hepatocellular carcinogenesis and progression in Toll-like receptor 2-deficient mice. *Hepatology* 2013;57:171–82.
- [269] Wang W, Li Q, Yamada T, Matsumoto K, Matsumoto I, Oda M, et al. Crosstalk to stromal fibroblasts induces resistance of lung cancer to epidermal growth factor receptor tyrosine kinase inhibitors. *Clin Cancer Res* 2009;15:6630–8.
- [270] Pontiggia O, Sampayo R, Raffo D, Motter A, Xu R, Bissell MJ, et al. The tumor microenvironment modulates tamoxifen resistance in breast cancer: a role for

- soluble stromal factors and fibronectin through beta1 integrin. *Breast Cancer Res Treat* 2012;133:459–71.
- [271] Madhok BM, Yeluri S, Perry SL, Hughes TA, Jayne DG. Dichloroacetate induces apoptosis and cell-cycle arrest in colorectal cancer cells. *Br J Cancer* 2010;102:1746–52.
- [272] Riemann A, Schneider B, Ihling A, Nowak M, Sauvant C, Thews O, et al. Acidic environment leads to ROS-induced MAPK signaling in cancer cells. *PLoS ONE* 2011;6:e22445.
- [273] Vangestel C, Van de Wiele C, Van Damme N, Staelens S, Pauwels P, Reutelingsperger CP, et al. (99mTc-(CO)(3) His-annexin A5 micro-SPECT demonstrates increased cell death by irinotecan during the vascular normalization window caused by bevacizumab. *J Nucl Med* 2011;52:1786–94.
- [274] Hernandez-Luna MA, Rocha-Zavaleta L, Vega MI, Huerta-Yepez S. Hypoxia inducible factor-1alpha induces chemoresistance phenotype in non-Hodgkin Lymphoma cell line via up-regulation of Bcl-xL. *Leuk Lymphoma* 2013;54:1048–55.
- [275] Islam N, Haqqi TM, Jepsen KJ, Kraay M, Welter JF, Goldberg VM, et al. Hydrostatic pressure induces apoptosis in human chondrocytes from osteoarthritic cartilage through up-regulation of tumor necrosis factor-alpha, inducible nitric oxide synthase, p53, c-myc, and bax-alpha, and suppression of bcl-2. *J Cell Biochem* 2002;87:266–78.
- [276] Campbell NE, Greenaway J, Henkin J, Moorehead RA, Petrik J. The thrombospondin-1 mimetic ABT-510 increases the uptake and effectiveness of cisplatin and paclitaxel in a mouse model of epithelial ovarian cancer. *Neoplasia* 2010;12:275–83.
- [277] Hua H, Wang Y, Wan C, Liu Y, Zhu B, Yang C, et al. Circadian gene mPer2 overexpression induces cancer cell apoptosis. *Cancer Sci* 2006;97:589–96.
- [278] Schafer H, Struck B, Feldmann EM, Bergmann F, Grage-Griebenow E, Geismann C, et al. TGF-beta1-dependent L1CAM expression has an essential role in macrophage-induced apoptosis resistance and cell migration of human intestinal epithelial cells. *Oncogene* 2013;32:180–9.
- [279] Kinugasa Y, Matsui T, Takakura N. CD44 expressed on cancer-associated fibroblasts is a functional molecule supporting the stemness and drug resistance of malignant cancer cells in the tumor microenvironment. *Stem Cells* 2014;32:145–56.
- [280] Golding JP, Wardhaugh T, Patrick L, Turner M, Phillips JB, Bruce JI, et al. Targeting tumour energy metabolism potentiates the cytotoxicity of 5-aminolevulinic acid photodynamic therapy. *Br J Cancer* 2013;109:976–82.
- [281] Nishi H, Nakada T, Kyo S, Inoue M, Shay JW, Isaka K. Hypoxia-inducible factor 1 mediates upregulation of telomerase (hTERT). *Mol Cell Biol* 2004;24:6076–83.
- [282] Heeg S, Hirt N, Queisser A, Schmiege H, Thaler M, Kunert H, et al. EGFR overexpression induces activation of telomerase via PI3K/AKT-mediated phosphorylation and transcriptional regulation through Hif1-alpha in a cellular model of oral-esophageal carcinogenesis. *Cancer Sci* 2011;102:351–60.
- [283] Anderson CJ, Hoare SF, Ashcroft M, Bilsland AE, Keith WN. Hypoxic regulation of telomerase gene expression by transcriptional and post-transcriptional mechanisms. *Oncogene* 2006;25:61–9.
- [284] Qu Y, Wang Z, Huang X, Wan C, Yang CL, Liu B, et al. Circadian telomerase activity and DNA synthesis for timing peptide administration. *Peptides* 2003;24:363–9.
- [285] Qu Y, Mao M, Li X, Liu Y, Ding J, Jiang Z, et al. Telomerase reconstitution contributes to resetting of circadian rhythm in fibroblasts. *Mol Cell Biochem* 2008;313:11–8.
- [286] Martinive P, Defresne F, Bouzin C, Saliez J, Lair F, Gregoire V, et al. Preconditioning of the tumor vasculature and tumor cells by intermittent hypoxia: implications for anticancer therapies. *Cancer Res* 2006;66:11736–44.
- [287] Mazzone M, Dettori D, Leite de Oliveira R, Loges S, Schmidt T, Jonckx B, et al. Heterozygous deficiency of PHD2 restores tumor oxygenation and inhibits metastasis via endothelial normalization. *Cell* 2009;136:839–51.
- [288] Kieda C, El Hafny-Rahbi B, Collet G, Lamerant-Fayel N, Grillon C, Guichard A, et al. Stable tumor vessel normalization with pO(2) increase and endothelial PTEN activation by inositol trispyrophosphate brings novel tumor treatment. *J Mol Med (Berl)* 2013;91:883–99.
- [289] Sun RC, Denko NC. Hypoxic regulation of glutamine metabolism through HIF1 and SIAH2 supports lipid synthesis that is necessary for tumor growth. *Cell Metab* 2014;19:285–92.
- [290] Liao J, Qian F, Tchabo N, Mhawech-Fauceglia P, Beck A, Qian Z, et al. Ovarian cancer spheroid cells with stem cell-like properties contribute to tumor generation, metastasis and chemotherapy resistance through hypoxia-resistant metabolism. *PLoS ONE* 2014;9:e84941.
- [291] Lamonte G, Tang X, Chen JL, Wu J, Ding CK, Keenan MM, et al. Acidosis induces reprogramming of cellular metabolism to mitigate oxidative stress. *Cancer Metab* 2013;1:23.
- [292] Matsubara T, Diresta GR, Kakunaga S, Li D, Healey JH. Additive influence of extracellular pH, oxygen tension, and pressure on invasiveness and survival of human osteosarcoma cells. *Front Oncol* 2013;3:199.
- [293] Kelly CJ, Hussien K, Fokas E, Kannan P, Shipley RJ, Ashton TM, et al. Regulation of O₂ consumption by the PI3K and mTOR pathways contributes to tumor hypoxia. *Radiother Oncol* 2014;111:72–80.
- [294] Seront E, Boidot R, Bouzin C, Karroum O, Jordan BF, Gallez B, et al. Tumour hypoxia determines the potential of combining mTOR and autophagy inhibitors to treat mammary tumours. *Br J Cancer* 2013;109:2597–606.
- [295] Okazaki H, Matsunaga N, Fujioka T, Okazaki F, Akagawa Y, Tsurudome Y, et al. Circadian regulation of mTOR by the ubiquitin pathway in renal cell carcinoma. *Cancer Res* 2014;74:543–51.
- [296] Rattigan YI, Patel BB, Ackerstaff E, Sukenick G, Koutcher JA, Glod JW, et al. Lactate is a mediator of metabolic cooperation between stromal carcinoma associated fibroblasts and glycolytic tumor cells in the tumor microenvironment. *Exp Cell Res* 2012;318:326–35.
- [297] Wojtkowiak JW, Rothberg JM, Kumar V, Schramm KJ, Haller E, Proemsey JB, et al. Chronic autophagy is a cellular adaptation to tumor acidic pH microenvironments. *Cancer Res* 2012;72:3938–47.
- [298] Belkin DA, Mitsui H, Wang CQ, Gonzalez J, Zhang S, Shah KR, et al. CD200 upregulation in vascular endothelium surrounding cutaneous squamous cell carcinoma. *JAMA Dermatol* 2013;149:178–86.
- [299] Hamzah J, Jugold M, Kiessling F, Rigby P, Manzur M, Marti HH, et al. Vascular normalization in Rgs5-deficient tumours promotes immune destruction. *Nature* 2008;453:410–4.
- [300] Chi Sabins N, Taylor JL, Fabian KP, Appleman LJ, Maranchie JK, Stolz DB, et al. DLK1: a novel target for immunotherapeutic remodeling of the tumor blood vasculature. *Mol Ther* 2013;21:1958–68.
- [301] Garuffi A, Pistrutto G, Ceci C, Di Renzo L, Santarelli R, Faggioni A, et al. Targeting COX-2/PGE(2) pathway in HIPK2 knockdown cancer cells: impact on dendritic cell maturation. *PLoS ONE* 2012;7:e48342.
- [302] Deng B, Zhu JM, Wang Y, Liu TT, Ding YB, Xiao WM, et al. Intratumor hypoxia promotes immune tolerance by inducing regulatory T cells via TGF-beta1 in gastric cancer. *PLoS ONE* 2013;8:e63777.
- [303] Lund AW, Duraes FV, Hirose S, Raghavan VR, Nembrini C, Thomas SN, et al. VEGF-C promotes immune tolerance in B16 melanomas and cross-presentation of tumor antigen by lymph node lymphatics. *Cell Rep* 2012;1:191–9.
- [304] Craig DH, Schaubert KL, Shiratsuchi H, Kan-Mitchell J, Basson MD. Increased pressure stimulates aberrant dendritic cell maturation. *Cell Mol Biol Lett* 2008;13:260–70.
- [305] Baronzio G, Schwartz L, Kiselevsky M, Guais A, Sanders E, Milanese G, et al. Tumor interstitial fluid as modulator of cancer inflammation, thrombosis, immunity and angiogenesis. *Anticancer Res* 2012;32:405–14.
- [306] Preise D, Scherz A, Salomon Y. Antitumor immunity promoted by vascular occluding therapy: lessons from vascular-targeted photodynamic therapy (VTP). *Photochem Photobiol Sci* 2011;10:681–8.
- [307] Miyazaki K, Wakabayashi M, Hara Y, Ishida N. Tumor growth suppression in vivo by overexpression of the circadian component, PER2. *Genes Cells* 2010;15:351–8.
- [308] Hsiao YW, Li CF, Chi JY, Tseng JT, Chang Y, Hsu LJ, et al. CCAAT/enhancer binding protein delta in macrophages contributes to immunosuppression and inhibits phagocytosis in nasopharyngeal carcinoma. *Sci Signal* 2013;6:ra59.
- [309] Daurkin I, Eruslanov E, Stoffs T, Perrin GQ, Algood C, Gilbert SM, et al. Tumor-associated macrophages mediate immunosuppression in the renal cancer microenvironment by activating the 15-lipoxygenase-2 pathway. *Cancer Res* 2011;71:6400–9.
- [310] Mace TA, Ameen Z, Collins A, Wojcik S, Mair M, Young GS, et al. Pancreatic cancer-associated stellate cells promote differentiation of myeloid-derived suppressor cells in a STAT3-dependent manner. *Cancer Res* 2013;73:3007–18.
- [311] Villalba M, Rathore MG, Lopez-Royuela N, Krzywinska E, Garaude J, Allende-Vega N. From tumor cell metabolism to tumor immune escape. *Int J Biochem Cell Biol* 2013;45:106–13.
- [312] Chirasani SR, Leukel P, Gottfried E, Hochrein J, Stadler K, Neumann B, et al. Diclofenac inhibits lactate formation and efficiently counteracts local immune suppression in a murine glioma model. *Int J Cancer* 2013;132:843–53.
- [313] Calcinotto A, Filipazzi P, Groni M, Iero M, De Milito A, Ricupito A, et al. Modulation of microenvironment acidity reverses energy in human and murine tumor-infiltrating T lymphocytes. *Cancer Res* 2012;72:2746–56.
- [314] James SE, Salsbury AJ. Effect of (plus or minus)-1,2-bis(3,5-dioxopiperazin-1-yl)propane on tumor blood vessels and its relationship to the antimetastatic effect in the Lewis lung carcinoma. *Cancer Res* 1974;34:839–42.
- [315] Lee SL, Rouhi P, Dahl Jensen L, Zhang D, Ji H, Hauptmann G, et al. Hypoxia-induced pathological angiogenesis mediates tumor cell dissemination, invasion, and metastasis in a zebrafish tumor model. *Proc Natl Acad Sci U S A* 2009;106:19485–90.
- [316] Cao Y. Opinion: emerging mechanisms of tumour lymphangiogenesis and lymphatic metastasis. *Nat Rev Cancer* 2005;5:735–43.
- [317] Lunt SJ, Kalliomaki TM, Brown A, Yang VX, Milosevic M, Hill RP. Interstitial fluid pressure, vascularity and metastasis in ectopic, orthotopic and spontaneous tumours. *BMC Cancer* 2008;8:2.
- [318] Rofstad EK, Tunheim SH, Mathiesen B, Graff BA, Halsor EF, Nilsen K, et al. Pulmonary and lymph node metastasis is associated with primary tumor interstitial fluid pressure in human melanoma xenografts. *Cancer Res* 2002;62:661–4.
- [319] Yu T, Wang Z, Liu K, Wu Y, Fan J, Chen J, et al. High interstitial fluid pressure promotes tumor progression through inducing lymphatic metastasis-related protein expressions in oral squamous cell carcinoma. *Clin Transl Oncol* 2014;16:539–47.
- [320] Rofstad EK, Gaustad JV, Egeland TA, Mathiesen B, Galappathi K. Tumors exposed to acute cyclic hypoxic stress show enhanced angiogenesis, perfusion and metastatic dissemination. *Int J Cancer* 2010;127:1535–46.
- [321] Oshima T, Takenoshita S, Akaike M, Kunisaki C, Fujii S, Nozaki A, et al. Expression of circadian genes correlates with liver metastasis and outcomes in colorectal cancer. *Oncol Rep* 2011;25:1439–46.
- [322] Wu M, Zeng J, Chen Y, Zeng Z, Zhang J, Cai Y, et al. Experimental chronic jet lag promotes growth and lung metastasis of Lewis lung carcinoma in C57BL/6 mice. *Oncol Rep* 2012;27:1417–28.

- [323] Zhang Y, Tang H, Cai J, Zhang T, Guo J, Feng D, et al. Ovarian cancer-associated fibroblasts contribute to epithelial ovarian carcinoma metastasis by promoting angiogenesis, lymphangiogenesis and tumor cell invasion. *Cancer Lett* 2011;303:47–55.
- [324] Qian BZ, Li J, Zhang H, Kitamura T, Zhang J, Campion LR, et al. CCL2 recruits inflammatory monocytes to facilitate breast-tumour metastasis. *Nature* 2011;475:222–5.
- [325] Rofstad EK, Mathiesen B, Kindem K, Galappathi K. Acidic extracellular pH promotes experimental metastasis of human melanoma cells in athymic nude mice. *Cancer Res* 2006;66:6699–707.
- [326] Peppicelli S, Bianchini F, Torre E, Calorini L. Contribution of acidic melanoma cells undergoing epithelial-to-mesenchymal transition to aggressiveness of non-acidic melanoma cells. *Clin Exp Metastasis* 2014;31(4):423–33.
- [327] Huang Y, Yuan J, Righi E, Kamoun WS, Ancukiewicz M, Nezivar J, et al. Vascular normalizing doses of antiangiogenic treatment reprogram the immunosuppressive tumor microenvironment and enhance immunotherapy. *Proc Natl Acad Sci U S A* 2012;109:17561–6.
- [328] Mo J, Sun B, Zhao X, Gu Q, Dong X, Liu Z, et al. Hypoxia-induced senescence contributes to the regulation of microenvironment in melanomas. *Pathol Res Pract* 2013;209:640–7.
- [329] Shin JY, Yoon IH, Kim JS, Kim B, Park CG. Vascular endothelial growth factor-induced chemotaxis and IL-10 from T cells. *Cell Immunol* 2009;256:72–8.
- [330] Aung KZ, Pereira BP, Tan PH, Han HC, Nathan SS. Interstitial fluid pressure as an alternate regulator of angiogenesis independent of hypoxia driven HIF-1 α in solid tumors. *J Orthop Res* 2012;30:2038–45.
- [331] Mao L, Dauchy RT, Blask DE, Slakey LM, Xiang S, Yuan L, et al. Circadian gating of epithelial-to-mesenchymal transition in breast cancer cells via melatonin-regulation of GSK3 β . *Mol Endocrinol* 2012;26:1808–20.
- [332] Kulbe H, Chakravarty P, Leinster DA, Charles KA, Kwong J, Thompson RG, et al. A dynamic inflammatory cytokine network in the human ovarian cancer microenvironment. *Cancer Res* 2012;72:66–75.
- [333] Dayan D, Salo T, Salo S, Nyberg P, Nurmenniemi S, Costea DE, et al. Molecular crosstalk between cancer cells and tumor microenvironment components suggests potential targets for new therapeutic approaches in mobile tongue cancer. *Cancer Med* 2012;1:128–40.
- [334] Karagiannis GS, Poutahidis T, Erdman SE, Kirsch R, Riddell RH, Diamandis EP. Cancer-associated fibroblasts drive the progression of metastasis through both paracrine and mechanical pressure on cancer tissue. *Mol Cancer Res* 2012;10:1403–18.
- [335] Estrella V, Chen T, Lloyd M, Wojtkowiak J, Cornnell HH, Ibrahim-Hashim A, et al. Acidity generated by the tumor microenvironment drives local invasion. *Cancer Res* 2013;73:1524–35.
- [336] Pratheeshkumar P, Kuttan G. Oleoic acid induces apoptosis by modulating p53, Bax, Bcl-2 and caspase-3 gene expression and regulates the activation of transcription factors and cytokine profile in B16F. *J Environ Pathol Toxicol Oncol* 2011;30:21–31.
- [337] Roy S, Deep G, Agarwal C, Agarwal R. Silibinin prevents ultraviolet B radiation-induced epidermal damages in JB6 cells and mouse skin in a p53-GADD45 α -dependent manner. *Carcinogenesis* 2012;33:629–36.
- [338] Thomas P, Wang YJ, Zhong JH, Kosaraju S, O'Callaghan NJ, Zhou XF, et al. Grape seed polyphenols and curcumin reduce genomic instability events in a transgenic mouse model for Alzheimer's disease. *Mutat Res* 2009;661:25–34.
- [339] Marden A, Walmsley RM, Schweizer LM, Schweizer M. Yeast-based assay for the measurement of positive and negative influences on microsatellite stability. *FEMS Yeast Res* 2006;6:716–25.
- [340] Lu LY, Ou N, Lu QB. Antioxidant induces DNA damage, cell death and mutagenicity in human lung and skin normal cells. *Sci Rep* 2013;3:169.
- [341] Liu R, Fu A, Hoffman AE, Zheng T, Zhu Y. Melatonin enhances DNA repair capacity possibly by affecting genes involved in DNA damage responsive pathways. *BMC Cell Biol* 2013;14:1.
- [342] Denissova NG, Nasello CM, Yeung PL, Tischfield JA, Brenneman MA. Resveratrol protects mouse embryonic stem cells from ionizing radiation by accelerating recovery from DNA strand breakage. *Carcinogenesis* 2012;33:149–55.
- [343] Mamede AC, Abrantes AM, Pedrosa L, Casalta-Lopes JE, Pires AS, Teixo RJ, et al. Beyond the limits of oxygen: effects of hypoxia in a hormone-independent prostate cancer cell line. *ISRN Oncol* 2013;2013:918207.
- [344] Wu J, Dauchy RT, Tirrell PC, Wu SS, Lynch DT, Jitawatanarat P, et al. Light at night activates IGF-1R/PDK1 signaling and accelerates tumor growth in human breast cancer xenografts. *Cancer Res* 2011;71:2622–31.
- [345] Elshazley M, Sato M, Hase T, Yamashita R, Yoshida K, Toyokuni S, et al. The circadian clock gene BMAL1 is a novel therapeutic target for malignant pleural mesothelioma. *Int J Cancer* 2012;131:2820–31.
- [346] Ryder M, Gild M, Hohl TM, Pamer E, Knauf J, Ghossein R, et al. Genetic and pharmacological targeting of CSF-1/CSF-1R inhibits tumor-associated macrophages and impairs BRAF-induced thyroid cancer progression. *PLoS ONE* 2013;8:e54302.
- [347] Gershon TR, Crowther AJ, Tikunov A, Garcia I, Annis R, Yuan H, et al. Hexokinase-2-mediated aerobic glycolysis is integral to cerebellar neurogenesis and pathogenesis of medulloblastoma. *Cancer Metab* 2013;1.
- [348] Onodera Y, Nam JM, Bissell MJ. Increased sugar uptake promotes oncogenesis via EPAC/RAP1 and O-GlcNAc pathways. *J Clin Invest* 2014;124:367–84.
- [349] Tyagi A, Singh RP, Ramasamy K, Raina K, Redente EF, Dwyer-Nield LD, et al. Growth inhibition and regression of lung tumors by silibinin: modulation of angiogenesis by macrophage-associated cytokines and nuclear factor-kappaB and signal transducers and activators of transcription 3. *Cancer Prev Res (Phila)* 2009;2:74–83.
- [350] Lin YG, Kunnumakkara AB, Nair A, Merritt WM, Han LY, Armaiz-Pena GN, et al. Curcumin inhibits tumor growth and angiogenesis in ovarian carcinoma by targeting the nuclear factor-kappaB pathway. *Clin Cancer Res* 2007;13:3423–30.
- [351] Moghaddam SJ, Barta P, Mirabolfathinejad SG, Ammar-Aouchiche Z, Garza NT, Vo TT, et al. Curcumin inhibits COPD-like airway inflammation and lung cancer progression in mice. *Carcinogenesis* 2009;30:1949–56.
- [352] Mineva ND, Paulson KE, Naber SP, Yee AS, Sonenshein GE. Epigallocatechin-3-gallate inhibits stem-like inflammatory breast cancer cells. *PLoS ONE* 2013;8:e73464.
- [353] Surh YJ, Chun KS, Cha HH, Han SS, Keum YS, Park KK, et al. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF-kappa B activation. *Mutat Res* 2001;480–481:243–68.
- [354] Corsini E, Dell'Agli M, Facchi A, De Fabiani E, Lucchi L, Boraso MS, et al. Enterodiol and enterolactone modulate the immune response by acting on nuclear factor-kappaB (NF-kappaB) signaling. *J Agric Food Chem* 2010;58:6678–84.
- [355] Jeong SK, Yang K, Park YS, Choi YJ, Oh SJ, Lee CW, et al. Interferon gamma induced by resveratrol analog, HS-1793, reverses the properties of tumor associated macrophages. *Int Immunopharmacol* 2014;22:303–10.
- [356] Salado C, Olaso E, Gallot N, Valcarcel M, Egilegor E, Mendoza L, et al. Resveratrol prevents inflammation-dependent hepatic melanoma metastasis by inhibiting the secretion and effects of interleukin-18. *J Transl Med* 2011;9:59.
- [357] Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, Beecher CW, et al. Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 1997;275:218–20.
- [358] Wang X, Bai H, Zhang X, Liu J, Cao P, Liao N, et al. Inhibitory effect of oleoic acid on hepatocellular carcinoma via ERK-p53-mediated cell cycle arrest and mitochondrial-dependent apoptosis. *Carcinogenesis* 2013;34:1323–30.
- [359] Tae N, Seo J, Min BS, Ryoo S, Lee JH. 3-Oxoolean-12-en-27-oic acid inhibits the proliferation of non-small cell lung carcinoma cells by inducing cell-cycle arrest at G0/G1 phase. *Anticancer Res* 2011;31:2179–85.
- [360] Jiang HL, Jin JZ, Wu D, Xu D, Lin GF, Yu H, et al. Celastrol exerts synergistic effects with PHA-665752 and inhibits tumor growth of c-Met-deficient hepatocellular carcinoma in vivo. *Mol Biol Rep* 2013;40:4203–9.
- [361] Wang WB, Feng LX, Yue QX, Wu WY, Guan SH, Jiang BH, et al. Paraptosis accompanied by autophagy and apoptosis was induced by celastrol, a natural compound with influence on proteasome, ER stress and Hsp90. *J Cell Physiol* 2012;227:2196–206.
- [362] Ge P, Ji X, Ding Y, Wang X, Fu S, Meng F, et al. Celastrol causes apoptosis and cell cycle arrest in rat glioma cells. *Neurol Res* 2010;32:94–100.
- [363] Tyagi A, Agarwal C, Agarwal R. Inhibition of retinoblastoma protein (Rb) phosphorylation at serine sites and an increase in Rb-E2F complex formation by silibinin in androgen-dependent human prostate carcinoma LNCaP cells: role in prostate cancer prevention. *Mol Cancer Ther* 2002;1:525–32.
- [364] Tyagi A, Agarwal C, Harrison G, Glode LM, Agarwal R. Silibinin causes cell cycle arrest and apoptosis in human bladder transitional cell carcinoma cells by regulating CDK1-CDK-cyclin cascade, and caspase 3 and PARP cleavages. *Carcinogenesis* 2004;25:1711–20.
- [365] Kaur M, Velmurugan B, Tyagi A, Deep G, Katiyar S, Agarwal C, et al. Silibinin suppresses growth and induces apoptotic death of human colorectal carcinoma LoVo cells in culture and tumor xenograft. *Mol Cancer Ther* 2009;8:2366–74.
- [366] Agarwal C, Singh RP, Dhanalakshmi S, Tyagi AK, Tecklenburg M, Sclafani RA, et al. Silibinin upregulates the expression of cyclin-dependent kinase inhibitors and causes cell cycle arrest and apoptosis in human colon carcinoma HT-29 cells. *Oncogene* 2003;22:8271–82.
- [367] Kang N, Wang MM, Wang YH, Zhang ZN, Cao HR, Lv YH, et al. Tetrahydrocurcumin induces G2/M cell cycle arrest and apoptosis involving p38 MAPK activation in human breast cancer cells. *Food Chem Toxicol* 2014;67:193–200.
- [368] Lim TG, Lee SY, Huang Z, Lim do Y, Chen H, Jung SK, et al. Curcumin suppresses proliferation of colon cancer cells by targeting CDK2. *Cancer Prev Res (Phila)* 2014;7:466–74.
- [369] Guo H, Xu YM, Ye ZQ, Yu JH, Hu XY. Curcumin induces cell cycle arrest and apoptosis of prostate cancer cells by regulating the expression of IkappaB α , c-Jun and androgen receptor. *Pharmazie* 2013;68:431–4.
- [370] Mukhopadhyay A, Banerjee S, Stafford LJ, Xia C, Liu M, Aggarwal BB. Curcumin-induced suppression of cell proliferation correlates with down-regulation of cyclin D1 expression and CDK4-mediated retinoblastoma protein phosphorylation. *Oncogene* 2002;21:8852–61.
- [371] Ahmad N, Adhmar VM, Gupta S, Cheng P, Mukhtar H. Role of the retinoblastoma (pRb)-E2F/DP pathway in cancer chemopreventive effects of green tea polyphenol epigallocatechin-3-gallate. *Arch Biochem Biophys* 2002;398:125–31.
- [372] Huh SW, Bae SM, Kim YW, Lee JM, Namkoong SE, Lee IP, et al. Anticancer effects of (–)-epigallocatechin-3-gallate on ovarian carcinoma cell lines. *Gynecol Oncol* 2004;94:760–8.
- [373] Saldanha SN, Kala R, Tollefsbol TO. Molecular mechanisms for inhibition of colon cancer cells by combined epigenetic-modulating epigallocatechin gallate and sodium butyrate. *Exp Cell Res* 2014;324:40–53.
- [374] Shen X, Zhang Y, Feng Y, Zhang L, Li J, Xie YA, et al. Epigallocatechin-3-gallate inhibits cell growth, induces apoptosis and causes S phase arrest in hepatocellular carcinoma by suppressing the AKT pathway. *Int J Oncol* 2014;44:791–6.

- [375] Huang WW, Tsai SC, Peng SF, Lin MW, Chiang JH, Chiu YJ, et al. Kaempferol induces autophagy through AMPK and AKT signaling molecules and causes G2/M arrest via downregulation of CDK1/cyclin B in SK-HEP-1 human hepatic cancer cells. *Int J Oncol* 2013;42:2069–77.
- [376] Xu W, Liu J, Li C, Wu HZ, Liu YW. Kaempferol-7-O-beta-D-glucoside (KG) isolated from *Smilax china* L. rhizome induces G2/M phase arrest and apoptosis on HeLa cells in a p53-independent manner. *Cancer Lett* 2008;264:229–40.
- [377] Choi EJ, Ahn WS. Kaempferol induced the apoptosis via cell cycle arrest in human breast cancer MDA-MB-453 cells. *Nutr Res Pract* 2008;2:322–5.
- [378] Casagrande F, Darbon JM. Effects of structurally related flavonoids on cell cycle progression of human melanoma cells: regulation of cyclin-dependent kinases CDK2 and CDK1. *Biochem Pharmacol* 2001;61:1205–15.
- [379] Hill SM, Blask DE, Xiang S, Yuan L, Mao L, Dauchy RT, et al. Melatonin and associated signaling pathways that control normal breast epithelium and breast cancer. *J Mammary Gland Biol Neoplasia* 2011;16:235–45.
- [380] McCann MJ, Gill CI, Linton T, Berrar D, McGlynn H, Rowland IR. Enterolactone restricts the proliferation of the LNCaP human prostate cancer cell line in vitro. *Mol Nutr Food Res* 2008;52:567–80.
- [381] Stan SD, Zeng Y, Singh SV. Ayurvedic medicine constituent withaferin A causes G2 and M phase cell cycle arrest in human breast cancer cells. *Nutr Cancer* 2008;60(Suppl. 1):51–60.
- [382] Roy RV, Suman S, Das TP, Luevano JE, Damodaran C. Withaferin A, a steroidal lactone from *Withania somnifera*, induces mitotic catastrophe and growth arrest in prostate cancer cells. *J Nat Prod* 2013;76:1909–15.
- [383] Grogan PT, Sleder KD, Samadi AK, Zhang H, Timmermann BN, Cohen MS. Cytotoxicity of withaferin A in glioblastoma involves induction of an oxidative stress-mediated heat shock response while altering Akt/mTOR and MAPK signaling pathways. *Invest New Drugs* 2013;31:545–57.
- [384] Adhami VM, Afaq F, Ahmad N. Involvement of the retinoblastoma (pRb)-E2F/DP pathway during antiproliferative effects of resveratrol in human epidermoid carcinoma (A431) cells. *Biochem Biophys Res Commun* 2001;288:579–85.
- [385] Wolter F, Akoglu B, Clausnitzer A, Stein J. Downregulation of the cyclin D1/Cdk4 complex occurs during resveratrol-induced cell cycle arrest in colon cancer cell lines. *J Nutr* 2001;131:2197–203.
- [386] Kim AL, Zhu Y, Zhu H, Han L, Kopelovich L, Bickers DR, et al. Resveratrol inhibits proliferation of human epidermoid carcinoma A431 cells by modulating MEK1 and AP-1 signalling pathways. *Exp Dermatol* 2006;15:538–46.
- [387] Yang Q, Wang B, Zang W, Wang X, Liu Z, Li W, et al. Resveratrol inhibits the growth of gastric cancer by inducing G1 phase arrest and senescence in a Sirt1-dependent manner. *PLOS ONE* 2013;8:e70627.
- [388] Tang S, Gao D, Zhao T, Zhou J, Zhao X. An evaluation of the anti-tumor efficacy of oleanolic acid-loaded PEGylated liposomes. *Nanotechnology* 2013;24:235102.
- [389] Li W, Ding Y, Sun YN, Yan XT, Yang SY, Choi CW, et al. Oleanane-type triterpenoid saponins from the roots of *Pulsatilla koreana* and their apoptosis-inducing effects on HL-60 human promyelocytic leukemia cells. *Arch Pharm Res* 2013;36:768–74.
- [390] Dizaji MZ, Malehmir M, Ghavamzadeh A, Alimoghaddam K, Ghaffari SH. Synergistic effects of arsenic trioxide and silibinin on apoptosis and invasion in human glioblastoma U87MG cell line. *Neurochem Res* 2012;37:370–80.
- [391] Fang HY, Chen SB, Guo DJ, Pan SY, Yu ZL. Proteomic identification of differentially expressed proteins in curcumin-treated MCF-7 cells. *Phytomedicine* 2011;18:697–703.
- [392] Lee WY, Chen YC, Shih CM, Lin CM, Cheng CH, Chen KC, et al. The induction of heme oxygenase-1 suppresses heat shock protein 90 and the proliferation of human breast cancer cells through its byproduct carbon monoxide. *Toxicol Appl Pharmacol* 2014;274:55–62.
- [393] Martin S, Lamb HK, Brady C, Lefkove B, Bonner MY, Thompson P, et al. Inducing apoptosis of cancer cells using small-molecule plant compounds that bind to GRP78. *Br J Cancer* 2013;109:433–43.
- [394] Song W, Dang Q, Xu D, Chen Y, Zhu G, Wu K, et al. Kaempferol induces cell cycle arrest and apoptosis in renal cell carcinoma through EGFR/p38 signaling. *Oncol Rep* 2014;31:1350–6.
- [395] Kim CH, Yoo YM. Melatonin induces apoptotic cell death via p53 in LNCaP cells. *Korean J Physiol Pharmacol* 2010;14:365–9.
- [396] Danbara N, Yuri T, Tsujita-Kyutoku M, Tsukamoto R, Uehara N, Tsubura A. Enterolactone induces apoptosis and inhibits growth of Colo 201 human colon cancer cells both in vitro and in vivo. *Anticancer Res* 2005;25:2269–76.
- [397] Hahm ER, Lee J, Singh SV. Role of mitogen-activated protein kinases and Mcl-1 in apoptosis induction by withaferin A in human breast cancer cells. *Mol Carcinog* 2013;53(11):907–16.
- [398] Wesolowska O, Wisniewski J, Bielawska-Pohl A, Paprocka M, Duarte N, Ferreira MJ, et al. Stilbenes as multidrug resistance modulators and apoptosis inducers in human adenocarcinoma cells. *Anticancer Res* 2010;30:4587–93.
- [399] Deeb D, Gao X, Liu Y, Kim SH, Pindolia KR, Arbab AS, et al. Inhibition of cell proliferation and induction of apoptosis by oleanane triterpenoid (CDDO-Me) in pancreatic cancer cells is associated with the suppression of hTERT gene expression and its telomerase activity. *Biochem Biophys Res Commun* 2012;422:561–7.
- [400] Deeb D, Gao X, Liu Y, Varma NR, Arbab AS, Gautam SC. Inhibition of telomerase activity by oleanane triterpenoid CDDO-Me in pancreatic cancer cells is ROS-dependent. *Molecules* 2013;18:3250–65.
- [401] Nasiri M, Zarghami N, Koshki KN, Mollazadeh M, Moghaddam MP, Yamchi MR, et al. Curcumin and silibinin inhibit telomerase expression in T47D human breast cancer cells. *Asian Pac J Cancer Prev* 2013;14:3449–53.
- [402] Ebrahimnezhad Z, Zarghami N, Keyhani M, Amirsaadat S, Akbarzadeh A, Rahmati M, et al. Inhibition of hTERT gene expression by silibinin-loaded PLGA-PEG-Fe₃O₄ in T47D breast cancer cell line. *Bioimpacts* 2013;3:67–74.
- [403] Thelen P, Wuttke W, Jarry H, Grzmil M, Ringert RH. Inhibition of telomerase activity and secretion of prostate specific antigen by silibinin in prostate cancer cells. *J Urol* 2004;171:1934–8.
- [404] Khaw AK, Hande MP, Kalthur G. Curcumin inhibits telomerase and induces telomere shortening and apoptosis in brain tumour cells. *J Cell Biochem* 2013;114:1257–70.
- [405] Wang X, Hao MW, Dong K, Lin F, Ren JH, Zhang HZ. Apoptosis induction effects of EGCG in laryngeal squamous cell carcinoma cells through telomerase repression. *Arch Pharm Res* 2009;32:1263–9.
- [406] Sadava D, Whitlock E, Kane SE. The green tea polyphenol, epigallocatechin-3-gallate inhibits telomerase and induces apoptosis in drug-resistant lung cancer cells. *Biochem Biophys Res Commun* 2007;360:233–7.
- [407] Berletch JB, Liu C, Love WK, Andrews LG, Katiyar SK, Tollefsbol TO. Epigenetic and genetic mechanisms contribute to telomerase inhibition by EGCG. *J Cell Biochem* 2008;103:509–19.
- [408] Martinez-Campa CM, Alonso-Gonzalez C, Mediavilla MD, Cos S, Gonzalez A, Sanchez-Barcelo EJ. Melatonin down-regulates hTERT expression induced by either natural estrogens (17beta-estradiol) or metalloestrogens (cadmium) in MCF-7 human breast cancer cells. *Cancer Lett* 2008;268:272–7.
- [409] Lanzilli G, Fuggetta MP, Tricarico M, Cottarelli A, Serafino A, Falchetti R, et al. Resveratrol down-regulates the growth and telomerase activity of breast cancer cells in vitro. *Int J Oncol* 2006;28:641–8.
- [410] Liu J, Wu N, Ma L, Liu M, Liu G, Zhang Y, et al. Oleanolic acid suppresses aerobic glycolysis in cancer cells by switching pyruvate kinase type M isoforms. *PLOS ONE* 2014;9:e91606.
- [411] Raina K, Agarwal C, Wadhwa R, Serkova NJ, Agarwal R. Energy deprivation by silibinin in colorectal cancer cells: a double-edged sword targeting both apoptotic and autophagic machineries. *Autophagy* 2013;9:697–713.
- [412] Yousefi M, Ghaffari SH, Soltani BM, Nafissi S, Momeny M, Zekri A, et al. Therapeutic efficacy of silibinin on human neuroblastoma cells: Akt and NF-kappaB expressions may play an important role in silibinin-induced response. *Neurochem Res* 2012;37:2053–63.
- [413] Vaughan RA, Garcia-Smith R, Dorsey J, Griffith JK, Bisoffi M, Trujillo KA. Tumor necrosis factor alpha induces Warburg-like metabolism and is reversed by anti-inflammatory curcumin in breast epithelial cells. *Int J Cancer* 2013;133:2504–10.
- [414] Moreira L, Araujo J, Costa T, Correia-Branco A, Faria A, Martel F, et al. Quercetin and epigallocatechin gallate inhibit glucose uptake and metabolism by breast cancer cells by an estrogen receptor-independent mechanism. *Exp Cell Res* 2013;319:1784–95.
- [415] Valenti D, de Bari L, Manente GA, Rossi L, Mutti L, Moro L, et al. Negative modulation of mitochondrial oxidative phosphorylation by epigallocatechin-3 gallate leads to growth arrest and apoptosis in human malignant pleural mesothelioma cells. *Biochim Biophys Acta* 2013;1832:2085–96.
- [416] Huang CH, Tsai SJ, Wang YJ, Pan MH, Kao JY, Way TD. EGCG inhibits protein synthesis, lipogenesis, and cell cycle progression through activation of AMPK in p53 positive and negative human hepatoma cells. *Mol Nutr Food Res* 2009;53:1156–65.
- [417] Filomeni G, Desideri E, Cardaci S, Graziani I, Piccirillo S, Rotilio G, et al. Carcinoma cells activate AMP-activated protein kinase-dependent autophagy as survival response to kaempferol-mediated energetic impairment. *Autophagy* 2010;6:202–16.
- [418] Thors L, Belghiti M, Fowler CJ. Inhibition of fatty acid amide hydrolase by kaempferol and related naturally occurring flavonoids. *Br J Pharmacol* 2008;155:244–52.
- [419] Wang BQ, Yang QH, Xu RK, Xu JN. Elevated levels of mitochondrial respiratory complexes activities and ATP production in 17-beta-estradiol-induced prolactin-secretory tumor cells in male rats are inhibited by melatonin in vivo and in vitro. *Chin Med J (Engl)* 2013;126:4724–30.
- [420] Sanchez-Sanchez AM, Martin V, Garcia-Santos G, Rodriguez-Blanco J, Casado-Zapico S, Suarez-Garnacho S, et al. Intracellular redox state as determinant for melatonin antiproliferative vs cytotoxic effects in cancer cells. *Free Radic Res* 2011;45:1333–41.
- [421] Gonzalez A, del Castillo-Vaquero A, Miro-Moran A, Tapia JA, Salido GM. Melatonin reduces pancreatic tumor cell viability by altering mitochondrial physiology. *J Pineal Res* 2011;50:250–60.
- [422] Hahm ER, Lee J, Kim SH, Sehwat A, Arlotti JA, Shiva SS, et al. Metabolic alterations in mammary cancer prevention by withaferin A in a clinically relevant mouse model. *J Natl Cancer Inst* 2013;105:1111–22.
- [423] Hahm ER, Moura MB, Kelley EE, Van Houten B, Shiva S, Singh SV. Withaferin A-induced apoptosis in human breast cancer cells is mediated by reactive oxygen species. *PLoS ONE* 2011;6:e23354.
- [424] Malik F, Kumar A, Bhushan S, Khan S, Bhatia A, Suri KA, et al. Reactive oxygen species generation and mitochondrial dysfunction in the apoptotic cell death of human myeloid leukemia HL-60 cells by a dietary compound withaferin A with concomitant protection by N-acetyl cysteine. *Apoptosis* 2007;12:2115–33.
- [425] Gomez LS, Zancan P, Marcondes MC, Ramos-Santos L, Meyer-Fernandes JR, Sola-Penna M, et al. Resveratrol decreases breast cancer cell viability and glucose metabolism by inhibiting 6-phosphofructo-1-kinase. *Biochimie* 2013;95:1336–43.

- [426] Fouad MA, Agha AM, Merzabani MM, Shouman SA. Resveratrol inhibits proliferation, angiogenesis and induces apoptosis in colon cancer cells: calorie restriction is the force to the cytotoxicity. *Hum Exp Toxicol* 2013;32:1067–80.
- [427] Filomeni G, Graziani I, Rotilio G, Ciriolo MR. *trans*-Resveratrol induces apoptosis in human breast cancer cells MCF-7 by the activation of MAP kinases pathways. *Genes Nutr* 2007;2:295–305.
- [428] Nagaraj S, Youn JI, Weber H, Iclozan C, Lu L, Cotter MJ, et al. Anti-inflammatory triterpenoid blocks immune suppressive function of MDSCs and improves immune response in cancer. *Clin Cancer Res* 2010;16:1812–23.
- [429] Deng W, Sun HX, Chen FY, Yao ML. Immunomodulatory activity of 3beta, 6beta-dihydroxyolean-12-en-27-oic acid in tumor-bearing mice. *Chem Biodivers* 2009;6:1243–53.
- [430] Forghani P, Khorramizadeh MR, Waller EK. Silibinin inhibits accumulation of myeloid-derived suppressor cells and tumor growth of murine breast cancer. *Cancer Med* 2014;3:215–24.
- [431] Bhattacharyya S, Md Sakib Hossain D, Mohanty S, Sankar Sen G, Chattopadhyay S, Banerjee S, et al. Curcumin reverses T cell-mediated adaptive immune dysfunctions in tumor-bearing hosts. *Cell Mol Immunol* 2010;7:306–15.
- [432] Meeran SM, Mantena SK, Katiyar SK. Prevention of ultraviolet radiation-induced immunosuppression by (–)-epigallocatechin-3-gallate in mice is mediated through interleukin 12-dependent DNA repair. *Clin Cancer Res* 2006;12:2272–80.
- [433] Jang JY, Lee JK, Jeon YK, Kim CW. Exosome derived from epigallocatechin gallate treated breast cancer cells suppresses tumor growth by inhibiting tumor-associated macrophage infiltration and M2 polarization. *BMC Cancer* 2013;13:421.
- [434] Lissoni P, Ardizzoia A, Barni S, Tancini G, Muttini M. Immunotherapy with subcutaneous low dose interleukin-2 plus melatonin as salvage therapy of heavily chemotherapy-pretreated ovarian cancer. *Oncol Rep* 1996;3:947–9.
- [435] Sinha P, Ostrand-Rosenberg S. Myeloid-derived suppressor cell function is reduced by Withaferin A, a potent and abundant component of *Withania somnifera* root extract. *Cancer Immunol Immunother* 2013;62:1663–73.
- [436] Hushmendy S, Jayakumar L, Hahn AB, Bhoiwala D, Bhoiwala DL, Crawford DR. Select phytochemicals suppress human T-lymphocytes and mouse splenocytes suggesting their use in autoimmunity and transplantation. *Nutr Res* 2009;29:568–78.
- [437] Wang B, Sun J, Li X, Zhou Q, Bai J, Shi Y, et al. Resveratrol prevents suppression of regulatory T-cell production, oxidative stress, and inflammation of mice prone or resistant to high-fat diet-induced obesity. *Nutr Res* 2013;33:971–81.
- [438] Yang Y, Paik JH, Cho D, Cho JA, Kim CW. Resveratrol induces the suppression of tumor-derived CD4+CD25+ regulatory T cells. *Int Immunopharmacol* 2008;8:542–7.
- [439] Deeb D, Gao X, Liu Y, Jiang D, Divine GW, Arbab AS, et al. Synthetic triterpenoid CDDO prevents the progression and metastasis of prostate cancer in TRAMP mice by inhibiting survival signaling. *Carcinogenesis* 2011;32:757–64.
- [440] Guo G, Yao W, Zhang Q, Bo Y. Oleanolic acid suppresses migration and invasion of malignant glioma cells by inactivating MAPK/ERK signaling pathway. *PLOS ONE* 2013;8:e72079.
- [441] Huang L, Zhang Z, Zhang S, Ren J, Zhang R, Zeng H, et al. Inhibitory action of Celastrol on hypoxia-mediated angiogenesis and metastasis via the HIF-1alpha pathway. *Int J Mol Med* 2011;27:407–15.
- [442] Singh RP, Raina K, Sharma G, Agarwal R. Silibinin inhibits established prostate tumor growth, progression, invasion, and metastasis and suppresses tumor angiogenesis and epithelial–mesenchymal transition in transgenic adenocarcinoma of the mouse prostate model mice. *Clin Cancer Res* 2008;14:7773–80.
- [443] Yang CL, Liu YY, Ma YG, Xue YX, Liu DG, Ren Y, et al. Curcumin blocks small cell lung cancer cells migration, invasion, angiogenesis, cell cycle and neoplasia through Janus kinase-STAT3 signalling pathway. *PLoS ONE* 2012;7:e37960.
- [444] Maruyama T, Murata S, Nakayama K, Sano N, Ogawa K, Nowatari T, et al. (–)-Epigallocatechin-3-gallate suppresses liver metastasis of human colorectal cancer. *Oncol Rep* 2014;31:625–33.
- [445] Sharma C, Nusri Qel A, Begum S, Javed E, Rizvi TA, Hussain A. (–)-Epigallocatechin-3-gallate induces apoptosis and inhibits invasion and migration of human cervical cancer cells. *Asian Pac J Cancer Prev* 2012;13:4815–22.
- [446] Park JH, Yoon JH, Kim SA, Ahn SG. (–)-Epigallocatechin-3-gallate inhibits invasion and migration of salivary gland adenocarcinoma cells. *Oncol Rep* 2010;23:585–90.
- [447] Astin JW, Jamieson SM, Eng TC, Flores MV, Misa JP, Chien A, et al. An in vivo antilymphatic screen in zebrafish identifies novel inhibitors of mammalian lymphangiogenesis and lymphatic-mediated metastasis. *Mol Cancer Ther* 2014;13:2450–62.
- [448] Chen HJ, Lin CM, Lee CY, Shih NC, Peng SF, Tsuzuki M, et al. Kaempferol suppresses cell metastasis via inhibition of the ERK-p38-JNK and AP-1 signaling pathways in U-2 OS human osteosarcoma cells. *Oncol Rep* 2013;30:925–32.
- [449] Rapozzi V, Perissin L, Zorzet S, Giraldo T. Effects of melatonin administration on tumor spread in mice bearing Lewis lung carcinoma. *Pharmacol Res* 1992;25(Suppl. 1):71–2.
- [450] Cos S, Fernandez R, Guezmes A, Sanchez-Barcelo EJ. Influence of melatonin on invasive and metastatic properties of MCF-7 human breast cancer cells. *Cancer Res* 1998;58:4383–90.
- [451] Miura D, Saarninen NM, Miura Y, Santti R, Yagasaki K. Hydroxymatairesinol and its mammalian metabolite enterolactone reduce the growth and metastasis of subcutaneous AH109A hepatomas in rats. *Nutr Cancer* 2007;58:49–59.
- [452] Thaiparambil JT, Bender L, Ganesh T, Kline E, Patel P, Liu Y, et al. Withaferin A inhibits breast cancer invasion and metastasis at sub-cytotoxic doses by inducing vimentin disassembly and serine 56 phosphorylation. *Int J Cancer* 2011;129:2744–55.
- [453] Ji Q, Liu X, Fu X, Zhang L, Sui H, Zhou L, et al. Resveratrol inhibits invasion and metastasis of colorectal cancer cells via MALAT1 mediated Wnt/beta-catenin signal pathway. *PLOS ONE* 2013;8:e78700.
- [454] Sheth S, Jajoo S, Kaur T, Mukherjee D, Sheehan K, Rybak LP, et al. Resveratrol reduces prostate cancer growth and metastasis by inhibiting the Akt/MicroRNA-21 pathway. *PLoS ONE* 2012;7:e51655.
- [455] Yeh CB, Hsieh MJ, Lin CW, Chiou HL, Lin PY, Chen TY, et al. The antimetastatic effects of resveratrol on hepatocellular carcinoma through the downregulation of a metastasis-associated protease by SP-1 modulation. *PLOS ONE* 2013;8:e56661.
- [456] Wu PK, Chi Shing Tai W, Liang ZT, Zhao ZZ, Hsiao WL. Oleanolic acid isolated from *Oldenlandia diffusa* exhibits a unique growth inhibitory effect against ras-transformed fibroblasts. *Life Sci* 2009;85:113–21.
- [457] Ni H, Zhao W, Kong X, Li H, Ouyang J. Celastrol inhibits lipopolysaccharide-induced angiogenesis by suppressing TLR4-triggered nuclear factor-kappa B activation. *Acta Haematol* 2014;131:102–11.
- [458] Xu P, Yin Q, Shen J, Chen L, Yu H, Zhang Z, et al. Synergistic inhibition of breast cancer metastasis by silibinin-loaded lipid nanoparticles containing TPGS. *Int J Pharm* 2013;454:21–30.
- [459] Xu Y, Zhang J, Han J, Pan X, Cao Y, Guo H, et al. Curcumin inhibits tumor proliferation induced by neutrophil elastase through the upregulation of alpha1-antitrypsin in lung cancer. *Mol Oncol* 2012;6:405–17.
- [460] Zgheib A, Lamy S, Annabi B. Epigallocatechin gallate targeting of membrane type 1 matrix metalloproteinase-mediated Src and Janus kinase/signal transducers and activators of transcription 3 signaling inhibits transcription of colony-stimulating factors 2 and 3 in mesenchymal stromal cells. *J Biol Chem* 2013;288:13378–86.
- [461] Luo H, Jiang BH, King SM, Chen YC. Inhibition of cell growth and VEGF expression in ovarian cancer cells by flavonoids. *Nutr Cancer* 2008;60:800–9.
- [462] Bandyopadhyay S, Romero JR, Chattopadhyay N. Kaempferol and quercetin stimulate granulocyte-macrophage colony-stimulating factor secretion in human prostate cancer cells. *Mol Cell Endocrinol* 2008;287:57–64.
- [463] Alvarez-Garcia V, Gonzalez A, Alonso-Gonzalez C, Martinez-Campa C, Cos S. Antiangiogenic effects of melatonin in endothelial cell cultures. *Microvasc Res* 2013;87:25–33.
- [464] Szarc vel Szic K, Op de Beeck K, Ratman D, Wouters A, Beck IM, Declerck K, et al. Pharmacological levels of Withaferin A (*Withania somnifera*) trigger clinically relevant anticancer effects specific to triple negative breast cancer cells. *PLOS ONE* 2014;9:e87850.
- [465] Shamim U, Hanif S, Albanyan A, Beck FW, Bao B, Wang Z, et al. Resveratrol-induced apoptosis is enhanced in low pH environments associated with cancer. *J Cell Physiol* 2012;227:1493–500.