



DIGITAL ACCESS TO
SCHOLARSHIP AT HARVARD
DASH.HARVARD.EDU



HARVARD LIBRARY
Office for Scholarly Communication

A Longitudinal Study of Peripubertal Serum Organochlorine Concentrations and Semen Parameters in Young Men: The Russian Children's Study

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Mínguez-Alarcón, Lidia, Oleg Sergeyeve, Jane S. Burns, Paige L. Williams, Mary M. Lee, Susan A. Korrick, Luidmila Smigulina, Boris Revich, and Russ Hauser. 2016. "A Longitudinal Study of Peripubertal Serum Organochlorine Concentrations and Semen Parameters in Young Men: The Russian Children's Study." <i>Environmental Health Perspectives</i> 125 (3): 460-466. doi:10.1289/EHP25. http://dx.doi.org/10.1289/EHP25 .
Published Version	doi:10.1289/EHP25
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:32072239
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

A Longitudinal Study of Peripubertal Serum Organochlorine Concentrations and Semen Parameters in Young Men: The Russian Children's Study

Lidia Mínguez-Alarcón,¹ Oleg Sergeev,^{2,3} Jane S. Burns,¹ Paige L. Williams,^{4,5} Mary M. Lee,⁶ Susan A. Korrick,^{1,7} Luidmila Smigulina,³ Boris Revich,⁸ and Russ Hauser^{1,5}

¹Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; ²Department of Genomics and Human Genetics Vavilov Institute of General Genetics, Russian Academy of Sciences, Moscow, Russia; ³Chapaevsk Medical Association, Chapaevsk, Samara Region, Russia; ⁴Department of Biostatistics, and ⁵Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; ⁶Pediatric Endocrine Division, Departments of Pediatrics and Cell & Developmental Biology, University of Massachusetts Medical School, Worcester, Massachusetts, USA; ⁷Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; ⁸Institute for Forecasting, Russian Academy of Sciences, Moscow, Russia

BACKGROUND: Exposures to endocrine-disrupting chemicals during critical phases of testicular development may be related to poorer semen parameters. However, few studies have assessed the association between childhood organochlorine (OC) exposure and adult semen parameters.

OBJECTIVE: We examined whether peripubertal serum OC concentrations are associated with semen parameters among young Russian men.

METHODS: From 2003 through 2005, 516 boys were enrolled at age 8–9 years and followed for up to 10 years. Serum OCs were measured in the enrollment samples using high-resolution mass spectrometry. At 18–19 years, 133 young men provided 1 or 2 semen samples (256 samples) collected approximately 1 week apart, which were analyzed for volume, sperm concentration, and motility. Unadjusted and adjusted linear mixed models were used to examine the associations of quartiles of lipid-standardized concentrations of dioxins [2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), polychlorinated dibenzo-*p*-dioxins (PCDDs)], furans, polychlorinated biphenyls (PCBs), and corresponding toxic equivalents (TEQs) with semen parameters.

RESULTS: The median (range) for TCDD was 2.9 (0.4–12.1) pg/g lipid and PCDD TEQ was 8.7 (1.0–36.0) pg TEQ/g lipid. Higher quartiles of TCDD and PCDD TEQs were associated with lower sperm concentration, total sperm count, and total motile sperm count (*p*-trends ≤ 0.05). The highest quartile of peripubertal serum TCDD concentrations was associated with a decrease (95% CI) of 40% (18, 66%), 29% (3, 64%), and 30% (2, 70%) in sperm concentration, total sperm count, and total motile sperm count, respectively, compared with the lowest quartile. Similar associations were observed for serum PCDD TEQs with semen parameters. Serum PCBs, furans, and total TEQs were not associated with semen parameters.

CONCLUSION: Higher peripubertal serum TCDD concentrations and PCDD TEQs were associated with poorer semen parameters.

CITATION: Mínguez-Alarcón L, Sergeev O, Burns JS, Williams PL, Lee MM, Korrick SA, Smigulina L, Revich B, Hauser R. 2017. A longitudinal study of peripubertal serum organochlorine concentrations and semen parameters in young men: the Russian Children's Study. *Environ Health Perspect* 125:460–466; <http://dx.doi.org/10.1289/EHP25>

Introduction

Over the past several decades, numerous studies have explored whether semen parameters have declined (Carlsen et al. 1992; Swan et al. 2000), and whether there are geographical differences in semen parameters both between (Jørgensen et al. 2001, 2002) and within countries (Swan et al. 2003). Recent literature has shown that serum concentrations of organochlorines (OCs), including dioxins, furans, and polychlorinated biphenyls (PCBs), are associated with decreased semen parameters (Faure et al. 2014; Meeker and Hauser 2010; Mocarelli et al. 2008, 2011; Paoli et al. 2015; Toft et al. 2006). Despite efforts to limit dioxin emissions and longstanding bans on PCB manufacture and use, there is still ongoing exposure through diet because these compounds bioconcentrate in the food chain due to their lipophilic properties and long half-lives (Schecter et al. 2001).

Among epidemiologic studies on OCs and semen parameters, the only one that explored childhood exposure and adult semen parameters was in Seveso, Italy, where an explosion in 1976 at a trichlorophenol manufacturing plant released up to 30 kg of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) (Mocarelli et al. 2008). The authors investigated the relationship of serum TCDD concentrations measured from blood samples taken in 1976 during childhood (1–9 years), puberty (10–17 years), or young adult life (18–26 years) with semen parameters and male reproductive hormones measured 22 years later. They did not measure other dioxins, furans or PCBs. Mocarelli and colleagues found that acute high exposure to TCDD in childhood (1–9 years), but not during puberty (10–17 years) or adulthood (18–26 years), was associated with poorer semen parameters later in adulthood. These compelling results were key data in

the U.S. Environmental Protection Agency (EPA) risk assessment for dioxins (U.S. EPA 2009). These results suggested that during childhood, when the testes are still immature, the activation of aryl hydrocarbon receptors (AhR) in the testes by TCDD may interfere with maturation of the seminiferous tubules and spermatogenesis and demonstrates that the juvenile reproductive system may be particularly vulnerable to TCDD exposure (Woodruff et al. 2010).

Given the importance of childhood exposures on reproductive health later in life, we conducted a prospective cohort study of Russian boys with a wide range of exposure to dioxins, furans, and PCBs stemming from environmental contamination of their community. Specifically we assessed the

Address correspondence to L. Mínguez-Alarcón, Department of Environmental Health, Harvard T.H. Chan School of Public Health, 665 Huntington Ave., Boston, MA 02115 USA. Telephone: (617) 432-4572. E-mail: lminguez@hsph.harvard.edu

Supplemental Material is available online (<http://dx.doi.org/10.1289/EHP25>).

The authors gratefully acknowledge all the children and adults involved in this study. We also acknowledge the Chapaevsk government (D. Blynsky and N. Malakhov), and the Chapaevsk Medical Association and Chapaevsk Central Hospital staff. We also thank our colleagues D.G. Patterson Jr. and W.E. Turner who were formerly at the Centers for Disease Control and Prevention (Atlanta, Georgia, USA) for their analysis of our biospecimens for organochlorine concentrations.

Funding was provided through grants R01ES0014370 and P30ES000002 from the National Institutes of Health/National Institute of Environmental Health Sciences, grant R82943701 from the U.S. Environmental Protection Agency, and grant 14-45-00065 from the Russian Science Foundation (O.S. and R.H.).

The authors declare they have no actual or potential competing financial interests.

Received: 24 February 2016; Revised: 25 July 2016; Accepted: 20 August 2016; Published: 7 October 2016.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

associations of peripubertal (measured at age 8–9 years) serum concentrations of dioxins, furans, and PCBs with semen parameters in young healthy men measured approximately 10 years later.

Methods

Study Population

The Russian Children's Study is an ongoing prospective study of 516 males (Hauser et al. 2008; Williams et al. 2010). Once enrolled at age 8–9 years, each boy underwent a physical examination, provided a blood sample for OC measurement, and together with his mother or guardian, completed health, lifestyle, and dietary questionnaires. Annual follow up examinations were conducted and questionnaires were completed. Of the original cohort of 516 boys, 124 (24%) were lost to follow-up by their 10th annual follow-up visit at age 18–19 years, 59 (11%) were too young for semen collection, 49 (15%) declined to participate in the semen study, 144 (28%) were pending (did not respond yet to invitation, temporarily relocated, or not yet sexually mature based on Tanner Stages and testicular volume), 4 had missing OC data, and 3 were excluded due to chronic disease. At ages 18–19 years, 133 young men who had serum OC concentrations measured at age 8–9 years and provided 1 or 2 semen samples collected approximately 1 week apart (256 samples) were included in this analysis (Figure 1).

The study was approved by the Human Studies Institutional Review Boards of the Chapaevsk Medical Association (Chapaevsk, Russia); Harvard T.H. Chan School of Public Health, Brigham and Women's Hospital (Boston, MA, USA), and University of Massachusetts Medical School (Worcester, MA, USA). At enrollment, the parent or guardian signed an informed consent, and each boy signed an assent before participation. At ≥ 18 years of age, the young man signed a consent form before providing the two semen samples.

Semen Parameters Assessment

The subjects' self-reported information about abstinence period, fever, and any illnesses within the previous month was collected before semen sampling. Semen samples were provided by masturbation in a study room near the Andrology Laboratory and kept at 37°C in an incubator until semen evaluation, which began within 1 hr after ejaculation (analysis for 88% of the samples began within 30 min). One hundred twenty-three men (92%) provided two semen samples collected approximately 1 week apart, and 10 men (8%) provided one semen sample. The actual abstinence period was calculated from the date and time of previous ejaculation and

the date and time of delivery of semen sample recorded by a technician.

Semen analysis was performed at the Andrology Laboratory according to the criteria recently updated (Björndahl et al. 2010) by the Nordic Association for Andrology (NAFA) and European Society of Human Reproduction and Embryology–Special Interest Group in Andrology (ESHRE-SIGA) (Kvist and Björndahl 2002). All samples were assessed by one technician (L.S.) who was blinded to the serum OC concentration. Semen volume was measured using a 1-, 5-, or 10-mL disposable pipette. For sperm motility assessment, 10 μ L of well-mixed semen was placed on a clean glass slide kept at 37°C and covered with a 22 \times 22 mm coverslip. The slide was placed on the heated stage of a microscope at 37°C and immediately examined at 400 \times magnification in duplicate. At least 200 sperm per slide were classified as the four World Health Organization (WHO) classes: rapidly progressive motile (class A), slowly progressive motile (class B), locally motile (class C) or immotile (class D), taking the average value for duplicate measures (WHO 1999). Percent motile sperm was defined as the sum of WHO classes A, B, and C. Sperm concentration was measured using an Improved Neubauer Chamber Hemacytometer viewed at phase contrast (200 \times magnification).

Organochlorine Exposure Assessment

Fasting blood samples were collected at the initial visit (when boys were 8–9 years old), and the serum fraction was stored at -35°C until shipment for analysis at the National Center for Environmental Health at the

Centers for Disease Control and Prevention (CDC; Atlanta, GA, USA). Analytes included 7 polychlorinated dibenzo-*p*-dioxins (PCDDs, or dioxins), 10 polychlorinated dibenzofurans (PCDFs, or furans), 4 co-planar PCBs (co-PCBs), 6 mono-*ortho*-substituted PCBs, and 31 other PCBs (non-dioxin-like PCBs) (Burns et al. 2009).

For dioxin-like analytes, sera, method blanks, and quality control samples (aliquots of pooled bovine sera) were spiked with a mixture of $^{13}\text{C}_{12}$ -labeled PCDDs/PCDFs and co-PCBs as internal standards, and serum analytes were isolated by solid phase extraction (SPE) followed by a multicolumn automated cleanup and enrichment procedure (Turner et al. 1997). Analytes were separated on a DB-5 MS capillary column (Phenomenex, Torrance, CA, USA) and quantified using selected-ion-monitoring (SIM) high-resolution (10,000 resolving power) mass spectrometry (HRGC-ID/HRMS; Thermo Electron North America, LLC, West Palm Beach, FL, USA) (Paterson et al. 1987). Quantification was by isotope dilution MS using calibration standards containing $^{13}\text{C}_{12}$ -labeled and unlabeled analytes. A similar approach was used for mono-*ortho* and non-dioxin-like PCBs (Barr et al. 2003). Samples were spiked with $^{13}\text{C}_{12}$ -labeled PCBs, extracted by either large (Turner et al. 1997) or small (Sjödin et al. 2004) volume SPE, and analyzed using HR GC/MS in SIM (Barr et al. 2003).

For all analyses, quality control sample coefficients of variation combining between-run and within-run reproducibility were generally $< 15\%$. All concentrations were expressed on a per-lipid basis, with serum total cholesterol and triglycerides measured enzymatically, and total lipids were calculated




Figure 1. Flow diagram of the Russian Children's Study.

Note: Information on BMI, smoking, and alcohol consumption was collected at the same visit year as the semen collection for 84 (63%) men, and within 3 years before semen collection for 49 (37%) men.

using the Phillips equation (Phillips et al. 1989). Congener concentrations below the limit of detection (LOD) were assigned the sample-specific LOD divided by the square root of 2 (Baccarelli et al. 2005).

Statistical Analysis

Dioxin toxic equivalents (TEQs) were calculated on a lipid basis using the 2005 WHO toxic equivalency factors to weigh the potency of each congener relative to TCDD before summation (Van den Berg et al. 2006). Although our *a priori* hypothesis focused on TCDD, we also explored the association of eight additional exposure metrics with semen parameters. These included 1) total (summed) TEQ measures (pg TEQ/g lipid) for combined dioxin, furan, co-planar PCB, and mono-ortho PCB congeners; 2–4) total (summed) TEQs (pg TEQ/g lipid) for each of the dioxins, furans, and co-PCBs; 5–7) total (summed) concentrations (pg/g lipid) for each of the dioxins (ΣPCDD), furans (ΣPCDF), and co-PCBs (ΣCo-PCB); and 8) total (summed) concentrations of non-dioxin-like PCBs, including mono-ortho-substituted PCBs (ΣPCBs) (ng/g lipid). OC measures were categorized into quartiles because of potential nonlinear associations.

We first summarized participant characteristics using medians and interquartile ranges (IQR) for continuous variables, and number and percentages for categorical variables. Linear mixed models were used to examine the relation between OC exposure

Table 1. Demographic characteristics and semen parameters of 133 young men contributing 256 semen samples in the Russian Children’s Study.

Characteristic	Median (IQR) or n (%)
Demographic characteristics^a	
Age (years)	18.3 (18.1–18.7)
Body mass index (kg/m ²)	21.0 (19.2–23.2)
Smoking status ^b	68 (51)
Alcohol consumption ^c	90 (68)
Semen parameters^d	
Volume (mL)	2.4 (1.8–3.5)
Sperm concentration (million/mL)	51.3 (26.6–78.8)
Total sperm count (million)	127 (61.0–222.0)
Sperm motility (A + B + C) ^e (%)	64.0 (57.0–68.0)
Total motile sperm count (million)	80.5 (35.8–141.0)
Abstinence time (days)	2.9 (2.0–6.0)

IQR, interquartile range.
^aAssessed at the time of semen collection (or at visit closest in time).
^bQuestion was “In the past year, have you smoked a cigarette, even a few puffs?” In some cases, the questionnaire was filled out up to 3 years before the semen sample was collected.
^cQuestion was “Have you drunk alcohol in the last year, including beer?” In some cases, the questionnaire was filled out up to 3 years before the semen sample was collected.
^dTwo semen samples were collected from 123 (93%) young men.
^eThis measure includes rapidly progressive motile (class A), slowly progressive motile (class B), and locally motile (class C).

and semen parameters with adjustment for potential confounders; within-person correlations in semen parameters across repeated samples were accounted for using random intercepts. We compared semen parameters (total sperm count, sperm concentration, percent motile sperm, total motile sperm count, and semen volume) for men with

higher quartiles of serum OC concentrations to those within the lowest quartile. Total sperm count (volume × sperm concentration) and total motile sperm count (total sperm count × percent motile sperm) were calculated. Total sperm count, sperm concentration, and total motile sperm count were log-transformed to approximate a normal distribution. Results

Table 2. Serum concentrations and TEQs for dioxins, furans, and PCBs measured at study enrollment (age 8–9 years of age) for 133 young men in the Russian Children’s Study.

Toxic equivalent/ concentration	Min	Percentile			Max
		25th	50th	75th	
TEQs (pg TEQ/g lipid)					
TCDD ^a	0.35	1.77	2.9	4.2	12.1
PCDD TEQ	0.95	5.69	8.7	13.3	36.0
PCDF TEQ	0.55	3.20	4.8	7.1	50.6
Co-PCB TEQ ^b	0.52	4.66	6.9	10.0	67.2
Total TEQ ^c	1.88	16.8	21.9	33.3	107
Concentration (pg/g lipid)					
PCDD	37.6	115	157	199	1,237
PCDF	14.4	29.4	44.5	63.3	406
Co-PCB ^d	62.5	131	188	273	965
Concentration (ng/g lipid)					
ΣPCBs ^e	58.3	152	235	352	1,500

^aAverage limit of detection (LOD) for TCDD was 0.60 (pg TEQ/g lipid); 16 samples (12%) were below LOD for TCDD.
^bSum of co-planar PCB TEQs [International Union of Pure and Applied Chemistry (IUPAC) congeners: 77, 81, 126, 169].
^cSum of TEQ measures for combined dioxin, furan, co-PCB and mono-ortho PCB congeners.
^dSum of co-planar PCB concentrations (IUPAC congeners: 77, 81, 126, 169).
^eSum of non-co-planar PCBs (IUPAC congeners: 18, 28, 52, 49, 44, 74, 66, 101, 99, 87, 110, 118, 105, 151, 149, 146, 153, 138/158, 128, 167, 156, 157, 178, 187, 183, 177, 172, 180, 170, 189, 201, 196/203, 195, 194, 206).




Figure 2. Adjusted mean semen parameters among 133 men (contributing 256 semen samples) from the Russian Children’s Study, by childhood serum TCDD concentrations. Data are presented as predicted marginal means (95% confidence intervals) by quartiles of TCDD concentrations (represented by the medians) adjusted for BMI, smoking status, alcohol drinker, season of sample collection, and abstinence time at the mean level of continuous covariates and adjusted for frequency of categorical measures. Motile sperm and total motile sperm count models were further adjusted by time elapsed between semen collection and analysis.

for these parameters were back-transformed to allow presentation of results in the original scale. Population marginal means (Searle et al. 1980) were utilized to present marginal population average semen parameters adjusted for the covariates (at the mean level for continuous variables and for categorical variables at a value weighted according to their frequencies) in the model. Tests for linear trends were conducted using quartile of serum OC concentrations as ordinal levels.

Potential confounding factors that were included in the models were selected primarily based on *a priori* evidence from the literature but supported empirically by associations with one or more of the semen parameters and/or serum OCs. In addition, we decided to include abstinence time regardless of statistical significance since this is a well-known predictor of most semen quality parameters, and thus can improve the precision of the exposure estimates in the model (Schisterman et al. 2009). Based on these criteria, all models were adjusted for body mass index (BMI) from the most recent physical examination, smoking status (yes vs. no, based on the response to the question “Have you smoked a cigarette, even a few puffs, within the past year?”), alcohol consumption (yes vs. no, based on the response to the question “Have you drunk alcohol in the last year, including beer?”), season of semen collection (autumn or winter vs. spring or summer), and abstinence time (< 2 days, 2–5 days, ≥ 5 days). Percent of motile sperm and total motile sperm count models were further adjusted for the time elapsed between semen collection and semen parameter analysis. Information on BMI, smoking status, and alcohol consumption was collected at the same visit year as the semen collection for 84 (63%) men, and within 3 years before semen collection for the remaining 49 (37%) men. BMI, smoking status, and alcohol consumption were unchanged between the two semen samples collected approximately 1 week apart; season, abstinence time, and time elapsed between semen collection and analysis were considered as time-varying measures for each semen sample. We analyzed the data using SAS (version 9.2; SAS Institute Inc., Cary, NC, USA), and two-sided *p*-values ≤ 0.05 were considered statistically significant.

Results

At the time of semen collection, study participants were young men with median age (IQR) = 18.3 (18.1–18.7) years, 100% Caucasian, and the median (IQR) BMI was 21.0 (19.2–23.2) kg/m² (Table 1). Fifty-one percent of the participants had smoked cigarettes (self-reported), and 68% had consumed alcohol (parental report) within the past year. One hundred thirty-three semen samples (52%) were above NAFA-ESHRE reference

values for sperm counts (≥ 80 million) and motility (≥ 60%) (Björndahl et al. 2010). The median (IQR) values for sperm parameters were 51.3 million/mL (26.6–78.8) for sperm concentration; 127 million (61.0–222) for total sperm count; and 64.0% (57.0–68.0) for sperm motility. Median (IQR) abstinence time was just under 3 days (2–6) (Table 1).

Serum concentrations of dioxins, furans, and PCBs among participants at ages 8–9 years are presented in Table 2. The median (range) values for TCDD and PCDD TEQs were 2.9 (0.4–12.1) pg/g lipid and 8.7 (1.0–36.0) pg TEQ/g lipid, respectively. Sixteen samples (12%) were below the LOD for TCDD. The median (range) of total serum TEQs was almost three times higher than levels among European children of similar age (Table 2) (Leijds et al. 2008; Link et al. 2005). The correlation between TCDD and PCDD TEQs was $r = 0.78$ ($p < 0.01$) and between PCDD TEQs and total TEQs was $r = 0.89$ ($p < 0.01$). The correlation between total TEQs and co-PCB TEQs was $r = 0.78$ ($p < 0.01$). Correlations among the dioxin and PCB congeners were lower ($r = 0.42$ – 0.57 , $p < 0.01$) (data not shown). When we compared baseline serum organochlorine concentrations adjusted by birth year between those young men who contributed semen samples and those who did

not, there were no significant differences (data not shown).

Higher serum TCDD and PCDD TEQs were associated with significantly lower semen parameters 10 years later in both unadjusted models (see Table S1) and models adjusted for BMI, smoking status, alcohol intake, season, and abstinence time (Figures 2 and 3 and Table 3). In adjusted models, on average, men in the highest quartile of serum TCDD TEQs had 40% lower sperm concentration (p -trend = 0.005), 29% lower total sperm count (p -trend = 0.05), and 30% lower total motile sperm count (p -trend = 0.05), compared to those in the lowest quartile (Figure 2). Similarly, men in the highest quartile of serum PCDD TEQs had a decrease of 39% in sperm concentration (p -trend = 0.02), 36% in total sperm count (p -trend = 0.04), and 40% in total motile sperm count (p -trend = 0.05), compared with the lowest quartile of PCDD TEQs (Figure 3).

There were no significant associations for summed concentrations of PCDDs, PCDFs, co-PCBs, or ΣPCBs with semen parameters in unadjusted (see Table S1) or adjusted models (Table 3). PCDF TEQs, co-PCB TEQs, or total TEQs were also not significantly associated with semen parameters in unadjusted (see Table S1) or adjusted models (Table 3).




Figure 3. Adjusted mean semen parameters among 133 men (contributing 256 semen samples) in the Russian Children’s Study, by childhood serum PCDD TEQs. Data are presented as predicted marginal means (95% confidence intervals) by quartiles of PCDD TEQs levels (represented by the medians) adjusted for BMI, smoking status, alcohol drinker, season of sample collection, and abstinence time at the mean level of continuous covariates and adjusted for frequency of categorical measures. Motile sperm and total motile sperm count models were further adjusted by time elapsed between semen collection and analysis.

Discussion

Our prospective cohort study showed that higher peripubertal serum TCDD and PCDD TEQs were associated with lower sperm concentration, total sperm count, and total motile sperm count measured 10 years later in healthy young men. Serum TCDD

and PCDD TEQs were not associated with percent motile sperm, so the association with total motile count was largely driven by the association with total sperm count. We did not observe associations of semen parameters with serum concentrations of PCDDs, PCDFs, co-PCBs, or ΣPCBs, nor with PCDF

TEQs, co-PCB TEQs, or total TEQs. The lack of association of semen parameters with total TEQs was surprising given the high correlation between PCDD TEQs and total TEQs. However, this might be explained by the fact that PCDDs account for slightly less than 40% of the total TEQs (Burns et al.

Table 3. Multivariable adjusted mean semen parameters by quartiles (Q)^a of serum dioxins, furans, and PCBs among 133 young men in the Russian Children's Study contributing 256 semen samples.

Toxic equivalent/ concentration	Volume (mL)	Sperm concentration (million/mL)	Total sperm count (million)	Motile sperm (%)	Total motile sperm count (million)
TEQs (pg TEQ/g lipid)					
TCDD					
Q1 (0.35–1.70)	2.7 (2.2, 3.2)	57.0 (45.0, 72.1)	128 (95.6, 173)	61.6 (58.6, 64.7)	78.0 (56.0, 109)
Q2 (1.77–2.45)	2.9 (2.5, 3.4)	51.8 (42.4, 63.3)	136 (105.0, 175)	65.4 (63.4, 67.4)	87.9 (67.1, 115)
Q3 (3.00–3.40)	2.6 (2.1, 2.9)	38.6 (28.2, 52.9)*	85.8 (60.4, 122)	59.5 (56.0, 62.9)	50.1 (33.5, 74.8)
Q4 (4.40–5.80)	3.1 (2.5, 3.7)	34.5 (25.0, 47.7)*	91.6 (63.5, 132)	60.1 (56.6, 63.7)	54.1 (36.0, 81.4)
<i>p</i> -trend	0.55	0.005	0.05	0.17	0.05
PCDD TEQ					
Q1 (0.95–5.62)	3.2 (2.7, 3.6)	64.7 (53.5, 78.2)	172 (136.0, 217)	63.4 (60.7, 66.1)	108.0 (82.5, 141)
Q2 (5.69–8.42)	2.6 (2.1, 3.1)	37.3 (27.6, 50.4)*	85.0 (58.9, 123)*	59.4 (56.1, 62.8)	49.2 (32.6, 73.5)*
Q3 (8.68–13.3)	2.4 (2.1, 2.8)*	41.9 (32.2, 54.8)*	87.7 (63.0, 122)*	63.3 (60.5, 66.1)	54.9 (38.1, 79.1)*
Q4 (13.7–36.0)	3.2 (2.6, 3.8)	39.4 (28.9, 53.6)*	109 (78.7, 150)*	60.7 (57.2, 64.3)	65.1 (45.1, 93.8)*
<i>p</i> -trend	0.89	0.02	0.04	0.55	0.05
PCDF TEQ					
Q1 (0.55–3.20)	2.9 (2.6, 3.4)	49.3 (36.4, 66.7)	128 (93.2, 176)	63.4 (60.8, 65.9)	80.3 (56.6, 114)
Q2 (3.29–4.66)	2.3 (1.9, 2.8)	43.3 (32.3, 58.0)	83.1 (57.2, 121)	59.3 (55.7, 62.9)	48.1 (31.8, 72.6)
Q3 (4.76–6.87)	3.1 (2.5, 3.6)	39.1 (30.9, 49.6)	103 (76.7, 140)	61.1 (58.2, 63.9)	62.3 (44.6, 87.2)
Q4 (7.10–50.6)	3.0 (2.5, 3.6)	47.8 (36.2, 63.1)	126 (94.5, 168)	63.0 (59.6, 66.5)	78.2 (56.5, 108)
<i>p</i> -trend	0.48	0.78	0.82	0.90	0.82
Co-PCB TEQ					
Q1 (0.52–4.63)	2.8 (2.3, 3.4)	56.5 (44.0, 72.6)	131 (97.6, 175)	63.1 (60.3, 66.0)	81.9 (59.6, 112)
Q2 (4.66–6.87)	2.9 (2.5, 3.3)	36.9 (26.2, 51.8)	95.6 (64.1, 142)	60.8 (57.8, 63.7)	57.0 (36.5, 89.0)
Q3 (6.88–9.97)	2.8 (2.2, 3.3)	37.4 (27.9, 50.2)	88.4 (62.2, 125)	62.1 (58.6, 65.6)	53.7 (36.0, 80.1)
Q4 (10.1–67.2)	2.9 (2.4, 3.4)	51.4 (40.1, 65.9)	127 (95.3, 168)	60.9 (57.3, 64.6)	76.0 (54.7, 106)
<i>p</i> -trend	0.89	0.73	0.88	0.47	0.77
Total TEQ					
Q1 (4.88–16.8)	3.0 (2.5, 3.5)	51.9 (38.3, 70.4)	131 (94.4, 181)	61.8 (58.7, 64.9)	80.4 (55.5, 116)
Q2 (17.0–21.4)	2.6 (2.2, 3.1)	38.9 (28.7, 52.6)	85.9 (57.9, 128)	61.4 (58.4, 64.3)	51.8 (33.8, 79.4)
Q3 (21.7–32.5)	2.9 (2.4, 3.5)	42.1 (33.9, 52.2)	102 (78.2, 132)	61.2 (58.1, 64.4)	60.8 (45.2, 82.0)
Q4 (33.3–107)	2.8 (2.3, 3.3)	44.8 (33.4, 60.2)	112 (82.4, 151)	61.9 (58.1, 65.6)	67.7 (47.8, 95.9)
<i>p</i> -trend	0.84	0.61	0.68	0.99	0.68
Concentration (pg/g lipid)					
PCDD					
Q1 (37.6–115)	2.9 (2.4, 3.3)	52.0 (39.4, 68.7)	130 (94.5, 180)	64.3 (61.9, 66.8)	83.1 (58.8, 118)
Q2 (118–157)	2.6 (2.0, 3.2)	43.2 (33.9, 55.0)	91.0 (65.8, 126)	58.9 (55.7, 62.0)	52.6 (37.0, 75.7)
Q3 (158–200)	3.3 (2.7, 3.8)	37.6 (28.0, 50.6)	108 (76.8, 151)	63.2 (60.4, 66.0)	67.2 (46.0, 98.3)
Q4 (201–1,237)	2.7 (2.2, 3.2)	47.2 (35.9, 62.1)	109 (81.3, 146)	60.4 (56.7, 64.0)	64.3 (46.3, 89.3)
<i>p</i> -trend	0.81	0.48	0.59	0.30	0.49
PCDF					
Q1 (14.4–29.2)	2.7 (2.2, 3.1)	53.1 (38.9, 72.5)	122 (86.2, 173)	63.7 (61.1, 66.3)	76.8 (53.1, 111)
Q2 (29.4–43.6)	2.6 (2.2, 3.0)	41.8 (32.1, 54.2)	89.6 (63.2, 127)	60.3 (57.0, 63.6)	53.0 (35.6, 78.9)
Q3 (44.5–63.0)	3.5 (2.9, 4.1)	37.2 (28.2, 48.9)	112 (80.8, 155)	60.0 (56.5, 63.3)	65.9 (45.4, 95.0)
Q4 (63.3–405)	2.6 (2.2, 3.0)	48.6 (37.2, 63.4)	115 (84.9, 155)	63.0 (59.8, 66.2)	71.3 (51.0, 100)
<i>p</i> -trend	0.47	0.60	0.93	0.79	0.98
Co-PCB					
Q1 (62.5–126)	2.6 (2.1, 3.1)	59.7 (45.8, 77.8)	128 (92.5, 179)	62.5 (59.4, 65.6)	79.1 (54.9, 114)
Q2 (130–184)	2.6 (2.2, 3.0)	39.3 (28.8, 53.7)	88.4 (61.8, 126)	61.5 (58.6, 64.4)	53.6 (36.0, 79.7)
Q3 (187–274)	3.3 (2.7, 3.8)	38.9 (29.7, 51.0)	108 (78.0, 149)	61.4 (58.6, 64.2)	65.0 (45.6, 92.7)
Q4 (275–965)	3.0 (2.5, 3.5)	43.9 (34.4, 56.1)	114 (86.3, 151)	61.5 (57.8, 65.1)	69.1 (49.9, 95.7)
<i>p</i> -trend	0.09	0.11	0.79	0.67	0.75
Concentration (ng/g lipid)					
ΣPCBs					
Q1 (58.3–151)	2.9 (2.4, 3.5)	52.5 (39.8, 69.3)	122 (87.4, 171)	62.6 (59.6, 65.7)	79.9 (55.9, 114)
Q2 (152–236)	2.6 (2.1, 3.0)	47.4 (34.8, 64.5)	103 (69.8, 152)	62.5 (59.9, 64.9)	65.1 (42.9, 98.8)
Q3 (239–352)	2.7 (2.3, 3.2)	33.8 (25.8, 44.3)*	84.3 (61.8, 115)*	61.5 (57.9, 65.1)	48.5 (33.4, 70.4)
Q4 (356–1,500)	3.0 (2.4, 3.6)	45.3 (34.3, 59.9)	110 (79.4, 152)	59.6 (55.9, 63.3)	68.6 (49.7, 94.7)
<i>p</i> -trend	0.81	0.24	0.47	0.19	0.36

^aData are presented as predicted estimates (95% confidence intervals) adjusted for BMI, smoking status, alcohol drinker, season, and abstinence time at the mean level of continuous covariates and adjusted for frequency of categorical measures. Motile sperm and total motile sperm count models were further adjusted by time to start semen analysis.

**p* < 0.05.

2009). This suggests that the associations we found may be more specific to PCDD TEQs than to overall TEQs, which also included contributions of PCDFs and co-planar- and mono-*ortho*-PCBs, which were not independently associated with semen parameters. Although cross-sectional studies on PCBs have reported negative associations with semen parameters (Meeker and Hauser 2010), we did not find longitudinal associations between childhood serum concentrations of PCBs and semen parameters in our cohort.

Similar to our TCDD results, those of Mocarelli et al. (2008) showed that men from the Seveso cohort who were acutely exposed to very high levels of TCDD during childhood (ages 1–9 years) had impaired semen parameters. Specifically, they had a 27% decrease in sperm concentration ($p = 0.025$), a 20% decrease in sperm motility ($p = 0.001$), and a 39% decrease in total motile sperm count ($p = 0.01$) 22 years later, compared with men in the control group without acute high exposure. In contrast, the Seveso boys exposed to high levels of TCDD during puberty (ages 10–17 years) had higher total sperm count and total motile sperm count than did men in the control group. These results suggest a differential effect of TCDD by age at exposure. The OC measurements in the Russian Children's Study reflect cumulative exposure up to age 8–9 years, whereas the boys in the Seveso cohort were exposed at a specific time point before age 10 years (mean age at exposure, 6.2 years); therefore, we can speculate that the Russian boys and this subset of Seveso boys were exposed before pubertal onset or very early during pubertal development. Both the Seveso study and our results suggest that the peripubertal period may be particularly susceptible to the deleterious effects of TCDD on adult semen parameters. In the Mocarelli et al. (2008) study, the median serum TCDD concentrations among the exposed group of children was 210 pg TEQ/g lipid and the control group had serum TCDD < 15 pg TEQ/g lipid. In contrast, for boys in our study, the median serum TCDD was 2.9 pg TEQ/g lipid, about 70-fold lower than exposed Seveso boys. Therefore, our results showed that childhood serum TCDD TEQ levels much lower than those measured in the Seveso study had a negative association with adult semen parameters. In addition, we found negative associations between childhood serum PCDD TEQs with sperm concentration, count and motile count, indicating that childhood exposure to other dioxins may also negatively affect semen parameters in adult life.

The period of sexual differentiation and reproductive tract organization during fetal development is highly sensitive to endocrine disrupting exposures which can

affect reproductive-tract development and subsequent pubertal timing (Sharpe 2006). However, childhood and adolescence may also be vulnerable to such exposures due to the developmental changes of pubertal maturation that occur at these ages (Bin-Abbas et al. 1999; Grumbach 2002). Previously, we reported that higher peripubertal serum dioxins were associated with delayed pubertal onset and sexual maturity in the Russian cohort (Burns et al. 2016; Korricks et al. 2011). The proliferation and differentiation of Sertoli cells, the support cells of the seminiferous tubules, are peripubertal androgen-dependent processes that are critical for spermatogenesis (Sharpe et al. 2003). Dioxins can inhibit testosterone biosynthesis (Svechnikov et al. 2010), and may have direct testicular actions as the AhR is widely expressed in the testes (Schultz et al. 2003). AhR-mediated disruption of androgen activity could affect proliferation of the Sertoli cells and their subsequent differentiation, and pubertal maturation of the seminiferous tubules (Sharpe et al. 2003; Woodruff et al. 2010). These mechanisms could contribute to the observed decrease in sperm count in adults who were exposed to TCDD and PCDD TEQs as young children (Mocarelli et al. 2008).

Our findings are in agreement with animal data showing TCDD inhibition of testicular development and function during critical periods of reproductive-tract development, including fetal, neonatal (Arima et al. 2009; Faqi et al. 1998), pubertal (el-Sabeawy et al. 1998), and adult stages (Oguz et al. 2013; Sönmez et al. 2011). Moreover, childhood exposures to dioxins, furans, and PCBs have been shown to adversely affect other key maturational processes, such as somatic growth and pubertal timing in our cohort (Burns et al. 2011, 2016; Korricks et al. 2011).

Our study has several potential limitations. First, we did not measure prenatal exposure to OCs, when sexual differentiation and reproductive tract organization occur. Nevertheless, childhood is also a vulnerable developmental period. Second, we excluded boys with severe chronic illnesses at study entry. If their diseases were caused by or at least partially attributable to pre- or perinatal exposure to dioxins, furans, and/or PCBs, the association of these exposures with semen parameters may be underestimated in our analyses. Third, in our study, the boys' median serum total TEQ concentrations were three times higher than the geometric mean in the U.S. National Health and Nutrition Examination Survey for males 12–19 years of age (no data were available for children < 12 years of age) (Patterson et al. 2008), and three times higher (using 1998 WHO total TEQs) than levels among similarly aged German boys (Link et al. 2005). This makes it difficult to investigate

the effects of very low exposures in our cohort. However, despite this, our concentrations of TCDD were much lower than those in the Seveso study, which was used by the U.S. EPA in their dioxin risk assessment document (U.S. EPA 2009).

The strengths of our study include its prospective design and long-term serial follow-up of participants which minimizes the risk of reverse causation, the consistency in analysis of semen samples by the same laboratory technician which prevents interobserver variation, the comprehensive adjustment for possible confounding variables collected using physical examination and questionnaire data, and the availability of two semen samples on almost all participants (93%).

Conclusions

Our results showed an association of peripubertal serum concentrations of TCDD with poorer semen parameters. Our results, along with toxicological evidence, suggest that peripubertal exposure to TCDD and dioxins may adversely impact adult semen parameters. We found this association at much lower TCDD concentrations than in the Seveso study, suggesting that moderate concentrations may also impact semen parameters and providing evidence that would be useful for risk assessment. Semen parameters are a marker of fertility and future studies on the impact of TCDD and dioxins on male fertility are warranted.

REFERENCES

- Arima A, Kato H, Ooshima Y, Tateishi T, Inoue A, Muneoka A, et al. 2009. *In utero* and lactational exposure to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) induces a reduction in epididymal and ejaculated sperm number in rhesus monkeys. *Reprod Toxicol* 28:495–502.
- Baccarelli A, Pfeiffer R, Consonni D, Pesatori AC, Bonzini M, Patterson DG Jr, et al. 2005. Handling of dioxin measurement data in the presence of non-detectable values: overview of available methods and their application in the Seveso chloracne study. *Chemosphere* 60:898–906.
- Barr JR, Maggio VL, Barr DB, Turner WE, Sjödin A, Sandau CD, et al. 2003. New high-resolution mass spectrometric approach for the measurement of polychlorinated biphenyls and organochlorine pesticides in human serum. *J Chromatogr B Analyt Technol Biomed Life Sci* 794:137–148.
- Bin-Abbas B, Conte FA, Grumbach MM, Kaplan SL. 1999. Congenital hypogonadotropic hypogonadism and micropenis: effect of testosterone treatment on adult penile size—why sex reversal is not indicated. *J Pediatr* 134:579–583.
- Björndahl L, Mortimer D, Barratt CLR, Castilla JA, Menkveld R, Kvist U, et al. 2010. *A Practical Guide to Basic Laboratory Andrology*. Cambridge, UK: Cambridge University Press.
- Burns JS, Lee MM, Williams PL, Korricks SA, Sergeev O, Lam T, et al. 2016. Associations of peripubertal serum dioxin and polychlorinated biphenyl concentrations with pubertal timing

- among Russian boys. *Environ Health Perspect* 124:1801–1180, doi: 10.1289/EHP154.
- Burns JS, Williams PL, Sergeev O, Korrick S, Lee MM, Revich B, et al. 2009. Predictors of serum dioxins and PCBs among peripubertal Russian boys. *Environ Health Perspect* 117:1593–1599, doi: 10.1289/ehp.0800223.
- Burns JS, Williams PL, Sergeev O, Korrick S, Lee MM, Revich B, et al. 2011. Serum dioxins and polychlorinated biphenyls are associated with growth among Russian boys. *Pediatrics* 127:e59–e68.
- Carlsen E, Giwercman A, Keiding N, Skakkebaek NE. 1992. Evidence for decreasing quality of semen during past 50 years. *BMJ* 305:609–613.
- el-Sabeawy F, Wang S, Overstreet J, Miller M, Lasley B, Enan E. 1998. Treatment of rats during pubertal development with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin alters both signaling kinase activities and epidermal growth factor receptor binding in the testis and the motility and acrosomal reaction of sperm. *Toxicol Appl Pharmacol* 150:427–442.
- Faqi AS, Dalsenter PR, Merker HJ, Chahoud I. 1998. Reproductive toxicity and tissue concentrations of low doses of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in male offspring rats exposed throughout pregnancy and lactation. *Toxicol Appl Pharmacol* 150:383–392.
- Faure AC, Viel JF, Bailly A, Blagosklonov O, Amiot C, Roux C. 2014. Evolution of sperm quality in men living in the vicinity of a municipal solid waste incinerator possibly correlated with decreasing dioxins emission levels. *Andrologia* 46:744–752.
- Grumbach MM. 2002. The neuroendocrinology of human puberty revisited. *Horm Res* 57(suppl 2):2–14.
- Hauser R, Sergeev O, Korrick S, Lee MM, Revich B, Gitin E, et al. 2008. Association of blood lead levels with onset of puberty in Russian boys. *Environ Health Perspect* 116:976–980, doi: 10.1289/ehp.10516.
- Jørgensen N, Andersen AG, Eustache F, Irvine DS, Suominen J, Petersen JH, et al. 2001. Regional differences in semen quality in Europe. *Hum Reprod* 16:1012–1019.
- Jørgensen N, Carlsen E, Neramo I, Punab M, Suominen J, Andersen AG, et al. 2002. East-West gradient in semen quality in the Nordic-Baltic area: a study of men from the general population in Denmark, Norway, Estonia and Finland. *Hum Reprod* 17:2199–2208.
- Korrick SA, Lee MM, Williams PL, Sergeev O, Burns JS, Patterson DG, et al. 2011. Dioxin exposure and age of pubertal onset among Russian boys. *Environ Health Perspect* 119:1339–1344, doi: 10.1289/ehp.1003102.
- Kvist U, Björndahl L. 2002. *Manual on Basic Semen Analysis*. Eshre Monographs. http://www.fivfrance.com/pro/pdf_NAFA_ESHRE2002.pdf [accessed 28 January 2016].
- Leijis MM, Koppe JG, Olie K, van Aalderen WM, Voogt P, Vulsma T, et al. 2008. Delayed initiation of breast development in girls with higher prenatal dioxin exposure; a longitudinal cohort study. *Chemosphere* 73:999–1004.
- Link B, Gabrio T, Zoellner I, Piechotowski I, Paepke O, Herrmann T, et al. 2005. Biomonitoring of persistent organochlorine pesticides, PCDD/PCDFs and dioxin-like PCBs in blood of children from South West Germany (Baden-Wuerttemberg) from 1993 to 2003. *Chemosphere* 58:1185–1201.
- Meeker JD, Hauser R. 2010. Exposure to polychlorinated biphenyls (PCBs) and male reproduction. *Syst Biol Reprod Med* 56:122–131.
- Mocarelli P, Gerthoux PM, Needham LL, Patterson DG Jr, Limonta G, Falbo R, et al. 2011. Perinatal exposure to low doses of dioxin can permanently impair human semen quality. *Environ Health Perspect* 119:713–718, doi: 10.1289/ehp.1002134.
- Mocarelli P, Gerthoux PM, Patterson DG Jr, Milani S, Limonta G, Bertona M, et al. 2008. Dioxin exposure, from infancy through puberty, produces endocrine disruption and affects human semen quality. *Environ Health Perspect* 116:70–77, doi: 10.1289/ehp.10399.
- Oguz F, Ciftci O, Aydin M, Timurkaan N, Beytur A, Altintas R, et al. 2013. Aminoguanidine prevents testicular damage-induced-2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) in male rats. *Andrologia* 45:225–231.
- Paoli D, Giannandrea F, Gallo M, Turci R, Cattaruzza MS, Lombardo F, et al. 2015. Exposure to polychlorinated biphenyls and hexachlorobenzene, semen quality and testicular cancer risk. *J Endocrinol Invest* 38:745–752.
- Patterson DG Jr, Hampton L, Lapeza CR Jr, Belser WT, Green V, Alexander L, et al. 1987. High-resolution gas chromatographic/high-resolution mass spectrometric analysis of human serum on a whole-weight and lipid basis for 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Anal Chem* 59:2000–2005.
- Patterson DG Jr, Turner WE, Caudill SP, Needham LL. 2008. Total TEQ reference range (PCDDs, PCDFs, cPCBs, mono-PCBs) for the US population 2001–2002. *Chemosphere* 73(1 suppl):S261–S277.
- Phillips DL, Pirkle JL, Burse VW, Bernert JT Jr, Henderson LO, Needham LL. 1989. Chlorinated hydrocarbon levels in human serum: effects of fasting and feeding. *Arch Environ Contam Toxicol* 18:495–500.
- Schechter A, Cramer P, Boggess K, Stanley J, Pöpke O, Olson J, et al. 2001. Intake of dioxins and related compounds from food in the U.S. population. *J Toxicol Environ Health A* 63:1–18.
- Schisterman EF, Cole SR, Platt RW. 2009. Over-adjustment bias and unnecessary adjustment in epidemiologic studies. *Epidemiology* 20:488–495.
- Schultz R, Suominen J, Värre T, Hakovirta H, Parvinen M, Toppari J, et al. 2003. Expression of aryl hydrocarbon receptor and aryl hydrocarbon receptor nuclear translocator messenger ribonucleic acids and proteins in rat and human testis. *Endocrinology* 144:767–776.
- Searle SR, Speed FM, Milliken GA. 1980. Population marginal means in the linear model: an alternative to least square means. *Am Stat* 34:216–221.
- Sharpe RM. 2006. Pathways of endocrine disruption during male sexual differentiation and masculinization. *Best Pract Res Clin Endocrinol Metab* 20:91–110.
- Sharpe RM, McKinnell C, Kivlin C, Fisher JS. 2003. Proliferation and functional maturation of Sertoli cells, and their relevance to disorders of testis function in adulthood. *Reproduction* 125:769–784.
- Sjödin A, Jones RS, Lapeza CR, Focant JF, McGahee EE III, Patterson DG Jr. 2004. Semiautomated high-throughput extraction and cleanup method for the measurement of polybrominated diphenyl ethers, polybrominated biphenyls, and polychlorinated biphenyls in human serum. *Anal Chem* 76:1921–1927.
- Sönmez M, Türk G, Çeribaşı AO, Sakin F, Ateşşahin A. 2011. Attenuating effect of lycopene and ellagic acid on 2,3,7,8-tetrachlorodibenzo-*p*-dioxin-induced spermiotoxicity and testicular apoptosis. *Drug Chem Toxicol* 34:347–356.
- Svechnikov K, Izzo G, Landreh L, Weisser J, Söder O. 2010. Endocrine disruptors and Leydig cell function. *J Biomed Biotechnol* 2010:684504, doi: 10.1155/2010/684504.
- Swan SH, Brazil C, Drobnis EZ, Liu F, Kruse RL, Hatch M, et al. 2003. Geographic differences in semen quality of fertile U.S. males. *Environ Health Perspect* 111:414–420, doi: 10.1289/ehp.5927.
- Swan SH, Elkin EP, Fenster L. 2000. The question of declining sperm density revisited: an analysis of 101 studies published 1934–1996. *Environ Health Perspect* 108:961–966.
- Toft G, Rignell-Hydbom A, Tyrkiel E, Shvets M, Giwercman A, Lindh CH, et al. 2006. Semen quality and exposure to persistent organochlorine pollutants. *Epidemiology* 17:450–458.
- Turner W, DiPietro E, Lapeza C, Green V, Gill J, Patterson DG Jr. 1997. A fast universal automated cleanup system for the isotope-dilution high-resolution mass spectrometric analysis of PCDDs, PCDFs, coplanar PCBs, PCB congeners, and persistent pesticides from the same serum sample. *Organohalogen Compounds* 31:26–31.
- U.S. EPA (U.S. Environmental Protection Agency). 2009. Summary of U.S. EPA Dioxin Workshop. February 18–20, 2009. Cincinnati, Ohio. EPA/600/R-09/027. Cincinnati, OH:U.S. EPA, Office of Research and Development. https://cfpub.epa.gov/si/si_public_file_download.cfm?p_download_id=489585 [accessed 28 January 2016].
- Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, et al. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 93:223–241.
- WHO (World Health Organization). 1999. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction*. 4th ed. Cambridge, UK:Cambridge University Press.
- Williams PL, Sergeev O, Lee MM, Korrick SA, Burns JS, Humblet O, et al. 2010. Blood lead levels and delayed onset of puberty in a longitudinal study of Russian boys. *Pediatrics* 125:e1088–e1096.
- Woodruff TJ, Janssen S, Guillette LJ Jr, Giudice LC. 2010. *Environmental Impacts on Reproductive Health and Fertility*. Cambridge, UK:Cambridge University Press.