

A DICHOTOMY THEOREM FOR THE GENERALIZED BAIRE SPACE AND ELEMENTARY EMBEDDABILITY AT UNCOUNTABLE CARDINALS

DOROTTYA SZIRÁKI AND JOUKO VÄÄNÄNEN

ABSTRACT. We consider the following dichotomy for Σ_2^0 finitary relations R on analytic subsets of the generalized Baire space for κ : either all R -independent sets are of size at most κ , or there is a κ -perfect R -independent set. This dichotomy is the uncountable version of a result found in (W. Kubiś, Proc. Amer. Math. Soc. 131 (2003), no 2.:619–623) and in (S. Shelah, Fund. Math. 159 (1999), no. 1:1–50). We prove that the above statement holds assuming \diamond_κ and the set theoretical hypothesis $I^-(\kappa)$, which is the modification of the hypothesis $I(\kappa)$ suitable for limit cardinals. When κ is inaccessible, or when R is a closed binary relation, the assumption \diamond_κ is not needed.

We obtain as a corollary the uncountable version of a result by G. Sági and the first author (Log. J. IGPL 20 (2012), no. 6:1064–1082) about the κ -sized models of a $\Sigma_1^1(L_{\kappa+\kappa})$ -sentence when considered up to isomorphism, or elementary embeddability, by elements of a K_κ subset of ${}^\kappa\kappa$. The role of elementary embeddings can be replaced by a more general notion that also includes embeddings, as well as the maps preserving $L_{\lambda\mu}$ for $\omega \leq \mu \leq \lambda \leq \kappa$ and the finite variable fragments of these logics.

1. INTRODUCTION

The domain of the *generalized Baire space for κ* , or the κ -Baire space for short, is the set ${}^\kappa\kappa$ of functions from κ to κ , and its topology is given by the basic open sets

$$N_p = \{x \in {}^\kappa\kappa : p \subseteq x\}$$

for all $p \in {}^{<\kappa}\kappa$. The *generalized Cantor space for κ* is ${}^\kappa 2$ with the topology induced by the κ -Baire space via the natural injection of ${}^\kappa 2$ into ${}^\kappa\kappa$. Unless otherwise stated, ${}^\kappa\kappa$ and ${}^\kappa 2$ are endowed with the above topologies, and for $2 \leq n < \omega$ the set ${}^n({}^\kappa\kappa)$ is endowed with the resulting product topology; as in classical descriptive set theory, this space is homeomorphic to ${}^\kappa\kappa$. The hypothesis $\kappa^{<\kappa} = \kappa$ is usually assumed when working with the κ -Baire and κ -Cantor space, because it implies that these spaces have some nice properties. Under it, for example, the standard bases of both spaces are of

2010 *Mathematics Subject Classification*. Primary 03E15; Secondary 03C45, 03C57, 03E55.

Key words and phrases. generalized Baire space, dichotomy theorem, Σ_2^0 relations, elementary embeddability, uncountable models.

size κ and consist of clopen sets, and, furthermore, the intersection (resp. union) of $< \kappa$ many open (closed) sets is open (closed).

A subset of a topological space X is defined to be $\Sigma_2^0(\kappa)$ (resp. $\Pi_2^0(\kappa)$), if it can be written as the union (intersection) of at most κ many closed (open) subsets of X . More generally, the collection of κ -Borel subsets of a topological space X is the smallest set of subsets of X which contains the open subsets and is closed under complementation and taking unions and intersections of at most κ many sets. A subset of X is κ -analytic, or $\Sigma_1^1(\kappa)$, if it can be obtained as a continuous image of a closed subset of the κ -Baire space ${}^\kappa\kappa$; this concept was introduced in [18] (for the κ -Baire space). In the case of the κ -Baire space ${}^\kappa\kappa$ (when $\kappa^{<\kappa} = \kappa$ holds), a subset is κ -analytic iff it is the image of a κ -Borel subset of ${}^\kappa\kappa$ under a κ -Borel map. When the topological space X is homeomorphic to a κ -analytic subset of the κ -Baire space, we will omit the “ κ ” when talking about $\Sigma_2^0(\kappa)$, $\Pi_2^0(\kappa)$, κ -Borel or κ -analytic subsets of X , i.e., we will call such subsets Σ_2^0 , Π_2^0 , Borel and analytic (or Σ_1^1) subsets, respectively.

We denote by $I^-(\kappa)$ the following set theoretical hypothesis:

there is a κ^+ -complete normal (non principal) ideal \mathcal{I} on κ^+ such that the set

$$\mathcal{I}^+ = \mathcal{P}(\kappa^+) - \mathcal{I}$$

contains a dense subset K such that every descending sequence of length $< \kappa$ of elements of K has a lower bound in K .

This hypothesis is the modification of the hypothesis $I(\kappa)$ (introduced in [9]) which is appropriate for limit cardinals κ (see also [7, 17, 26] and [27] where the specific case of $I(\omega)$ is considered).

If κ is a regular cardinal and $\lambda > \kappa$ is measurable, then Lévy-collapsing λ to κ^+ yields a model of ZFC in which $I^-(\kappa)$ holds. The corresponding statement for $I(\kappa)$ is an unpublished result of Richard Laver; the proofs can be reconstructed from the $I(\omega)$ case, which is described in [7]. Note that \diamond_κ will also hold in the model obtained in the above way, and therefore the consistency of $I^-(\kappa)$ together with \diamond_κ follows from the existence of a measurable $\lambda > \kappa$. A model in which $I^-(\kappa)$ holds and κ is inaccessible can be obtained in the same way, if we start out from a situation where κ is inaccessible and there exists a measurable $\lambda > \kappa$. The following two facts were suggested to us by Menachem Magidor.

1. Given any cardinal κ , and assuming the existence of a measurable $\lambda > \kappa$, one can also obtain a model in which $2^\kappa > \kappa^+$ holds together with $I^-(\kappa)$ (and also \diamond_κ), by forcing with the product of the Lévy-collapse of λ to

κ^+ and $Add(\kappa, \mu)$ for any $\mu > \lambda$ (the standard forcing notion that adds μ -many Cohen subsets of κ).

2. It is also possible for a supercompact κ to satisfy $I^-(\kappa)$ and even $2^\kappa > \kappa^+$ (see Remark 2.11).

We note that the hypothesis $\kappa^{<\kappa} = \kappa$, which is useful when considering the κ -Baire space, follows from our assumption $I^-(\kappa)$. This implication can be proven by a straightforward generalization of the proof of the fact that $I(\omega)$ implies CH [26, p. 13]. Using this fact, and a result of Shelah [25] that \diamond_κ holds when $\kappa = \mu^+ = 2^\mu > \aleph_1$, we have that $I^-(\kappa)$ implies \diamond_κ whenever $\kappa > \aleph_1$ is a successor cardinal.

A subset Y of the generalized Baire space ${}^\kappa\kappa$ is defined to be *perfect* if Y consists of the κ -branches $[T]$ of a *perfect tree* T , i.e., a subtree $T \subseteq {}^{<\kappa}\kappa$ whose set of splitting nodes is cofinal and which is $< \kappa$ -closed (i.e., every increasing sequence in T of length $< \kappa$ has an upper bound in T); this concept was introduced in [27]. Note that $X \subseteq {}^\kappa\kappa$ contains a perfect subset iff there is a continuous injection of ${}^\kappa 2$ into X iff there is such a Borel injection; see, e.g., [4, Proposition 2].

Let X be an analytic subset of the κ -Baire space ${}^\kappa\kappa$. We say that R is a Σ_2^0 (*closed, Borel, etc.*) *relation on X* if, for some $1 \leq n < \omega$, R is a $\Sigma_2^0(\kappa)$ (*closed, κ -Borel, etc.*) subset of the product space ${}^n X$, where X is endowed with the subspace topology induced by the κ -Baire space. A subset Y of X is *R -independent* iff for all pairwise distinct $y_0, \dots, y_{n-1} \in Y$ we have $(y_0, \dots, y_{n-1}) \notin R$.

Our main result, whose proof is detailed in Section 2, is as follows.

Theorem 2.4 *Assume $I^-(\kappa)$ and either that \diamond_κ or that κ is inaccessible.*

Suppose R is a Σ_2^0 relation on an analytic subset of the κ -Baire space ${}^\kappa\kappa$. Then either all R -independent sets are of size $\leq \kappa$, or there exists a perfect R -independent set.

We note that in the case that R is a *closed binary relation* on an analytic subset of ${}^\kappa\kappa$, only the hypothesis $I^-(\kappa)$ is needed for the above dichotomy to hold; see Remark 2.5.

This dichotomy is the uncountable generalization of [12, Corollary 2.13]. (There, the result is formulated in terms of homogeneous sets of G_δ colorings on analytic spaces; see Corollary 2.7 for this formulation in the uncountable case.) The special case of the result for binary relations on Polish spaces is also mentioned in [23, Remark 1.14].

The above dichotomy for uncountable κ does not follow from ZFC alone; if there exists a weak κ -Kurepa tree, for example, then it can not hold for κ . This further implies that when $V = L$, the dichotomy fails for all uncountable κ , and that the consistency of this dichotomy is at least that of an inaccessible cardinal. (See Remark 2.6 for details.) By Theorem 2.4, we have that the consistency of a measurable cardinal implies the consistency of the above dichotomy; however we do not yet know its exact consistency strength.

A specific case of Theorem 2.4 is that the (κ) -Silver dichotomy holds for Σ_2^0 equivalence relations on the κ -Baire space under the assumption $I^-(\kappa)$ together with either \diamond_κ or the inaccessibility of κ . Recently, a considerable effort has been made to investigate set theoretical conditions implying (the consistency of) the satisfaction or the failure of the Silver dichotomy for Borel equivalence relations on the generalized Baire space, see, e.g., [4, 6] and [5, Section 4.2]. It would be worth investigating whether our hypotheses imply this more general case as well.

In Section 3, we use the results of the previous section to obtain model theoretic dichotomies motivated by the spectrum problem.

Suppose κ is a cardinal and ψ is a sentence in $\Sigma_1^1(L_{\kappa+\kappa})$ (i.e., it is a second order sentence of the form $\exists \bar{R} \varphi(\bar{R})$ where \bar{R} is a set of $\leq \kappa$ many symbols disjoint from the original vocabulary and $\varphi(\bar{R})$ is an $L_{\kappa+\kappa}$ sentence in the expanded language). One obtains interesting questions by considering, instead of the number of non-isomorphic κ -sized models of ψ , the possible sizes of sets of such models which are pairwise non-elementarily embeddable, as in for example [1, 22]. More generally, the role of elementary embeddings may be replaced by embeddings preserving (in the sense of (1) in Definition 3.2) “nice” sets of formulas, possibly of some extension of first order logic.

We consider the case when the “nice” sets of formulas to be preserved are what we call *fragments* of $L_{\kappa+\kappa}$ (see Definition 3.1). Examples of fragments of $L_{\kappa+\kappa}$ include not only the set of all first order formulas and the set At of all atomic formulas and their negations (the maps preserving these sets of formulas are elementary embeddings and embeddings, respectively), but also the infinitary logics $L_{\lambda\mu}$, where $\omega \leq \mu \leq \lambda \leq \kappa$, and n -variable fragments of these logics. In the case of fragments $F \subseteq L_{\kappa+\omega}$ and sentences $\psi \in F$, the set of models of ψ together with the embeddings preserving F forms an abstract elementary class, and the corresponding version of the above question has been studied in, e.g., [24]. To the best knowledge of the

first author, this question has not been studied yet in the case fragments of $L_{\kappa+\kappa}$ which are not subsets of $L_{\kappa+\omega}$.

Since we are dealing with models of size κ up to elementary embeddability or, more generally, up to embeddability by maps preserving fragments of $L_{\kappa+\kappa}$ (and isomorphisms are a special case of such embeddings), we may assume that all models have domain κ . Accordingly, let Mod_κ^ψ denote the set of models of ψ with domain κ , and denote by $\text{Inj}(\kappa)$ the set of injective functions in ${}^\kappa\kappa$. Then any embedding between elements of Mod_κ^ψ (preserving a fragment of $L_{\kappa+\kappa}$) is an element of $\text{Inj}(\kappa)$. It is natural to ask what happens when, in the above questions, the role of $\text{Inj}(\kappa)$ is replaced by a certain subset H of $\text{Inj}(\kappa)$, i.e., when Mod_κ^ψ is considered up to only the embeddings which are in H (and preserve the given fragment of $L_{\kappa+\kappa}$). Notice that when H is a subgroup of $\text{Sym}(\kappa)$, the above question reduces to considering models up to isomorphisms in H .

More precisely, let \mathcal{A} and \mathcal{B} be models with domain κ . For a subset H of $\text{Inj}(\kappa)$, we say that \mathcal{A} is *H-elementarily embeddable* into \mathcal{B} iff there is some function h in H that embeds \mathcal{A} elementarily into \mathcal{B} . (In the special case that H is a subgroup of $\text{Sym}(\kappa)$, we say that the two structures are *H-isomorphic*.) Analogously, we define the more general notion of *(F, H)-elementary embeddability* for fragments F of $L_{\kappa+\kappa}$ (see Definition 3.2). We are interested in the possible sizes of pairwise non *(F, H)-elementarily embeddable* subsets of Mod_κ^ψ . By introducing the set H of “allowed embeddings” as an extra parameter, we may study explicitly the role the topological properties of H play in the above question.

A subset C of a topological space is defined to be *κ -compact* if any open cover of C has a subcover of size $< \kappa$, and C is K_κ if it can be written as the union of at most κ many κ -compact subsets. A topological space is K_κ if it is a K_κ subset of itself.

The next dichotomy theorem, which is the main result of Section 3, gives an answer to the above question when H is a K_κ subset of the κ -Baire space, or, for certain fragments, when H is a K_κ subset of the product space $({}^\kappa\kappa, \tau_{\text{pr}})$, where τ_{pr} is the product topology on the set ${}^\kappa\kappa$ obtained by equipping κ with the discrete topology.

Theorem 3.8 *Assume the set theoretical hypotheses of Theorem 2.4. Suppose that $H \subseteq \text{Inj}(\kappa)$, F is a fragment of $L_{\kappa+\kappa}$ and ψ is a sentence of $\Sigma_1^1(L_{\kappa+\kappa})$. Suppose that either*

1. H is a K_κ subset of the κ -Baire space, or
2. H is a K_κ subset of the product space $({}^\kappa\kappa, \tau_{\text{pr}})$ and $F \subseteq L_{\kappa+\omega}$.

If there are at least κ^+ many pairwise non (F, H) -elementarily embeddable models in Mod_κ^ψ , then there are perfectly many such models.

Theorem 3.8 can be seen as uncountable version of [21, Theorems 5.8 and 5.9]. We note that these two cited theorems of [21] also follow from [12, Corollary 2.13] or from [23, Remark 1.14].

In order to prove Theorem 3.8, we have to first show in the beginning of Section 3 that for any fragment F and K_κ subset H of the κ -Baire space, Mod_κ^ψ can be viewed as an analytic subset of the κ -Cantor space on which (F, H) -elementary embeddability is a Σ_2^0 binary relation (and when $F \subseteq L_{\kappa+\omega}$, this holds even when H is K_κ only in the product topology τ_{pr}). This is done by considering a Borel refinement (the Borel refinement t_F induced by F) of the canonical topology used to study the deep connections between model theory and generalized descriptive set theory (see [18] and, e.g., [28] and [5]), and generalizing to the uncountable case an argument in [19]. These arguments allow us to obtain Theorem 3.8 as a special case of our more general dichotomy result, Theorem 2.4. In Theorem 3.8, the word “perfect” may refer to the topology $t_{F'}$ induced by *any* fragment F' of $L_{\kappa+\kappa}$ (see Proposition 3.7).

We remark that when κ is a non-weakly compact cardinal, the assumption $I^-(\kappa)$ implies that there are no K_κ subsets of the κ -Baire space other than those of size $\leq \kappa$. However, K_κ sets of size $> \kappa$ exist in the case of the product space $({}^\kappa\kappa, \tau_{\text{pr}})$, or in the case of the κ -Baire space when κ is weakly compact. (See Remark 2.10.)

One possible motivation for investigating the above questions for K_κ subsets H , even in the case of the κ -Baire space for κ non-weakly compact, is the following. Let F be a fixed fragment of $L_{\kappa+\kappa}$ and equip Mod_κ^ψ with the topology t_F described above. Consider, for each $H \subseteq \text{Inj}(\kappa)$, the (F, H) -elementary embeddability relation R_H^F viewed as a subset of $\text{Mod}_\kappa^\psi \times \text{Mod}_\kappa^\psi$. Specifically, the relation $R_{\text{Inj}(\kappa)}^F$ (of embeddability by any map preserving F) corresponds to the original question where the set of “allowed” embeddings “has not been restricted”. Because the standard base of the space $(\text{Mod}_\kappa^\psi, t_F)$ is of size κ , it is possible to construct a subset (even a submonoid) H of $\text{Inj}(\kappa)$ of size $\leq \kappa$ such that R_H^F is dense in $R_{\text{Inj}(\kappa)}^F$. On the one hand, the density of R_H^F in $R_{\text{Inj}(\kappa)}^F$ may be interpreted, on an intuitive level, to mean that “the action of H on Mod_κ^ψ is locally similar to the action of $\text{Inj}(\kappa)$ ”. On the other hand, $|H| \leq \kappa$ and is therefore a K_κ subset of the κ -Baire space, which implies that our model theoretic dichotomy result Theorem 3.8 is applicable in this case as well.

Notation. The notation we use is mostly standard. In particular, for sets X, Y and ordinals λ , ${}^X Y$ denotes the set of functions from X into Y , and $[X]^\lambda$ denotes the set of subsets of X which are of size λ . Furthermore, id_X denotes the identity function on X . When $n \in \omega$, we denote elements of ${}^n X$ by \bar{x} , and x_i denotes the i^{th} coordinate of \bar{x} . When $f \in {}^X Y$, we use the notation $f(\bar{x})$ for the element of ${}^n Y$ whose i^{th} coordinate is $f(x_i)$.

Unless otherwise mentioned, we assume the set ${}^\kappa \kappa$ is equipped with the κ -Baire topology. However, we will sometimes consider the product topology on the set ${}^\kappa \kappa$, where κ is equipped with the discrete topology; we denote this space by $({}^\kappa \kappa, \tau_{\text{pr}})$.

2. INDEPENDENT SETS OF Σ_2^0 RELATIONS

In this section, we prove the main result of the paper, Theorem 2.4. Some corollaries of this theorem are stated at the end of the section.

Accordingly, we assume throughout this section that the hypothesis $I^-(\kappa)$ holds. We also assume that R is a Σ_2^0 relation on an analytic subset X of the κ -Baire space and that $B \subseteq X$ is an R -independent set of size κ^+ . Our goal is to show that under the additional hypothesis of \diamond_κ or when κ is inaccessible, X has a perfect R -independent subset.

We start by describing the basic idea of the proof in the special case when $X = {}^\kappa \kappa$; the more general version will follow easily from this one. We are going to construct functions $(p_\xi \in {}^{<\kappa} \kappa : \xi \in {}^{<\kappa} 2)$ in such a way that for all $\eta \subseteq \xi \in {}^{<\kappa} 2$ we have that $p_\eta \subseteq p_\xi$ and that $p_{\xi \smallfrown 0}$ and $p_{\xi \smallfrown 1}$ are incomparable; closing the set of these p_ξ 's under initial subsequences will give us a perfect tree T . In the case that \diamond_κ holds or when κ is inaccessible, further conditions can be made on the p_ξ 's to ensure that the κ -branches of T form an R -independent set. In order to be able to construct all the p_ξ 's, we also make sure that the basic open subsets N_{p_ξ} are "big" in the following sense. By the assumption $I^-(\kappa)$, we can fix a κ^+ -complete normal ideal \mathcal{I} on the set B and a subset K of \mathcal{I}^+ which is dense and in which every descending chain of length $< \kappa$ has a lower bound. We will guarantee the existence of sets $B_\xi \in K$ (for all $\xi \in {}^{<\kappa} 2$) such that $B_\xi \subseteq N_{p_\xi}$.

Lemmas 2.1 to 2.3 below are needed to ensure that these sets and functions can be constructed. When stating these lemmas, we assume that B, \mathcal{I} and K are fixed and satisfy the requirements described in the above paragraph. Note again that our assumption $I^-(\kappa)$ implies $\kappa^{<\kappa} = \kappa$.

Lemma 2.1. *Suppose that $p \in {}^{<\kappa} \kappa$ and $B' \in \mathcal{I}^+$ are such that $B' \subseteq N_p$. Then there exist $p_0, p_1 \supseteq p$ such that*

1. $N_{p_0} \cap N_{p_1} = \emptyset$ (i.e., $p_0 \not\subseteq p_1$ and $p_1 \not\subseteq p_0$), and
2. $N_{p_i} \cap B' \in \mathcal{I}^+$ for $i = 0, 1$.

Proof. Assume, seeking a contradiction, that for p and B' as above, no such p_0, p_1 exist. For any $\alpha < \kappa$ such that $\text{dom } p < \alpha$, we have, by $B' \subseteq N_p$, that

$$B' = \bigcup \{N_s \cap B' : p \subset s \in {}^\alpha \kappa\}.$$

Thus, because \mathcal{I} is a κ^+ -complete ideal, $B' \in \mathcal{I}^+$ and $|\alpha \kappa| \leq \kappa$, our assumption implies that there exists exactly one $s_\alpha \in {}^\alpha 2$ extending p such that

$$N_{s_\alpha} \cap B' \in \mathcal{I}^+ \text{ and } B' - N_{s_\alpha} \in \mathcal{I}.$$

Our assumption also implies that we must have $s_\beta \subset s_\alpha$ for all $\text{dom } p < \beta < \alpha < \kappa$, and therefore $s = \bigcup_{\text{dom } p < \alpha < \kappa} s_\alpha$ is an element of ${}^\kappa \kappa$. Then $\bigcap_{\text{dom } p < \alpha < \kappa} N_{s_\alpha} = \{s\}$, and so,

$$B' \subseteq \{s\} \cup \bigcup_{\text{dom } p < \alpha < \kappa} (B' - N_{s_\alpha}) \in \mathcal{I},$$

by the κ^+ -completeness and the non-principality of the ideal \mathcal{I} , contradicting the assumptions of the lemma. \square

In the next two lemmas, $0 < n < \omega$ and S denotes a closed subset of $n({}^\kappa \kappa)$ such that B is S -independent.

Lemma 2.2. *Suppose that $p_0, \dots, p_{n-1} \in {}^{<\kappa} \kappa$ and $B_0, \dots, B_{n-1} \in \mathcal{I}^+$ are such that*

$$B_i \subseteq N_{p_i} \text{ and } p_i \not\subseteq p_j \text{ for all } i, j < n \text{ with } i \neq j.$$

Then there exist $q_0, \dots, q_{n-1} \in {}^{<\kappa} \kappa$ and $B'_0, \dots, B'_{n-1} \in K$ for which

1. $(N_{q_0} \times \dots \times N_{q_{n-1}}) \cap S = \emptyset$,
2. $p_i \subseteq q_i$ for all $i < n$, and
3. $B'_i \subseteq N_{q_i} \cap B_i$ for all $i < n$.

Proof. Take an arbitrary $\bar{x} = (x_0, \dots, x_{n-1}) \in B_0 \times \dots \times B_{n-1}$. The x_i are pairwise distinct elements of the S -independent set B , and therefore $\bar{x} \notin S$. Thus, since S is closed and $x_i \in B_i \subseteq N_{p_i}$, there exists a $\bar{q}(\bar{x}) = (q_0(\bar{x}), \dots, q_{n-1}(\bar{x})) \in n({}^{<\kappa} \kappa)$, such that $p_i \subseteq q_i(\bar{x}) \subseteq x_i$, and

$$N_{q_0(\bar{x})} \times \dots \times N_{q_{n-1}(\bar{x})} \subseteq n({}^\kappa \kappa) - S.$$

Fix $\bar{y} \in B_0 \times \dots \times B_{n-2}$. Since the ideal \mathcal{I} is κ^+ -complete and does not contain B_{n-1} , and also because there are $\kappa^{<\kappa} = \kappa$ many possibilities for the $\bar{q}_i(\bar{x})$'s, there exists $\bar{q}(\bar{y}) \in n({}^{<\kappa} \kappa)$ such that the set

$$A(\bar{y}) = \{x \in B_{n-1} : \bar{q}(\bar{y}, x) = \bar{q}(\bar{y})\} \text{ is in } \mathcal{I}^+.$$

Similarly, using induction on $k \leq n$, we can define, for all $\bar{y} \in B_0 \times \dots \times B_{n-(k+1)}$ elements $\bar{q}(\bar{y})$ of ${}^n(<^\kappa \kappa)$ for which

$$A(\bar{y}) = \{x \in B_{n-k} : \bar{q}(\bar{y}, x) = \bar{q}(\bar{y})\} \text{ is in } \mathcal{I}^+.$$

Finally, for $k = n$, we obtain $\bar{q} = (q_0, \dots, q_{n-1}) \in {}^n(<^\kappa \kappa)$ and $A = \{x \in B_0 : \bar{q}(x) = \bar{q}\}$.

To see that q_0, \dots, q_{n-1} will satisfy the requirements of the lemma, define an element $\bar{y} = (y_0, \dots, y_{n-1}) \in B_0 \times \dots \times B_{n-1}$ such that $\bar{q} = \bar{q}(\bar{y})$, as follows: let $y_0 \in A$ be arbitrary, and if $1 \leq i \leq n-1$ and y_0, \dots, y_{i-1} have been defined, then let y_i be an arbitrary element of $A(y_0, \dots, y_{i-1})$. Then, using that $\bar{q} = \bar{q}(\bar{y})$, we have $p_i \subseteq q_i$ and $N_{q_0} \times \dots \times N_{q_{n-1}} \subseteq {}^n(\kappa \kappa) - S$.

Furthermore, for $0 \leq i \leq n-1$, the set $A(y_0, \dots, y_{i-1})$ is a subset of B_i by definition, and also of N_{q_i} , because of the following: for all $x \in A(y_0, \dots, y_{i-1})$, we can find $\bar{z} \in B_{i+1} \times \dots \times B_{n-1}$ such that

$$\bar{q}(y_0, \dots, y_{i-1}, x, \bar{z}) = \bar{q}(y_0, \dots, y_{i-1}, x) = \bar{q},$$

implying that $q_i \subseteq x$. Therefore, by $A(y_0, \dots, y_{i-1}) \in \mathcal{I}^+$ and the density of K in \mathcal{I}^+ , there exists a $B'_i \in K$ such that $B'_i \subseteq B_i \cap N_{q_i}$. \square

The next lemma will aid us in proving the theorem for Σ_2^0 relations in the case κ is inaccessible.

Lemma 2.3. *Suppose $\mu < \kappa$ is a cardinal, and $(p_\alpha \in {}^{<\kappa}\kappa : \alpha < \mu)$ and $(B_\alpha \in \mathcal{I}^+ : \alpha < \mu)$ are such that*

$$B_\alpha \subseteq N_{p_\alpha} \text{ and } p_\alpha \not\subseteq p_\beta \text{ for all } \alpha, \beta < \mu \text{ with } \alpha \neq \beta.$$

Then there exist, for all $\alpha < \mu$, elements $q_\alpha \in {}^{<\kappa}\kappa$ and $B'_\alpha \in K$ for which

1. $(N_{q_{\alpha_0}} \times \dots \times N_{q_{\alpha_{n-1}}}) \cap S = \emptyset$ for all pairwise different $\alpha_0, \dots, \alpha_{n-1} < \mu$,
2. $p_\alpha \subseteq q_\alpha$ for all $\alpha < \mu$, and
3. $B'_\alpha \subseteq N_{q_\alpha} \cap B_\alpha$ for all $\alpha < \mu$.

Proof. To prove this lemma, we basically iterate the application of the previous one as many times as necessary (being careful that the iteration goes through at limit stages); the details are below. To simplify notation, we assume $n = 2$.

Let $((\alpha_\sigma, \beta_\sigma) : \sigma < \mu')$ be an enumeration of $\{(\alpha, \beta) : \alpha, \beta \in \mu, \alpha \neq \beta\}$, where $\mu' = \mu$ in the case that $\mu \geq \omega$, and $\mu' = \mu^2 - \mu$ for $\mu < \omega$. By transfinite induction for $\sigma \leq \mu'$, we define

$$p_\alpha^\sigma \in {}^{<\kappa}\kappa \text{ and } B_\alpha^\sigma \in K \text{ for all } \alpha < \mu \text{ simultaneously,}$$

in such a way that the three properties below are true for all $\alpha < \mu$ and $\tau < \sigma \leq \mu'$. For simplicity, we use the notation $P_\alpha^\sigma = N_{p_\alpha^\sigma}$.

- (i.) $(P_{\alpha_\tau}^{\tau+1} \times P_{\beta_\tau}^{\tau+1}) \cap S = \emptyset$.
- (ii.) $p_\alpha \subseteq p_\alpha^\tau \subseteq p_\alpha^\sigma$ and $B_\alpha \supseteq B_\alpha^\tau \supseteq B_\alpha^\sigma$,
- (iii.) $B_\alpha^\sigma \subseteq P_\alpha^\sigma$.

This is enough, because once these sequences have been defined, $q_\alpha = p_\alpha^{\mu'}$ and $B'_\alpha = B_\alpha^{\mu'}$ will satisfy the conclusions of the lemma. To see that item 1 of the conclusion of the lemma is satisfied, note that we have $N_{q_\alpha} = P_\alpha^{\mu'} \subseteq P_\alpha^{\tau+1}$ for all $\alpha < \mu, \tau < \mu'$, and use property (i.) above.

To build the sequences, we start from $p_\alpha^0 = p_\alpha$ (and $P_\alpha^0 = N_{p_\alpha}$) and some $B_\alpha^0 \in K$ such that $B_\alpha^0 \subseteq B_\alpha$. For limit ordinals σ , we define

$$p_\alpha^\sigma = \bigcup_{\tau < \sigma} p_\alpha^\tau \text{ (and thus } P_\alpha^\sigma = \bigcap_{\tau < \sigma} P_\alpha^\tau \text{)}.$$

Then, by $B_\alpha^\tau \in K$ and the closure property of K , there exists $B_\alpha^\sigma \in K$ such that $B_\alpha^\sigma \subseteq \bigcap_{\tau < \sigma} B_\alpha^\tau$. Therefore, using that $B_\alpha^\tau \subseteq P_\alpha^\tau$ for all $\tau < \sigma$ and the definition of p_α^σ , we also have $B_\alpha^\sigma \subseteq P_\alpha^\sigma$.

In the case of a successor ordinal $\sigma + 1$, we define the elements $p_{\alpha_\sigma}^{\sigma+1}, p_{\beta_\sigma}^{\sigma+1}$ of ${}^{<\kappa}\kappa$ and the sets $B_{\alpha_\sigma}^{\sigma+1}, B_{\beta_\sigma}^{\sigma+1}$ to be those obtained from an application of Lemma 2.2 to the functions $p_{\alpha_\sigma}^\sigma, p_{\beta_\sigma}^\sigma$ and the sets $B_{\alpha_\sigma}^\sigma, B_{\beta_\sigma}^\sigma$. For $\alpha_\sigma \neq \gamma \neq \beta_\sigma$, we leave everything unchanged, i.e., we let $p_\gamma^{\sigma+1} = p_\gamma^\sigma$ and $B_\gamma^{\sigma+1} = B_\gamma^\sigma$. Then these definitions will satisfy all three required properties. \square

We now state and prove the main result of our paper.

Theorem 2.4. *Assume $I^-(\kappa)$ and either that \diamond_κ or that κ is inaccessible.*

Suppose R is a Σ_2^0 relation on an analytic subset of the κ -Baire space ${}^\kappa\kappa$. Then either all R -independent sets are of size $\leq \kappa$, or there exists a perfect R -independent set.

We note that, as was mentioned in the Introduction, $I^-(\kappa)$ implies \diamond_κ for successor cardinals $\kappa > \aleph_1$. Furthermore, when R is a closed binary relation on X , the above dichotomy follows from $I^-(\kappa)$ alone; see Remark 2.5 below.

Proof. As throughout the rest of this section, we assume $I^-(\kappa)$. We start by proving the theorem in the case that R is a Σ_2^0 relation on the whole space ${}^\kappa\kappa$. Suppose R is n -ary (where $0 < n < \omega$), and that $(R_\alpha : \alpha < \kappa)$ are closed subsets of ${}^n({}^\kappa\kappa)$ such that $R = \bigcup_{\alpha < \kappa} R_\alpha$. Let B be an independent set of R of size κ^+ . By the hypothesis $I^-(\kappa)$, there is a κ^+ -complete normal ideal \mathcal{I} on B and a subset K of \mathcal{I}^+ which is dense and in which every descending chain of length $< \kappa$ has a lower bound.

First, assume \diamond_κ . Then, the following combinatorial principle also holds:

there exists a sequence of n -tuples

$$((x_\alpha^0, \dots, x_\alpha^{n-1}) \in {}^n(\alpha 2) : \alpha < \kappa)$$

such that $x_\alpha^i \neq x_\alpha^j$ for all $i < j < n$, $\alpha < \kappa$, and for all $(x_0, \dots, x_{n-1}) \in {}^n(\kappa 2)$ with $x_i \neq x_j$ for all $i < j < n$ we have that the set $\{\alpha \in \kappa : x_\alpha^i = x_i \upharpoonright \alpha \text{ for all } i < n\}$ is cofinal in κ .

We note that whenever κ is not inaccessible, the above combinatorial principle is equivalent to \diamond_{κ} , by the Proposition in [16]. (See also [2] and [20], from which this equivalence follows when $\kappa = \omega_1$ and when κ is a successor cardinal, respectively.)

We define sets $(B_\xi \in K : \xi \in {}^{<\kappa}2)$ and functions $(p_\xi \in {}^{<\kappa}\kappa : \xi \in {}^{<\kappa}2)$ in such a way that the items below are satisfied for all $\xi, \eta \in {}^{<\kappa}2$ and $\alpha < \kappa$. We use the notation $P_\xi = N_{p_\xi}$.

1. $B_\xi \subseteq P_\xi$;
2. if $\eta \subseteq \xi$, then $B_\eta \supseteq B_\xi$ and $p_\eta \subseteq p_\xi$;
3. if $\xi \neq \eta \in {}^\alpha 2$, then $P_\xi \cap P_\eta = \emptyset$ (i.e., p_ξ and p_η are incomparable);
4. for all $\gamma < \alpha$ we have

$$(P_{x_\alpha^0} \times \dots \times P_{x_\alpha^{n-1}}) \cap R_\gamma = \emptyset$$

(i.e., for any $\bar{t} \in {}^n(\kappa\kappa)$ such that $t_i \supseteq p_{x_\alpha^i}$ for all $i < n$, we have $\bar{t} \notin R_\gamma$).

By the second and third conditions, the map $t : {}^\kappa 2 \rightarrow {}^\kappa \kappa$; $x \mapsto \bigcup_{\alpha < \kappa} p_{x \upharpoonright \alpha}$ is a continuous injection whose image $Im(t)$ is a perfect subset of ${}^\kappa \kappa$. Furthermore, item 4 guarantees that $Im(t)$ will be R -independent: if $x_0, \dots, x_{n-1} \in {}^\kappa 2$ are pairwise distinct and $\gamma < \kappa$ is arbitrary, then there exists a $\gamma < \alpha < \kappa$ such that $x_\alpha^i = x_i \upharpoonright \alpha$ for all $i < n$, and therefore, by 4, we have that $(t(x_0), \dots, t(x_{n-1})) \notin R_\gamma$.

Thus, it is enough to see that sets B_ξ and functions p_ξ satisfying the conditions above can indeed be constructed. To guarantee the first three conditions, we will use Lemma 2.1 and the density of K at successor stages, and we will use the closure property of $K \subseteq \mathcal{I}^+$ at limit stages. Then, the fourth property can be guaranteed using Lemma 2.2. The details are below.

Let $p_\emptyset = \emptyset$ and let $B_\emptyset \in K$ be arbitrary. Now, fix $\alpha < \kappa$ and suppose that p_η and B_η have been defined for all $\eta \in {}^{<\alpha}2$ so that conditions 1 to 4 hold. We first construct elements $p'_\xi \in {}^{<\kappa}\kappa$ and $B'_\xi \in K$, for all $\xi \in {}^\alpha 2$ which satisfy the first three required conditions, i.e., for all $\xi, \eta \in {}^\alpha 2$ with $\xi \neq \eta$,

$$p'_\xi \not\subseteq p'_\eta \text{ and } B'_\xi \subseteq N_{p'_\xi},$$

and for all $\beta < \alpha$ we have $p'_\xi \supseteq p_{\xi \upharpoonright \beta}$ and $B'_\xi \subseteq B_{\xi \upharpoonright \beta}$.

If $\alpha = \beta + 1$, then, for each $\eta \in {}^\beta 2$, apply Lemma 2.1 for p_η and B_η to obtain $p'_{\eta \smallfrown 0}$ and $p'_{\eta \smallfrown 1}$. By the density of K in \mathcal{I}^+ , we can find $B'_{\eta \smallfrown 0}, B'_{\eta \smallfrown 1} \in K$ such that $B'_{\eta \smallfrown i} \subseteq N_{p'_{\eta \smallfrown i}} \cap B_\eta$ for $i = 0, 1$. If α is a limit ordinal, then for all $\xi \in {}^\alpha 2$, let $p'_\xi = \bigcup_{\beta < \alpha} p_{\xi \upharpoonright \beta}$, and, using the closure property of K , choose sets $B'_\xi \in K$ such that $B'_\xi \subseteq \bigcap_{\beta < \alpha} B_{\xi \upharpoonright \beta}$. We have $B'_\xi \subseteq N_{p'_\xi}$ because $B_{\xi \upharpoonright \beta} \subseteq P_{\xi \upharpoonright \beta}$ for all $\beta < \alpha$ and by the definition of p'_ξ .

After p'_ξ and B'_ξ have been constructed for all $\xi \in {}^\alpha 2$, we apply Lemma 2.2 for the closed subset $\bigcup_{\gamma < \alpha} R_\gamma$ of ${}^n({}^\kappa \kappa)$ and for $p'_{x_\alpha^0}, \dots, p'_{x_\alpha^{n-1}}$ and $B'_{x_\alpha^0}, \dots, B'_{x_\alpha^{n-1}}$ in order to obtain the elements $p_{x_\alpha^0}, \dots, p_{x_\alpha^{n-1}}$ and $B_{x_\alpha^0}, \dots, B_{x_\alpha^{n-1}}$. In the case that $\xi \notin \{x_\alpha^0, \dots, x_\alpha^{n-1}\}$, we leave everything unchanged, i.e., we let $p_\xi = p'_\xi$ and $B_\xi = B'_\xi$. This choice of the p_ξ 's and B_ξ 's will guarantee that all four required properties are satisfied.

Now, assume that κ is inaccessible. We modify the previous argument in every step of the construction, by requiring, instead of condition 4, that the following property is satisfied for all $\alpha < \kappa$:

4'. for all pairwise distinct $\xi_0, \dots, \xi_{n-1} \in {}^\alpha 2$ and $\gamma < \alpha$, we have

$$(P_{\xi_0} \times \dots \times P_{\xi_{n-1}}) \cap R_\gamma = \emptyset.$$

This property will ensure that the perfect set $Im(t)$ is an independent set of R_γ for each $\gamma < \kappa$ and therefore of R . Condition 4' can be guaranteed in the α^{th} step of the construction by applying Lemma 2.3 for the closed subset $\bigcup_{\gamma < \alpha} R_\gamma$ of ${}^n({}^\kappa \kappa)$ and for the collections $(p'_\xi : \xi \in {}^\alpha 2)$ and $(B'_\xi : \xi \in {}^\alpha 2)$ of the previous argument; note that $|{}^\alpha 2| < \kappa$ by the inaccessibility of κ . This completes the proof of the theorem in the case that R is a Σ_2^0 relation on the whole space ${}^\kappa \kappa$.

Now, assume that R is an n -ary Σ_2^0 relation on a closed subset X of the κ -Baire space ${}^\kappa \kappa$. Then $R' = R \cup ({}^n({}^\kappa \kappa) - {}^n X)$ is an n -ary Σ_2^0 relation on ${}^\kappa \kappa$ (note that the assumption $\kappa^{< \kappa} = \kappa$ implies that all open subsets of ${}^\kappa \kappa$ are also Σ_2^0). Furthermore, the R' -independent sets of size at least n are subsets of X and are therefore also R -independent. Hence, this case follows by applying the previous case for R' .

Finally, suppose that X is an arbitrary analytic subset of ${}^\kappa \kappa$ and R is an n -ary Σ_2^0 relation on X . We can assume that $\Delta_n = \{\bar{x} \in {}^n X : x_i = x_j \text{ for some } i < j < n\}$ is a subset of R (we can take $R \cup \Delta_n$ instead of R if necessary, because Δ_n is a closed relation on X and because, by definition, a set $Y \subseteq X$ is $(R \cup \Delta_n)$ -independent iff it is R -independent).

Because X is analytic, there exists a closed subset X'' of ${}^\kappa \kappa$ and a continuous function $f : {}^\kappa \kappa \rightarrow {}^\kappa \kappa$ such that $X = f[X'']$. Denoting by $f(\bar{x})$ the

vector $(f(x_0), \dots, f(x_{n-1}))$ (for $\bar{x} \in {}^n X$), we let

$$R'' = \{\bar{x} \in {}^n X'' : f(\bar{x}) \in R\}.$$

That is, R'' is the inverse image of R under the continuous function ${}^n X'' \longrightarrow {}^n X$; $\bar{x} \mapsto f(\bar{x})$, and is therefore a Σ_2^0 n -ary relation on X'' .

If B is an R -independent set of size κ^+ , then any $B'' \subseteq X''$ such that $f[B''] = B$ and $f \upharpoonright B''$ is injective is an R'' -independent set (by the definition of R'' and the injectivity of $f \upharpoonright B''$), and B'' has cardinality κ^+ . Therefore, by the previous case, R'' has a perfect independent set, i.e., there is a continuous injection t of ${}^\kappa 2$ into X'' whose image is R'' -independent. Then, by the definition of R'' and because $\Delta_n \subseteq R$, the function $f \circ t$ is a continuous injection of ${}^\kappa 2$ into X whose image is R -independent, and thus, X has a perfect R -independent subset. \square

Remark 2.5. When R is a *closed binary* relation on an analytic subset X of the κ -Baire space, the dichotomy in Theorem 2.4 already follows from the hypothesis $I^-(\kappa)$, even when neither \diamond_κ nor the inaccessibility of κ is assumed. This can be seen by a slight modification of the proof of Theorem 2.4, which we give below.

We first describe the proof for a closed subset X of the κ -Baire space; the more general case of analytic subsets follows just as in the proof of Theorem 2.4. (However, the argument for obtaining the statement for closed sets X from the $X = {}^\kappa \kappa$ case no longer works, since $({}^\kappa \kappa \times {}^\kappa \kappa) - (X \times X)$ is not necessarily closed when X is closed.)

Suppose that R is a closed subset of $X \times X$ and $B \subseteq X$ is an R -independent set of size κ^+ , and let \mathcal{I} and K be as in the proof of Theorem 2.4. As before, we define, for all $\xi \in {}^{<\kappa} 2$, functions $p_\xi \in {}^{<\kappa} \kappa$ and sets $B_\xi \in K$ satisfying certain conditions. We require items 1 to 3 of the proof of Theorem 2.4; these conditions can be guaranteed exactly as in that proof. For R -independence, it is enough this time to ensure, using Lemma 2.2 (for $n = 2$), that

$$4''. (P_{\xi \smallfrown 0} \times P_{\xi \smallfrown 1}) \cap R = \emptyset \text{ for all } \xi \in {}^{<\kappa} 2$$

(where $P_\nu = N_{p_\nu}$). That the κ -branches defined by two distinct $x, y \in {}^\kappa 2$ will not be R -related follows from applying item 4'' to the node $\xi \in {}^{<\kappa} 2$ at which x and y split. The sufficiency of item 4'' is the reason the extra assumption of either \diamond_κ or the inaccessibility of κ is unnecessary.

We have to add a fifth condition to guarantee that the perfect set defined by the p_ξ 's is a subset of X . Since X is closed and $\kappa^{<\kappa} = \kappa$ holds, there exist open sets X_α such that $X = \bigcap_{\alpha < \kappa} X_\alpha$. We require that

5. $P_\xi \subseteq X_\alpha$ for all $\alpha < \kappa$ and $\xi \in {}^\alpha 2$.

Suppose $\xi \in {}^\alpha 2$ and a p'_ξ and B'_ξ has been defined satisfying items 1-3 and 4''. Then $B'_\xi \subseteq B \subseteq X \subseteq X_\alpha$, and so, (using item 1) we have $B'_\xi \subseteq N_{p'_\xi} \cap X_\alpha$. Therefore, using the facts that $B'_\xi \in \mathcal{I}^+$, that \mathcal{I} is κ^+ -complete and that $N_{p'_\xi} \cap X_\alpha$ is the union of at most κ many basic open sets, we can find an extension $p_\xi \supseteq p'_\xi$ such that $N_{p_\xi} \subseteq X_\alpha$ and $N_{p_\xi} \cap B'_\xi \in \mathcal{I}^+$. By the density of K in \mathcal{I}^+ , there exists $B_\xi \in K$ which is a subset of $N_{p_\xi} \cap B'_\xi$, and thus, p_ξ and B_ξ satisfy all 5 required conditions. This completes the proof of the case when X is a closed subset of the κ -Baire space.

To obtain from this case the dichotomy for closed binary relations R on arbitrary analytic subsets X of ${}^\kappa \kappa$, we employ exactly the same argument (for $n = 2$) as the one that was used at end of the proof of Theorem 2.4 to show the corresponding implication for Σ_2^0 relations. Note that, using the notation at the end of the proof of Theorem 2.4, R'' is now a closed binary relation on a closed subset X'' of ${}^\kappa \kappa$, because it is the inverse image of the closed subset R of $X \times X$ under a continuous function.

Remark 2.6. The special case of the dichotomy in Theorem 2.4 for the empty binary relation $R = \emptyset$ is that the κ -perfect set property holds for closed (and even analytic) subsets of the κ -Baire space, i.e., any closed (analytic) subset is either of size $\leq \kappa$ or contains a perfect subset. (In fact, by a generalization of [27, Theorem 1], or as a special case of Remark 2.5, the κ -perfect set property for closed sets follows already from the assumption $\Gamma^-(\kappa)$.)

This implies that our dichotomy does not hold for κ if there exists a weak κ -Kurepa tree, i.e., a tree of height κ with at least κ^+ many κ -branches (of length κ) whose α^{th} level has size $\leq |\alpha|$ for stationarily many $\alpha \in \kappa$. If T is a weak κ -Kurepa tree, then the set $[T]$ of κ -branches of T is a closed subset of the κ -Baire space of size $\geq \kappa^+$ that does not contain a perfect subset; see [5, Section 4] and [4]. The idea of using Kurepa trees to obtain counterexamples to the \aleph_1 -perfect set property had already appeared in [27] and [18].

We therefore have the following two facts. Firstly, in $V = L$, our dichotomy does not hold for any uncountable regular κ , by [4, Lemma 4]. Secondly, our dichotomy for κ implies, by a result of Robert Solovay [10, Sections 3 and 4], that κ^+ is an inaccessible cardinal in L . Note that by Theorem 2.4, the consistency of our dichotomy follows from the existence of a measurable $\lambda > \kappa$; however, we do not yet know its exact consistency strength.

As mentioned in the Introduction, Theorem 2.4 can also be seen as the uncountable version of a result [12, Corollary 2.13] about homogeneous subsets of G_δ colorings on analytic spaces. A *coloring* of a set X is any subset C of $[X]^n$ for some $1 < n \in \omega$, and a set $B \subseteq X$ is *C-homogeneous* if $[B]^n \subseteq C$. Notice that $[X]^n$ can be identified with an open subset of ${}^n X$, namely $\{\bar{x} \in {}^n X : x_i \neq x_j \text{ for all } i < j < n\}$. Thus, when X is an analytic subset of the κ -Baire space, we may speak about $\mathbf{\Pi}_2^0$ colorings on X . The “complement” of such colorings can be considered as Σ_2^0 relations on X , and furthermore, the concept of homogeneity translates to that of independence. Hence, Theorem 2.4 yields the following.

Corollary 2.7. *Assume $\Gamma^-(\kappa)$ and either that \diamond_κ or that κ is inaccessible.*

Suppose that C is a $\mathbf{\Pi}_2^0$ coloring on an analytic subset of ${}^\kappa \kappa$. Then either all C -homogeneous sets are of size $\leq \kappa$, or there is a perfect C -homogeneous set.

In the remainder of the section, we state some corollaries of the $n = 2$ case of Theorem 2.4.

Corollary 2.8. *Assume $\Gamma^-(\kappa)$ and either that \diamond_κ or that κ is inaccessible.*

Then the κ -Silver dichotomy holds for Σ_2^0 equivalence relations on analytic subsets of ${}^\kappa \kappa$.

We use the following notation in the next corollary. For H a topological space, X any set and $S \subseteq H \times X \times X$, let R_S be the projection of S onto $X \times X$, i.e.,

$$R_S = \{(x, y) : (h, x, y) \in S \text{ for some } h \in H\}.$$

Specifically, for the action a of a group H on X , R_a is the orbit equivalence relation.

Recall from the introduction that a topological space is κ -compact if any of its open covers has a subcover of size $< \kappa$, and is K_κ , if it can be written as the union of at most κ many κ -compact subsets. A topological group is K_κ if it is K_κ as a topological space. The first item of the next corollary is stated because we will need to use this more general form later.

Corollary 2.9. *Assume $\Gamma^-(\kappa)$ and either that \diamond_κ or that κ is inaccessible.*

Let X be an analytic subset of the κ -Baire space (equipped with the subspace topology) and let H be an arbitrary K_κ topological space.

1. If $S \subseteq H \times X \times X$ is closed, then either all R_S -independent sets have size $\leq \kappa$ or there is a perfect R_S -independent set.
2. If H is a group that acts continuously on X , then there are either $\leq \kappa$ many or perfectly many orbits.

Proof. A generalization of a standard argument from the countable case [8, Exercise 3.4.2] shows that if H is a κ -compact topological space, then R_S is a closed subset of $X \times X$. In more detail, let $(x, y) \in X \times X - R_S$ be arbitrary. Because S is closed, we can choose for all $h \in H$ open sets $U_h \subseteq H$ and $V_h \subseteq X \times X$ such that

$$(h, x, y) \in U_h \times V_h \subseteq H \times X \times X - S.$$

By the κ -compactness of H , there exists a set $I \in [H]^{<\kappa}$ such that $H = \bigcup_{h \in I} U_h$. Then $V = \bigcap_{h \in I} V_h$ is an open subset of $X \times X$ such that $(x, y) \in V \subseteq X \times X - R_S$.

Thus R_S is indeed closed in the case H is κ -compact, implying that if H is K_κ , then R_S is a Σ_2^0 binary relation on X . An application of Theorem 2.4 completes the proof of 1, of which the second item is a special case. \square

Remark 2.10. In the rest of the paper, we will be considering the special cases of K_κ topological spaces obtained by endowing K_κ subsets of the κ -Baire space with the subspace topology (induced by the κ -Baire topology). We also consider K_κ subsets of the product space $({}^\kappa\kappa, \tau_{\text{pr}})$ equipped with the subspace topology induced by τ_{pr} . (Recall from the introduction that τ_{pr} is the product topology on the set ${}^\kappa\kappa$ obtained by endowing κ with the discrete topology.) We therefore give characterizations of, or some examples for, such sets below.

Notice that K_κ subsets of the κ -Baire space are always K_κ subsets of $({}^\kappa\kappa, \tau_{\text{pr}})$ as well, due to the fact that the κ -Baire topology on the set ${}^\kappa\kappa$ is finer than the product topology τ_{pr} .

When κ is a weakly compact cardinal, the converse also holds; in this case, the following are equivalent for any $H \subseteq {}^\kappa\kappa$.

1. H is a K_κ subset of the κ -Baire space.
2. H is a K_κ subset of the product space $({}^\kappa\kappa, \tau_{\text{pr}})$.
3. H is a Σ_2^0 subset of the κ -Baire space which is *eventually bounded* (this means, by definition, that there exists an $x \in {}^\kappa\kappa$ such that for all $h \in H$, we have $h \leq^* x$, i.e., there exists an $\alpha < \kappa$ such that $h(\beta) \leq x(\beta)$ for all $\alpha \leq \beta < \kappa$).

The equivalence of the first and third items follows from [15, Lemma 2.6], and, as just noted, the second item is implied by the first one.

The third item can be obtained from the second using the fact that a κ -compact subset C of $({}^\kappa\kappa, \tau_{\text{pr}})$ is closed in the κ -Baire topology and is bounded (i.e., there exists an $x \in {}^\kappa\kappa$ such that $c(\beta) \leq x(\beta)$ for all $c \in C$ and $\beta < \kappa$). The above fact can be proven by modifying standard arguments from the countable case (see [11, Exercise 4.11]). In more detail, suppose that $C \subseteq {}^\kappa\kappa$ is κ -compact in the product topology τ_{pr} . If we take any $\alpha < \kappa$ (and we denote by $N_{\{(\alpha, \gamma)\}}$ the set $\{y \in {}^\kappa\kappa : y(\alpha) = \gamma\} \in \tau_{\text{pr}}$ for all $\gamma < \kappa$), the family

$$\{N_{\{(\alpha, \gamma)\}} : \gamma < \kappa \text{ and } C \cap N_{\{(\alpha, \gamma)\}} \neq \emptyset\}$$

is a disjoint τ_{pr} -open cover of C , and must therefore be of size $< \kappa$. Thus, we can define $x \in {}^\kappa\kappa$ by letting $x(\alpha) = \sup\{\gamma < \kappa : C \cap N_{\{(\alpha, \gamma)\}} \neq \emptyset\}$ for all $\alpha < \kappa$, and x witnesses that C is bounded.

To see that C is closed in the κ -Baire topology, let $z \in {}^\kappa\kappa - C$. We can choose, for all $y \in C$, disjoint neighborhoods $U_y, V_y \in \tau_{\text{pr}}$ of z and y respectively. By the κ -compactness of C , the τ_{pr} -open cover $\{V_y : y \in C\}$ of C can be refined to a subcover $\{V_y : y \in I\}$ of size $< \kappa$. Then the intersection $U = \bigcap_{y \in I} U_y$ is disjoint from C , contains z , and is open in the κ -Baire topology (because the κ -Baire topology is finer than τ_{pr} and is closed under intersections of size $< \kappa$).

Now, consider the case when $\kappa^{<\kappa} = \kappa$ is not weakly compact. If $I^-(\kappa)$ holds, the K_κ subsets of the κ -Baire space are exactly those of size at most κ . This is because by [15, Corollary 2.8], the above equivalence holds when $\kappa^{<\kappa} = \kappa$ is not weakly compact and the closed subsets of the κ -Baire space have the κ -perfect set property (this latter requirement is implied by $I^-(\kappa)$; see Remark 2.6). Examples of subsets H of the set ${}^\kappa\kappa$ which are κ -compact subsets of the *product space* $({}^\kappa\kappa, \tau_{\text{pr}})$ but are not K_κ subsets of the κ -Baire space, even when $I^-(\kappa)$ is not assumed, include by [15, Lemma 2.2] (and Tychonoff's theorem) the set $H = {}^\kappa 2$ and, more generally, sets of the form $H = \prod_{\alpha < \kappa} I_\alpha$, where $I_\alpha \in [\kappa]^{<\kappa}$ and I_α is finite except for $< \kappa$ many $\alpha < \kappa$. (See [14] and the references therein for when this last assumption can be weakened.)

Remark 2.11. In light of the previous remark, especially the fact that K_κ subsets of the κ -Baire space are of size $\leq \kappa$ when κ is not weakly compact and $I^-(\kappa)$ holds, we mention the following observation.

A model of ZFC in which κ is weakly compact (in fact, supercompact) and $I^-(\kappa)$ holds can be obtained, starting out from a situation in which κ is supercompact and there exists a measurable $\lambda > \kappa$, in the following way. Before Lévy-collapsing λ to κ^+ , one first applies the Laver preparation [13] to

make the supercompactness of κ indestructible by any $< \kappa$ -directed closed forcing. If we would also like to have $2^\kappa > \kappa^+$ together with $I^-(\kappa)$ for a supercompact κ , we force after the Laver-preparation with the product of the Lévy-collapse and $Add(\kappa, \mu)$ for some $\mu > \lambda$. We would like to thank Menachem Magidor for suggesting the arguments found in this remark.

3. ELEMENTARY EMBEDDABILITY ON MODELS OF SIZE κ

In this section, we use our previous results to obtain dichotomy properties for the set of models with domain κ of some $\Sigma_1^1(L_{\kappa+\kappa})$ -sentence, when considered up to isomorphism, embeddability or elementary embeddability by elements of some K_κ subset H of the κ -Baire space. More generally, we will be interested in these models up to maps (in H) that preserve certain subsets of $L_{\kappa+\kappa}$ -formulas, which we will call *fragments* (see Definition 3.1 below). In the case of some fragments, it will be enough to assume that H is a K_κ subset of the product space $({}^\kappa\kappa, \tau_{\text{pr}})$; (recall from the introduction that τ_{pr} is the product topology on the set ${}^\kappa\kappa$, where κ is given the discrete topology).

Notation. We will use the following notation in this section. As usual, $Sym(\kappa)$ denotes the permutation group of κ , and we write $Inj(\kappa)$ for the monoid of all injective functions of ${}^\kappa\kappa$.

The symbol L denotes a fixed first order language which contains only relation symbols and is of size at most κ . However, the arguments below also work in the case of languages which have infinitary relations of arity $< \kappa$. We assume the language L has κ many variables, the sequence of which is denoted by $(v_i : i \in \kappa)$. The symbols \mathcal{A} , \mathcal{B} , etc. are used to denote L -structures whose domains are A , B , etc. The set of all L -structures with domain κ is denoted by Mod_κ^L . Given a structure $\mathcal{A} \in \text{Mod}_\kappa^L$, we identify \mathcal{A} -valuations with elements of ${}^\kappa\kappa$.

As usual, $L_{\lambda\mu}$ (where $\omega \leq \mu \leq \lambda \leq \kappa^+$ and $\mu \leq \kappa$) denotes the infinitary language which allows conjunctions and disjunctions of $< \lambda$ many formulas and quantification over $< \mu$ many variables. See, e.g., Definitions 1.1.2 and 1.1.3 of [3] for the precise definition of $L_{\lambda\mu}$ -formulas or, alternatively, Definitions 9.12 and 9.13 of [29]. In particular, note that by definition, an $L_{\lambda\mu}$ -formula contains $< \mu$ many free variables, from $\{v_i : i \in \kappa\}$. The concept of the *subformulas* of a formula $\varphi \in L_{\kappa+\kappa}$ is defined by induction on the complexity of φ as usual (see, e.g., Definition 1.3.1 of [3] or [29, p. 234]. Note that if φ is obtained as $\varphi = \bigwedge \Phi$, then any $\phi \in \Phi$ is defined to be a subformula of φ , but $\bigwedge \Phi'$, where $\Phi' \subset \Phi$, is not a subformula).

For $\varphi \in L_{\kappa+\kappa}$ and $h \in {}^\kappa\kappa$, we denote by $s_h\varphi$ the formula obtained from φ by simultaneously substituting, for all $i \in \kappa$, the variable $v_{h(i)}$ for the variable v_i . We say that a set F of $L_{\kappa+\kappa}$ -formulas is *closed under substitution* if for any $\varphi \in F$ and $h \in {}^\kappa\kappa$ we have $s_h\varphi \in F$. Note that since $\kappa^{<\kappa} = \kappa$, closing a nonempty set of formulas of $L_{\kappa+\kappa}$ of size $\leq \kappa$ under substitution leads to a set of formulas of size κ .

Definition 3.1. We define a subset F of the infinitary language $L_{\kappa+\kappa}$ to be a *fragment of $L_{\kappa+\kappa}$* if $|F| = \kappa$,

1. F contains all atomic formulas,
2. F is closed under negation and taking subformulas, and
3. F is closed under substitution of variables.

Definition 3.2. Suppose F is a fragment of $L_{\kappa+\kappa}$, $H \subseteq \text{Inj}(\kappa)$, and $\mathcal{A}, \mathcal{B} \in \text{Mod}_\kappa^L$. We say that \mathcal{A} is (F, H) -*elementarily embeddable* into \mathcal{B} iff there is a map $h \in H$ such that

- (1) $\mathcal{A} \models \varphi[a]$ iff $\mathcal{B} \models \varphi[h \circ a]$ for all $\varphi \in F$ and valuations $a \in {}^\kappa\kappa$.

Note that any function $h \in {}^\kappa\kappa$ satisfying (1), where F is any fragment, must be an embedding of \mathcal{A} into \mathcal{B} (by item 1 of Definition 3.1), and in particular, we must have $h \in \text{Inj}(\kappa)$. This is the reason that we have chosen to define the above concept only for subsets H of $\text{Inj}(\kappa)$ (instead of arbitrary subsets of ${}^\kappa\kappa$). We remark that if H is a *submonoid* of $\text{Inj}(\kappa)$, then the binary relation of (F, H) -elementary embeddability on Mod_κ^L is a partial order (but there is no reason for this to hold when H is an arbitrary subset of $\text{Inj}(\kappa)$).

Example 3.3. Many interesting notions are special cases of (F, H) -elementary embeddability, including

- H -embeddability (when F is the set At of atomic formulas and their negations),
- H -elementary embeddability (when $F = L_{\omega\omega}$), and
- H -isomorphism, when H is a subgroup of $\text{Sym}(\kappa)$ (and F is any fragment. Two models in Mod_κ^L are defined to be *H -isomorphic* iff there exists a $h \in H$ which is an isomorphism between them.) Note that H -isomorphism is an equivalence relation on Mod_κ^L when H is a subgroup of $\text{Sym}(\kappa)$.

Other maps that may be of interest are

- those obtained when $F = L_{\lambda\mu}$ (where $\omega \leq \mu \leq \lambda \leq \kappa$), or
- those preserving the n -variable fragment $L_{\lambda\omega}^n$ of this logic. (By definition, the n variable fragment of $L_{\lambda\mu}$, or equivalently of $L_{\lambda\omega}$, consists

of those formulas which use only the variables v_0, \dots, v_{n-1} . In this case, the corresponding fragment F is the set of those $L_{\lambda\omega}$ -formulas which contain at most n (arbitrary) variables from $\{v_i : i \in \kappa\}$. It is this fragment F that we will denote by $L_{\lambda\omega}^n$.

A fragment $F \subseteq L_{\kappa+\kappa}$ induces a topology t_F on the set Mod_{κ}^L in a natural way. To a formula φ and a valuation $a \in {}^{\kappa}\kappa$, we correlate the basic clopen set $\text{Mod}_{\kappa}(\varphi, a) = \{\mathcal{A} \in \text{Mod}_{\kappa}^L : \mathcal{A} \models \varphi[a]\}$. The topology t_F on Mod_{κ}^L is obtained by taking arbitrary unions of intersections of $< \kappa$ many sets from the collection

$$b_F = \{\text{Mod}_{\kappa}(\varphi, a) : \varphi \in F, a \in {}^{\kappa}\kappa\},$$

and we denote by $\mathcal{M}od_F$ the topological space $(\text{Mod}_{\kappa}^L, t_F)$. The canonical topological space used to study the connections between model theory and the generalized Baire space is $\mathcal{M}od_{\text{At}}$, and it is homeomorphic to the Cantor space ${}^{\kappa}2$; see [18, 28, 5]. An advantage of working with t_F instead of t_{At} is that (F, H) -elementary embeddability induces a “ t_F -continuous action” of H on $\text{Mod}_{\kappa}^{\psi}$ (in the more general sense of item 1 of Corollary 2.9; see the proof of Theorem 3.8 below). The above fact is needed in order for us to be able to use the results of the previous section.

Firstly, we show that for an arbitrary fragment F , the space $\mathcal{M}od_F$ is homeomorphic to a $\mathbf{\Pi}_2^0$ subset X_F of ${}^{\kappa}2$. Our proof is basically a generalization from the countable case of a proof in [19]. To start, note that a bijection between κ and F allows us to define the generalized Cantor topology on ${}^F 2$. In fact, since κ is regular (by $\kappa^{<\kappa} = \kappa$), another basis for this topology is $\{N_p : p \in {}^{\Phi}2 \text{ for some } \Phi \in [F]^{<\kappa}\}$, where $N_p = \{x \in {}^F 2 : p \subseteq x\}$, and therefore this topology does not depend on the chosen bijection.

For a fragment F of $L_{\kappa+\kappa}$, define an injection $i_F : \text{Mod}_{\kappa}^L \rightarrow {}^F 2$ as follows: if $\mathcal{A} \in \text{Mod}_{\kappa}^L$, then let $i_F(\mathcal{A}) \in {}^F 2$ be such that

$$i_F(\mathcal{A})(\varphi) = 1 \text{ iff } \mathcal{A} \models \varphi[\text{id}_{\kappa}]$$

for all $\varphi \in F$. The function i_F is in fact injective, because if \mathcal{A}, \mathcal{B} are different structures in Mod_{κ}^L , then there exists a formula φ in At (the set of all atomic formulas and their negations) such that $\mathcal{A} \models \varphi[\text{id}_{\kappa}]$ and $\mathcal{B} \not\models \varphi[\text{id}_{\kappa}]$, i.e.,

$$i_F(\mathcal{A})(\varphi) = 1, \text{ and } i_F(\mathcal{B})(\varphi) = 0.$$

This implies, using the fact that $\text{At} \subseteq F$ by Definition 3.1, that $i_F(\mathcal{A}) \neq i_F(\mathcal{B})$.

We denote by X_F the image of the injection i_F .

Proposition 3.4. *If F is a fragment of $L_{\kappa+\kappa}$, then X_F is a $\mathbf{\Pi}_2^0$ subset of the Cantor space ${}^F 2$, and i_F is a homeomorphism from $\mathcal{M}od_F$ onto its image X_F , where X_F is equipped with the subspace topology.*

Proof. Because F is closed under substitution, the collection b_F from which the topology t_F is obtained is actually equal to $\{\text{Mod}_\kappa(\varphi, \text{id}_\kappa) : \varphi \in F\}$. Using this fact, it is not hard to see that the injection i_F is a homeomorphism between $\mathcal{M}od_F$ and its image X_F .

To see that $X_F \subseteq {}^F 2$ is $\mathbf{\Pi}_2^0$, we define the following subsets of ${}^F 2$. For any $h \in {}^\kappa \kappa$, we denote by $\text{supp}(h)$ the set of those $\alpha \in \kappa$ for which $h(\alpha) \neq \alpha$.

$$X_0 = \{x \in {}^F 2 : x(\psi) = 1 \text{ iff } x(\neg\psi) = 0 \text{ for all } \psi \in F\};$$

$$X_1 = \{x \in {}^F 2 : \text{if } \psi \in F \text{ and } \psi = \bigwedge \Phi \text{ for some } \Phi \in [F]^{<\kappa},$$

$$\text{then } x(\psi) = 1 \text{ iff for all } \varphi \in \Phi \text{ we have } x(\varphi) = 1\};$$

$$X_2 = \{x \in {}^F 2 : \text{if } \psi \in F \text{ and } \psi = \exists(v_\beta : \beta \in I)\varphi \text{ where } \varphi \in F \text{ and } I \in [\kappa]^{<\kappa},$$

$$\text{then } x(\psi) = 1 \text{ iff } x(s_h\varphi) = 1$$

$$\text{for some } h \in {}^\kappa \kappa \text{ such that } \text{supp}(h) \subseteq I\};$$

$$X_3 = \{x \in {}^F 2 : \text{for all } i, j \in \kappa \text{ we have } x(v_i = v_j) = 1 \text{ iff } i = j\},$$

and we let X denote their intersection. We claim that the X_i 's are all $\mathbf{\Pi}_2^0$ subsets of ${}^F 2$ and show this in detail for X_1 . For $\psi = \bigwedge \Phi \in F$, the set

$$X_{1,\psi}^1 = \{x \in {}^F 2 : x(\psi) = 1 \text{ and } x(\varphi) = 1 \text{ for all } \varphi \in \Phi\} = \bigcap_{\varphi \in \Phi} N_{\{(\psi,1),(\varphi,1)\}}$$

is $\mathbf{\Pi}_2^0$, while the set

$$X_{1,\psi}^0 = \{x \in {}^F 2 : x(\psi) = 0 \text{ and } x(\varphi) = 0 \text{ for some } \varphi \in \Phi\} = \bigcup_{\varphi \in \Phi} N_{\{(\psi,0),(\varphi,0)\}}$$

is open. Therefore $X_1 = \bigcap \{X_{1,\psi}^0 \cup X_{1,\psi}^1 : \psi = \bigwedge \Phi \in F\}$ is also a $\mathbf{\Pi}_2^0$ subset of the Cantor space ${}^F 2$. That X_0 , X_2 and X_3 are also $\mathbf{\Pi}_2^0$ can be seen similarly; in the case of X_2 , one has to use the fact that $|\{s_h\varphi : h \in {}^\kappa \kappa\}| \leq \kappa$ holds, because φ has $< \kappa$ variables and $\kappa^{<\kappa} = \kappa$.

Therefore the intersection X of the X_i 's is also $\mathbf{\Pi}_2^0$, and it remains to see that $X_F = X$. It is straightforward to show that for any $\mathcal{A} \in \text{Mod}_\kappa^L$, we have $i_F(\mathcal{A}) \in X$, and so $X_F \subseteq X$. For the other direction, first observe that if $x, y \in X$ and $x \upharpoonright \text{At} = y \upharpoonright \text{At}$, then $x = y$ also holds, by an easy induction on the complexity of formulas. Suppose $x \in X$ is arbitrary. We wish to define a model $\mathcal{A} \in \text{Mod}_\kappa^L$ such that $x = i_F(\mathcal{A})$; by the above observation, it is enough to require that $x \upharpoonright \text{At} = i_F(\mathcal{A}) \upharpoonright \text{At}$. Clearly, the L -model \mathcal{A} whose domain is κ and whose relations are defined by letting, for

each n -ary relation symbol R of L and $\alpha_1, \dots, \alpha_n \in \kappa$,

$$(\alpha_1, \dots, \alpha_n) \in R^A \text{ iff } x(R(v_{\alpha_1}, \dots, v_{\alpha_n})) = 1,$$

satisfies these requirements. \square

Let ψ be an arbitrary sentence in $\Sigma_1^1(L_{\kappa+\kappa})$ (recall that this means ψ is a second order sentence of the form $\exists \bar{R} \varphi(\bar{R})$ where \bar{R} is a set of $\leq \kappa$ many symbols disjoint from L and $\varphi(\bar{R})$ is an $L_{\kappa+\kappa}$ -sentence in the language expanded by \bar{R}). We let Mod_κ^ψ be the set of elements of Mod_κ^L which are models of ψ . If F is a fragment of $L_{\kappa+\kappa}$, we denote by $\mathcal{M}od_F^\psi$ the corresponding subspace of the topological space $\mathcal{M}od_F$, or in other words, $\mathcal{M}od_F^\psi$ is obtained by endowing Mod_κ^ψ with the topology t_F . We furthermore define

$$X_F^\psi = \{i_F(\mathcal{A}) : \mathcal{A} \in \text{Mod}_\kappa^\psi\},$$

the set of elements of X_F corresponding to models of ψ .

Corollary 3.5. *If ψ is a sentence of $\Sigma_1^1(L_{\kappa+\kappa})$ and F is a fragment of $L_{\kappa+\kappa}$, then X_F^ψ is an analytic subset of ${}^F 2$. Furthermore, the map $i_F \upharpoonright \mathcal{M}od_F^\psi$ is a homeomorphism from $\mathcal{M}od_F^\psi$ onto its image X_F^ψ equipped with the subspace topology.*

Proof. As we have seen at the beginning of the previous proof, it is enough to show that X_F^ψ is analytic. First, in the case when $\psi \in F$ (and therefore is an $L_{\kappa+\kappa}$ -sentence), we have $X_F^\psi = X_F \cap N_{\{(\psi,1)\}}$. Thus, X_F^ψ is a $\mathbf{\Pi}_2^0$ subset of ${}^F 2$ by Proposition 3.4.

Now, in general, suppose that ψ is the sentence $\exists \bar{R} \varphi(\bar{R})$. Let F' be the fragment generated (in the expanded language) by $F \cup \{\varphi(\bar{R})\} \cup \bar{R}$. Then we have that X_F^ψ is the image of the $\mathbf{\Pi}_2^0$ set $X_{F'}^\varphi$ under the continuous map ${}^{F'} 2 \rightarrow {}^F 2$, $x \mapsto x \upharpoonright F$ and is therefore analytic. (Equivalently, $\mathcal{M}od_F^\psi$ is the image of $\mathcal{M}od_{F'}^\varphi$ under the continuous map defined by taking the L -reducts of models for the expanded language.) \square

We now turn to the “number” of pairwise non (F, H) -elementarily embeddable models of an arbitrary sentence $\psi \in \Sigma_1^1(L_{\kappa+\kappa})$, where $H \subseteq \text{Inj}(\kappa)$ and F is a fragment of $L_{\kappa+\kappa}$. We will say that ψ has *perfectly many non (F, H) -elementarily embeddable models* if the binary relation of (F, H) -elementary embeddability on Mod_κ^ψ has a t_F -perfect independent set. (Note that we may speak about t_F -perfect sets since $\mathcal{M}od_F^\psi$ is homeomorphic to a subset of the κ -Baire space.)

We remark that, as Proposition 3.7 below shows, the choice of the fragment that generates the topology on Mod_κ^ψ is actually irrelevant in the above definition. That is, given any fragment F' of $L_{\kappa+\kappa}$, the sentence ψ

has perfectly many non (F, H) -elementarily embeddable models iff the binary relation of (F, H) -elementary embeddability on Mod_κ^ψ has a $t_{F'}$ -perfect independent set.

Below, by a t_F -Borel subset of Mod_κ^ψ , we mean a κ -Borel subset of $\mathcal{M}od_F^\psi$, and a map $f : X \rightarrow \text{Mod}_\kappa^\psi$ (where X is a topological space) is t_F -Borel iff the inverse images of t_F -Borel subsets of Mod_κ^ψ are κ -Borel subsets of X .

Proposition 3.6. *Let F and F' be arbitrary fragments of $L_{\kappa+\kappa}$.*

1. *A subset of Mod_κ^ψ is t_F -Borel iff it is $t_{F'}$ -Borel.*
2. *A map $f : {}^\kappa 2 \rightarrow \text{Mod}_\kappa^\psi$ is t_F -Borel iff it is $t_{F'}$ -Borel.*

Proof. An easy induction shows that for any $\varphi \in L_{\kappa+\kappa}$ (and therefore for any $\varphi \in F'$) and valuation $a \in {}^\kappa \kappa$, the set $\text{Mod}_\kappa(\varphi, a)$ is t_F -Borel. Consequently, all $t_{F'}$ -Borel sets are t_F -Borel sets as well. By symmetry, we have item 1, of which item 2 is a direct consequence. \square

Proposition 3.7. *Let F and F' be arbitrary fragments of $L_{\kappa+\kappa}$, and suppose R is a binary relation on Mod_κ^ψ . Then R has a t_F -perfect independent set iff it has a $t_{F'}$ -perfect independent set.*

Proof. The binary relation R has a t_F -perfect independent set iff there is a t_F -Borel (instead of “ t_F -continuous”) injection of ${}^\kappa 2$ into Mod_κ^ψ such that

$$(*) \quad \text{for all } x \neq y \in {}^\kappa 2, \text{ we have } (r(x), r(y)) \notin R;$$

see for example [4, Proposition 2] for a proof. (A map satisfying $(*)$ is called a *reduction* of $\text{id}_{{}^\kappa 2}$ to R .) Therefore, using item 2 of Proposition 3.6 above, we obtain the conclusion of this proposition immediately. \square

We now state and prove the main theorem of this section.

Theorem 3.8. *Assume $I^-(\kappa)$ and either that \diamond_κ holds or that κ is inaccessible. Suppose that $H \subseteq \text{Inj}(\kappa)$, F is a fragment of $L_{\kappa+\kappa}$ and ψ is a sentence of $\Sigma_1^1(L_{\kappa+\kappa})$. Suppose that either*

1. *H is a K_κ subset of the κ -Baire space, or*
2. *H is a K_κ subset of the product space $({}^\kappa \kappa, \tau_{\text{pr}})$ and $F \subseteq L_{\kappa+\omega}$.*

If there are at least κ^+ many pairwise non (F, H) -elementarily embeddable models in Mod_κ^ψ , then there are perfectly many such models.

Note that since the κ -Baire topology is finer than the product topology τ_{pr} , being a K_κ subset of the κ -Baire space implies being a K_κ subset of the product space $({}^\kappa \kappa, \tau_{\text{pr}})$. This implication is strict if and only if κ is not weakly compact. Moreover, if κ is not weakly compact and $I^-(\kappa)$ holds, then

the K_κ subsets of the κ -Baire space are exactly the ones of size $\leq \kappa$ (see Remark 2.10).

We note that, (as mentioned in the Introduction), it is consistent to also have $2^\kappa > \kappa^+$ together with the set theoretical hypothesis of the above theorem, relative to the consistency of a measurable $\lambda > \kappa$. Furthermore, starting from a stronger situation, we can also assume that κ is a weakly compact cardinal such that $I^-(\kappa)$ and $2^\kappa > \kappa^+$ (see Remark 2.11).

Proof. We start with the proof of item 1. Assume, as usual, that H is given the subspace topology induced by the κ -Baire space and X_F^ψ is equipped with the subspace topology induced by the generalized Cantor space ${}^F 2$.

Define the subset S of $H \times X_F^\psi \times X_F^\psi$ by letting, for any $h \in H$ and $\mathcal{A}, \mathcal{B} \in \text{Mod}_\kappa^\psi$,

$$(h, i_F(\mathcal{B}), i_F(\mathcal{A})) \in S \text{ iff } h \text{ witnesses that } \mathcal{A} \text{ is} \\ (F, H)\text{-elementarily embeddable into } \mathcal{B}.$$

Then, since F is closed under substitution and negation, $(h, i_F(\mathcal{B}), i_F(\mathcal{A})) \in S$ iff for all formulas φ in F , $\mathcal{A} \models \varphi[\text{id}_\kappa]$ implies that $\mathcal{B} \models \varphi[h \circ \text{id}_\kappa]$, or equivalently that $\mathcal{B} \models s_h \varphi[\text{id}_\kappa]$. Therefore

$$S = \{(h, y, x) \in H \times X_F^\psi \times X_F^\psi : \text{for all } \varphi \in F, x(\varphi) = 1 \text{ implies } y(s_h \varphi) = 1\}.$$

Claim 3.9. *S is a closed subset of $H \times X_F^\psi \times X_F^\psi$.*

Proof of Claim 3.9. We prove that the complement $U = H \times X_F^\psi \times X_F^\psi - S$ is open. Suppose that $(h, y, x) \in U$, or in other words, there exists $\varphi \in F$ such that $x(\varphi) = 1$ and $y(s_h \varphi) = 0$. Then, since the set $\Delta(\varphi)$ of free variables of φ is of size $< \kappa$, the set $N_1 = N_{h \upharpoonright \Delta(\varphi)} \cap H$ is an open subset of H . Furthermore, $h' \in N_1$ implies that for all $x \in X_F^\psi$, we have $x(s_{h'} \varphi) = x(s_h \varphi)$. Thus, denoting by N_2 and N_3 the open subsets of X_F^ψ determined by the conditions $z(\varphi) = 1$ and $z(s_h \varphi) = 0$, respectively, we obtain an open neighborhood $N_1 \times N_2 \times N_3$ of (h, y, x) which is also a subset of U . This completes the proof of Claim 3.9.

Clearly, the projection R_S of S onto $X_F^\psi \times X_F^\psi$ is the relation corresponding to (F, H) -elementary embeddability on Mod_κ^ψ (i.e., $(i_F(\mathcal{B}), i_F(\mathcal{A})) \in R_S$ iff \mathcal{A} is F -elementarily embeddable into \mathcal{B} by H). By Corollary 3.5, X_F^ψ is an analytic subset of the generalized Cantor space ${}^F 2$, and H is a K_κ topological space by the assumption of item 1. Thus, by Corollary 2.9, we have the required conclusion.

To see item 2, we equip H with the subspace topology induced by the *product topology* τ_{pr} on the set ${}^\kappa \kappa$. As before, X_F^ψ is given the subspace

topology induced by the generalized Cantor space ${}^F 2$, and the set S is defined as above. Then, using the assumption $F \subseteq L_{\kappa+\omega}$ of item 2, one can show that

$$S \text{ is a closed subset of the space } H \times X_F^\psi \times X_F^\psi.$$

This can be seen by using the argument in the proof of Claim 3.9 and taking note of the fact that, since the set of free variables of any $\varphi \in F$ is finite by the assumption $F \subseteq L_{\kappa+\omega}$, the set denoted by N_1 in the proof of Claim 3.9 is an open subset of H even when the topology on H is inherited from the product space $({}^\kappa \kappa, \tau_{\text{pr}})$.

Furthermore, H is a K_κ topological space by the first assumption of item 2, and X_F^ψ is an analytic subset of the generalized Cantor space ${}^F 2$ by Corollary 3.5. Therefore Corollary 2.9 can again be applied to obtain the required conclusion. \square

Remark 3.10. Suppose that Φ is an arbitrary κ -sized subset of $L_{\kappa+\kappa}$ -formulas which is closed under substitution. For a subset H of ${}^\kappa \kappa$, consider the models in Mod_κ^ψ up to maps $h \in H$ which preserve the formulas in Φ (i.e., maps $h : \mathcal{A} \rightarrow \mathcal{B}$ such that for all $\varphi \in \Phi$ and valuations $a \in {}^\kappa \mathcal{A}$, if $\mathcal{A} \models \varphi[a]$ then $\mathcal{B} \models \varphi[h \circ a]$. Note that in this case, such maps h need not be injective). Using the topology t_F , where F is the fragment generated by Φ , the proof of Theorem 3.8 can be generalized to yield an analogous statement about the “number of models” up to such maps. This version seems to cover all natural generalizations of Theorem 3.8.

Specifically, when Φ is the set of those atomic formulas which do not contain the $=$ symbol, a map h preserves Φ iff it is a homomorphism. Therefore, when H is a K_κ subset of the product space $({}^\kappa \kappa, \tau_{\text{pr}})$, we have that the following holds under the assumptions of Theorem 2.4: *if there are at least κ^+ many models in Mod_κ^ψ such that no $h \in H$ is a homomorphism from one into another, then there are perfectly many such models.*

We conclude this section by mentioning some interesting special cases of Theorem 3.8.

Corollary 3.11. *Assume that $\Gamma^-(\kappa)$ and either \diamond_κ or that κ is inaccessible. Let $H \subseteq \text{Inj}(\kappa)$ and let ψ be a sentence of $\Sigma_1^1(L_{\kappa+\kappa})$.*

1. *Suppose H is a K_κ subset of the product space $({}^\kappa \kappa, \tau_{\text{pr}})$. If there are at least κ^+ many pairwise non H -elementarily embeddable models in Mod_κ^ψ , then there is a perfect set of such models.*
2. *The above also holds for H -embeddability, as well as (F, H) -elementary embeddability when F is either $L_{\lambda\omega}$ or $L_{\lambda\omega}^n$, and $n < \omega \leq \lambda \leq \kappa$.*

3. Suppose H is a K_κ subset of the κ -Baire space. Then the same statement holds for $(L_{\lambda\mu}, H)$ -elementary embeddability, where $\omega < \mu \leq \lambda \leq \kappa$.
4. Now, suppose that H is a subgroup of $\text{Sym}(\kappa)$ which is K_κ again in the product topology τ_{pr} . If there are κ^+ many pairwise non H -isomorphic models in Mod_κ^ψ , then there is a perfect set of such models.

Acknowledgements. The research of the first author was partially supported by the Central European University Budapest Foundation (CEUBPF) and by the Hungarian National Research, Development and Innovation Office grant no. 113017. The authors would like to thank Menachem Magidor for suggesting arguments found in the Introduction and in Remark 2.11. The first author would like to thank Lajos Soukup, Tapani Hyttinen, Vadim Kulikov, Philipp Schlicht and Boban Veličković for valuable discussions and Gábor Sági for reading and making useful comments on this manuscript.

REFERENCES

- [1] J. Baldwin. Diverse classes. *J. Symb. Log.*, 54(3):857–893, 1989.
- [2] K. J. Devlin. Variations on \diamond . *J. Symb. Log.*, 44(1):51–58, 1979.
- [3] M. A. Dickmann. *Large Infinitary Languages*, volume 83 of *Model theory, Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Company, Amsterdam, 1975.
- [4] S. D. Friedman. Consistency of the Silver dichotomy in generalised Baire space. *Fund. Math.*, 227(2):179–186, 2014.
- [5] S. D. Friedman, T. Hyttinen, and V. Kulikov. *Generalized Descriptive Set Theory and Classification Theory*. American Mathematical Society, Providence, 2014.
- [6] S. D. Friedman and V. Kulikov. Failures of the Silver dichotomy in the generalised Baire space. *J. Symb. Log.*, 80(2):661–670, 2015.
- [7] F. Galvin, T. Jech, and M. Magidor. An ideal game. *J. Symb. Log.*, 43(2):284–292, 1978.
- [8] S. Gao. *Invariant Descriptive Set Theory*, volume 293 of *Pure and Applied Mathematics, A Series of Monographs and Textbooks*. Taylor & Francis Group, 2009.
- [9] W. Hodges and S. Shelah. Infinite games and reduced products. *Ann. Math. Log.*, 20(1):77–108, 1981.
- [10] T. Jech. Trees. *J. Symb. Log.*, 36(1):1–14, 1971.
- [11] A. S. Kechris. *Classical Descriptive Set Theory*, volume 165 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [12] W. Kubiś. Perfect cliques and G_δ colorings of Polish spaces. *Proc. Amer. Math. Soc.*, 131(2):619–623, 2003.
- [13] R. Laver. Making the supercompactness of κ indestructible under κ -directed closed forcing. *Israel J. Math.*, 29(4):385–388, 1978.
- [14] P. Lipparini. Compactness of powers of ω . *Bull. Pol. Acad. Sci. Math.*, 61(3-4):239–246, 2013.
- [15] P. Lücke, L. Motto Ros, and P. Schlicht. The Hurewicz dichotomy for generalized Baire spaces. Preprint, available at <http://arxiv.org/abs/1506.03364>.
- [16] P. Matet. Corrigendum to “Partitions and Diamond”. *Proc. Amer. Math. Soc.*, 101(3):519–520, 1987.
- [17] A. Mekler, S. Shelah, and J. Väänänen. The Ehrenfeucht-Fraïssé game of length ω_1 . *Trans. Amer. Math. Soc.*, 339:567–580, 1993.

- [18] A. Mekler and J. Väänänen. Trees and Π_1^1 subsets of $\omega_1\omega_1$. *J. Symb. Log.*, 58(3):1052–1070, 1993.
- [19] M. Morley. The number of countable models. *J. Symb. Log.*, 35(1):14–18, 1970.
- [20] A. Piolunowicz. On the \diamond principle. *Fund. Math.*, 122(2):149–157, 1984.
- [21] G. Sági and D. Sziráki. Some variants of Vaught’s conjecture from the perspective of algebraic logic. *Log. J. IGPL*, 20(6):1064–1082, 2012.
- [22] S. Shelah. The number of pairwise non-elementarily-embeddable models. *J. Symb. Log.*, 54(4):1431–1455, 1989.
- [23] S. Shelah. Borel sets with large squares. *Fund. Math.*, 159(1):1–50, 1999.
- [24] S. Shelah. *Classification theory for abstract elementary classes*, volume 18 of *Studies in Logic*. College Publications, London, 2009.
- [25] S. Shelah. Diamonds. *Proc. Amer. Math. Soc.*, 138(6):2151–2161, 2010.
- [26] S. Shelah, H. Tuuri, and J. Väänänen. On the number of automorphisms of uncountable models. *J. Symb. Log.*, 58(4):1402–1418, 1993.
- [27] J. Väänänen. A Cantor-Bendixson theorem for the space $\omega_1^{\omega_1}$. *Fund. Math.*, 137:187–199, 1991.
- [28] J. Väänänen. Games and trees in infinitary logic: A Survey. In M. Krynicki, M. Mostowski, and L. Szczerba, editors, *Quantifiers*, pages 105–138. Kluwer Academic Publishers, 1995.
- [29] J. Väänänen. *Models and Games*, volume 132 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2011.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES, REÁLTANODA U. 13–15, H–1053 BUDAPEST, HUNGARY, AND DEPARTMENT OF MATHEMATICS AND ITS APPLICATIONS, CENTRAL EUROPEAN UNIVERSITY, NÁDOR U. 9, H–1051 BUDAPEST, HUNGARY

E-mail address: sziraki_dorottya@phd.ceu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, GUSTAF HÄLLSTRÖMIN KATU 2B, FI–00014 UNIVERSITY OF HELSINKI, FINLAND, AND INSTITUTE FOR LOGIC, LANGUAGE AND COMPUTATION, UNIVERSITY OF AMSTERDAM, 1090 GE AMSTERDAM, NETHERLANDS

E-mail address: jouko.vaananen@helsinki.fi