



DIGITAL ACCESS TO  
SCHOLARSHIP AT HARVARD  
DASH.HARVARD.EDU



HARVARD LIBRARY  
Office for Scholarly Communication

# Pyrazines from bacteria and ants: convergent chemistry within an ecological niche

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Silva-Junior, Eduardo A., Antonio C. Ruzzini, Camila R. Paludo, Fabio S. Nascimento, Cameron R. Currie, Jon Clardy, and Mônica T. Pupo. 2018. "Pyrazines from bacteria and ants: convergent chemistry within an ecological niche." <i>Scientific Reports</i> 8 (1): 2595. doi:10.1038/s41598-018-20953-6. <a href="http://dx.doi.org/10.1038/s41598-018-20953-6">http://dx.doi.org/10.1038/s41598-018-20953-6</a> .
Published Version	<a href="https://doi.org/10.1038/s41598-018-20953-6">doi:10.1038/s41598-018-20953-6</a>
Citable link	<a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:35014955">http://nrs.harvard.edu/urn-3:HUL.InstRepos:35014955</a>
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA">http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA</a>

# SCIENTIFIC REPORTS

OPEN

## Pyrazines from bacteria and ants: convergent chemistry within an ecological niche

Eduardo A. Silva-Junior<sup>1</sup>, Antonio C. Ruzzini<sup>2,5</sup>, Camila R. Paludo<sup>1</sup>, Fabio S. Nascimento<sup>3</sup>, Cameron R. Currie<sup>4</sup>, Jon Clardy<sup>2</sup> & Mônica T. Pupo<sup>1</sup>

Ants use pheromones to coordinate their communal activity. Volatile pyrazines, for instance, mediate food resource gathering and alarm behaviors in different ant species. Here we report that leaf-cutter ant-associated bacteria produce a family of pyrazines that includes members previously identified as ant trail and alarm pheromones. We found that L-threonine induces the bacterial production of the trail pheromone pyrazines, which are common for the host leaf-cutter ants. Isotope feeding experiments revealed that L-threonine along with sodium acetate were the biosynthetic precursors of these natural products and a biosynthetic pathway was proposed.

Leaf-cutter ants are major herbivores in the New World Tropics where they provide essential ecosystem services<sup>1</sup>. These ants collect fresh plant material to cultivate a symbiotic fungus, *Leucoagaricus gongylophorus*, that serves as their food source<sup>2</sup>. The scale at which these ants collect leaves makes them major agricultural pests in temperate regions of the New World<sup>3,4</sup>. Colonies of leaf-cutter ants in the genus *Atta* can have populations in the millions<sup>5</sup> housed in subterranean nests that include an extensive network of tunnels and chambers which contain their fungal gardens<sup>3</sup>. To accomplish the task of keeping such massive populations fed, the leaf-cutter ants coordinate their harvesting behavior by using trail pheromones that are secreted from specialized poison glands<sup>6–8</sup>. In the case of *Atta sexdens*<sup>8,9</sup>, 2,5-dimethylpyrazine (1) and 3-ethyl-2,5-dimethylpyrazine (2) have been identified both as components of poison gland secretions and trail pheromones.


Pyrazines are heterocyclic natural products commonly used by ants to communicate. Mandibular glands secretions from Ponerini, Odontomachini and Ectatommini ants contain a mixture of 3-alkyl-2,5-dimethylpyrazines and 3-alkyl-2,6-dimethylpyrazines, compounds that mediate alarm behavior in *Odontomachus* workers<sup>10,11</sup>. Pyrazines are also components of the trail pheromones produced by the poison glands of the leaf-cutter ants *Acromyrmex octospinosus*, *Atta bisphaerica*, *Atta cephalotes*, and *Atta sexdens*<sup>7,12,13</sup>. These natural products likewise mediate communication in bacteria: 3,5-dimethylpyrazin-2-ol is a quorum sensing regulator implicated in biofilm formation by *Vibrio cholerae*<sup>14</sup>.

The bacterial associates of ants in the tribe Attine have recently been exploited as a promising source of antibiotics<sup>15–17</sup> thought to protect their hosts. The production of volatiles – a critical class of molecules that guides ant behavior – by these bacteria, however, remain largely unknown. There is some evidence that volatile organic compounds (VOCs) produced by bacterial insect symbionts are sources of pheromones utilized by their hosts<sup>18,19</sup>. Herein, we provide the first report of a leaf-cutter ant-associated bacterium, *Serratia marcescens* 3B2, that is capable of producing pyrazines previously identified as trail pheromones (1 and 2) of the host ant *Atta sexdens rubropilosa* and alarm pheromones (4 and 5) of other ants<sup>8,9,11,20</sup>. The biosynthetic precursors of the trail pheromone pyrazines were unveiled by isotope feeding experiments.


### Results

**Ant associated bacteria produce a small family of pyrazines.** Eight bacteria with similar morphology were isolated from *A. sexdens rubropilosa* worker ants belonging to two different colonies. The isolates were

<sup>1</sup>Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, 14040903, Brazil. <sup>2</sup>Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA, 02115, United States of America. <sup>3</sup>Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, 14040901, Brazil. <sup>4</sup>Department of Bacteriology, University of Wisconsin, Madison, WI, 53706, United States of America. <sup>5</sup>Present address: Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon, SK, S7N 5B4, Canada. Correspondence and requests for materials should be addressed to M.T.P. (email: [mtpupo@fcrp.usp.br](mailto:mtpupo@fcrp.usp.br))



**Figure 1.** Pyrazines produced by *Serratia marcescens* 3B2.




**Figure 2.** Structure, <sup>13</sup>C (red) and <sup>1</sup>H (blue) NMR data of compound **6** in (A). Key HMBC and COSY correlations (B).

derived from two sources: whole ant bodies that had been washed with a sterile aqueous solution<sup>21,22</sup> or worker ants that had been sectioned after a surface sterilization<sup>23</sup>. The VOCs from the eight isolates grown on agar ISP-2 medium were extracted using a SPME polydimethylsiloxane fiber and analyzed by GC-MS (Figs S1–S5, ESI<sup>†</sup>).


The main compounds produced by the isolates were pyrazines **3–6** (Fig. 1). Compound **3** is a methoxylated analogue of 3,5-dimethylpyrazin-2-ol, a recently identified bacterial quorum sensing regulator<sup>14</sup>. Compounds **4** and **5** were previously reported as alarm pheromones of non-fungus growing ants<sup>10,11</sup>. Compound **6** (Fig. 2) is an unreported pyrazine and its structure was determined based on HRESIMS (C<sub>12</sub>H<sub>20</sub>N<sub>2</sub>, *m/z* 193.1694 [M + H]<sup>+</sup>; Fig. S10, ESI<sup>†</sup>) and NMR data (Table S1, ESI<sup>†</sup>). More specifically, the 1D and 2D NMR spectra indicated that the structure is 2,5-dimethyl-3-(3'-methylpentyl)pyrazine (Figs S6–S9, ESI<sup>†</sup>). gHMBC correlations and a key gCOSY coupling between the methyl hydrogens (δ 0.95) with the hydrogen at δ 1.66 (H3') established the position of this methyl group at C3'. The GC-MS fragmentation profile of **6** corroborates with NMR data, indicating a methyl group at C3' (Fig. S3, ESI<sup>†</sup>).

**L-threonine induces the bacterial production of trail pheromone pyrazines.** In order to investigate if the trail pheromone pyrazines **1** and **2**, found in the hosts ants (Figs S11–S13, ESI<sup>†</sup>), could also be produced by the bacterium, we examined the metabolites from a single isolate – *S. marcescens* 3B2 – using minimal and defined growth media. This strain was identified by sequencing the 16S rRNA gene using the universal primers 27F and 1492R, and the resulting sequence was compared with GenBank sequences. We cultured *S. marcescens* 3B2 using M9 agar supplemented with glucose and a single amino acid: L-alanine, L-serine, L-threonine or L-valine, (Figs S14–S16, ESI<sup>†</sup>). Importantly, filtered sterilized amino acids were added to autoclaved media to avoid heat-induced pyrazine formation<sup>24</sup>. Compounds **1** and **2** were observed only in the M9 agar supplemented with glucose and L-threonine, and could also be detected in bulk extraction of the host, *Atta sexdens rubropilosa* (Fig. 3). The bacterial production of **1** and **2** was dependent on the concentration of L-threonine in the culture media, tested at 1.0, 1.5 and 2.0% (w/v; Fig. S17–S20, ESI<sup>†</sup>). We did not detect **1** and **2** from cultures with low L-threonine concentrations (0–0.5% w/v; Figs S17–S20, ESI<sup>†</sup>). In addition, L-threonine supplementation resulted in reduced yields of the more abundant pyrazines **4–6**, and the production of **3** was also upregulated (Fig. S17, ESI<sup>†</sup>).

**L-threonine and acetate are the biosynthetic precursors of the trail pheromone pyrazines.** To study the role of L-threonine in the production of **1** and **2**, we grew *S. marcescens* 3B2 on M9 agar supplemented with 0.2% of glucose, 0.5% of L-alanine and 1.5% of L-threonine or L-[U-<sup>13</sup>C,<sup>15</sup>N]-threonine (Figs S21–S30, ESI<sup>†</sup>). GC-MS analysis showed the incorporation of eight mass units into both **1** and **2** (Figs S27 and S28, ESI<sup>†</sup>). The ethyl group of **2** is eliminated during GC-MS analysis and produces a fragment ion with the charge at the 2,5-dimethylpyrazine ring (Fig. S4, ESI<sup>†</sup>). The fragment ion of **2**, produced in the medium supplemented with L-[U-<sup>13</sup>C,<sup>15</sup>N]-threonine, has the expected M + 8 shift (*m/z* 116) compared to the control (*m/z* 108), showing that the incorporation occurred just in the pyrazine ring. High-resolution electrospray ionization mass spectrometry (HR-ESIMS) analysis corroborated with the GC-MS data. The major ions observed for both **1** and **2** were consistent with the incorporation of six <sup>13</sup>C and two <sup>15</sup>N atoms into each (Fig. 4; Figs S29 and S30; Tables S2 and S3, ESI<sup>†</sup>). Minor fermentation products contained three <sup>13</sup>C atoms and a single <sup>15</sup>N atom, suggesting that endogenously produced L-threonine was also incorporated during the feeding experiment.



**Figure 3.** GC-MS chromatograms showing the production of the trail pheromone pyrazines **1** and **2** by *S. marcescens* 3B2 cultivated on M9 agar medium supplemented with 0.2% of glucose and 2% of L-threonine in (A) and their presence in gasters of *Atta sexdens rubropilosa* in (B).




**Figure 4.** Labelling patterns of pyrazines **1** and **2** obtained after feeding the biosynthetic precursors L-[U-<sup>13</sup>C, <sup>15</sup>N]-threonine and U-<sup>13</sup>C-sodium acetate in *S. marcescens* 3B2 culture.

To investigate the origin of the ethyl-substitution in **2**, we added U-<sup>13</sup>C-sodium acetate to the growth medium. Under these conditions, no incorporation was observed into **1** whereas two additional mass units were incorporated into **2**, as evidenced by GC-MS and HR-ESI-MS (Figs S31–S35, ESI<sup>†</sup>). To better establish the positions of <sup>13</sup>C acetate-derived atoms we took advantage of the elimination process that occurs during GC-MS of **2**. Equivalent fragment ions ( $m/z$  108) were detected in both labelled (molecular ion with  $m/z$  138) and control sample (molecular ion with  $m/z$  136), indicating that acetate was incorporated into the eliminated ethyl group. *Serratia marcescens* 3B2 was also cultivated with <sup>15</sup>N-L-alanine and 3-<sup>13</sup>C-L-alanine; however, no significant isotopic labelling of the 3,5-dimethylpyrazine rings of compounds **1** and **2** were observed from these amino acids (Figs S36–S45, ESI<sup>†</sup>). Instead, we observed low levels of a single <sup>13</sup>C incorporation into the ethyl group of compound **2** when the media was supplemented with 3-<sup>13</sup>C-L-alanine, which is most likely due to the metabolism of L-alanine to acetate<sup>25</sup>. Much higher levels of M + 2 ions were observed using uniformly <sup>13</sup>C labelled acetate alone or in combination with an equimolar concentration of unlabeled L-alanine (Table S3, Figs S46 and S47, ESI<sup>†</sup>). Altogether, the results demonstrate that acetate is the precursor of the compound **2** ethyl group. In contrast of **1** and **2**, the 2,5-dimethylpyrazine rings of **4**, **5** and **6** were labelled with <sup>15</sup>N-L-alanine and 3-<sup>13</sup>C-L-alanine (Figs S48–S50, ESI<sup>†</sup>). As reported for 3,5-dimethylpyrazin-2-ol<sup>14</sup>, isotope labelled L-alanine and L-threonine were incorporated into the 3,5-dimethylpyrazine ring of **3** (Fig. S51, ESI<sup>†</sup>). These results indicate that *S. marcescens* 3B2 produces the same structural ring using different biosynthetic precursors.

We proposed the *S. marcescens* 3B2 biosynthetic pathways of **1** and **2** starting with L-threonine, which is converted to aminoacetone (Fig. 5). Condensation of two units of aminoacetone, followed by dehydration, yields the 2,5-dimethyl-3,6-dihydropyrazine (DHP), which is aromatized by ring oxidation forming **1**. Compound **2** can be produced by the same biosynthetic pathway, with an addition of acetate at C3. The resulting carbonyl group is reduced to alcohol, followed by water elimination and spontaneous tautomerization to originate **2**.

## Discussion

The discovery of bacterially-produced pyrazines from isolates of leaf-cutter ants presents an interesting example of niche-specific chemistry. While ant-derived microbes have been chemically investigated<sup>15,16,26,27</sup>, this study is the first to report convergent chemistry between ants and their associated bacteria. Moreover, biological roles of pyrazines and their effects on ant behavior are well-established. VOCs are commonly used by insects for



**Figure 5.** Proposed biosynthesis of pyrazines **1** and **2**, involving oxidations (ox.), reductions (red.), spontaneous tautomerization (taut.) and dehydration ( $-H_2O$ ).

communication<sup>18,28,29</sup>. Pyrazines, for instance, have been found in ants, phasmids and wasps, playing important roles as alarm, defense, sex and trail pheromones<sup>9,11,30–33</sup>.

Insects can use compounds from their diet as pheromone precursors, as occur with the bark beetles *Ips paraconfusus*, which acquire pheromone precursors from plants<sup>34,35</sup>. Arthropod-associated bacteria can also provide VOCs for insects: the German cockroach *Blattella germanica*<sup>18</sup> and the locust *Schistocerca gregaria*<sup>19</sup> use VOCs produced by gut symbiotic bacteria. The pyrazines produced by *S. marcescens* ant-associated bacteria likely contribute to the complex molecular cocktail of pheromones that are used by *A. sexdens rubropilosa* ants. An interesting secondary hypothesis, and one that takes a bacterial perspective, posits that bacteria have evolved to attract their hosts. The trail pheromones are essentially attractants, and free-living forms of the bacteria may produce the pyrazines to initiate a symbiotic relationship with the leaf-cutters. Bacteria in the genus *Serratia* were also identified in colonies of the fungus-growing ants *Atta colombica*, *Apterostigma dentigerum*, *Cyphomyrmex longiscapus* and *Trachymyrmex zeteki*, but their VOCs were not investigated<sup>36,37</sup>. The producer(s) of volatile pyrazines in the *A. sexdens rubropilosa* ants merit subsequent field and laboratory investigations in order to better understand and delineate the insect and possible bacterial contribution to pheromonal communication in leaf-cutter ants.

Despite their important role as trail pheromones, very little is known about the biosynthesis of pyrazines in ants and bacteria. The utilization of L-threonine by *Serratia* results in the transient accumulation of aminoacetone<sup>38</sup>, a metabolite that may represent a direct precursor to **1** and **2**. An aminoacetone oxidase from the human oral microbiome isolate *Streptococcus oligofermentans* has been reported to catalyze the formation of **1** from two equivalents of aminoacetone<sup>39</sup>. Moreover, in a closely related molecule, 3,5-dimethylpyrazin-2-ol (DPO), a threonine dehydrogenase, which catalyzes the formation of 2-amino-3-ketobutyric acid from L-threonine was implicated<sup>14</sup>. It is the spontaneous decarboxylation of 2-amino-3-ketobutyric acid that results in the aforementioned pool of aminoacetone. An enzyme that utilizes aminoacetone and L-alanine to form the pyrazine DPO, however, is unknown.

Finding that L-threonine induces the ant-isolate bacterium *S. marcescens* 3B2 to produce the trail pheromone pyrazines **1** and **2** is consistent with the high level of incorporation observed in these products. Amino acids often trigger significant physiological responses in bacteria and higher organisms<sup>40–43</sup>. The amino acid L-isoleucine induces epithelial cells to produce  $\beta$ -defensin, which belongs to a family of antimicrobial peptides important to the mammalian innate immune system to combat pathogens<sup>40</sup>. L-proline is an amino acid component of the insect hemolymph, and contributes to the upregulation of the bacterial production of small molecules essential for the virulence of the nematode-associated bacteria<sup>41</sup>. Finally, L-tryptophan produced by a marine algae induce symbiotic bacteria to produce the growth promoter auxin indole-3-acetic acid<sup>42,43</sup>. The role and regulation of L-threonine levels in the poison glands of *Atta* are currently unknown, but clearly amino acid availability shapes the distribution of pyrazines produced by the ant-associated *Serratia*.

In summary, *Serratia marcescens* 3B2 isolated from the leaf-cutter ant *Atta sexdens rubropilosa* produces volatile pyrazines, which can vary according the composition of the culturing media. Pyrazines **1** and **2** are known components of its host's trail pheromone, and pyrazines **4** and **5** are likewise alarm pheromones of other ants<sup>10,11</sup>. The bacterial biosynthesis of **1** and **2** is stimulated by L-threonine, which was found to be the precursor of their 2,5-dimethylpyrazine rings, while the ethyl group of **2** is derived from acetate. The physiological and ecological relevance of these unprecedented findings merits to be further explored in the future.

## Methods

**General method for bacterial isolation and identification.** Ant-associated bacteria were isolated by washing ants and plating the aqueous cell suspensions on selective medium<sup>21,22</sup>. We performed two independent isolation experiments, using two *Atta sexdens rubropilosa* nests, collected in the campus of University of São Paulo, Ribeirão Preto, SP, Brazil and maintained in laboratorial conditions (colony 1 and colony 2), according



to SISBIO authorization 46555-5, and CNPq process 010936/2014-9. For the first isolation process, three *Atta sexdens rubropilosa* ants from colony 1 were transferred to a sterile 1.5 mL Eppendorf tube containing 750  $\mu\text{L}$  of 0.9% NaCl isotonic aqueous solution. The tube was vortexed during approximately 1 min and 100  $\mu\text{L}$  of washing solution were inoculated on ISP-2 agar plates (80 mm; 4% of yeast extract, 10% of malt extract, 4% of glucose and 20% of agar in distilled water) added with antifungal agents (nystatin 0.04  $\text{g}\cdot\text{L}^{-1}$  and cycloheximide 0.05  $\text{g}\cdot\text{L}^{-1}$ ). The plates were incubated at 30 °C and the grown bacterial colonies were isolated on ISP-2 agar plates.

For the second isolation procedure, ants from colony 2 were surveyed for bacterial strains. Three ants were surface sterilized using a modified version of a previously methodology used by our group<sup>23</sup>. Each ant was submerged in 70% ethanol during two minutes, followed by 5% of sodium hypochlorite for one minute, 70% of ethanol during one minute and rinsed three times with sterile water. The ants were crushed and transferred to ISP-2 agar plates added with antifungal agents. A non-surface sterilized ant was transferred into a sterile 1.5 mL Eppendorf tube containing 750  $\mu\text{L}$  of 0.9% NaCl isotonic aqueous solution, the tube was vortexed during approximately 1 min and 100  $\mu\text{L}$  of washing solution were inoculated on ISP-2 agar plate with antifungal agents. Additionally, three ants were also surface sterilized and the poison glands were dissected in aseptic conditions. The poison glands were inoculated on ISP-2 agar plates to isolate bacterial colonies. The bacterial isolation was performed according the procedure done for colony 1. Strains 3B2 and 3B4 were isolated in the first isolation procedure, while in the second experiment, strains ASLIM1, ASLIM5, ASLIM6, ASLIM7 were isolated from sterile ants, strain ASLFLB2 from a non-sterile ant, and strain AS1GV1B1 from a poison gland. Stocks of the eight bacterial isolates were prepared in ISP-2 or M9 liquid media with 20% of glycerol and maintained in -80 °C freezer.

The bacterial identification was carried out by sequencing the 16S rRNA, using the primers 27F and 1492R<sup>44</sup>, and the sequences were compared with GenBank sequences by BLASTn analysis. The *S. marcescens* 3B2 16S rRNA sequence has been deposited in the GenBank, and can be accessed using the GenBank accession number MF450442.

**General procedure for bacterial cultivation.** Aliquots of glycerol stocks (100  $\mu\text{L}$ ) were inoculated on ISP-2 agar plates (80 mm) containing approximately 25 mL of medium. Empty Petri plates were sealed on top of the inoculated plates, and incubated for three days at 30 °C. The polydimethylsiloxane GC-MS fibber was exposed inside of these plates during approximately 60 min to adsorb the volatile organic compounds (VOCs), which were analyzed by GC-MS.

*Serratia marcescens* 3B2 was also cultivated using the synthetic medium M9 added with 0.2% of glucose and 1.5% of the amino acids L-threonine, L-serine, L-valine and L-alanine in independent experiments. The amino acid L-threonine was also added in the M9 medium at 0.5%, 1.0%, 1.5% and 2.0%. All amino acids were dissolved with distilled water and filter sterilized before addition to the culture medium.

M9 medium (100 mL): 20 mL of M9 5X, 2 mL of glucose 20%, 200  $\mu\text{L}$  of  $\text{MgSO}_4$  1 M, 10  $\mu\text{L}$  of  $\text{CaCl}_2$  1 M, the final volume was adjusted with 100 mL of distilled water and the solution was filter sterilized. To prepare M9 agar medium, 20 g of agar in 40 mL of distilled water were autoclaved and mixed with filter sterilized: 20 mL of M9 5X, 2 mL of glucose 20%, 200  $\mu\text{L}$  of  $\text{MgSO}_4$  1 M, 10  $\mu\text{L}$  of  $\text{CaCl}_2$  1 M, and the final volume was adjusted with 100 mL of sterile distilled water.

**Chemical analysis.** The GC-MS analyses were performed on a GC-2010 Plus Shimadzu coupled with DB-5 MS column (30 m  $\times$  0.25 mm  $\times$  0.25  $\mu\text{m}$ ). The polydimethylsiloxane fibber with the VOCs was exposed during three minutes into the GC-MS injector at 250 °C. The oven temperature started with 50 °C during 5 min, followed by an increase of 3 °C/min to 180 °C and held for 10 min. Analyses were carried out in splitless injection mode, with helium as carrier gas, ion source temperature of 250 °C and 70 eV for sample ionization. VOCs were identified by comparing the GC-MS spectra with those of databases Wiley 7 and NIST 08, analyzing fragmentation patterns and comparing retention times and mass spectra with Sigma-Aldrich standards of pyrazines 1 and 2.

The LC-MS analyses were performed on an Accela UHPLC (Thermo Scientific, Carlsbad, CA, USA) interfaced with an Exactive<sup>TM</sup> Plus Orbitrap mass spectrometer (MS) (Thermo Scientific). The Kinetex  $\text{C}_{18}$  (150  $\times$  2.10 mm, 1.7  $\mu\text{m}$ ) column was eluted with mixtures of acetonitrile ( $\text{CH}_3\text{CN}$ ) and water, both add with 0.1% of formic acid, with the following gradient at flow rate of 400  $\mu\text{L}/\text{min}$ : 0–15 min of gradient from 5% of  $\text{CH}_3\text{CN}$  to 50% of  $\text{CH}_3\text{CN}$ , 15–20 min of isocratic 50% of  $\text{CH}_3\text{CN}$ , 20–30 min of gradient from 50% of  $\text{CH}_3\text{CN}$  to 100% of  $\text{CH}_3\text{CN}$ , 30–35 min of isocratic 100% of  $\text{CH}_3\text{CN}$  and 35–40 min of gradient from 100% of  $\text{CH}_3\text{CN}$  to 5% of  $\text{CH}_3\text{CN}$ . Samples were analyzed in positive ESI mode over the mass range 100 to 1500 Da, mass accuracy was below 3 ppm, capillary temperature was 320 °C and spray voltage was 3.6 kV.

**Isotope feeding experiments.** The bacterial culturing for isotope feeding experiments were performed using 10 mL GC-MS vial added with 2 mL of culture media. The medium M9 agar supplemented with 0.2% of glucose, 1.5% of L-threonine and 0.5% of L-alanine was mixed with labelled or non-labelled precursors dissolved in MilliQ water. To prepare the culture media, 0.04 g of agar were mixed with 700  $\mu\text{L}$  of MilliQ water into 10 mL SPME vials and they were sterile autoclaved. After that, the precursors were dissolved with 900  $\mu\text{L}$  of sterile water and they were mixed with 400  $\mu\text{L}$  of M9 5X salts, 40  $\mu\text{L}$  of 20% aqueous glucose solution, 4  $\mu\text{L}$  of  $\text{MgSO}_4$  1 M and 0.2  $\mu\text{L}$  of  $\text{CaCl}_2$  1 M, which were finally filter sterilized and mixed with the sterile agar. Aliquots (100  $\mu\text{L}$ ) of *Serratia marcescens* 3B2 glycerol stocks on M9 were inoculated into each 10 mL GC-MS vials containing the culture media, which were sealed and incubated for three days at 30 °C. The VOCs were extracted using a polydimethylsiloxane GC-MS fibber exposed inside of these vials during approximately 60 min, and the VOCs were analyzed by GC-MS. Isotope-labelled precursors L-[U-<sup>13</sup>C, <sup>15</sup>N]-threonine, U-<sup>13</sup>C-sodium acetate, 3-<sup>13</sup>C-L-alanine and <sup>15</sup>N-L-alanine were purchased from Sigma-Aldrich.

**Purification of compound 6.** *S. marcescens* 3B2 was cultivated onto 20 plates of ISP-2 agar (100 mL per 150 × 15 mm Petri dish) for 14 days at 30 °C and the culture medium was extracted with dichloromethane. The resulting extract was fractionated using solid-phase extraction (silica 2 g, Fisher cartridge) eluted with the following solvent gradient: 100% of hexane, 80% of hexane and 20% of ethyl acetate, 60% of hexane and 40% of ethyl acetate, 40% of hexane and 60% of ethyl acetate, 20% of hexane and 80% of ethyl acetate and 100% of ethyl acetate. Compound 6 (3.8 mg) was isolated from the fraction with 80% of hexane and 20% of ethyl acetate, which was purified using semi-preparative HPLC (Agilent 1200 series HPLC-DAD) coupled to a normal phase column (Phenomenex Luna 5 µm silica, 250 × 10 mm) and eluted with the isocratic mobile phase of 98% hexane and 2% of isopropanol (3 mL/min).

## References

- Hölldobler, B. & Wilson, E. O. *The Ants*. (Harvard University Press, 1990).
- De Fine Licht, H. H., Boomsma, J. J. & Tunlid, A. Symbiotic adaptations in the fungal cultivar of leaf-cutting ants. *Nat. Commun.* **5**, 5675, <https://doi.org/10.1038/ncomms5675> (2014).
- Weber, N. A. Fungus-growing ants. *Science* **153**, 587–604, <https://doi.org/10.1126/science.153.3736.587> (1966).
- Licht, H. H. D. & Boomsma, J. J. Forage collection, substrate preparation, and diet composition in fungus-growing ants. *Ecol. Entomol.* **35**, 259–269, <https://doi.org/10.1111/j.1365-2311.2010.01193.x> (2010).
- Fernández-Marín, H. *et al.* Functional role of phenylacetic acid from metapleural gland secretions in controlling fungal pathogens in evolutionarily derived leaf-cutting ants. *Proc. R. Soc. B* **282**, 20150212 (2015).
- Jackson, B. D. & Morgan, E. D. Insect chemical communication: pheromones and exocrine glands of ants. *Chemoecology* **4**, 125–144 (1993).
- Evershed, R. & Morgan, E. The amounts of trail pheromone substances in the venom of workers of four species of attine ants. *Insect Biochem.* **13**, 469–474 (1983).
- Morgan, E. D., Keegans, S. J., Tits, J., Wenseleers, T. & Billen, J. Preferences and differences in the trail pheromone of the leaf-cutting ant *Atta sexdens sexdens* (Hymenoptera: Formicidae). *Eur. J. Entomol.* **103**, 553–558 (2006).
- Cross, J. H. *et al.* The major component of the trail pheromone of the leaf-cutting ant, *Atta sexdens rubropilosa* Forel. *J. Chem. Ecol.* **5**, 187–203 (1979).
- Morgan, E. D., Do Nascimento, R. R., Keegans, S. J. & Billen, J. Comparative study of mandibular gland secretions of workers of ponerine ants. *J. Chem. Ecol.* **25**, 1395–1409 (1999).
- Wheeler, J. W. & Blum, M. S. Alkylpyrazine alarm pheromones in ponerine ants. *Science* **182**, 501–503 (1973).
- Cross, J. H., West, J. R., Silverstein, R. M., Jutsum, A. R. & Cherrett, J. M. Trail pheromone of the leaf-cutting ant, *Acromyrmex octospinosus* (Reich), (Formicidae: Myrmicinae). *J. Chem. Ecol.* **8**, 1119–1124 (1982).
- De Oliveira, J., Martinez, O., Carnieri, N., Vilela, E. & Reis, H. Components of the trail pheromone of the leaf-cutting ants *Atta laevigata* F. Smith and *Atta bisphaerica* Forel (Formicidae: Attini). *An. Soc. Entomol. Bras.* **19**, 145–154 (1990).
- Papenfort, K. *et al.* A *Vibrio cholerae* autoinducer-receptor pair that controls biofilm formation. *Nat. Chem. Biol.* **13**, 551–557 (2017).
- Oh, D.-C., Poulsen, M., Currie, C. R. & Clardy, J. Dentigerumycin: a bacterial mediator of an ant-fungus symbiosis. *Nat. Chem. Biol.* **5**, 391–393 (2009).
- Van Arnam, E. B., Ruzzini, A. C., Sit, C. S., Currie, C. R. & Clardy, J. A rebeccamycin analog provides plasmid-encoded niche defense. *J. Am. Chem. Soc.* **137**, 14272–14274 (2015).
- Van Arnam, E. B. *et al.* Selvamycin, an atypical antifungal polyene from two alternative genomic contexts. *Proc. Natl. Acad. Sci. USA* **113**, 12940–12945 (2016).
- Wada-Katsumata, A. *et al.* Gut bacteria mediate aggregation in the German cockroach. *Proc. Natl. Acad. Sci. USA* **112**, 15678–15683, <https://doi.org/10.1073/pnas.1504031112> (2015).
- Dillon, R. J., Vennard, C. T. & Charnley, A. K. Pheromones: exploitation of gut bacteria in the locust. *Nature* **403**, 851 (2000).
- Brown, W. & Moore, B. Volatile secretory products of an Australian formicine ant of the genus *Calomyrmex* (Hymenoptera: Formicidae). *Insect Biochem.* **9**, 451–460 (1979).
- Poulsen, M. & Currie, C. R. Symbiont interactions in a tripartite mutualism: exploring the presence and impact of antagonism between two fungus-growing ant mutualists. *PLoS One* **5**, e8748 (2010).
- Marsh, S. E. *et al.* Association between *Pseudonocardia* symbionts and *Atta* leaf-cutting ants suggested by improved isolation methods. *Int. Microbiol.* **16**, 17–25 (2013).
- Paludo, C. R. *et al.* Whole-Genome Sequence of *Bacillus* sp. SDLI1, Isolated from the Social Bee *Scaptotrigona depilis*. *Genome Announc.* **4**, e00174–00116 (2016).
- Shu, C. K. Pyrazine formation from serine and threonine. *J. Agr. Food Chem.* **47**, 4332–4335, <https://doi.org/10.1021/jf9813687> (1999).
- Nelson, D. L. & Cox, M. M. *Lehninger Principles of Biochemistry*. (W. H. Freeman and Company, 2004).
- Sit, C. S. *et al.* Variable genetic architectures produce virtually identical molecules in bacterial symbionts of fungus-growing ants. *Proc. Natl. Acad. Sci. USA* **112**, 13150–13154 (2015).
- Schoenian, I. *et al.* Chemical basis of the synergism and antagonism in microbial communities in the nests of leaf-cutting ants. *Proc. Natl. Acad. Sci. USA* **108**, 1955–1960 (2011).
- Gary, N. E. Chemical mating attractants in the queen honey bee. *Science* **136**, 773–774 (1962).
- David Morgan, E. Trail pheromones of ants. *Physiological Entomology* **34**, 1–17 (2009).
- Morgan, E. D. Chemical sorcery for sociality: exocrine secretions of ants (Hymenoptera: Formicidae). *Myrmecol. News* **11**, 79–90 (2008).
- Showalter, D. N., Troyer, E. J., Aklu, M., Jang, E. B. & Siderhurst, M. S. Alkylpyrazines: alarm pheromone components of the little fire ant, *Wasmannia auropunctata* (Roger) (Hymenoptera, Formicidae). *Insect. Soc.* **57**, 223–232, <https://doi.org/10.1007/s00040-010-0075-4> (2010).
- Dossey, A. T., Gottardo, M., Whitaker, J. M., Roush, W. R. & Edison, A. S. Alkylidimethylpyrazines in the defensive spray of *Phyllium westwoodii*: a first for order Phasmatodea. *J. Chem. Ecol.* **35**, 861–870 (2009).
- Bohman, B. *et al.* Discovery of pyrazines as pollinator sex pheromones and orchid semiochemicals: implications for the evolution of sexual deception. *New Phytol.* **203**, 939–952 (2014).
- Byers, J. Pheromone biosynthesis in the bark beetle, *Ips paraconfusus*, during feeding or exposure to vapours of host plant precursors. *Insect Biochem.* **11**, 563–569 (1981).
- Reddy, G. V. & Guerrero, A. Interactions of insect pheromones and plant semiochemicals. *Trends Plant Sci.* **9**, 253–261 (2004).
- Aylward, F. O. *et al.* Complete genome of *Serratia* sp. strain FGI 94, a strain associated with leaf-cutter ant fungus gardens. *Genome Announc.* **1**, e00239–00212 (2013).
- Aylward, F. O. *et al.* Convergent bacterial microbiotas in the fungal agricultural systems of insects. *MBio* **5**, e02077–02014 (2014).
- Komatsubara, S., Murata, K., Kisumi, M. & Chibata, I. Threonine degradation by *Serratia marcescens*. *J. Bacteriol.* **135**, 318–323 (1978).

39. Molla, G. *et al.* Aminoacetone oxidase from *Streptococcus oligofermentans* belongs to a new three-domain family of bacterial flavoproteins. *Biochem. J.* **464**, 387–399, <https://doi.org/10.1042/bj20140972> (2014).
40. Fehlbaum, P., Rao, M., Zasloff, M. & Anderson, G. M. An essential amino acid induces epithelial  $\beta$ -defensin expression. *Proc. Natl. Acad. Sci. USA* **97**, 12723–12728 (2000).
41. Crawford, J. M., Kontnik, R. & Clardy, J. Regulating alternative lifestyles in entomopathogenic bacteria. *Curr. Biol.* **20**, 69–74 (2010).
42. Amin, S. *et al.* Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* **522**, 98 (2015).
43. Segev, E. *et al.* Dynamic metabolic exchange governs a marine algal-bacterial interaction. *Elife* **5**, e17473 (2016).
44. Cafaro, M. J. *et al.* Specificity in the symbiotic association between fungus-growing ants and protective *Pseudonocardia* bacteria. *Proc. R. Soc. B* **278**, 1814–1822 (2011).

## Acknowledgements

The authors thank the Fogarty International Center, National Institutes of Health (NIH), USA (grant U19TW009872); São Paulo Research Foundation (FAPESP), Brazil (grants #2013/50954-0, 2013/04092-7, 2012/24204-1, 2012/22487-6, 2014/24330-2, 2015/25301-9); CAPES and CNPq for funding and research support; and Norberto Lopes and Izabel Turatti for helping with GC-MS analysis; Emily Mevers for helping with NMR analysis. ACR was supported by a postdoctoral fellowship from the Canadian Institutes of Health Sciences.

## Author Contributions

E.A.S.J., A.C.R., C.R.P., F.S.N., C.R.C., J.C., and M.T.P. designed research; E.A.S.J., A.C.R. and C.R.P. performed research; E.A.S.J., A.C.R., C.R.P., F.S.N., C.R.C., J.C., and M.T.P. analyzed data; E.A.S.J., A.C.R., C.R.P., J.C. and M.T.P. wrote the paper. All authors reviewed the manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <https://doi.org/10.1038/s41598-018-20953-6>.

**Competing Interests:** The authors declare no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018