

**Serveur Académique Lausannois SERVAL [serval.unil.ch](https://serval.unil.ch)**

## **Author Manuscript**

**Faculty of Biology and Medicine Publication**

**This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.**

Published in final edited form as:

**Title:** Chaperones convert the energy from ATP into the nonequilibrium stabilization of native proteins.

**Authors:** Goloubinoff P., Sassi A.S., Fauvet B., Barducci A., De Los Rios P.

**Journal:** Nature Chemical Biology

**Year:** 2018

**Volume:** 14(4)

**Pages:** 388-395

**DOI:** [10.1038/s41589-018-0013-8](https://doi.org/10.1038/s41589-018-0013-8)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

# **Molecular chaperones convert the energy from the ATPase cycle into the non-equilibrium stabilisation of native proteins**

Pierre Goloubinoff<sup>1\*</sup>, Alberto S. Sassi<sup>2</sup>, Bruno Fauvet<sup>1</sup>, Alessandro Barducci<sup>3,4</sup>, and Paolo De Los Rios<sup>2\*</sup>

<sup>1</sup>Department of Plant Molecular Biology, Université de Lausanne, 1015 Lausanne, Switzerland

<sup>2</sup>Institute of Physics, School of Basic Sciences and Institute of Bioengineering, School of Life Sciences, École Polytechnique Fédérale de Lausanne – EPFL, 1015 Lausanne, Switzerland.

<sup>3</sup>Inserm, U1054, Montpellier, France

<sup>4</sup>Université de Montpellier, CNRS, UMR 5048, Centre de Biochimie Structurale, Montpellier, France

\*Corresponding authors

## **Abstract**

Protein homeostasis, namely the ensemble of cellular mechanisms collectively controlling the activity, stability and conformational states of proteins, depends on energy-consuming processes. *De novo* protein synthesis requires ATP hydrolysis for peptide bond formation. Controlled degradation by the chaperone-gated proteases requires ATP hydrolysis to unfold target proteins and render their peptide bonds accessible to hydrolysis. During and following translation, different classes of molecular chaperones require ATP hydrolysis to control the conformational state of proteins, favor their folding into their active conformation and avoid, under stress, their conversion into potentially harmful aggregates. Furthermore, specific ATP-fueled unfolding chaperones can dynamically revert aggregation itself. Why do molecular chaperones need ATP and what are the consequences on protein homeostasis of the energy contributed by the ATPase cycle? Here we used various biochemical assays and physical modeling and showed that both bacterial chaperones GroEL (HSP60) and DnaK (HSP70) can use part of the energy liberated by ATP to preserve the native state population of their substrate proteins, even under conditions in which the native state is, according to equilibrium thermodynamics, unstable.

Expectedly, upon ATP exhaustion the chaperone substrates spontaneously lose their activity, reverting to their equilibrium non-native state. In the presence of chaperones, proteins are thus open, driven, *non-equilibrium* systems, whose behavior is only partially determined by equilibrium thermodynamics, with intriguing and far-reaching consequences for our understanding of protein homeostasis, protein biophysics and, more generally, protein evolution.

## Introduction

Anfinsen *et al.*<sup>1</sup> demonstrated that artificially unfolded proteins can refold spontaneously into their native, active state without assistance from other macromolecules, solely guided by their primary amino-acid sequences. These seminal *in vitro* refolding experiments implied that, compared to unfolded nascent polypeptides, natively folded proteins are thermodynamically more stable conformations positioned at the bottom of the free energy landscape. This can be reasonably expected from 3.5 billion years of evolution that strived to produce optimally active proteins capable of sustaining long-lasting cellular functions in fluctuating stressful environments. Nonetheless, Anfinsen also noticed that, *in vitro*, the spontaneous folding process can be prone to errors as not all artificially unfolded proteins could efficiently reach their native state, and formed instead inactive insoluble aggregates. Indeed, although the funnel-like structure of the free-energy landscape of proteins should in general favour their native folding<sup>2,3</sup>, local minima, corresponding to metastable misfolded conformations, can transiently halt polypeptides on their way to the native state. There, the kinetically trapped species might pursue further stability by coalescing into stable inactive aggregates, as in the case of alpha-synuclein fibrils and beta-amyloids. Aggregates can be stabilised by intra- and inter-molecular contacts of otherwise inappropriately exposed hydrophobic surfaces and are typically enriched in cross  $\beta$ -sheets<sup>4</sup>, and at sufficiently high protein concentrations their free-energy might be lower than the native state<sup>5-7</sup>. Moreover, mutations and environmental stresses, such as heat-shock, can perturb the free-energy landscape and render the native state intrinsically less stable than other partially compact, misfolded and aggregated conformations<sup>8</sup>. Because protein aggregates and

their misfolded precursors appear to be the primary cause for various degenerative diseases, they have gained great interest in the last decades<sup>9,10</sup>.

Molecular chaperones have come to prominence because of their role in assisting *de novo* protein folding and in preventing under stress the formation of toxic aggregates, by several complementary mechanisms involving protein binding and unfolding<sup>5,11,12</sup>. Attesting for their foremost function in sustaining cellular life, up to 3-5 % of the total protein mass of unstressed non-cancerous metazoan cells may be constituted from the conserved families of ATPase chaperones, namely the 60 kDa Heat Shock Protein, Hsp60 (GroEL, the bacterial chaperones in parenthesis), Hsp70 (DnaK), Hsp100 (ClpB) and Hsp90 (HtpG)<sup>13</sup>. This amount may further increase following heat stress<sup>14</sup>, or in immortalized cancer cells, which, owing to their abnormal elevated chaperone load, resist apoptosis and protein damage under various chemotherapeutic and physical stresses<sup>15</sup>. Because of their affinity for water-exposed hydrophobic amino acids, chaperones preferentially bind non-native proteins exposing hydrophobic surfaces, thereby passively preventing protein aggregation<sup>16-18</sup>. The role of the chaperone ATPase cycle has been traditionally, but possibly oversimplistically, associated with the directional cycling through several different molecular conformations, each optimally primed for one of the various steps of the chaperone action: binding, “processing” and release of their to-be refolded, or to-be prevented from aggregating<sup>19</sup>, polypeptide substrates.

Quite surprisingly, a fundamental consequence of the ATPase activity of chaperones, which stems from fundamental laws of physics and remains largely under-appreciated, is that the constant contribution of energy from ATP, aside from generating heat, could also drive the system into a *non-equilibrium steady state*, where proteins interacting with the chaperones would distribute over their

conformational ensemble differently than at equilibrium<sup>20-23</sup>. A testable prediction in such scenario is that, using the energy available from ATP, chaperones should be able to favour the native state of their substrates even under adverse denaturing conditions, where the native state is thermodynamically less stable than the various unfolded, misfolded and aggregated species.

Here, we primarily focussed on bacterial GroEL, which was the first chaperone to be studied *in vivo*<sup>24</sup> and *in vitro*<sup>16</sup> and thereafter thoroughly characterized structurally and functionally<sup>25,26</sup> and on malate dehydrogenase (MDH), which is an extensively studied natural substrate of GroEL in mitochondria and bacteria, as well as on citrate synthase, another well-studied model protein. Further, we show that the same non-equilibrium principles likely apply both to the network of chaperones, composed by GroEL and GroES, the disaggregase ClpB and DnaK, that in cells supervises protein homeostasis, and to Hsp70 (DnaK) alone, which is an abundant ATPase chaperone representing at least 0.5% of the total protein mass of animal cells<sup>13,15</sup>. We find that similarly to GroEL, these systems can also convert the free energy from the ATPase cycle into the *non-equilibrium* stabilisation of the native state of labile protein substrates, by mechanisms that, although very different at the molecular level from cage sequestration by GroEL, are likely analogous in their overarching principle.

### **The free-energy landscape of proteins in non-native conditions**

Purified MDH from porcine mitochondria, which is the model protein that we used in most of the experiments, is stable at 25°C, while it spontaneously loses its activity at more elevated temperatures<sup>27</sup>, such as 37°C<sup>28</sup>. These results suggest a picture of the

free-energy landscape of MDH at 37°C as represented in Fig.1, where the native, active dimer (ND) is in kinetic equilibrium with the compact, native monomer (NM), which is in turn in kinetic equilibrium with a less-structured intermediate (I) characterized by a higher free-energy. The intermediate is likely short-lived, and it is in kinetic equilibrium also with a collapsed but non-native misfolded state (M)<sup>29-31</sup>, which is more stable (*i.e.* lower in free-energy) even than the native dimer.

Aggregation, which is a concentration dependent reaction, further proceeds by the binding of non-native monomers to other non-native monomers or to already formed aggregates (A). Since it has been noted that non-native MDH remains in conformations that, when brought back into native conditions, could refold spontaneously in native conditions even after incubation at denaturing temperatures for tens of minutes<sup>32</sup>, the aggregation is very likely a slow process compared to denaturation at micro-molar, or lower, concentrations. These findings also agree with observations that several non-native proteins can remain soluble for long periods of time, even under adverse conditions that do not favour the native state<sup>8,33,34</sup>.

### **Under denaturing conditions GroEL and ATP maintain MDH in a *non-equilibrium* native state**

Hartman *et al.*<sup>28</sup> remarkably found that at 37°C the rate of spontaneous denaturation of MDH decreased in the presence of substoichiometric amount of GroEL and GroES 7-mers and ATP. These intriguing observations did not trigger a more fundamental issue: does GroEL use ATP hydrolysis to seemingly “preserve”, under a denaturing temperature, a conformational state that is not the most stable one

as dictated by thermodynamic equilibrium? We address here this crucial question, which has remained to date unanswered.

At 25°C, as expected, native MDH remained stably active with or without GroELS and ATP (Fig.S1A). At 37°C, MDH spontaneously lost its active form at an approximate rate of  $0.007 \text{ min}^{-1}$  (Fig.2A), compatibly with Hartman *et al.*<sup>28</sup> and with the scheme in Fig.1. Addition of a molar excess of GroELS (3.5  $\mu\text{M}$  protomers of GroEL and GroES) (but without ATP) over MDH did not affect the rate of denaturation (Fig.S1B). In contrast, when GroELS, ATP and a pyruvate kinase/phosphoenolpyruvate-based regeneration system were supplemented at  $t=0'$ , they caused a net reactivation of nearly 75 nM MDH on top of the initial amount of active enzyme. This implied that in the initial stock of nominally native MDH there was at most 80-85% of truly native, active MDH species, alongside ~15-20% inactive yet soluble and GroELS-amenable species, in keeping with the results of Peralta *et al.*<sup>32</sup>, since no protein aggregate has yet been shown to be solubilized by GroELS+ATP<sup>12,35</sup>.

Surprisingly, following reactivation MDH remained active and apparently stable, despite the denaturing conditions in which the active state should not have persisted. When GroELS and ATP were added later after the start of thermal denaturation, native state maintenance again followed the net reactivation of some MDH (Fig.2B), which had remained chaperone-amenable during 30-60 minutes of the heat treatment. The amounts of recovered MDH significantly decreased only when GroELS and ATP were added at much later times, once more suggesting that the formation of aggregates (Fig.2B, non-recoverable fraction) was much slower than the denaturation process in our assay conditions.



Our observations highlighted that at 37°C, where the non-native ensemble is more stable than the native ensemble (Fig.1), the primary ATP-dependent action of GroELS was not limited to the mere prevention of aggregation of the non-native MDH. Instead, as represented in Fig.1, GroELS behaved as if it actively rescued the recoverable misfolded MDH conformers, pushing them up the free-energy landscape towards the native active state, where, despite being only metastable, they were maintained for extended periods. These conclusions were further strengthened by a similar action of GroELS+ATP during heat denaturation of citrate synthase, another classical GroEL substrate (Fig.S2)<sup>36</sup>.

Basic physics principles dictate that the steady maintenance of MDH in its otherwise unstable native state must be strictly ATP-dependent. Indeed, if no energy could be extracted from the ATPase cycle, equilibrium thermodynamics would inescapably govern the system, and GroELS could not oppose the thermodynamically inevitable denaturation of the MDH. We thus devised experimental setups where we could control whether the system was at, or away from, equilibrium.

As in Fig.2A, an excess of GroELS over MDH, supplemented with ATP and a regeneration system, was expectedly able to maintain the native MDH level when added from the start and, when added at a later stage, to recover some activity and then maintain it constant despite the denaturing temperature (Fig.2C). In both cases, the subsequent addition of apyrase, which readily hydrolysed ATP into AMP, deprived GroELS of its energy source, inhibiting its ability to maintain the non-equilibrium active stabilisation of the metastable native MDH at 37°C. As a consequence, the system spontaneously sought the MDH inactive denatured state, which is thermodynamically the most stable (by contrast, at 25°C, where native MDH is intrinsically stable, addition of apyrase had no effect, see Fig.S1A).

Using MDH that was first artificially unfolded by urea, Anfinsen-type *in vitro* refolding experiments showed that, following removal of urea by abrupt dilution, there was no spontaneous MDH refolding at 37°C (Fig.2D), consistent with the thermodynamic instability of the native state at that temperature. In contrast, we found that the presence of GroELS+ATP at the time of the urea dilution resulted in the transient major accumulation of native MDH at 37°C, and in its subsequent maintenance over time. Apyrase addition readily brought the level of native MDH back under the control of equilibrium thermodynamics, and led to its spontaneous denaturation despite the presence of the chaperones (Fig.2D).

The amount of energy that is liberated by the ATPase cycle,  $\Delta G_{ATP}$ , and that is available for GroELS to inject into maintaining the metastable MDH into its native state, depends on the concentrations of ATP, ADP and inorganic phosphate Pi through the relation:

$$\Delta G_{ATP} = k_B T \left\{ \ln([ATP]/([ADP][Pi])) - \ln([ATP]_{eq}/([ADP]_{eq}[Pi]_{eq})) \right\}.$$

Here,  $k_B$  is the Boltzmann constant,  $T$  is the absolute temperature and the *eq* subscript marks the concentrations that the different molecules would have if the hydrolysis reaction could run to completion (*i.e.*  $[ADP]_{eq}/[ATP]_{eq} \approx 10^6$ - $10^8$  depending on the solution conditions<sup>37</sup>). To analyse the role of  $\Delta G_{ATP}$  in more detail, we changed the concentrations of ADP in the GroELS-mediated refolding experiments of urea pre-unfolded MDH at 37°, while keeping ATP constant at a sub-physiological concentration (400  $\mu$ M), which was, however, still in excess over GroELS (3.5  $\mu$ M). Increasing amounts of ADP, from 0  $\mu$ M to 1600  $\mu$ M, were added at the start of the reaction, with the goal of progressively reducing the value of  $\Delta G_{ATP}$ , as predicted from the above formula. Expectedly, larger ADP concentrations inhibited the initial

effectiveness of the chaperone mediated refolding reaction of the MDH (Fig.2E). Later, the reaction was further hindered by the decreased concentrations of ATP that was consumed, and by the reciprocal increased concentration of ADP. This brought the system back towards equilibrium conditions, namely  $\Delta G_{\text{ATP}}=0$ .

Together, the results in Figs.2C-E confirmed that the effect of GroELS is strictly dependent, and tuneable, by the amount of energy available from ATP, which thus drove and maintained the system in a non-equilibrium state.

### **Non-equilibrium thermodynamics model of chaperone action**

We next rationalized these observations with a model that captured the main properties of the free-energy landscape of MDH and of the ATP-fuelled chaperone action as represented in Fig.1. We chose to describe the action of chaperones at a minimalistic level of detail to focus on the broader and general thermodynamic implications of the energy consumed during their cycle. The explicit reactions of chaperone-substrate association and dissociation allowed accounting for the chaperone concentrations that, when needed, we chose to vary in order to serve our experimental demonstration. When bound to chaperones, substrates could further convert between more and less structured non-native conformations, based on the compelling experimental evidence for Hsp60s (and for Hsp70s and Hsp100s) that following binding, misfolded substrates generally undergo a loss of secondary structure (unfolding) and an expansion, while still in complex with the chaperones<sup>34,38-46</sup>, and as emphasized in other modelling schemes for the action of GroELS<sup>47-49</sup>. This assumption was further strengthened by observing that the compacting action of a crowding agent (15% PEG 6000)<sup>50,51</sup>, which inhibited both the misfolded-to-native

and native-to-misfolded conversions of MDH at 25°C and at 37°C (Fig.S5A), was antagonized by GroELS+ATP at 37°C (Fig.S5B). In other words, the energy of ATP was used by GroELS to facilitate the de-compaction and/or unfolding of its substrates.

In this model, the native species do not interact with the chaperones, in agreement with the generally-accepted view that in order to act efficiently in the crowded environment of cells, molecular chaperones must have a high affinity for non-native, aggregation-prone *de novo* synthesized, *de novo* translocated or stress-misfolded polypeptides, while keeping an extremely low affinity for the very dense population of surrounding proteins that are native<sup>52</sup>.

The detailed internal movements of GroEL, as well as GroES binding and release, the precise steps of ATP hydrolysis and nucleotide exchange, and their cooperative and/or anti-cooperative nature are usually taken into account when addressing the precise molecular steps of the cycle of the action of GroEL<sup>53-56</sup>. Here, they needed not to be explicitly represented in the model, as they were not necessary to extract the effects of the energy consumed by the chaperones on the behaviour of the system. Indeed, according to the laws of non-equilibrium thermodynamics, which must be valid at any coarse-grained level of description, the effect of the energy  $\Delta G_{\text{ATP}}$  contributed to the system by the ATPase activity could be rigorously integrated in the relations between the association and dissociation rates of the chaperone-substrate complexes (see SI).

The rates of the transitions between the different MDH states, the rate of aggregation and the rates of the chaperone action have been chosen to reasonably reproduce the measured spontaneous denaturation of native MDH dimers at 37°C and the third reactivation curve (when GroELS and ATP were added at  $t=30'$ ) observed in

Fig.2A (see SI for the full list of rates). All model results presented in this work have been obtained with this same set of intrinsic parameters.

Once the parameters had been set, we found that the model closely reproduced the other data in Fig.2A, when the GroELS was added either earlier or later during the thermal denaturation (*cf.* Fig.3A). More importantly, the model gave access to estimates of the various populations of MDH species that were not directly detectable experimentally (Fig.3B). The chaperone-resistant non-recoverable fraction was heterogeneous: it comprised an irreversibly inactive component, likely aggregates, which expectedly grew over time at 37°C, and a proportionally diminished soluble component, which was mostly composed of near-native inactive monomers.

In the model, the switch from non-equilibrium to equilibrium conditions induced by apyrase could be captured by sharply setting  $\Delta G_{\text{ATP}}=0$  at the time of apyrase addition, and the results agreed with the experiments (compare Fig.3C with Fig.2C). Similarly, we could reproduce the results from Anfinsen-type refolding of artificially unfolded MDH experiments (Fig.3D). In the model, we could also control the amount of energy available to chaperones by changing the [ATP]/[ADP] ratio and taking into account its variation in time (see SI for the implementation in the model). The same behaviour observed experimentally in Fig.2E. was reproduced by the model (Fig.3E), further confirming that the energy of ATP was transduced into the stability of the native state of MDH.

### **GroELS functions through iterative cycles fuelled by the ATPase activity.**

We then investigated whether the observed ATP-dependent ability of GroELS to regenerate and transiently maintain native MDH under denaturing conditions (Fig.1A)

was possibly due to the stabilisation during long-lived in-cage binding, of a minority of denaturing polypeptide substrates by a molar excess of GroELS.

We thus exposed during 8 hours, an MDH concentration higher than in previous experiments ( $2.5 \mu\text{M}$ ) to decreasing substoichiometric amounts of GroELS in the presence of a non-limiting amount of ATP (backed by a long-lasting ATP regeneration system) (Fig.4A). Following 8 hours at  $25^\circ\text{C}$ , the native MDH expectedly remained nearly fully native, independently of the presence or absence of GroELS. Without ATP, all GroELS concentrations remained ineffective at preserving at  $37^\circ\text{C}$  the MDH activity and 85% of the total MDH, *i.e.* about  $2.1 \mu\text{M}$ , became inactive in 8 hours. However, in the presence of ATP, as little as  $357 \text{ nM}$  of GroEL<sub>7</sub> rings ( $2.5 \mu\text{M}$  of GroEL protomers) could actively maintain a 7-fold excess of MDH in the native state during the 8 hours at  $37^\circ\text{C}$ . At  $\text{EC}_{50}$ , as little as  $56 \text{ nM}$  of GroEL<sub>7</sub> rings ( $392 \text{ nM}$  of GroEL protomers) could effectively maintain in the native state as many as  $923 \text{ nM}$  of MDH, implying that on average, each GroEL<sub>7</sub> cavity had actively and iteratively converted at least 17 molecules of inactive MDH monomers into active dimers, more than it could bind at any given time during the 8 hours (Fig.4A, inset). This ruled out the possibility that the stabilisation of the native species resulted from the lengthy association with the chaperone cage. Without changing parameters, the model could closely reproduce these experimental data (Fig.4B), including the minimal number of recovered MDH molecules, which was about 20 per GroEL<sub>7</sub> ring at  $\text{EC}_{50}$  (Fig.4B inset, red squares). Moreover, because at any moment of the 8 hours reaction each MDH monomer that was released from the chaperone cavity and that had natively refolded and possibly transiently reassembled into an active dimer could denature again, the model further unravelled that each GroEL<sub>7</sub> cavity had to undergo at least 400 ATP-fuelled cycles of MDH binding, processing and release in order to

yield the measured elevated end-product amounts of native MDH after 8 hours at 37°C (Fig.4B inset, blue squares).

These results imply that continuously misfolding polypeptides must have been effectively converted into native complexes against thermodynamic equilibrium through a highly iterative process which is similar to catalysis, because the conversion between the misfolded and the intermediate state is much faster than it would be spontaneously, and where the catalyst molecule (here, GroELS) is available for multiple turnovers of the reaction. Yet, at variance with catalysis, this process is not accelerated in both directions because of energy consumption, leading to the observed change of the native state fraction in steady-state.

### **GroELS uses the energy of ATP to shape the effective free-energy landscape of proteins**

The results in Figs.2 and 4 indicate that by readily processing the thermally misfolded proteins as they formed and by releasing the processed species to transiently form native complexes, GroELS and ATP were in fact maintaining the concentration of the aggregation-prone misfolded proteins very low, and consequently dramatically reduced the process of aggregation that strongly depends on the concentration of misfolded precursors. This justified our further simplification of the model by plainly neglecting the contribution of aggregation. It then became possible to formally define the steady-state *effective* free-energy difference between the native and non-native ensembles,  $\Delta G_{N,nN}^{\text{eff}} = -k_B T \ln([N]/[nN])$  (see SI for a discussion) as a function of  $\Delta G_{\text{ATP}}$ . Here,  $[N]$  was the total concentration of native conformers ( $[N]=[NM]+[ND]$ ) and  $[nN]$  the total concentration of non-native conformers

([nN]=[I]+[M]+[CI]+[CM]) (see Fig.1). Expectedly, at 37°C, in equilibrium conditions ( $\Delta G_{\text{ATP}}=0$ ) the non-native ensemble was more stable than the native ensemble ( $\Delta G_{\text{N,nN}}^{\text{eff}}=\Delta G_{\text{N,nN}}^{\text{eq}}>0$ ), driving the steady spontaneous denaturation of native MDH (Fig.5A). As  $\Delta G_{\text{ATP}}$  increased by injecting progressively more energy in the reaction, the native ensemble became stabilized. For  $\Delta G_{\text{ATP}}> 2.7$  kcal/mol, the native ensemble became more stable than the non-native ensemble. For  $\Delta G_{\text{ATP}}=7.7$  kcal/mol (the value used in the model in Figs.2, 3A-D and 5C), the native-ensemble was found to be apparently more stable than the non-native ensemble by about 1.8 kcal/mol, corresponding to a total energy transfer by the chaperone of about 2 kcal/mol from the ATPase cycle to the steady-state stability of the native state product. Pictorially, the effect of the action of GroELS thus corresponded to a remodelling of the free-energy landscape of the substrate, where the misfolded state was destabilised and raised above the native state, as it would be expected under physiological conditions (Fig.5B).

The model also rationalized the role that an elevated cellular concentration of GroELS (an estimation from quantitative proteomics of a rat liver cells indicates that in the mitochondrial stroma there are about 130  $\mu\text{M}$  HSPD1 (GroEL) protomers, as many HSPE1 (GroES) protomers, for about  $\sim 160$   $\mu\text{M}$  of MDH2 protomers<sup>13</sup>), might have for the evolution of proteins carrying cellular functions necessitating high flexibility at an unavoidable cost of decreased overall intrinsic stability (Fig. 5C). As shown by Anfinsen, intrinsically stable native proteins ( $\Delta G_{\text{N,nN}}^{\text{eq}}<0$ ) may not need assistance from other proteins. Here, we found, however, that as the thermodynamic stability of labile proteins was progressively challenged by mild stress or mutations (corresponding to increasing values of  $\Delta G_{\text{N,nN}}^{\text{eq}}>0$ ), molecular chaperones could



recruit the energy of ATP to maintain in their native state, proteins that are intrinsically metastable, thereby extending their effective range of activity within stressed cells (the orange region in Fig.5C represents the net gain of native proteins over the amount that should be present according to equilibrium thermodynamics). Noticeably, even under harsh denaturing conditions, where an excess of GroELS is unable to maintain MDH in the native state, in the model, it was still able to bind the non-native aggregation-prone species (blue region). When GroELS was in excess over the substrate, it could reduce the amount of free non-native species (green region), thereby impeding their irreversible aggregation and justifying, in part, the somewhat overstated characterizing of molecular chaperones as being able to prevent the aggregation of other proteins<sup>57</sup>.

### **The full power of energy transduction is unleashed by the entire chaperone network**

In bacterial cells, protein homeostasis depends on several ATPase chaperones, comprising GroELS, DnaK (and its co-chaperones), HtpG and the disaggregase ClpB<sup>58</sup>. We thus tested the effects of a more physiological multi-chaperone composition, for various concentrations of DnaK, on MDH at 37°C (Fig.S4). At 37°C, when urea-unfolded MDH was pre-incubated for 70 minutes with ATP, ClpB, limiting amounts of GroELS, and the DnaK co-chaperones DnaJ and GrpE, but without DnaK, no significant native MDH refolding was observed. The subsequent addition of increasing amounts of DnaK caused the increasing accumulation of native MDH, despite the non-native temperature, demonstrating that DnaK, which can act both upstream and downstream of ClpB<sup>59</sup> and of GroELS<sup>60,61</sup>, can facilitate the conversion of the energy of ATP into increased stability of labile native proteins.

As matter of fact, early *in vitro* experiments with DnaK already indicated that also Hsp70 chaperones are able to convert the energy of ATP into a non-equilibrium native state stabilisation of the thermolabile luciferase at denaturing temperature<sup>62</sup>. We further demonstrated here that this is an intrinsically non-equilibrium effect, by showing that upon the addition of apyrase, spontaneous denaturation of luciferase was restored (Fig.S3A). We further tested the ability of DnaK to stabilize native MDH at 37°C, and observed that the addition of KJE and ATP produced a significant reactivation of native MDH, although only short-lived likely because the rate of MDH denaturation was faster than the slow pace of DnaK-mediated reactivation (Fig.S3B). We conclude thus that the KJE+ATP chaperone system is also able to transduce the energy of ATP into the non-equilibrium stabilisation of the native state of both Luciferase and MDH at 37°C, even if, in the case of MDH, less efficiently than GroELS.

Together, these results suggest that distinct ATPase chaperones may act individually as well as cooperatively at the effective injection of energy from ATP to artificially maintain labile proteins in their native states under stressful conditions where they spontaneously tend to misfold and aggregate, likely because aggregates represent the true free-energy minimum and are thus thermodynamically most stable<sup>5,6</sup>. In the cell, the conformational landscape could thus be modified even more dramatically than by GroELS alone, by raising the apparent free-energy, not only of misfolded monomeric conformers, as in Fig.5B, but also of already formed stable aggregates, thereby further extending the conditions in which labile proteins can remain mostly native.

## Conclusions

Our parallel experimental and theoretical findings have revealed that, at its most fundamental level, the role of ATPase chaperones such as GroEL, DnaK and ClpB, is to transduce the energy liberated by the ATPase cycle into an enhanced *non-equilibrium* steady-state stability of otherwise labile native protein substrates, even under conditions where their native states are unstable. In this respect, the function of chaperones is more deeply rooted in non-equilibrium thermodynamics than in the molecular aspects of the ATP-driven conformational transitions in the chaperone complexes, that we have neglected here in favour of a simplified model in which we considered only the constraints imposed by thermodynamics, that mandatorily apply independently of the level of simplification of the chaperone cycle. Thus, at the price of missing part of the molecular accuracy characterizing more complex simulation schemes<sup>63,64</sup>, our simpler strategy provided a general rigorous thermodynamic perspective that broadly applies to many ATPase chaperones.

In a cellular perspective, we may thus propose that ATPase chaperones in general supply cells with an energy consuming mechanism to withstand prolonged destabilizing stresses, such as the diurnal cycle of temperatures, by maintaining the most labile members of the proteome in a soluble functional state until the environmental stress is over. Chaperones might thus increase the freedom of evolution to accumulate functionally favourable but destabilizing mutations<sup>65</sup>, by artificially maintaining mutated proteins active and functional in the cell, against their thermodynamic tendency to spontaneously denature. This view is compatible with directed evolution experiments that have demonstrated that, when chaperones are

over-expressed, enzymes can evolve faster, remain active and soluble even when challenged by slightly destabilizing mutations<sup>66-69</sup>.

Our observation with labile proteins, together with the emerging evidence that ATP-dependent mechanisms can also actively maintain RNA molecules in steady-state conformations that are different from the ones they seek to adopt at equilibrium<sup>70-72</sup>, lends further support to the general concept of the cytoplasm being an “active milieu”<sup>73,74</sup>. Indeed, when ATP becomes limiting, essential biological functions, such as the maintaining of steep ion gradients across membranes, collapse and lead to cell death. Thus, energy-consuming nanomachines need to be constantly at work to maintain cells in a metabolically active state. In particular, stress-labile macromolecules, such as DNA, RNA and proteins damaged by exposures to various physical and chemical stresses, need to be constantly proofread by enzymes, such as topoisomerases, helicases, RNA chaperones and now, as we show, also by protein chaperones, which, at an energy cost, can repair and convert them to their active, functional conformations. These ATP-fuelled enzymes can thus remodel the free-energy landscape of biological macromolecules, which consequently become less limited by the dictates of equilibrium thermodynamics. This calls for questioning the correlation between the behaviour of marginally stable biological macromolecules observed *in vitro*, as compared to their effective behaviour in the highly crowded, chaperone-rich environment of the cell. From the perspective of the landscape theory of protein folding, which posits that the native state sits at the bottom of a funnel-shaped free-energy basin<sup>2,3</sup>, our results suggest that the presence of other minima, possibly lower in energy than the native state, is not going to affect the correct the folding as long as the corresponding conformations can be processed by chaperones.

The present results are part of a growing recognition that all aspects of life, from its origins billions of years ago to its present most intimate workings, depend on a constant influx and subsequent dissipation of energy. Only the accounting of this energy budget, and of the way it is used to offset very complex biological systems away from their natural tendency to reach thermodynamic equilibrium, can lead to a correct quantitative description of biological systems. As Erwin Schrödinger stated, “Living matter evades the decay to equilibrium”<sup>75</sup>. Chaperones represent a vivid incarnation of Schrödinger’s fundamental insight.

## **Materials and Methods**

### **Proteins:**

GroEL, GroES were purified according to Torok *et al.*<sup>76</sup>. DnaK, DnaJ were purified as according to Gur *et al.*<sup>77</sup>. GrpE was a gift from H.-J. Schönfeld, F. Hoffmann-La Roche, Basel, Switzerland. ClpB was purified according to Woo *et al.*<sup>78</sup>. Firefly luciferase (Luc) was purified and assayed in as described in Sharma *et al.*<sup>42</sup>. Mitochondrial malate dehydrogenase (MDH) was from pig heart mitochondria (Boehringer Mannheim) and assayed according to Diamant *et al.*<sup>79</sup>. Citrate synthase (CS) was from Sigma and assayed according to Buchner *et al.*<sup>36</sup>.

### **Chaperone assays:**

The chaperone-refolding buffer was 100 mM Tris 7.5, 150 mM KCl, 20 mM MgCl<sub>2</sub>, 10 mM Dithiothreitol. Unless mentioned otherwise, all chaperone assays used 4 mM ATP 5 mM 2-phosphoenolpyruvate and pyruvate kinase (Sigma) to regenerate the ATP.

Concentrations of chaperones, cochaperones and reporter protein: Unless mentioned otherwise, all protein concentrations were expressed as protomers. Hence, an assay containing 1  $\mu$ M GroEL and 1  $\mu$ M GroES, contained 71.4 nM of tetradecameric GroEL<sub>14</sub> complexes, corresponding to 142.8 nM of active heptameric catalytic sites (GroEL<sub>7</sub> cavities), which could be transiently covered by as many (142.8 nM) functional GroES<sub>7</sub> heptamers during the ATP-driven catalytic cycle.

Chaperone stoichiometries in assays: GroEL and GroES protomers were always equimolar, as in bacteria and mitochondria<sup>15</sup>. For DnaK, DnaJ, GrpE and ClpB molar ratios were 5:1:1:2

Native reporter enzymes (MDH, CS, Luc) from stocks were diluted to the indicated final concentrations (0.25-2.5  $\mu\text{M}$ ) and incubated in thin-walled PCR tubes in a thermocycler at constant indicated temperatures, in the presence of the chaperones and ATP, typically for about two hours, but in the experiment in Fig.4, for up to 8 hours. At indicated times, aliquots were assayed in a spectrophotometer or a luminometer. Linear enzymatic rates were converted into molar concentrations (nM) of natively refolded chaperone products as explained below.

For experiments following the native refolding of pre-unfolded reporter enzymes, a 40-fold excess of MDH reporter enzyme was preincubated in 7 M urea and 20 mM DTT for 5 min at 25°C. Typically, 2  $\mu\text{l}$  were then flush-diluted and mixed into 78  $\mu\text{l}$  of pre-warmed refolding buffer containing chaperones and ATP, as indicated, and further incubated at the indicated temperatures. In contrast to Guanidium HCl that affects GroEL-GroES activity, the 175 mM traces of Urea were inconsequential for the chaperonin assay<sup>80</sup>.

#### Calculation of the scale for graphs where enzymatic activity is reported:

For a given total concentration [E] of the enzyme E in solution, (E being MDH, citrate synthase or luciferase), the maximal activity measured throughout the whole set of experiments represented in a figure is considered to represent [E], and all other activities are thus proportionally scaled when expressed as a concentration.

#### **Acknowledgments**

We thank summer project students Patrick Inns and Ivanoé Koog for setting up the Citrate Synthase enzymatic assay and performing key preliminary and control chaperone assays. This project was financed in part by Swiss National Science

Foundation Grants 140512/1, 31003A\_156948 and by Grant C15.0042 from the Swiss State Secretariat for Education Research and Innovation to P.G. and B.F. Grant 200020\_163042 from the Swiss National Science Foundation to P.D.L.R and A.S.S. A.B. acknowledges the support of the French Agence Nationale de la Recherche (ANR), under grant ANR-14-ACHN-0016.



## Figure captions

### **Figure 1. Free energy landscape of MDH at 37°C and schematic action of chaperones.**

The free-energy landscape of MDH at 37°C is represented by the native (active) dimer (ND), the native (inactive) monomer (NM), a less structured intermediate (I) which is higher in free energy and a misfolded state (M), which is the lowest in free-energy. Non-native conformations can further lead to the irreversible formation of aggregates (A), although likely at much slower rates. The spontaneous conversion reactions between the different states are represented with black arrows (dashed arrows for aggregation to highlight the different timescale of this process). Chaperones act on the non-native ensemble by associating to and dissociating from misfolded (CM) and intermediate (CI) conformations. While bound to chaperones, misfolded and intermediate conformations can interconvert into each other. Reactions driven by chaperones are represented by red arrows. The energy driving the chaperone action is graphically indicated with the yellow-to-red closed-circle arrow.

### **Figure 2. GroELS recovers and then maintain MDH activity in denaturing conditions, in a stringently ATP-dependent, non-equilibrium, reaction.**

A) Time course of the activity of MDH (0.5  $\mu\text{M}$ ) at 37°C in the absence (black circles) and presence (red circles) of GroEL (3.5  $\mu\text{M}$ ) and GroES (3.5  $\mu\text{M}$ ) and ATP regeneration system (see Methods). In the absence of GroELS MDH spontaneously lost activity at an apparent rate of about 0.007  $\text{min}^{-1}$ . GroELS was supplemented at different  $t=0'$ , 15', 30', 45' and 70' (see Methods for the choice of the scale on the y-

axis; the same procedure applies to all graphs where activity is expressed as a concentration). B) Final MDH fractions (at 130') for GroELS addition at different times from the experiments in Fig.1A. White bars represent the concentration of active MDH at the time of GroELS addition, whereas the recovered fractions (blue bars) are the differences between the active protein concentrations at 130' and at the time of GroELS addition. The concentration of proteins that cannot be recovered by GroELS is represented with red bars. C) GroELS (3.5  $\mu\text{M}$ ) and ATP regeneration system were added to 0.25  $\mu\text{M}$  MDH at 37°C (denaturing conditions) at t=0' (red filled circles) or at t=60' (red empty circles). Apyrase was added at t=105'. D) Refolding of urea-pre-denatured 0.5  $\mu\text{M}$  MDH at 37°C in the initial presence of different GroELS concentrations (as indicated) and ATP regeneration system. Apyrase was added at t=60'. E) Refolding of urea-pre-denatured 0.5  $\mu\text{M}$  MDH at 37°C in the initial presence of GroELS (3.5  $\mu\text{M}$ ), 400  $\mu\text{M}$  ATP and various concentrations of ADP (as indicated). As predicted by the correct accounting of the energy available for the chaperone action (see main text), increasing concentrations of ADP progressively inhibit the reaction.

### **Figure 3. Results from the model.**

The panels here are in one-to-one correspondence to the panels in Fig.2, and have been obtained for a single set of intrinsic parameters (the rates of the reactions). Concentrations of chaperones and substrates are the same as in the corresponding experiments. A) and B) Model results for the same setups as in Figs.2A and 2B respectively. C) Results from the model for the same conditions as in Fig.1C. D) Results from the model for the same conditions as in Fig.2D. E) Results from the

model for the same conditions as in Fig.2E (see SI for details on the modelling of ATP consumption).

**Figure 4. ATP-driven catalytic action of substoichiometric amounts of GroELS on MDH at 37°C.**

A) Activity of 2.5  $\mu\text{M}$  MDH after incubation for 8 hours in the presence of increasing amounts of GroEL (and equimolar GroES) at 25°C (black circles) and at 37°C in the presence (full red squares) and absence (empty red squares) of ATP and regeneration system. The ratio of the concentration of GroEL<sub>7</sub> rings (considered as the minimal active unit of GroEL) to the concentration of MDH protomers is given. Inset) The minimal number of non-native MDH protomers that should be rescued by each GroEL<sub>7</sub> ring to explain the observed activity after 8h. B) results of the model in the same conditions as in panel A. Inset) The minimal number of non-native MDH protomers that should be rescued by each GroEL<sub>7</sub> ring in order to explain the observed activity after 8h (red squares) is similar to the experimental one (~15 protomers per GroEL<sub>7</sub>). The model gives also access to the effective number of MDH protomers that had to be processed by each ring (blue squares), which is more than one order of magnitude larger (~300 per GroEL<sub>7</sub> at EC50), highlighting the iterative, catalytic action of GroELS on its substrates.

**Figure 5. GroELS turn the energy of ATP into a non-equilibrium stabilisation of the native state of MDH.**

A) The *effective* free-energy difference between the native and non-native two ensembles  $\Delta G_{N,nN}^{\text{eff}} = -k_B T \ln\{[N]/[nN]\}$  (as defined in the main text) as a function of

$\Delta G_{\text{ATP}}$ , from equilibrium conditions ( $\Delta G_{\text{ATP}}=0$ ) to conditions exceeding the ones that, in the model, reproduce the experiments.  $\Delta G_{\text{N,nN}}^{\text{eff}}$  is positive at equilibrium at 37°C, attesting for the thermodynamic instability of the native ensemble. The native ensemble of MDH is stabilized as progressively more energy is injected into the system. B) The action of GroELS on the non-native ensemble in non-equilibrium results can be visualized as a modification of the free-energy landscape of non-native conformations, whereby misfolded proteins have a much higher free-energy than native conformations. C) The relevance of the action of GroELS as a function of the equilibrium stability of their substrates ( $\Delta G_{\text{N,nN}}^{\text{eq}}$ ). When  $\Delta G_{\text{N,nN}}^{\text{eq}} < 0$ , substrates benefit very little from the help of GroELS because they are already stable. As the equilibrium stability of the native state is challenged (increasing  $\Delta G_{\text{N,nN}}^{\text{eq}}$ ), energy injection in the system helps maintaining the native state populated against its thermodynamic tendency (the orange region represents the net gain of native proteins thanks to chaperones). Extreme instability resulting from strong stresses or highly deleterious mutations cannot be compensated by chaperones (blue region), which can anyway avoid protein aggregation by binding the non-native conformations. Non-native but unbound, and thus aggregation-prone conformations thus represent a minor part of the solution (green region).







Figure 1



**Figure 2**




**Figure 3**



**Figure 4**





**Figure 5**

## BIBLIOGRAPHY

1. Anfinsen, C. B., Haber, E., Sela, M. & White, F. H. The kinetics of formation of native ribonuclease during oxidation of the reduced polypeptide chain. *Proc. Natl. Acad. Sci. USA* **47**, 1309–1314 (1961).
2. Onuchic, J. N., Luthey-Schulten, Z. & Wolynes, P. G. Theory of protein folding: the energy landscape perspective. *Annual review of physical chemistry* **48**, 545–600 (1997).
3. Onuchic, J. N. & Wolynes, P. G. Theory of protein folding. *Curr. Opin. Struct. Biol.* **14**, 70–75 (2004).
4. Dobson, C. M. Protein folding and misfolding. *Nature* **426**, 884–890 (2003).
5. Hartl, F. U., Bracher, A. & Hayer-Hartl, M. Molecular chaperones in protein folding and proteostasis. *Nature* **475**, 324–332 (2011).
6. Knowles, T. P. J., Vendruscolo, M. & Dobson, C. M. The amyloid state and its association with protein misfolding diseases. *Nat Rev Mol Cell Bio* **15**, 384–396 (2014).
7. Gershenson, A., Gierasch, L. M., Pastore, A. & Radford, S. E. Energy landscapes of functional proteins are inherently risky. *Nat Chem Biol* **10**, 884–891 (2014).
8. Natalello, A. *et al.* Biophysical characterization of two different stable misfolded monomeric polypeptides that are chaperone-amenable substrates. *J. Mol. Biol.* **425**, 1158–1171 (2013).
9. Hipp, M. S., Park, S.-H. & Hartl, F. U. Proteostasis impairment in protein-misfolding and -aggregation diseases. *Trends Cell Biol* **24**, 506–514 (2014).
10. Goloubinoff, P. Mechanisms of protein homeostasis in health, aging and disease. *Swiss Med Wkly* **146**, (2016).
11. Goloubinoff, P., Mogk, A., Zvi, A. P., Tomoyasu, T. & Bukau, B. Sequential mechanism of solubilization and refolding of stable protein aggregates by a bichaperone network. *Proc. Natl. Acad. Sci. USA* **96**, 13732–13737 (1999).
12. Finka, A., Mattoo, R. U. H. & Goloubinoff, P. Experimental Milestones in the Discovery of Molecular Chaperones as Polypeptide Unfolding Enzymes. *Annu Rev Biochem* (2016). doi:10.1146/annurev-biochem-060815-014124
13. Gat-Yablonski, G. *et al.* Quantitative proteomics of rat livers shows that unrestricted feeding is stressful for proteostasis with implications on life span. *Aging (Albany NY)* **8**, 1735–1758 (2016).
14. Finka, A., Sood, V., Quadroni, M., Rios, P. D. L. & Goloubinoff, P. Quantitative proteomics of heat-treated human cells show an across-the-board mild depletion of housekeeping proteins to massively accumulate few HSPs. *Cell Stress Chaperon* **20**, 605–620 (2015).
15. Finka, A. & Goloubinoff, P. Proteomic data from human cell cultures refine mechanisms of chaperone-mediated protein homeostasis. *Cell Stress Chaperon* (2013). doi:10.1007/s12192-013-0413-3

16. Goloubinoff, P., Christeller, J. T., Gatenby, A. A. & Lorimer, G. H. Reconstitution of active dimeric ribulose biphosphate carboxylase from an unfolded state depends on two chaperonin proteins and Mg-ATP. *Nature* **342**, 884–889 (1989).
17. Wiech, H., Buchner, J., Zimmermann, R. & Jakob, U. Hsp90 Chaperones Protein Folding In vitro. *Nature* **358**, 169–170 (1992).
18. Jakob, U., Gaestel, M., Engel, K. & Buchner, J. Small Heat-Shock Proteins Are Molecular Chaperones. *Journal of Biological Chemistry* **268**, 1517–1520 (1993).
19. Clare, D. K. & Saibil, H. R. ATP-driven molecular chaperone machines. *Biopolymers* **99**, 846–859 (2013).
20. Seifert, U. Stochastic thermodynamics, fluctuation theorems and molecular machines. *Rep Prog Phys* **75**, 126001 (2012).
21. Frank, G. A. *et al.* Out-of-equilibrium conformational cycling of GroEL under saturating ATP concentrations. *Proceedings of the National Academy of Sciences* **107**, 6270–6274 (2010).
22. De Los Rios, P. & Barducci, A. Hsp70 chaperones are non-equilibrium machines that achieve ultra-affinity by energy consumption. *Elife* **3**, e02218 (2014).
23. Barducci, A. & De Los Rios, P. Non-equilibrium conformational dynamics in the function of molecular chaperones. *Curr. Opin. Struct. Biol.* **30**, 161–169 (2015).
24. Goloubinoff, P., Gatenby, A. A. & Lorimer, G. H. GroE heat-shock proteins promote assembly of foreign prokaryotic ribulose biphosphate carboxylase oligomers in *Escherichia coli*. *Nature* **337**, 44–47 (1989).
25. Sigler, P. B. *et al.* Structure and function in GroEL-mediated protein folding. *Annu Rev Biochem* **67**, 581–608 (1998).
26. Hayer-Hartl, M., Bracher, A. & Hartl, F. U. The GroEL-GroES Chaperonin Machine: A Nano-Cage for Protein Folding. *Trends Biochem Sci* **41**, 62–76 (2016).
27. Jurgensen, S. R., Wood, D. C., Mahler, J. C. & Harrison, J. H. The immobilization of mitochondrial malate dehydrogenase on Sepharose beads and the demonstration of catalytically active subunits. *Journal of Biological Chemistry* **256**, 2383–2388 (1981).
28. Hartman, D. J., Surin, B. P., Dixon, N. E., Hoogenraad, N. J. & Høj, P. B. Substoichiometric amounts of the molecular chaperones GroEL and GroES prevent thermal denaturation and aggregation of mammalian mitochondrial malate dehydrogenase in vitro. *Proc. Natl. Acad. Sci. USA* **90**, 2276–2280 (1993).
29. Agashe, V. R., Shastry, M. C. & Udgaonkar, J. B. Initial hydrophobic collapse in the folding of barstar. *Nature* **377**, 754–757 (1995).
30. Dobson, C. M. & Karplus, M. The fundamentals of protein folding: bringing together theory and experiment. *Curr. Opin. Struct. Biol.* **9**, 92–101 (1999).

31. Walters, B. T., Mayne, L., Hinshaw, J. R., Sosnick, T. R. & Englander, S. W. Folding of a large protein at high structural resolution. *Proceedings of the National Academy of Sciences* **110**, 18898–18903 (2013).
32. Peralta, D., Hartman, D. J., Hoogenraad, N. J. & Høj, P. B. Generation of a stable folding intermediate which can be rescued by the chaperonins GroEL and GroES. *FEBS Lett.* **339**, 45–49 (1994).
33. Horwich, A. L., LOW, K. B., Fenton, W. A., HIRSHFIELD, I. N. & Furtak, K. Folding in-Vivo of Bacterial Cytoplasmic Proteins - Role of Groel. *Cell* **74**, 909–917 (1993).
34. Priya, S. *et al.* GroEL and CCT are catalytic unfoldases mediating out-of-cage polypeptide refolding without ATP. *Proc. Natl. Acad. Sci. USA* **110**, 7199–7204 (2013).
35. Goloubinoff, P., Christeller, J. T., Gatenby, A. A. & Lorimer, G. H. Reconstitution of active dimeric ribulose biphosphate carboxylase from an unfolded state depends on two chaperonin proteins and Mg-ATP. *Nature* **342**, 884–889 (1989).
36. Buchner, J. *et al.* GroE facilitates refolding of citrate synthase by suppressing aggregation. *Biochemistry* **30**, 1586–1591 (1991).
37. Moran, U., Phillips, R. & Milo, R. Snapshot: Key Numbers in Biology. *Cell* **141**, 1262–1262 (2010).
38. Horst, R. *et al.* Direct NMR observation of a substrate protein bound to the chaperonin GroEL. *Proc. Natl. Acad. Sci. USA* **102**, 12748–12753 (2005).
39. Lin, Z., Madan, D. & Rye, H. S. GroEL stimulates protein folding through forced unfolding. *Nat Struct Mol Biol* **15**, 303–311 (2008).
40. Sharma, S. *et al.* Monitoring protein conformation along the pathway of chaperonin-assisted folding. *Cell* **133**, 142–153 (2008).
41. Libich, D. S., Tugarinov, V. & Clore, G. M. Intrinsic unfoldase/foldase activity of the chaperonin GroEL directly demonstrated using multinuclear relaxation-based NMR. *Proceedings of the National Academy of Sciences* **112**, 8817–8823 (2015).
42. Sharma, S. K., De Los Rios, P., Christen, P., Lustig, A. & Goloubinoff, P. The kinetic parameters and energy cost of the Hsp70 chaperone as a polypeptide unfoldase. *Nat Chem Biol* **6**, 914–920 (2010).
43. Kellner, R. *et al.* Single-molecule spectroscopy reveals chaperone-mediated expansion of substrate protein. *Proceedings of the National Academy of Sciences* **111**, 13355–13360 (2014).
44. Lee, J. H. *et al.* Heterogeneous binding of the SH3 client protein to the DnaK molecular chaperone. *Proceedings of the National Academy of Sciences* **112**, E4206–15 (2015).
45. Sekhar, A., Rosenzweig, R., Bouvignies, G. & Kay, L. E. Mapping the conformation of a client protein through the Hsp70 functional cycle. *Proceedings of the National Academy of Sciences* **112**, 10395–10400 (2015).
46. Sekhar, A., Rosenzweig, R., Bouvignies, G. & Kay, L. E. Hsp70 biases the

- folding pathways of client proteins. *Proceedings of the National Academy of Sciences* **113**, E2794–801 (2016).
47. Todd, M. J., Lorimer, G. H. & Thirumalai, D. Chaperonin-facilitated protein folding: optimization of rate and yield by an iterative annealing mechanism. *Proc. Natl. Acad. Sci. USA* **93**, 4030–4035 (1996).
  48. Tehver, R. & Thirumalai, D. Kinetic model for the coupling between allosteric transitions in GroEL and substrate protein folding and aggregation. *J. Mol. Biol.* **377**, 1279–1295 (2008).
  49. Santra, M., Farrell, D. W. & Dill, K. A. Bacterial proteostasis balances energy and chaperone utilization efficiently. *Proceedings of the National Academy of Sciences* **114**, E2654–E2661 (2017).
  50. Mittal, J. & Best, R. B. Dependence of protein folding stability and dynamics on the density and composition of macromolecular crowders. *Biophys J* **98**, 315–320 (2010).
  51. Hayer-Hartl, M. & Minton, A. P. A simple semiempirical model for the effect of molecular confinement upon the rate of protein folding. *Biochemistry* **45**, 13356–13360 (2006).
  52. Priya, S., Sharma, S. K. & Goloubinoff, P. Molecular chaperones as enzymes that catalytically unfold misfolded polypeptides. *FEBS Lett.* **587**, 1981–1987 (2013).
  53. Yifrach, O. & Horovitz, A. Nested Cooperativity in the Atpase Activity of the Oligomeric Chaperonin Groel. *Biochemistry* **34**, 5303–5308 (1995).
  54. Corrales, F. J. & Fersht, A. Toward a mechanism for GroEL center dot GroES chaperone activity: An ATPase-gated and -pulsed folding and annealing cage. *Proc. Natl. Acad. Sci. USA* **93**, 4509–4512 (1996).
  55. Kipnis, Y., Papo, N., Haran, G. & Horovitz, A. Concerted ATP-induced allosteric transitions in GroEL facilitate release of protein substrate domains in an all-or-none manner. *Proc. Natl. Acad. Sci. USA* **104**, 3119–3124 (2007).
  56. Clare, D. K. *et al.* ATP-Triggered Conformational Changes Delineate Substrate-Binding and -Folding Mechanics of the GroEL Chaperonin. *Cell* 1–11 (2012). doi:10.1016/j.cell.2012.02.047
  57. Horwich, A. L., Apetri, A. C. & Fenton, W. A. The GroEL/GroES cis cavity as a passive anti-aggregation device. *FEBS Lett.* **583**, 2654–2662 (2009).
  58. Doyle, S. M. & Wickner, S. Hsp104 and ClpB: protein disaggregating machines. *Trends Biochem Sci* **34**, 40–48 (2009).
  59. Haslberger, T. *et al.* M domains couple the ClpB threading motor with the DnaK chaperone activity. *Mol. Cell* **25**, 247–260 (2007).
  60. Langer, T. *et al.* Successive Action of Dnak, Dnaj and Groel Along the Pathway of Chaperone-Mediated Protein Folding. *Nature* **356**, 683–689 (1992).
  61. Buchberger, A., Schroder, H., Hesterkamp, T., Schönfeld, H. J. & Bukau, B. Substrate shuttling between the DnaK and GroEL systems indicates a chaperone network promoting protein folding. *J. Mol. Biol.* **261**, 328–333 (1996).

62. Sharma, S. K., De Los Rios, P. & Goloubinoff, P. Probing the different chaperone activities of the bacterial HSP70-HSP40 system using a thermolabile luciferase substrate. *Proteins* **79**, 1991–1998 (2011).
63. Hu, B., Mayer, M. P. & Tomita, M. Modeling Hsp70-mediated protein folding. *Biophys J* **91**, 496–507 (2006).
64. Powers, E. T., Powers, D. L. & Gierasch, L. M. FoldEco: a model for proteostasis in *E. coli*. *Cell Rep* **1**, 265–276 (2012).
65. De Los Rios, P. & Goloubinoff, P. Protein folding: Chaperoning protein evolution. *Nat Chem Biol* **8**, 226–228 (2012).
66. Lindquist, S. Protein folding sculpting evolutionary change. *Cold Spring Harb. Symp. Quant. Biol.* **74**, 103–108 (2009).
67. Williams, T. A. & Fares, M. A. The Effect of Chaperonin Buffering on Protein Evolution. *Genome Biol Evol* **2**, 609–619 (2010).
68. Tokuriki, N. & Tawfik, D. S. Chaperonin overexpression promotes genetic variation and enzyme evolution. *Nature* **459**, 668–673 (2009).
69. Tokuriki, N. & Tawfik, D. S. Stability effects of mutations and protein evolvability. *Curr. Opin. Struct. Biol.* **19**, 596–604 (2009).
70. Bhaskaran, H. & Russell, R. Kinetic redistribution of native and misfolded RNAs by a DEAD-box chaperone. *Nature* **449**, 1014–U2 (2007).
71. Yang, Q., Fairman, M. E. & Jankowsky, E. DEAD-box-protein-assisted RNA structure conversion towards and against thermodynamic equilibrium values. *J. Mol. Biol.* **368**, 1087–1100 (2007).
72. Rouskin, S., Zubradt, M., Washietl, S., Kellis, M. & Weissman, J. S. Genome-wide probing of RNA structure reveals active unfolding of mRNA structures in vivo. *Nature* **505**, 701–705 (2014).
73. Bershtein, S., Mu, W., Serohijos, A. W. R., Zhou, J. & Shakhnovich, E. I. Protein quality control acts on folding intermediates to shape the effects of mutations on organismal fitness. *Mol. Cell* **49**, 133–144 (2013).
74. Rodrigues, J. V. *et al.* Biophysical principles predict fitness landscapes of drug resistance. *Proceedings of the National Academy of Sciences* (2016). doi:10.1073/pnas.1601441113
75. Schrödinger, E. 1.-1. What is life? The physical aspect of the living cell. (1945).
76. Torok, Z. *et al.* Evidence for a lipochaperonin: association of active protein-folding GroESL oligomers with lipids can stabilize membranes under heat shock conditions. *Proc. Natl. Acad. Sci. USA* **94**, 2192–2197 (1997).
77. Gur, E. *et al.* The *Escherichia coli* DjlA and CbpA proteins can substitute for DnaJ in DnaK-mediated protein disaggregation. *J Bacteriol* **186**, 7236–7242 (2004).
78. Woo, K. M., Kim, K. I., Goldberg, A. L., Ha, D. B. & Chung, C. H. The heat-shock protein ClpB in *Escherichia coli* is a protein-activated ATPase. *Journal of Biological Chemistry* **267**, 20429–20434 (1992).
79. Diamant, S., Azem, A., Weiss, C. & Goloubinoff, P. Increased efficiency of

- GroE-assisted protein folding by manganese ions. *J Biol Chem* **270**, 28387–28391 (1995).
80. Todd, M. J. & Lorimer, G. H. Stability of the asymmetric Escherichia coli chaperonin complex. Guanidine chloride causes rapid dissociation. *Journal of Biological Chemistry* **270**, 5388–5394 (1995).