



# PROCEEDINGS

VOLUME II

The 2<sup>nd</sup> International Conference  
on Animal Nutrition and Environment  
(ANI-NUE2017)

*“Towards the Betterment of Animal  
Productivity, Conserving Resources  
and Environment”*

November 1-4,  
**2017**

Pullman Raja Orchid Hotel,  
Khon Kaen, Thailand



**EDITORS:** Cherdthong, A., Foiklang, S., Mapato, C., Pilajun, R., Kang, SC. and Wanapat, M.

**ISBN:** 978-616-438-084-4

Organized by

Tropical Feed Resources Research and Development Center (TROFREC), Department of Animal Science,  
Faculty of Agriculture, Khon Kaen University





**In Recognition for**

**Professor Dr. Metha Wanapat**

**“The Khon Kaen University Distinguished  
Research Professor”**

**for**

**His 37 years of continuous serving in  
teaching, research, education, technology  
development, and transfer in Animal  
Science nationally and internationally.**

**The 2<sup>nd</sup> International Conference on Animal Nutrition and  
Environment (ANI-NUE2017) Organizing Committee,**

**November 1, 2017**

**PROCEEDINGS**  
*of*  
**The 2<sup>nd</sup> International Conference on**  
**Animal Nutrition and Environment**  
**(ANI-NUE2017)**  
**Volume II**

**Editors:** Cherdthong, A., Foiklang, S., Mapato, C., Pilajun, R., Kang, SC. and Wanapat, M.

**ISBN:** 978-616-438-084-4

**November 1-4, 2017**

**Pullman Raja Orchid Hotel, Khon Kaen, Thailand**

## Contents (Continued)

<b>ANN-01-0071: “Effects of supplementation of Piper sarmentosum leaves powder in concentrates on feed efficiency, rumen fermentation and protozoa in Thai native beef cattle”</b> <i>Anusorn Cherdthong and Benjamad Khonkhaeng-Thailand</i>	400
<b>Session 9</b>	
<b>ANN-01-0061: “Determination of level protein intake to control fat and protein in carcass of fattened lambs”</b> <i>Rizky Choirunnisa, Mukh Arifin, Robert Kusetyawan, Febrian Rhamadya, Ari Prima, Vita Restitrisnani, Nadlirotun Luthfi, Sutaryo and Agung Purnomoadi-Indonesia</i>	408
<b>ANN-01-0044: “Effect of yeast-fermented cassava pulp levels on growth performance of growing goats fed Napier Pakchong 1 grass”</b> <i>Saran Parisuthikul, Tichakorn Thumtong, Theerasant Phothee, Panuwat Teebklang, Wunchai Inthiseang, Wichan Kaewluan and Ruangyote Pilajun-Thailand</i>	415
<b>ANN-01-0102: “Effect of mangosteen peel liquid protected soybean meal on methonogen and microbial population using in in vitro gas fermentation technique”</b> <i>Kampanat Phesatcha, Burarat Phesatcha, Suban Foiklang, Metha Wanapat-Thailand</i>	422
<b>ANN-01-0082: “Fermentation quality and in vitro digestibility of fermented total mix ration with difference roughage and fermentation period”</b> <i>Pichad Khejornart, Teerayoot Jantanam and Metha Wanapat-Thailand</i>	429
<b>ANN-01-0076: “Effect of replacement soybean meal by yeast waste on feed intake and rumen ecology in Thai native beef cattle”</b> <i>Anusorn Cherdthong, Phussorn Sumadong, Suban Foiklang, and Nipa Milintawisamai-Thailand</i>	436
<b>ANN-01-0077: “Chemical composition and In vitro gas production of the local Thai and India Moringa (<i>Moringa oleifera</i> Lam.) for ruminant”</b> <i>Chaiyawan Wattanachant, Nattha Rattanacosol, Chaermpon Yuangklang, and Apichat Loopachr-Thailand</i>	441
<b>Session 10</b>	
<b>ANN-01-0013: “Use of Body Measurements to Predict Intermuscular Fat in Thin-tailed Lambs”</b> <i>Ulia Renfelia Baysi, Ari Prima, Farah Nabila, Pradhipta Hersandika, Endang Purbowati, Christina Maria Sri Lestari, and Agung Purnomoadi-Indonesia</i>	447
<b>ANN-01-0099: “Effect of inclusion of enzyme from fermented tomato pomace with <i>Aspergillus niger</i> (FETPAN) in total mixed ration on feed intake and growth performance in beef cattle”</b> <i>Chalermpon Yuangklang, Jiravan Khotsakdee, and Krisit Vasupen-Thailand</i>	452
<b>ANN-01-0100: “Influence of Tropical Roughages Combined with Urea and Bamboo Grass (<i>Tiliacora Triandra</i>, Diels) Supplementation on Gas Production and In Vitro Degradability”</b> <i>Chaichana Suriyapha, Metha Wanapat and Chinda Wann-Thailand</i>	457



## Contents (Continued)

ANN-01-0087: “Effect of Bamboo grass ( <i>Tiliacora Triandra</i> ) pellet supplementation on feed intake, nutrient digestibility and rumen microbial population in Thai native beef” <i>Chinda Wann, Metha Wanapat, Choawarit Mapato and Chaichana Suriyapha-Thailand</i>	463
ANN-01-0105: “ <i>In Vitro</i> Fermentation and Methane Production Influenced by Leucaena Silage and Mangosteen Peel Powder” <i>Sungchhang Kang, Metha Wanapat, Ratana Khun, Laiheang Chhen, Piseth Try, Kimsan Sok, Ratha Long and Vannak Un-Cambodia</i>	469
ANN-01-0106: “Influence of Bamboo Leaf Meal Supplementation on <i>In Vitro</i> Gas Production and Digestibility” <i>Sungchhang Kang and Metha Wanapat-Cambodia</i>	475
<b>Session 11</b>	
ANN-01-0075: “Effect of Banana flower powder contained in high quality feed block and roughage to concentrate ratio on in vitro gas production kinetics, digestibility and fermentation” <i>Suban Foiklang, Metha Wanapat, Bounnaxay Viennasay, and Thitima Norrapoke-Thailand</i>	481
ANN-01-0047: “Feed intake and blood metabolite of goats fed urea-calcium hydroxide treated oil palm frond” <i>Pin Chanjula, Suradech phetarwut, and Anusorn Cherdthong-Thailand</i>	486
ANN-01-0070: “Blood chemistries of dairy cow during pre-calving, at calving and post-calving period” <i>Chamnanwit Promkot, Pitukpol Porn-aneke, and Teerawat Srinukool-Thailand</i>	491
ANN-01-0060: “The study on determination of feed digestibility using frequency of defecation on Thin Tailed lamb” <i>Febrianto Dwi Nugroho, Ari Prima, Edy Rianto, Agung Purnomoadi-Indonesia</i>	494
ANN-01-0062: “Effect of <i>Terminalia Chebula</i> RETZ. meal on nutrient intake, digestibility and microbial population of goats” <i>Nirawan Gunun, Pongsatorn Gunun, Anusorn Cherdthong, Sineenart Polyorach and Metha Wanapat-Thailand</i>	497
ANN-01-0090: “Effect of bamboo grass pellet (Bamboo-cass) levels on gas production kinetics and in vitro degradability” <i>Bounnaxay Viennasay and Metha Wanapat-Thailand</i>	500
ANN-01-0084: “Effect of fresh cassava root with feed block containing high sulfur on gas kinetics and rumen fermentation using <i>in vitro</i> gas production technique” <i>Gamonmas Dageaw and Anusorn Cherdthong-Thailand</i>	507
<b>Session 12</b>	
ANN-01-0072: “Inclusion of yeast waste as protein source to replace soybean meal in concentrate diet on ruminal fermentation and kinetics of gas using a gas production technique” <i>Anusorn Cherdthong, Rittikeard Prachumchai, Chanadol Supapong, Metha Wanapat, Suban Foiklang, and Nipa Milintawisamai-Thailand</i>	514

## Contents (Continued)

<b>ANN-01-0055: “Using of Urea and Molasses Fermented Cassava Pulp on Rumen Fermentation and Methane Production”</b> <i>Thitima Norrapoke and Tanitpan Phongchongmit-Thailand</i>	519
<b>ANN-01-0046: “The Meat Quality of Bali Beef fed with Supplement Blocks with Different Liquid Smoke Levels as Antioxidant and Binder”</b> <i>Effendi Abustam, Muhammad Irfan Said, and Muhammad Yusuf-Indonesia</i>	526
<b>ANN-01-0094: “Effect of Flemingia (<i>Flemingia macrophylla</i>) as a protein replacement of soybean meal on feed intake, digestibility of nutrients and microbial population in Thai native beef cattle”</b> <i>Burarat Phesatcha and Metha Wanapat-Thailand</i>	532
<b>ANN-01-0068: “Nutrient utilization and rumen ecology of Thai indigenous cattle given hay and sago palm pith with different levels of soybean meal”</b> <i>Bunseelarp Wiyada and Ngampongsai Wanwisa-Thailand</i>	540
<b>ANN-01-0085: “Comparison between Hay and Silage of <i>Pennisetum purpurium</i> cv. Mahasarakham feeding on feed intake, nutrient digestibility, and rumen fermentation in Thai native beef bulls”</b> <i>Chaowarit Mapato and Metha Wanapat-Thailand</i>	545
<b>ANN-01-0078: “Increasing Productive Performance of Native Chickens by Herbs in Rural Community”</b> <i>Narumon Somkuna, Eakkasit Somkuna, Jarous Sawangtap and Phinithi Ratchwicha-Thailand</i>	549
<b>Session 13</b>	
<b>ANN-01-0059: “Regression Models for Estimating Fat Carcass Percentage Using Chest Measurement in Thin Tailed Lambs”</b> <i>Farah Nabila, Pradhipta Hersandika, Ari Prima, Vita Restitrisnani, Nadlirotun Luthfi, Endang Purbowati, Sutaryo and Agung Purnomoadi-Indonesia</i>	553
<b>ANN-01-0065: “Development of Near Infrared Spectroscopy for nondestructive and rapid measurement of chemical compositions and somatic cell counts in raw milk”</b> <i>Onanong Pongchompu, Metha Wanapat, Chaluntorn Vichasilp, Yaungyote Jindatajak and Pongsagorn Pongchompu-Thailand</i>	558
<b>ANN-01-0014: “Effect of Hydroponic Maize Fodder Supplementation on Production Performance in Graded Murrah Buffaloes of Scarce Rainfall Zone”</b> <i>Atturi Krishna Murthy, Dhanalakshmi Guduru, Y.G.Prasad, and Sarjan Reddy Kapa-India</i>	563
<b>ANN-01-0081: “Effect of fermented total mixed ration with microbial culture on fermentation quality and <i>in vitro</i> digestibility”</b> <i>Pichad Khejornsart and Metha Wanapat-Thailand</i>	567
<b>ANN-01-0104- Effect of Addition of Siamese Neem Foliage on pH and Number of Lactic Acid Bacteria in Napier Grass Silage</b> <i>Anan Petlum, Sukanya Kamphayae, Pramote Paengkoum, Walailuck Kaewwongsa, Smerjai Bureenok, Tanaporn Plong-uan, and Thanyalak Theppaw-Thailand</i>	574



## **Regression Models for Estimating Fat Carcass Percentage Using Chest Measurement in Thin Tailed Lambs**

**Farah Nabila, Pradhipta Hersandika, Ari Prima, Vita Restitrisnani, Nadlirotun Luthfi, Endang Purbowati, Sutaryo, and Agung Purnomoadi\***

*Department of Animal Science, Faculty of Animal and Agricultural Sciences,  
Diponegoro University, Semarang, Indonesia 50275  
\*Corresponding email: agung194@yahoo.com*

### **Abstract**

This study was conducted to evaluate the possibility of using chest measurement for subcutaneous, intermuscular, and total fat percentage in Thin Tailed Lambs. Twenty one heads of three months old male thin tailed lambs with initial body weight (BW)  $14.57 \pm 2.19$  kg were raised up to 3 months fattening period. Chest girth (CG), chest depth (CD), and chest width (CW) were measured before slaughtered. The fat of the carcass was separated into subcutaneous, intermuscular, and total fat, then weighed. The data was analyzed by linear regression to determine the correlation and equation between chest measurement and fat carcass percentage, then was evaluated by t-test, standard error (SE) and the differences were measured to predict the accuracy of the equation. The results showed that the correlation between CG, CW and CG/CD in intermuscular and total fat percentage was positive and significant ( $P < 0.05$ ) with correlation value moderate to high ( $r = 0.510 - 0.664$ ), while subcutaneous were not significant. SE of each variable showed a low value (0.002-0.020) that indicate the prediction is close to the actual fat percentage value. The lowest differences between prediction and actual value could be found in CG regression equation in total fat (0.368%), while the highest value was 8.918% in intermuscular fat in CW/CG regression. Based on the results of this study, it can be concluded that CG regression is the best equation for estimating intermuscular, and total fat percentage using chest measurement in Thin Tailed Lambs.

**Keywords:** chest measurement, fat percentage, regression, lambs

### **Introduction**

Fat is one of factor that determine carcass grading qualification in animals. Fat carcass consist of subcutaneous, intermuscular and intramuscular. The minimum requirement for either extractable fat in order to achieve acceptable consumer satisfaction for grilling red meat at 5% for sheepmeat (Hopkins et al., 2006). There are some methods to predict the fat before slaughtering the lambs. The ultrasound scanning is one of the methods that can be used for estimating fat carcass percentage of live animals (Grill et al., 2015).

Body Condition Score (BCS) have been used as reflection of animal's fat reserve. However, BCS is a subjective means of assessing an animal's lean body mess and body fat (Summers et al., 2012). Therefore there is a need to have an accurate and objective measurement for estimating fat while the animal is still alive. Body measurement can be used to predict body



weight and carcass characteristics (Agamy et al., 2015). Chest girth is one of variable that have been used to predict body weight. Agamy et al. (2015) reported that chest girth had the highest correlation with body weight in Barki Lambs. The fat thickness of lambs can also indicated in chest depth. This study was conducted to evaluate the possibility of using chest measurement for subcutaneous, intermuscular, and total fat percentage in Thin Tailed Lambs.

## Materials and Methods

Twenty one heads of male thin tailed lambs ( $\pm$  3 months old) with initial body weight (BW)  $14.57 \pm 2.19$  kg (CV= 15.03%) were used in this study. They were fattened by fed the complete feed contained three levels of crude protein (CP; 14, 16 and 18%) and two levels of total digestible nutrients (TDN; 60, 70%) that composed of rice bran, cassava meal, sugar cane top, cassava peel, soybean meal, fish meal, molasses and mineral and was given in pelleted form. All lambs were housed in individual pen and given freely access to feed and water throughout the experimental period.


Parameters measured were CG (cm), CW (cm), CD (cm), CG/CD, CW/CD, CW/CG, subcutaneous, intermuscular, and total fat percentage of carcass weight. Those measurements were taken before slaughtered that used tape measures and elaborate calipers. All lambs were slaughtered randomly after 3 months of feeding. Lambs were fasted for 6 hours before slaughtered. The slaughter method was done follow Islamic methods. The carcass was kept in a cold room at 18°C for 10 hours. The fat of carcass were separated into subcutaneous, intermuscular, and total fat (comprises of subcutaneous and intermuscular), then were weighed.

The relationship between body measurements and fat carcass percentage were analyzed by correlation regression analysis. The strength of correlation coefficient was evaluated by the value described by Steel and Torrie (1960), while the accuracy of the equation of regression was evaluated by t-test, standard error (SE) and the differences between predicted value and actual value. The accuracy was also determined if the difference value was low and statistically significant.

## Results and Discussion

The correlation between chest measurement and fat carcass percentage are presented in Figure 1. The percentage of subcutaneous fat in CG, CW, CD, CG/CD, and CW/CD has a positive correlation value (0.163, 0.074, 0.135, 0.076, and 0.017, respectively) while in CW/CG was negative (-0.019). The negative correlation in CW/CG is caused by the growth of subcutaneous fat might be maximum, so it will decrease the growth rate. Owens et al. (1993) reported that fat deposition is starting from intermuscular, subcutaneous and intramuscular. Intermuscular and total fat percentage showed a positive correlation in all chest measurements. The correlation between CG in intermuscular and total fat percentage showed a strong correlation value, being 0.582 and 0.664, respectively, while, in CG/CD showed a medium correlation value, being 0.510 and 0.533, respectively. The bigger chest measurement value will give the more fat carcass percentage. This result agreed to the report of Abd-Alla (2014) that total fat stores weight of carcass were positively correlated with chest girth in Barki Lambs.





**Figure 1.** The correlation between chest measurements and fat carcass percentage; % subcutaneous (●); % intermuscular (○) and % total fat (□).

Table 1 shows the equation regression of chest measurement and fat carcass percentage. The table showed the correlation of intermuscular and total fat in CG, CW and CG/CD were statistically significant ( $P < 0.05$ ), while subcutaneous fat was not significant ( $P > 0.05$ ). The growth of subcutaneous fat might be stable, so it's statistically not significant with chest measurements. Fat deposition is starting from intermuscular, subcutaneous and intramuscular (Owens et al., 1993). Chest girth were reported to have the highest correlation with body weight in Barki Lambs (Agamy et al., 2015). The SE of both variables showed a low value (from 0.002 to 0.020) and have a small differences of the actual value (0.368 - 7.520%). These result means either intermuscular or total fat regression equation in CG and CG/CD can be used. The CG regression is the best indicator of overall value, because it has the lowest differences and strong correlation in intermuscular and total fat percentage. The results of this study are in agreement with Nigm et al. (1995) who concluded that heart girth (or chest girth) was the best single measurement for predicting different carcass traits of Merino sheep.

## Conclusion

This study concluded that chest girth (CG) has a strong correlation to intermuscular and total fat carcass percentage, so that the regression equation using CG has a good accuracy to estimate intermuscular and total fat carcass percentage in thin tailed lambs as shown by the low differences between the predicted and actual value.



## Acknowledgements

The authors thank to the members of “Domba Cemara Team” of Faculty of Animal and Agricultural Sciences, Diponegoro University, for their kind supports and help in this study.

## References

- Abd-Alla, M. S. 2014. A comparative study on body measurements and carcass characteristics in Egyptian Sheep and Goats. *Asian J. Anim. Vet. Adv.* 9: 292-301.
- Agamy, R., A.Y. Abdel-Moneim, M. S. Abd-Alla, I.I. Abdel-Mageed, and G. M. Ashmawi. 2015. Using linear body measurements to predict body weight and carcass characteristics of three Egyptian Fat-Tailed Sheep breeds. *Asian J. Anim. Vet. Adv.* 10 (7): 335-344.
- Grill, L., F. Ringdorfer, R. Baumung, and B. Fuerst-Waltl. 2015. Evaluation of ultrasound scanning to predict carcass composition of Austrian meat sheep. *Small Rumin. Res.* 123, 260-268.
- Hopkins, D. L., R. S. Hegarty, P. J. Walker, and D.W. Pethick. 2006. Relationship between animal age, intramuscular fat, cooking loss, pH, shear force and eating quality of aged meat from sheep. *Aust. J. Exp. Agric.* 46: 879-884.
- Nigm, A. A., O. M. Abdalla, M.B. Aboul-Ela, H.M. Kamel, and M.A. Ahmed, 1995. Meat characteristics of sheep and goat breeds commonly consumed in U.A.E. 2. Use of body dimensions for predicting body and carcass weights. *Emir. J. Agric. Sci.*, 7: 39-54.
- Owens, F.N., P. Dubeski, and C.F. Hanson. 1993. Factors that alter the growth and development of ruminants. *J. Anim. Sci.* 71 (11): 3138-3150.
- Steel, R. G. D., and J .H. Torrie. 1960. *Principles and Procedures of Statistics*. McGraw-Hill Book Co., New York.
- Summers, L., K. J. Clingerman, and X. Yang. 2012. Validation of a body condition scoring system in Rhesus Macaques (*Macaca mulatta*): assesment of body composition by using dual-energy X-ray absorptiometry. *J. Anim. Assoc. Lab. Anim. Sci.* 51 (1): 88-93.