

Journal of the Arkansas Academy of Science

Volume 71

Article 41

2017

Anatomical Distribution of Clinostomum Metacercariae in the Tissues of Pond-Raised Channel Catfish (*Ictalurus punctatus*)

J. Singleton

University of Maryland Eastern Shore, Princes Anne, MD


James J. Daly Sr.

University of Arkansas for Medical Sciences, Little Rock, AR, jamesdalysr@yahoo.com

K. Wagner

University of Arkansas for Medical Sciences, Little Rock, AR.

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

 Part of the [Animal Diseases Commons](#), [Aquaculture and Fisheries Commons](#), and the [Parasitic Diseases Commons](#)

Recommended Citation

Singleton, J.; Daly, James J. Sr.; and Wagner, K. (2017) "Anatomical Distribution of Clinostomum Metacercariae in the Tissues of Pond-Raised Channel Catfish (*Ictalurus punctatus*)," *Journal of the Arkansas Academy of Science*: Vol. 71 , Article 41.

Available at: <http://scholarworks.uark.edu/jaas/vol71/iss1/41>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, ccmiddle@uark.edu.

Anatomical Distribution of *Clinostomum* Metacercariae in the Tissues of Pond-Raised Channel Catfish (*Ictalurus punctatus*)

J. Singleton¹, J.J. Daly Sr², and K. Wagner²

¹University of Maryland Eastern Shore (retired), Princes Anne, MD 21853

²University of Arkansas for Medical Sciences (retired), Little Rock, AR 72205

Correspondence: jamesdalysr@yahoo.com

Running title: *Clinostomum* in Channel Catfish

Previously Daly *et al.* (2007) found that the distribution of *Clinostomum marginatum* (“yellow grub”) metacercariae in the mouth and gills (orobranchial cavity) of smallmouth bass (*Micropterus dolomieu*) was highly proportional to the total body metacercariae. One could use this relationship to estimate the *Clinostomum* larval abundance in a smallmouth population by counting only the number in the mouth and gills without lethal necropsy. Lorio (1989) pointed out that yellow grub in channel catfish could cause a marketing problem for catfish farmers. A simple examination of visible anatomic sites (orobranchial areas) would be helpful for catfish growers as a tool for monitoring yellow grub in their stock. An infection of yellow grub in catfish (*Ictalurus punctatus*) in a pond in Northwest Arkansas offered the opportunity to see if such an approach would be feasible and worthwhile and to see if similar tissue distribution of proportionality existed with another fish host other than smallmouth bass.

Fifty- four catfish of similar age and size (35±3.9 cm SL; range 28-45; weight 326±169 g; range 190-1215) were taken from a pond in Washington Co. in 1995 and necropsied. The recovered yellow grubs (1712 from 54 hosts) into groupings of mouth, muscles, gills, fins and internal sites and counted. Descriptive statistics and regression analysis were done with Microsoft excel 2010.

In Fig. 1. the percentage of cysts in each of the anatomical sites are seen. The majority of the cysts, (59%), are in the orobranchial visible areas of the fish (gills + mouth). The population descriptors for yellow grub in pond-raised catfish are found in Table 1. All sites but one, muscle, have SD/Mean (Index of Dispersion) ratios of much less than one indicating a random infection process. This is unusual since most helminth infections have shown a stochastic and overdispersion of cysts in a few hosts and fewer worms in most of the other hosts. The simplest explanation for this would be that the commercial pond environment

would favor random association with snail-released cercariae because there are few or no areas that the host fish can establish territorial dominance that would otherwise stratify the host-parasite relationship. The muscle SD/Mean data would indicate a different infection route for that particular anatomical site. Mean intensity, i.e. removing zero infections from the calculations, did not show much difference from total population data due to the relatively small number of zero infections. Regression analysis of mean abundance with total population versus other sites were found to be highly correlative (Table 2, Fig. 2). Importantly, total population versus the visible sites (gill-mouth) showed high correlation with $r = 0.89$, $p = 3.7E-19$. This data (gill + mouth = 59% of the cysts) somewhat agreed with that of Vianna *et al.* (2005), with *C. complanatum* in *Rhamdea quellan* (a Brazilian catfish) which showed 42 % of metacercariae in the head region of the host but differed from 16 different Ouachita and Ozark smallmouth infections where less grubs were found in the head region: 14 % with a range of 5-16%, (Daly *et al.* 2014). and 19% (Taber 1972).




Figure 1. Percent distribution of *Clinostomum* sp. metacercarial cysts in different anatomical sites of pond-raised channel catfish (*Ictalurus punctatus*).

J. Singleton, J.J. Daly Sr, and K Wagner

Table 1. Population descriptors (Bush *et al.* 1997) of *Clinostomum* sp. metacercarial cysts in different anatomical sites in channel catfish (*Ictalurus punctatus*) from a pond in Northwest Arkansas.

Mean abundance							
	Total	Gill	Mouth	Mouth+Gill	Fin	Muscle	Internal
Mean	31.7	5.7	13.0	18.7	5.8	3.9	3.3
SD	21.4	5.5	8.7	11.8	5.6	5.8	3.6
Max	92	19	34	48	25	28	16
%	100	83	98	100	91	70	67
SD/Mean	0.68	0.78	0.64	0.63	0.85	1.1	0.68

Mean Intensity							
	Total	Gill	Mouth	Mouth+Gill	Fin	Muscle	Internal
Mean	31.7	5.7	13.0	18.7	6.5	5.7	5.1
SD	21.4	5.3	8.6	11.8	5.5	6.2	3.4
Count	54	45	53	54	48	37	35
SD/Mean	0.68	0.78	0.5	0.63	0.85	1.1	0.67

Table 2. Regression analyses for key population descriptors of *Clinostomum* sp cysts in *Ictalurus punctatus*.

Independent Variable	Dependent Variable	X	Intercept	r	P
Total	Gill	0.16	0.7	0.60	8.3E-08
Total	Mouth	0.33	2.5	0.82	5.3E-14
Total	Gill + Mouth	1.60	1.5	0.89	3.7E-19
Total	Fins	3.20	13.1	0.83	7.6E-12
Total	Muscle	0.22	1.1	0.78	5.2E-12
Total	Internal Sites	0.09	0.6	0.50	1.0E-04




Figure 2. Regression analysis of total yellow grubs versus gill-mouth yellow grubs. Dashed line is actual total number of grubs and the circles represent the predicted total grubs calculated from regression coefficients.

***Clinostomum* in Channel Catfish**

In conclusion, this study shows that proportionality of *Clinostomum* larval infections exists between anatomical sites in commercial catfish as well as in smallmouth bass and also in an acanthocephalan infection of a microcrustacean. (Daly *et al.* 2014; Daly and Wagner 2016). Furthermore, McAllister *et al.* 2010 used this technique for estimating yellow grub in largemouth bass from a pond that did not require lethal necropsying of a highly valued host. Thus, counting visible grubs in the head region without necropsy gives a good estimate of the total worm burden and can be a useful tool for survey work and for catfish farmers who would not have to sacrifice economically valuable stock in order to monitor for yellow grub infections.

Literature Cited

- Bush AO, KD Lafferty, JF Lotz, and AW Shostak.** 1997. Parasitology meets ecology on its own terms. Margolis *et al.* revisited. *Journal of Parasitology* 83:575-583.
- Daly JJ Sr. and K Wagner.** 2016. Proportionality between population descriptors (covariables) in an aggregated acanthocephalan (helminth) infection of a microcrustacean, *Journal of the Arkansas Academy of Science* 70:82-87.
- Daly JJ Sr.** 2014. Proportionality of population descriptors of metacercariae of *Clinostomum marginatum* in the orobranchial cavity of black bass (*Micropterus* spp.) from Arkansas Ozark and Ouachita streams. *Journal of the Arkansas Academy of Science* 68:57-64
- Daly JJ Sr, RJ Keller, and B DeYoung.** 2007. A non-invasive technique for assessing the population parameters of metacercariae of *Clinostomum marginatum* in smallmouth bass (*Micropterus dolomieu*). *Journal of the Arkansas Academy of Science* 61:37-43.
- Lorio WJ.** 1989. Experimental control of metacercariae of the yellow grub *Clinostomum marginatum* in channel catfish. *Journal of Aquatic Animal Health*. 1:269-271.
- McAllister CT, CR Bursey, JA Crawford, AR Kuhns, C Shaffer, and SE Trauth.** 2010. Metacercariae of *Clinostomum* (Trematoda: Digenea) from three species of *Ambystoma* (Caudata: Ambystomatidae) from Arkansas and Illinois. *Comparative Parasitology* 77:25-30.
- Taber CA.** 1972. The yellow grub in centarchids of Southwest Missouri streams. *Progressive Fish-Culturist* 34:119.
- Vianna RT, JP Junior, and SA Brandon.** 2005. *Clinostomum complanatum* (Digenea: Clinostomatidae) density in *Rhamdea quellen* (Siluriformes: Pimelodidae) from South Brazil. *Brazilian Archives of Biology and Technology* 48:635-642.