

CONVERGENCE OF HENCKY-TYPE DISCRETE BEAM MODEL TO EULER INEXTENSIBLE *ELASTICA* IN LARGE DEFORMATION: RIGOROUS PROOF

JEAN-JACQUES ALIBERT¹, ALESSANDRO DELLA CORTE²,
AND PIERRE SEPPECHER^{1,3}

1: IMATH, Université du Sud Toulon-Var, France

2: DIMA, Sapienza Università di Roma, Italy

3: L.M.A CNRS AMU Marseille, France

ABSTRACT. The present paper concerns rigorous homogenization of a Hencky-type discrete beam model, which is useful for the numerical study of complex fibrous systems as pantographic sheets as well as woven fabrics. Γ -convergence of the discrete model towards the inextensible Euler's beam model is proven and the result is established for placements in \mathbb{R}^d in large deformation regime. KEYWORDS: Γ -convergence; Inextensible Euler beam; Hencky-type beam discretization.

1. INTRODUCTION

Rigorous results on homogenization are very important for today's theoretical and applied mechanics. This is especially true for the numerical investigation of very complex systems, as even with today's computational tools they may require a long computation time, and thus the *a priori* reliability of the results is of course desirable. The investigation of metamaterials (see [11] for a review of recent results) is among the topical research directions in which one often deals with very expensive numerical simulations, as the implementation of the desired (often exotic) properties at the macro-scale are usually realized by means of a very complex microstructure ([18, 13]). The theory of microstructured/micromorphic continua is by now well developed, with several sound and interesting results (see e.g. [5, 19] as general references on Cosserat continua, [33, 48, 4] for related results and [40, 23, 47, 29, 9, 42, 36, 38] for different kinds of applications of microstructured models). Still, it is necessary to develop suitable convergence arguments if one wants to solidly rely on the numerical simulations based on the solution of the simplified equations coming from the micromorphic/generalized continuum model used for the description of the metamaterial.

In the present paper we focus on special micro-structured systems which can be described as discrete systems. In this case, the reliability of the homogenization has to be intended in two ways:

- (1) Real world micro-structured systems with suitably small characteristic lengths have to be well described by the homogenized continuum model;
- (2) The numerical simulation of the equations coming from the homogenized model (that are usually way simpler than the ones coming from the discrete model) has to converge in a suitable sense.

In principle, there is no reason to believe that the ordinary assumptions made for classical (Cauchy) continuum models are suitable for models describing objects that are so different from the phenomenology originally motivating them. Indeed, very often generalized continuum models are called for, and in particular higher gradient theories (see e.g. [12, 6, 16, 41, 17, 34]) are being successfully employed in a number of cases for the homogenization of systems with complex geometry at the micro-scale ([2, 8, 35, 37, 39]). In the present paper we address this kind of question for Euler's beam model (also known as *Elastica*), which is the elementary constituent for a large class of complex fibrous systems, including the promising case of pantographic sheets (see [44, 22, 10, 15] for theoretical and numerical results and [46] for experimental ones in this direction). Specifically, we want to provide a rigorous justification for the discrete approximation by Heinrich Hencky (1885-1951) [24] of Euler's beam model in large deformation, which is becoming increasingly topical in today's research in structural and computational mechanics ([20, 21, 1]) and metamaterials ([28]). In particular, we address here the ideal case in which the beam is perfectly inextensible, while future investigation will be devoted to the more general extensible case.

2. CONVERGENCE OF MEASURE FUNCTIONALS

Before setting the mechanical problem we are interested in, we need to recall some (well known) mathematical tools for describing the placement and the energy of the discrete beam model and define a suitable convergence for the sequence of the discrete energy functionals.

Let $(C[0, 1])^d$ be the space of vector valued continuous functions on $[0, 1]$ endowed with the uniform norm $\|\varphi\|_\infty := \sup\{\|\varphi(t)\| : t \in [0, 1]\}$ and $(\mathcal{M}[0, 1])^d$ the set of vector valued bounded measures on $[0, 1]$ endowed with the norm

$$\|\mu\|_{\mathcal{M}} := \sup\{\langle \mu, \varphi \rangle : \varphi \in (C[0, 1])^d, \|\varphi\|_\infty = 1\}$$

where $\langle \cdot, \cdot \rangle$ stands for the duality bracket between $(\mathcal{M}[0, 1])^d$ and $(C[0, 1])^d$. Recall that if a sequence of vector valued bounded measures (μ_n) satisfies $\sup_n \|\mu_n\|_{\mathcal{M}} < +\infty$ then there exists a vector valued bounded measure μ and a subsequence (μ_{n_k}) which converges to μ with respect to the weak*-topology of $(\mathcal{M}([0, 1])^d)$ i.e.

$$\lim_{k \rightarrow \infty} \langle \mu_{n_k}, \varphi \rangle = \langle \mu, \varphi \rangle$$

for every $\varphi \in (C([0, 1])^d)$.

Let (F_n) and F be functionals on $(\mathcal{M}[0, 1])^d$ with values in $\mathbb{R} \cup \{+\infty\}$. We say that F_n Γ -converges to F if the following holds ([7]):

- (i) *Upper bound inequality.* For every $\mu \in (\mathcal{M}([0, 1])^d)$, there exists a sequence (μ_n) in $(\mathcal{M}[0, 1])^d$ weak*-converging to μ for which

$$\limsup_{n \rightarrow \infty} F_n(\mu_n) \leq F(\mu).$$

- (ii) *Lower bound inequality.* For every $\mu \in (\mathcal{M}[0, 1])^d$ and every sequence (μ_n) in $(\mathcal{M}[0, 1])^d$ weak*-converging to μ ,

$$\liminf_{n \rightarrow \infty} F_n(\mu_n) \geq F(\mu).$$

Such a Γ -convergence result is efficient when the following property of the sequence (F_n) holds:

- (iii) *Relative compactness.* For every sequence (μ_n) in $(\mathcal{M}[0, 1])^d$

$$\sup_n F_n(\mu_n) < +\infty \implies \sup_n \|\mu_n\|_{\mathcal{M}} < +\infty.$$

Informally speaking, relative compactness ensures that controlling the deformation energy is enough to control the norm of the measure employed for the description of the current configuration of the discrete model.

3. MICRO-MODEL FOR NON-LINEAR BEAMS

3.1. Discrete configurations and operators. Let δ_t denote the Dirac measure at the point $t \in [0, 1]$. The reference configuration of the discrete micro-system is constituted by $n + 1$ nodes placed at the points $\frac{i}{n}$, $i = 0, \dots, n$. Therefore it can be identified with the positive Radon measure $\frac{1}{n}$ placed at the points $\frac{i}{n}$ where $i = 0, 1, \dots, n$. Therefore, it will be described by the positive Radon measure on $[0, 1]$

$$\bar{\nu}_n := \frac{1}{n} \sum_{i=0}^n \delta_{\frac{i}{n}} \quad (1)$$

We assume that the reference (unstressed) configuration of the beam is straight, has unitary length and lays parallel to e_1 , i.e. the first vector of the canonical base of \mathbb{R}^d . The current configuration of the beam can be described by a vector bounded measure μ on $[0, 1]$ of the form $\mu(dt) := u(t)\bar{\nu}_n(dt)$ where the placement function $u : [0, 1] \rightarrow \mathbf{R}^d$ is defined $\bar{\nu}_n$ -almost everywhere i.e. at the points $\frac{i}{n}$ where $i = 0, 1, \dots, n$ (see Fig. 1 for a graphical representation of the discrete model).

In what follows, we will use the following notations.

$$\nu_n^+ := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{\frac{i}{n}} \quad \nu_n^- := \frac{1}{n} \sum_{i=1}^n \delta_{\frac{i}{n}} \quad \nu_n := \frac{1}{n} \sum_{i=1}^{n-1} \delta_{\frac{i}{n}} \quad (2)$$

$$D_n^+ u(t) := n(u(t + \frac{1}{n}) - u(t)) \quad D_n^- u(t) := n(u(t) - u(t - \frac{1}{n})) \quad (3)$$




FIGURE 1. Graphical representation of Hencky discrete model consisting of inextensible bars and rotational springs. In the graph $\theta_i := \theta_n(u)(\frac{i}{n})$.

$$D_n^2 u := n(D_n^+ u - D_n^- u) \quad (4)$$

Note that, if u is a placement function, $D_n^+ u$ is defined ν_n^+ -almost everywhere, $D_n^- u$ is defined ν_n^- -almost everywhere and $D_n^2 u$ is defined ν_n -almost everywhere.

3.2. Left hand side clamped inextensible beam. A placement function u is said to be admissible for a left hand side clamped beam if the following condition holds:

$$u(0) = 0 \quad \text{and} \quad D_n^+ u(0) = e_1 \quad (5)$$

It is said to be admissible for an inextensible beam if the following condition holds: $\|u(\frac{i+1}{n}) - u(\frac{i}{n})\| = \frac{1}{n}$ for $i = 0, 1, \dots, n-1$. This condition can be written

$$\|D_n^+ u\| = 1 \quad \nu_n^+ \text{-almost everywhere} \quad (6)$$

3.3. Deformation energy associated with three points interactions. At each node $\frac{i}{n}$, for $i = 1, \dots, n-1$, a rotational spring is placed, whose deformation energy depends on the angle $\theta_n(u)(\frac{i}{n}) \in (-\pi, +\pi)$ formed by the vectors $u(\frac{i+1}{n}) - u(\frac{i}{n})$ and $u(\frac{i}{n}) - u(\frac{i-1}{n})$. This energy must vanish when the angle is zero. We assume, following [14, 45], that this energy is proportional to $1 - \cos(\theta_n(u)(\frac{i}{n}))$. Hence, when the discrete system is in the configuration described by the bounded measure $\mu(dt) = u(t)\bar{\nu}_n(dt)$, its energy is given by

$$E_n^3(\mu) := \frac{1}{n} \sum_{i=1}^{n-1} n^2 (1 - \cos \theta_n(u)(\frac{i}{n})) \quad \text{where} \quad \cos \theta_n(u)(\frac{i}{n}) = \frac{D_n^+ u(\frac{i}{n}) \cdot D_n^- u(\frac{i}{n})}{\|D_n^+ u(\frac{i}{n})\| \|D_n^- u(\frac{i}{n})\|}$$

or equivalently

$$E_n^3(\mu) = \frac{1}{2} \int \left\| n \left(\frac{D_n^+ u(t)}{\|D_n^+ u(t)\|} - \frac{D_n^- u(t)}{\|D_n^- u(t)\|} \right) \right\|^2 \nu_n(dt)$$

The above energy is well defined if the placement function u is such that $D_n^+ u \neq 0$ ν_n^+ -almost everywhere. This is clearly the case when u is admissible for an inextensible beam. In this case, the discrete energy has the reduced form

$$E_n^3(\mu) = \frac{1}{2} \int \|D_n^2 u(t)\|^2 \nu_n(dt) \quad (7)$$

4. FROM MICRO TO MACRO MODEL - Γ -CONVERGENCE RESULT.

This section is devoted to left hand side clamped inextensible beam.

4.1. Functionals associated to the micro model. Let \mathcal{M}_n denote the set of those vector bounded measures of the form $\mu(dt) = u(t)\bar{\nu}_n(dt) \in (\mathcal{M}[0, 1])^d$ and such that

$$u(0) = 0 \quad \text{and} \quad D_n^+ u(0) = e_1 \quad \text{and} \quad \|D_n^+ u\| = 1 \quad \nu_n^+ \text{ - almost everywhere} \quad (8)$$

The total energy functional (associated to the discrete model) is given by

$$E_n(\mu) := \begin{cases} \frac{1}{2} \int \|D_n^2 u(t)\|^2 \nu_n(dt) & \text{if } \mu(dt) = u(t)\bar{\nu}_n(dt) \in \mathcal{M}_n \\ +\infty & \text{otherwise} \end{cases} \quad (9)$$

4.2. Functional associated to the macro model. Let $H^2(0, 1)$ denote the usual Sobolev space. Relying on well-known embedding theorems, any function $u \in H^2(0, 1)$ will be considered as a $C^1[0, 1]$ -function. Let \mathcal{M} be the set of those vector bounded measures of the form $\mu(dt) = u(t)dt \in (\mathcal{M}[0, 1])^d$ with $u \in (H^2((0, 1))^d)$ and such that

$$u(0) = 0 \quad \text{and} \quad u'(0) = e_1 \quad \text{and} \quad \|u'(t)\| = 1 \text{ for every } t \in [0, 1] \quad (10)$$

The total energy functional (associated to the continuous model) is given by

$$E(\mu) := \begin{cases} \frac{1}{2} \int_0^1 \|u''(t)\|^2 dt & \text{if } \mu(dt) = u(t)dt \in \mathcal{M} \\ +\infty & \text{otherwise} \end{cases} \quad (11)$$

4.3. Γ -convergence result. Our main result is the following :

Theorem 4.1. *The sequence (E_n) satisfies the relative compactness property and Γ -converges to the functional E .*

If we compare Theorem 4.1 with the results proved in [3], the difficulty relies on the fact that the beam is inextensible, which corresponds to a nonlinear constraint.

5. PROOF OF THE MAIN RESULT

5.1. Approximation of a sequence with bounded energy. Let (μ_n) be a sequence in $(\mathcal{M}(0, 1])^d$ with bounded energy. This means that there exists some positive real number M such that

$$\mu_n(dt) = u_n(t)\bar{\nu}_n(dt) \in \mathcal{M}_n \quad \text{and} \quad \int \|D_n^2 u_n(t)\|^2 \nu_n(dt) \leq M. \quad (12)$$

for every integer n . Let us define the sequence $(\bar{\mu}_n)$ by setting $\bar{\mu}_n(dt) = \bar{u}_n(t)dt$, with \bar{u}_n piecewise C^2 in $(0, 1)$ satisfying:

$$\begin{aligned} \bar{u}_n(0) &= 0 \quad , \quad \bar{u}'_n(0) = e_1 \\ \bar{u}''_n(t) &= D^2 u_n\left(\frac{i}{n}\right) \quad \text{as soon as } t \in \left(\frac{i}{n} - \frac{1}{2n}, \frac{i}{n} + \frac{1}{2n}\right). \end{aligned}$$

Notice that $\bar{u}_n \in (H^2(0, 1))^d$ but in general $\bar{u}_n \notin \mathcal{M}$ because $\|\bar{u}'_n(t)\|$ is not necessarily equal to 1. The following result will be used to establish the lower bound inequality.

Lemma 5.1. *Let (μ_n) be a sequence in $(\mathcal{M}(0, 1])^d$ with bounded energy. Then, the sequence (\bar{u}_n) defined above is bounded with respect to the usual H^2 -norm and satisfies the following properties.*

$$\int_0^1 \|\bar{u}''_n(t)\|^2 dt = \int \|D_n^2 u_n(t)\|^2 \nu_n(dt) \quad \text{for every } n, \quad (13)$$

$$\lim_{n \rightarrow \infty} \|\bar{u}'_n(t)\| = 1 \quad \text{for every } t \in [0, 1], \quad (14)$$

$$\bar{\mu}_n - \mu_n \text{ converges to 0 with respect to the weak* topology.} \quad (15)$$

Proof. One has $\bar{u}_n(0) = 0$, $\bar{u}'_n(0) = e_1$ and

$$\int_0^1 \|\bar{u}''_n(t)\|^2 dt = \sum_{i=1}^{n-1} \int_{\frac{i}{n} - \frac{1}{2n}}^{\frac{i}{n} + \frac{1}{2n}} \|\bar{u}''_n(t)\|^2 dt = \int \|D_n^2 u_n(t)\|^2 \nu_n(dt) \leq M$$

which implies that the sequence (\bar{u}_n) is bounded with respect to the usual H^2 -norm. Hence, the two sequences (\bar{u}'_n) and (\bar{u}_n) are equicontinuous on $[0, 1]$ and uniformly bounded on $[0, 1]$. More precisely, for any $s, t \in [0, 1]$,

$$\|\bar{u}'_n(t) - \bar{u}'_n(s)\| \leq \sqrt{M} \sqrt{|t - s|} \quad \text{and} \quad \|\bar{u}'_n(t)\| \leq 1 + \sqrt{M} \quad (16)$$

$$\|\bar{u}_n(t) - \bar{u}_n(s)\| \leq (1 + \sqrt{M})|t - s| \quad \text{and} \quad \|\bar{u}_n(t)\| \leq 1 + \sqrt{M} \quad (17)$$

On the other hand, a first computation gives that for any $i = 1, \dots, n - 1$,

$$\bar{u}'_n\left(\frac{i}{n} + \frac{1}{2n}\right) = e_1 + \int_0^{\frac{i}{n} + \frac{1}{2n}} \bar{u}''_n(t) dt = e_1 + \frac{1}{n} \sum_{k=1}^i D_n^2 u_n\left(\frac{k}{n}\right) = D_n^+ u_n\left(\frac{i}{n}\right)$$

Since $\|D_n^+ u_n\| = 1$ ν_n^+ -almost everywhere and the sequence (\bar{u}'_n) is equicontinuous on $[0, 1]$, we obtain (15). A second computation gives $\bar{u}_n(1) = u_n(1)$ and

$$\begin{aligned}\bar{u}_n\left(\frac{i}{n}\right) &= \frac{i}{n}e_1 + \int_0^{\frac{i}{n}} \left(\frac{i}{n} - s\right) \bar{u}_n''(s) ds \\ &= \frac{i}{n}e_1 + \frac{1}{n} \sum_{k=1}^{i-1} \left(n \int_{\frac{k}{n} - \frac{1}{2n}}^{\frac{k}{n} + \frac{1}{2n}} \left(\frac{i}{n} - s\right) ds \right) D_n^2 u_n\left(\frac{k}{n}\right) + \left(\int_{\frac{i}{n} - \frac{1}{2n}}^{\frac{i}{n}} \left(\frac{i}{n} - s\right) ds \right) D_n^2 u_n\left(\frac{i}{n}\right) \\ &= \frac{i}{n}e_1 + \frac{1}{n} \sum_{k=1}^{i-1} \left(\frac{i-k}{n}\right) D_n^2 u_n\left(\frac{k}{n}\right) + \frac{1}{8n^2} D_n^2 u_n\left(\frac{i}{n}\right) \\ &= u_n\left(\frac{i}{n}\right) + \frac{1}{8n^2} D_n^2 u_n\left(\frac{i}{n}\right)\end{aligned}$$

for every $i = 1, \dots, n-1$. As a consequence, the inequality $\|\bar{u}_n - u_n\| \leq \frac{\sqrt{M}}{8n}$ holds $\bar{\nu}_n$ -almost everywhere. Let $\varphi \in C([0, 1])^2$. A third computation gives

$$\begin{aligned}|\langle \bar{\mu}_n - \mu_n, \varphi \rangle| &= \left| \sum_{i=1}^n \int_{\frac{i-1}{n}}^{\frac{i}{n}} (\varphi(t) \cdot \bar{u}_n(t) - \varphi\left(\frac{i}{n}\right) \cdot u_n\left(\frac{i}{n}\right)) dt \right| \\ &\leq \sum_{i=1}^n \int_{\frac{i-1}{n}}^{\frac{i}{n}} \|\varphi(t)\| \|\bar{u}_n(t) - \bar{u}_n\left(\frac{i}{n}\right)\| dt + \sum_{i=1}^n \int_{\frac{i-1}{n}}^{\frac{i}{n}} \|\varphi(t) - \varphi\left(\frac{i}{n}\right)\| \|\bar{u}_n\left(\frac{i}{n}\right)\| dt \\ &\quad + \int \|\varphi(t)\| \|\bar{u}_n(t) - u_n(t)\| \nu_n(dt) \\ &\leq \frac{1 + \sqrt{M}}{n} \int_0^1 \|\varphi(t)\| dt + (1 + \sqrt{M}) \sum_{i=1}^n \int_{\frac{i-1}{n}}^{\frac{i}{n}} \|\varphi(t) - \varphi\left(\frac{i}{n}\right)\| dt \\ &\quad + \frac{\sqrt{M}}{8n} \int \|\varphi(t)\| \nu_n(dt)\end{aligned}$$

Since

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \int_{\frac{i-1}{n}}^{\frac{i}{n}} \|\varphi(t) - \varphi\left(\frac{i}{n}\right)\| dt = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \int \|\varphi(t)\| \nu_n(dt) = \int_0^1 \|\varphi(t)\| dt$$

we conclude that the sequence $(\bar{\mu}_n - \mu_n)$ converges to 0 with respect to the weak*-topology of $(\mathcal{M}[0, 1])^d$. The proof is complete. \square

5.2. The Proof of Theorem 4.1. We divide this proof in three steps.

Step 1. (*Relative compactness*). Let $\mu(dt) := u(t) \bar{\nu}_n(dt) \in \mathcal{M}_n$. Since $\|u(0)\| = 0$ and $\|D_n^+ u\| = 1$ ν_n^+ -almost everywhere, one has $\|u\| \leq 1$ $\bar{\nu}_n$ -almost everywhere, hence

$$\|\mu\|_{\mathcal{M}} = \int \|u(t)\| \bar{\nu}_n(dt) \leq 1.$$

Step 2. (*Upper bound inequality*). Let $\mu(dt) := u(t)dt \in \mathcal{M}$. Since $u \in (C^1[0, 1])^d$, we define $\mu_n(dt) = u_n(t)\bar{\nu}_n(dt)$ by setting

$$u_n(0) = 0 \quad \text{and} \quad u_n\left(\frac{i}{n}\right) = \frac{1}{n} \sum_{k=0}^{i-1} u'\left(\frac{k}{n}\right) \quad (\text{for } i = 1, \dots, n)$$

Note that $D_n^+ u_n\left(\frac{i}{n}\right) = u'\left(\frac{i}{n}\right)$. Then $D_n^+ u_n(0) = e_1$ and $\|D_n^+ u_n\| = 1$ ν_n^+ -almost everywhere. Hence one has $\mu_n \in \mathcal{M}_n$ and

$$D_n^2 u_n\left(\frac{i}{n}\right) = n\left(D_n^+ u_n\left(\frac{i}{n}\right) - D_n^+ u_n\left(\frac{i-1}{n}\right)\right) = n\left(u'\left(\frac{i}{n}\right) - u'\left(\frac{i-1}{n}\right)\right) = n \int_{\frac{i-1}{n}}^{\frac{i}{n}} u''(t) dt$$

then, using Jensen inequality we obtain

$$\limsup_{n \rightarrow \infty} \int \|D_n^2 u_n\|^2 d\nu_n = \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^{n-1} \left\| n \int_{\frac{k}{n}}^{\frac{k+1}{n}} u''(t) dt \right\|^2 \leq \int_0^1 \|u''(t)\|^2 dt.$$

Let $\varphi \in (C[0, 1])^d$. Since u' is continuous on $[0, 1]$ we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \langle \mu_n, \varphi \rangle &:= \lim_{n \rightarrow \infty} \int \varphi(t) \cdot u_n(t) \bar{\nu}_n(dt) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \varphi\left(\frac{i}{n}\right) \cdot \left(\frac{1}{n} \sum_{k=0}^{i-1} u'\left(\frac{k}{n}\right) \right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \varphi\left(\frac{i}{n}\right) \cdot \left(u\left(\frac{i}{n}\right) + \sum_{k=0}^{i-1} \int_{\frac{k}{n}}^{\frac{k+1}{n}} (u'\left(\frac{k}{n}\right) - u'(t)) dt \right) \\ &= \lim_{n \rightarrow \infty} \int \varphi(t) \cdot u(t) \bar{\nu}_n(dt) \end{aligned}$$

Hence, Riemann's Theorem implies that the sequence (μ_n) converges to μ with respect to the weak*-topology of $(\mathcal{M}[0, 1])^d$.

Step 3. (*Lower bound inequality*). Let $\mu, \mu_n \in (\mathcal{M}[0, 1])^d$ such that (μ_n) converges to μ with respect to the weak*-topology of $(\mathcal{M}[0, 1])^d$. Without loss of generality we may assume that $\mu_n(dt) = u_n(t)\bar{\nu}_n(dt) \in \mathcal{M}_n$ and there exists a nonnegative real number M such that for every n

$$\int \|D_n^2 u_n(t)\|^2 \nu_n(dt) \leq M.$$

Let $(\bar{\mu}_n)$ be the sequence of measures defined in subsection 5.1. By Lemma 5.1, this sequence converges to μ with respect to the weak*-topology of $(\mathcal{M}[0, 1])^d$. Since $\bar{\mu}_n(dt) = \bar{u}_n(t)dt$ and the sequence (\bar{u}_n) is bounded with respect to the usual H^2 -norm, there exists $u \in (H^2(0, 1])^d$ such that $\mu(dt) = u(t)dt$ and

$$\liminf_{n \rightarrow \infty} \int \|D_n^2 u_n(t)\|^2 \nu_n(dt) = \liminf_{n \rightarrow \infty} \int_0^1 \|\bar{u}_n''(t)\|^2 dt \geq \int_0^1 \|u''(t)\|^2 dt$$

Since the space $H^2(0, 1)$ is compactly embedded on $C^1[0, 1]$, the sequence (\bar{u}'_n) converges to (u') with respect to the uniform norm over $[0, 1]$. Hence, using Lemma 5.1, We obtain

$$\|u'(t)\| = 1 \quad \text{for every } t \in [0, 1]$$

then $u \in \mathcal{M}$. The proof is complete.

6. CONCLUSIONS

We proved a Γ -convergence result for a Hencky-type discretization of an inextensible Euler beam in large deformation regime. Future investigations should generalize the result (in a suitable form) for extensible beam models; moreover, it will be interesting to extend the convergence argument to Generalized Beam Models ([32, 31, 30, 25]) and also to the dynamics of the discrete system, which should of course take into account the possibility of various kinds of dynamic instabilities ([43, 27, 26]). Finally, it has to be remarked that Hencky-type discretization for *Elastica* has proven to be very effective, and is in fact used by several computational software packages (as for instance by MATLAB[®]). The present result gives a sound mathematical argument which this kind of numerical evidence can be based on.

REFERENCES

- [1] B Emek Abali, Wolfgang H Müller, and Victor A Eremeyev. Strain gradient elasticity with geometric nonlinearities and its computational evaluation. *Mechanics of Advanced Materials and Modern Processes*, 1(1):1, 2015.
- [2] Jean-Jacques Alibert and Alessandro Della Corte. Second-gradient continua as homogenized limit of pantographic microstructured plates: a rigorous proof. *Zeitschrift für angewandte Mathematik und Physik*, 66(5):2855–2870, 2015.
- [3] Jean-Jacques Alibert, Pierre Seppecher, and Francesco dell’Isola. Truss modular beams with deformation energy depending on higher displacement gradients. *Mathematics and Mechanics of Solids*, 8(1):51–73, 2003.
- [4] Holm Altenbach and Victor A Eremeyev. On the linear theory of micropolar plates. *ZAMM-Journal of Applied Mathematics and Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik*, 89(4):242–256, 2009.
- [5] Johannes Altenbach, Holm Altenbach, and Victor A Eremeyev. On generalized cosserat-type theories of plates and shells: a short review and bibliography. *Archive of Applied Mechanics*, 80(1):73–92, 2010.
- [6] Ugo Andreaus, Francesco dell’Isola, Ivan Giorgio, Luca Placidi, Tomasz Lekszycki, and Nicola Luigi Rizzi. Numerical simulations of classical problems in two-dimensional (non) linear second gradient elasticity. *International Journal of Engineering Science*, 108:34–50, 2016.
- [7] Andrea Braides. *Gamma-convergence for Beginners*, volume 22. Clarendon Press, 2002.
- [8] Antonio Carcaterra, Francesco dell’Isola, R Esposito, and M Pulvirenti. Macroscopic description of microscopically strongly inhomogenous systems: A mathematical basis for the synthesis of higher gradients metamaterials. *Archive for Rational Mechanics and Analysis*, 218(3):1239–1262, 2015.

- [9] Ching S Chang and Anil Misra. Packing structure and mechanical properties of granulates. *Journal of engineering mechanics*, 116(5):1077–1093, 1990.
- [10] M Cuomo, F dell’Isola, and Leopoldo Greco. Simplified analysis of a generalized bias test for fabrics with two families of inextensible fibres. *Zeitschrift für angewandte Mathematik und Physik*, 67(3):1–23, 2016.
- [11] Dionisio Del Vescovo and Ivan Giorgio. Dynamic problems for metamaterials: review of existing models and ideas for further research. *International Journal of Engineering Science*, 80:153–172, 2014.
- [12] F. dell’Isola, G. Sciarra, and S. Vidoli. Generalized Hooke’s law for isotropic second gradient materials. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 2009.
- [13] Francesco dell’Isola, Sara Bucci, and Antonio Battista. Against the fragmentation of knowledge: The power of multidisciplinary research for the design of metamaterials. In *Advanced Methods of Continuum Mechanics for Materials and Structures*, pages 523–545. Springer, 2016.
- [14] Francesco dell’Isola, I Giorgio, M Pawlikowski, and NL Rizzi. Large deformations of planar extensible beams and pantographic lattices: heuristic homogenization, experimental and numerical examples of equilibrium. In *Proc. R. Soc. A*, volume 472, page 20150790. The Royal Society, 2016.
- [15] Francesco dell’Isola, Ivan Giorgio, and Ugo Andreaus. Elastic pantographic 2d lattices: a numerical analysis on the static response and wave propagation. *Proceedings of the Estonian Academy of Sciences*, 64(3):219, 2015.
- [16] Francesco dell’Isola and Pierre Seppecher. The relationship between edge contact forces, double forces and interstitial working allowed by the principle of virtual power. *Comptes rendus de l’Académie des sciences. Série IIB, Mécanique, physique, astronomie*, page 7, 1995.
- [17] Francesco dell’Isola and David Steigmann. A two-dimensional gradient-elasticity theory for woven fabrics. *Journal of Elasticity*, 118(1):113–125, 2015.
- [18] Francesco dell’Isola, David Steigmann, and Alessandro Della Corte. Synthesis of fibrous complex structures: Designing microstructure to deliver targeted macroscale response. *Applied Mechanics Reviews*, 67(6):060804, 2015.
- [19] Victor A Eremeyev, Leonid P Lebedev, and Holm Altenbach. Foundations of micropolar mechanics, 2012.
- [20] Demeter G Fertis. *Nonlinear structural engineering*. Springer, 2006.
- [21] Samuel Forest and Rainer Sievert. Nonlinear microstrain theories. *International Journal of Solids and Structures*, 43(24):7224–7245, 2006.
- [22] Ivan Giorgio, Alessandro Della Corte, Francesco dell’Isola, and David J Steigmann. Buckling modes in pantographic lattices. *Comptes rendus Mecanique*, 344(7):487–501, 2016.
- [23] Ivan Giorgio and Daria Scerrato. Multi-scale concrete model with rate-dependent internal friction. *European Journal of Environmental and Civil Engineering*, pages 1–19, 2016.
- [24] Heinrich Hencky. *Über die angenäherte Lösung von Stabilitätsproblemen im Raum mittels der elastischen Gelenkkette*. PhD thesis, Engelmann, 1921.
- [25] A Luongo and D Zulli. A non-linear one-dimensional model of cross-deformable tubular beam. *International Journal of Non-Linear Mechanics*, 66:33–42, 2014.
- [26] Angelo Luongo and Francesco D’Annibale. Double zero bifurcation of non-linear viscoelastic beams under conservative and non-conservative loads. *International Journal of Non-Linear Mechanics*, 55:128–139, 2013.
- [27] Angelo Luongo and Francesco D’Annibale. A paradigmatic minimal system to explain the ziegler paradox. *Continuum Mechanics and Thermodynamics*, 27(1-2):211–222, 2015.

- [28] Graeme Walter Milton. Adaptable nonlinear bimode metamaterials using rigid bars, pivots, and actuators. *Journal of the Mechanics and Physics of Solids*, 61(7):1561–1568, 2013.
- [29] Anil Misra and Yang Yang. Micromechanical model for cohesive materials based upon pseudo-granular structure. *International Journal of Solids and Structures*, 47(21):2970–2981, 2010.
- [30] Giuseppe Piccardo, Alberto Ferrarotti, and Angelo Luongo. Nonlinear generalized beam theory for open thin-walled members. *Mathematics and Mechanics of Solids*, page 1081286516649990, 2016.
- [31] Giuseppe Piccardo, Gianluca Ranzi, and Angelo Luongo. A complete dynamic approach to the generalized beam theory cross-section analysis including extension and shear modes. *Mathematics and Mechanics of Solids*, 19(8):900–924, 2014.
- [32] Giuseppe Piccardo, Gianluca Ranzi, and Angelo Luongo. A direct approach for the evaluation of the conventional modes within the gbt formulation. *Thin-Walled Structures*, 74:133–145, 2014.
- [33] W Pietraszkiewicz and VA Eremeyev. On natural strain measures of the non-linear micropolar continuum. *International Journal of Solids and Structures*, 46(3):774–787, 2009.
- [34] Luca Placidi. A variational approach for a nonlinear one-dimensional damage-elasto-plastic second-gradient continuum model. *Continuum Mechanics and Thermodynamics*, 28(1-2):119–137, 2016.
- [35] Luca Placidi, Ugo Andreaus, and Ivan Giorgio. Identification of two-dimensional pantographic structure via a linear d4 orthotropic second gradient elastic model. *Journal of Engineering Mathematics*, DOI: 10.1007/s10665-016-9856-8, 2016.
- [36] Luca Placidi, Sérgio H Faria, and Kolumban Hutter. On the role of grain growth, recrystallization and polygonization in a continuum theory for anisotropic ice sheets. *Annals of Glaciology*, 39(1):49–52, 2004.
- [37] Luca Placidi, Leopoldo Greco, Sara Bucci, Emilio Turco, and Nicola Luigi Rizzi. A second gradient formulation for a 2d fabric sheet with inextensible fibres. *Zeitschrift für angewandte Mathematik und Physik*, 67(5):114, 2016.
- [38] Luca Placidi, Ralf Greve, Hakime Seddik, and Sérgio H Faria. Continuum-mechanical, anisotropic flow model for polar ice masses, based on an anisotropic flow enhancement factor. *Continuum Mechanics and Thermodynamics*, 22(3):221–237, 2010.
- [39] Y Rahali, I Giorgio, JF Ganghoffer, and Francesco dell’Isola. Homogenization à la piola produces second gradient continuum models for linear pantographic lattices. *International Journal of Engineering Science*, 97:148–172, 2015.
- [40] Daria Scerrato, Ivan Giorgio, Angela Madeo, Ali Limam, and Felix Darve. A simple nonlinear model for internal friction in modified concrete. *International Journal of Engineering Science*, 80:136–152, 2014.
- [41] Giulio Sciarra, Nicoletta Ianiro, Angela Madeo, et al. A variational deduction of second gradient poroelasticity i: general theory. *Journal of Mechanics of Materials and Structures*, 3(3):507–526, 2008.
- [42] Hakime Seddik, Ralf Greve, Luca Placidi, Ilka Hamann, and Olivier Gagliardini. Application of a continuum-mechanical model for the flow of anisotropic polar ice to the edml core, antarctica. *Journal of Glaciology*, 54(187):631–642, 2008.
- [43] Gerard Taig, Gianluca Ranzi, and Angelo Luongo. Gbt pre-buckling and buckling analyses of thin-walled members under axial and transverse loads. *Continuum Mechanics and Thermodynamics*, 28(1-2):41–66, 2016.

- [44] Emilio Turco, Francesco dell’Isola, Nicola Luigi Rizzi, Roman Grygoruk, Wolfgang H Müller, and Christian Liebold. Fiber rupture in sheared planar pantographic sheets: Numerical and experimental evidence. *Mechanics Research Communications*, 76:86–90, 2016.
- [45] Emilio Turco, Francesco dell’Isola, Antonio Cazzani, and Nicola Luigi Rizzi. Hencky-type discrete model for pantographic structures: numerical comparison with second gradient continuum models. *Zeitschrift für angewandte Mathematik und Physik*, 67(4):1–28, 2016.
- [46] Emilio Turco, Maciej Golaszewski, Antonio Cazzani, and Nicola Luigi Rizzi. Large deformations induced in planar pantographic sheets by loads applied on fibers: experimental validation of a discrete lagrangian model. *Mechanics Research Communications*, 76:51–56, 2016.
- [47] Yang Yang and Anil Misra. Micromechanics based second gradient continuum theory for shear band modeling in cohesive granular materials following damage elasticity. *International Journal of Solids and Structures*, 49(18):2500–2514, 2012.
- [48] VA Yeremeyev and LM Zubov. The theory of elastic and viscoelastic micropolar liquids. *Journal of Applied Mathematics and Mechanics*, 63(5):755–767, 1999.