

Relationship between blood remifentanyl concentration and stress hormone levels during pneumoperitoneum in patients undergoing laparoscopic cholecystectomy

P. ACETO¹, C. DELLO RUSSO², C. LAI³, V. PERILLI¹, N. FUCCI⁴,
N. DE GIOVANNI⁴, A. PIRAS⁵, P. NAVARRA², L. SOLLAZZI¹

¹Department of Anesthesiology and Intensive Care, A. Gemelli University Hospital Foundation, Rome, Italy

²Institute of Pharmacology, Catholic University Medical School, Rome, Italy

³Department of Dynamic and Clinical Psychology, "Sapienza" University of Rome, Italy

⁴Institute of Public Health, Section of Legal Medicine, Catholic University Medical School, Rome, Italy

⁵Institute of Clinical Pediatrics, Catholic University Medical School, Rome, Italy

Abstract. – OBJECTIVE: The effect of remifentanyl on stress response to surgery is unclear. However, there are not clinical studies investigating the relationship between blood remifentanyl concentrations and stress hormones. Therefore, the aim of the present study was to assess the association between blood remifentanyl concentrations measured after pneumoperitoneum and cortisol (CORT) or prolactin (PRL) ratio (intraoperative/preoperative value), in patients undergoing laparoscopic cholecystectomy.

PATIENTS AND METHODS: Patients did not receive any pre-anesthetic medication. Anesthesia induction was standardized. Anesthesia maintenance was performed with inhaled sevoflurane at age-adjusted 1.0 minimum alveolar concentration and intravenous remifentanyl at infusion rate ranging from 0.1 to 0.4 mcg/kg/min. Blood samples were withdrawn before anesthesia induction and 5 min after achieving a pneumoperitoneum pressure of 12 mmHg. Correlation analyses were performed to evaluate the relationship between measured blood remifentanyl concentrations, CORT or PRL ratio (intraoperative/preoperative value) and remifentanyl dose delivered by the pump.

RESULTS: A significant inverse correlation was found between CORT ratio and measured blood remifentanyl concentration ($p=0.03$) or planned remifentanyl dose ($p=0.04$). No correlations were found between blood remifentanyl concentration and PRL ratio ($p=0.83$).

CONCLUSIONS: Our data suggest that the CORT response to surgical stress is more efficiently counteracted by increased blood remifentanyl concentration.

Key Words

Remifentanyl, Cortisol, Prolactin, Stress, Pneumoperitoneum, General anesthesia.

Introduction

It has been demonstrated that laparoscopic surgery may produce a moderate stress response although this approach is considered as minimally invasive¹. Intravenously administered remifentanyl, combined with general anesthetics, provides stable hemodynamic during laparoscopic surgery^{2,3}. Previous studies regarding remifentanyl effect on stress response in laparoscopic surgery are conflicting. Some authors reported remifentanyl dose-independent stress hormone variations in response to pneumoperitoneum, whereas other studies demonstrated reduced hormone release only at high doses^{1,4}. In this regard, it may be suggested that remifentanyl, as other opioids, could mitigate stress response activation in a dose-dependent manner^{5,6}. Currently, there are not clinical studies investigating the relationship between blood concentrations of remifentanyl and stress hormones during surgery. In this scenario, the aim of the present study was to assess the association between blood remifentanyl concentration measured after pneumoperitoneum and stress hormones (cortisol and prolactin) ratio (intraoperative/preoperative value), in patients undergoing laparoscopic cholecystectomy. Also, we investigated the relationship between blood remifentanyl concentration and the remifentanyl dose delivered by pump.

Patients and Methods

After local Ethics Committee approval and patients' informed consent, 30 adult patients with an American Society of Anesthesiologists phy-

sical status I, scheduled for laparoscopic cholecystectomy, were recruited. Exclusion criteria were: age >70 years; Body Mass Index >30 kg/m²; chronic use of medications. In the pre-anesthetic room, blood samples were collected to measure the baseline levels of cortisol (CORT) and prolactin (PRL). Patients did not receive any pre-anesthetic medication, whereas anesthesia induction was standardized using propofol 2 mg/kg, remifentanyl 0.25 mcg/kg/min and cisatracurium 0.15 mg/kg. Anesthesia maintenance was performed with inhaled sevoflurane at 1.0 age-adjusted minimum alveolar concentration and intravenous remifentanyl at infusion rate ranging from 0.1 to 0.4 mcg/kg/min, to maintain non-invasive mean arterial pressure and heart rate within a range of 20% more or less than the basal values. Intravenous paracetamol 1 g and morphine 0.06 mg/kg were administered to patients and local anesthetic infiltration (ropivacaine 4.75%) of trocar insertion sites was performed at the end of surgery, accordingly to local multimodal analgesia protocol. The same surgeon performed all the procedures by using Hasson's open access technique. Intraoperative blood samples were collected 5 min after achieving a pneumoperitoneum pressure of 12 mmHg (carbon-dioxide flow rate of 1.2 L/min) to assess both remifentanyl blood concentration and intraoperative levels of stress hormones. Blood samples were stored in capped vacutainer tubes at low temperatures, -20°C and -80°C for stress hormone and remifentanyl evaluations respectively. Samples were analyzed within a maximum period of 90 days from surgery. To prevent remifentanyl degradation, collection tubes containing EDTA whole blood were placed in ice water immediately after sampling. CORT and PRL serum levels were measured by using a chemiluminescence assay developed for the Abbott Architect i1000sr immunologic analyzer (Abbott Laboratories, Chicago, IL, USA). Remifentanyl concentrations were measured via a laboratory-developed assay, according to the following protocol. Briefly, 1 mL aliquots of samples, containing an internal standard (alfentanil), were mixed with 2 mL acetonitrile. After centrifugation, the supernatant was evaporated and reconstituted with 1 mL buffer phosphate 0.1 M at pH=8.5. A liquid/liquid extraction with dichloromethane was performed. After evaporation of the solvent, the dry extracts were reconstituted with 25 mL ethyl acetate and analyzed by GC/MS (gas chromatography/mass spectrometry) technology. A focus

GC couplet with DSQ operative in electron impact mode (70eV) was employed. An Equity 5 capillary column, 30 m-0.25 mm-0.25 mm film thickness, was used for 1 min isotherm at 70°C with further 15°C/min linear increment up to 290°C (5 min final isotherm). The acquisition in selected ion monitoring mode was performed by choosing 3 ions for remifentanyl (m/z 227-303-319), and internal standard (alfentanil m/z 289-268-222). Bispectral index (BIS; BIS Vista™ monitoring system, Aspect Medical System Inc., Norwood, MA, USA) and cardiovascular parameters (non-invasive blood pressure and heart rate) were recorded throughout anesthesia.

Statistical Analysis

Correlation analyses were performed to assess the relationship between measured blood remifentanyl concentration, planned remifentanyl dose (delivered by pump) and CORT or PRL ratio (intraoperative/preoperative value) and between planned and measured remifentanyl concentration. $p < 0.05$ was considered statistically significant. All the analyses were performed using the Statistical Version 10.0 software (StatSoft, Tulsa, OK, USA).


Results

Out of the 30 patients originally enrolled in the study, only 24 patients (mean age: 48.7±12.9 years; gender (M/F): 11/13; weight: 73.6±12.4 kg) with completed case report form were included in the final analysis (Table I). 6 patients were excluded for the following reasons: one patient did not meet the exclusion criteria; another patient due to clotted blood withdrawal and 4 patients due to the abnormal values of remifentanyl blood concentrations (i.e. out of validated range, 0.1-20 ng/ml). A significant inverse correlation was found between CORT ratio and measured blood remifentanyl

Table I. Values are calculated from five consecutive readings, each one registered every minute after establishment of pneumoperitoneum.

Patients (n=24)	
BIS value	43.2±6.5
Heart rate	63.8±8.8
Systolic pressure	109±8.3
Diastolic pressure	66±5.1

Figure 1. Scatterplot showing an inverse correlation between remifentanyl measured concentration and CORT ratio.



concentration (Pearson's $r=-0.45$; $p=0.03$) or planned remifentanyl dose (Pearson's $r=-0.42$; $p=0.04$) (Figure 1). On the other hand, no correlation was found between blood remifentanyl concentration and PRL ratio (Pearson's $r=-0.05$; $p=0.83$). Also, we observed a significant correlation between planned remifentanyl dose (delivered by pump) and blood remifentanyl concentration (Pearson's $r=0.72$; $p=0.0001$).

Discussion

The main result of this study was that CORT ratio (intraoperative/preoperative blood levels) was lower in patients showing higher remifentanyl blood concentrations, thus indicating that surgical stress was counteracted more efficiently. Previous studies regarding the effect of remifentanyl on stress response in laparoscopic surgery are conflicting, including not clear evidence on dose-dependent hormonal response to pneumoperitoneum¹⁻⁴. It is essential to blunt stress hormone secretion to prevent adverse postoperative events, including cognitive and psychological complications⁷⁻⁹. The physiological feedback mechanism that leads to the inhibition of adrenocorticotrophic hormone synthesis due to increased CORT concentration is disrupted during surgery⁶. Therefore, the anesthetic drugs are essential to prevent an excessive hypothalamic-pituitary-adrenal (HPA) axis activation in response to surgical trauma. From this work, it emerges that remifentanyl can mitigate

cortisol response in a dose-dependent manner. On the other hand, most of the anesthetic drugs, including opioids, have been found to stimulate PRL release during anaesthesia¹⁰. It has been demonstrated that patients undergoing endovascular abdominal aortic aneurysm experienced less pain but showed a more intense PRL response in the remifentanyl group when compared to the midazolam-fentanyl group¹¹. However, in our study, no relationship was found between blood remifentanyl concentration and PRL release. Our findings also show a significant correlation between remifentanyl blood concentration and target infusion concentration, albeit the methodology used to assess remifentanyl blood level needs additional validation data to meet the Food and Drug Administration requirements for pharmacokinetic studies.

Conclusions

CORT response to surgical stress seemed to be more efficiently counteracted by increased blood remifentanyl concentration in patients undergoing laparoscopic cholecystectomy. Despite the small size of the analyzed sample, this study may contribute to understanding the complex interaction between the effects of surgical stimulus and remifentanyl on stress response.

Conflict of Interest

The authors declared no conflict of interest.

References

- 1) WATANABE K, KASHIWAGI K, KAMIYAMA T, YAMAMOTO M, FUKUNAGA M, INADA E, KAMIYAMA Y. High-dose remifentanil suppresses stress response associated with pneumoperitoneum during laparoscopic colectomy. *J Anesth* 2014; 28: 334-340.
- 2) ACETO P, PERILLI V, LAI C, SACCO T, MODESTI C, LUCA E, DE SANTIS P, SOLLAZZI L, ANTONELLI M. Minimum alveolar concentration threshold of sevoflurane for postoperative dream recall. *Minerva Anestesiol* 2015; 81: 1201-1209.
- 3) ACETO P, LAI C, PERILLI V, DELLO RUSSO C, FEDERICO B, NAVARRA P, PROIETTI R, SOLLAZZI L. Stress-related biomarkers of dream recall and implicit memory under anaesthesia. *Anaesthesia* 2013; 68: 1141-1147.
- 4) MYRE K, RAEDER J, ROSTRUP M, BUANES T, STOKLAND O. Catecholamine release during laparoscopic fundoplication with high and low doses of remifentanil. *Acta Anaesthesiol Scand* 2003; 47: 267-273.
- 5) DESBOROUGH JP. The stress response to trauma and surgery. *Br J Anaesth* 2000; 85: 109-117.
- 6) ACETO P, LAI C, DELLO RUSSO C, PERILLI V, NAVARRA P, SOLLAZZI L. Stress response to surgery, anaesthetics role and impact on cognition. *J Anesth Clin Res* 2015; 6: 539.
- 7) ACETO P, PERILLI V, LAI C, SACCO T, ANCONA P, GASPERIN E, SOLLAZZI L. Update on post-traumatic stress syndrome after anesthesia. *Eur Rev Med Pharmacol Sci* 2013; 17: 1730-1737.
- 8) CASATI A, FANELLI G, PIETROPAOLI P, PROIETTI R, TUFANO R, MONTANINI S, DANELLI G, NUZZI M, MENTEGAZZI F, TORRI G, MARTANI C, SPREAFICO E, FIERRO G, PUGLIESE F, DE COSMO G, ACETO P, SERVILLO G, MONACO F. Monitoring cerebral oxygen saturation in elderly patients undergoing general abdominal surgery: a prospective cohort study. *Eur J Anaesthesiol* 2007; 24: 59-65.
- 9) ZHI XL, LI CY, XUE M, HU Y, JI Y. Changes in cognitive function due to combined propofol and remifentanil treatment are associated with phosphorylation of Tau in the hippocampus, abnormal total water and calcium contents of the brain, and elevated serum S100 β levels. *Eur Rev Med Pharmacol Sci* 2016; 20: 2156-2162.
- 10) MUJAGIC Z, CICKO E, VERAG-BROZOVIC V, PRAŠO M. Serum level of cortisol and prolactin in patients treated under total intravenous anesthesia with propofol-fentanyl and under balanced anesthesia with isoflurane-fentanyl. *Cent Eur J Med* 2008; 3: 459-463.
- 11) BARBIERI A, GIULIANI E, GENAZZANI A, BARALDI E, FERRARI A, D'AMICO R, COPPI G. Analgesia and endocrine surgical stress: effect of two analgesia protocols on cortisol and prolactin levels during abdominal aortic aneurysm endovascular repair. *Neuro Endocrinol Lett* 2011; 32: 526-529.