

1 **Actin proteolysis during ripening of dry fermented sausages at different pH values**

2

3 A. Berardo^a, B. Devreese^b, H. De Maere^c, D.A. Stavropoulou^d, G. Van Royen^e, F. Leroy^d, S. De
4 Smet^{a*}

5

6 ^aLaboratory of Animal Nutrition and Animal Product Quality, Faculty of Bioscience Engineering,
7 Ghent University, Proefhoevestraat 10, 9090 Melle, Belgium

8 ^bLaboratory for Protein Biochemistry and Biomolecular Engineering, Department of Biochemistry
9 and Microbiology, Ghent University, K.L. Ledeganckstraat 35, B-9000 Ghent, Belgium

10 ^c Research Group for Technology and Quality of Animal Products, Department M²S, member of
11 LForCe, KU Leuven Technology Campus, Gebroeders De Smetstraat 1, B-9000 Ghent, Belgium

12 ^dResearch Group of Industrial Microbiology and Food Biotechnology (IMDO), Faculty of Sciences
13 and Bio-engineering Sciences, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium

14 ^e Technology and Food Science Unit, Institute for Agricultural and Fisheries Research (ILVO),
15 Brusselsesteenweg 370, 9090 Melle, Belgium

16

17

18

19

20 *Corresponding author:

21 Stefaan De Smet

22 Proefhoevestraat 10 9090 Melle, Belgium

23 Tel. ++ 32 (0)9 264 90 01

24 Fax ++ 32 (0)9 264 90 99

25 E-mail stefaan.desmet@ugent.be

26 **ABSTRACT**

27 In dry fermented sausages, myofibrillar proteins undergo intense proteolysis generating small
28 peptides and free amino acids that play a role in flavour generation. This study aimed to identify
29 small peptides arising from actin proteolysis, as influenced by the type of processing. Therefore,
30 two acidification profiles were imposed by adding a different dose of dextrose, in order to mimic
31 the pH normally obtained in southern-type (“high pH”; pH 5.2 after fermentation and pH 6.0 after
32 ripening) and northern-type (“low pH”; pH 4.9 after fermentation and pH 5.0 after ripening) dry
33 fermented sausages. The identification of peptides within the raw meat was done by liquid
34 chromatography coupled to mass spectrometry in a data-independent positive mode of acquisition
35 (LCMS^E), both after fermentation (4 days) and at the end of ripening (28 days). During
36 manufacturing of the dry fermented sausages, actin was highly proteolysed, especially in nine
37 regions of the sequence. After fermentation, 52 and 42 actin-derived peptides were identified at
38 high and low pH, respectively, which further increased to 66 and 144 peptides, respectively, at the
39 end of ripening. Looking at the cleavage sites, cathepsins B and D likely played an important role.
40 The activity of exopeptidases was also evident since many peptides differed by a single amino acid.

41

42 **Keywords:** actin, proteolysis, dry fermented sausages, mass spectrometry.

43

44 1. INTRODUCTION

45

46 Dry fermented sausages are processed meat products that undergo a fermentation and drying period
47 before consumption. During processing, quality-affecting physico-chemical changes occur of which
48 several are driven by proteolysis (Spaziani et al., 2009; Ordóñez et al., 1999). Proteolysis is carried
49 out by enzymes that are endogenous to the meat as well as by enzymes from microbial origin
50 (Molly et al., 1997). This process is of fundamental importance because it influences the flavour of
51 dry fermented sausages (Ordóñez et al., 1999). The resulting small peptides and amino acids not
52 only directly influence taste but also serve as substrates for microorganisms that further convert
53 them into flavour compounds, as is the case for coagulase-negative staphylococci (Stavropoulou et
54 al., 2015; Sánchez Mainar et al., 2016).

55 Proteolysis in dry fermented sausages is usually studied by determining protein and non-
56 protein nitrogen (Defernando et al., 1991) and free α -NH₂-N (Verplaetse et al., 1992), by
57 electrophoresis techniques (Hughes et al., 2002), and by measuring the release of amino acids
58 during ripening (Defernando et al., 1991). In addition, the use of mass spectrometry-based
59 proteomic techniques now also allows for the identification of peptides generated during ripening
60 (Mora et al., 2015). As a result, cleavage sites of proteolytic enzymes can be determined to better
61 unravel the proteolytic mechanisms at play during ripening. Proteomic approaches also have the
62 potential to generate a more accurate view on the effect of processing factors on the dynamics of
63 proteolysis. It is for instance known that proteolytic enzymes have different pH optima (Hughes et
64 al., 1999, 2000). Therefore, the prevailing acidification profile during processing is expected to
65 affect proteolysis and thus flavour formation. Dry fermented sausages can be roughly divided in
66 two groups with different acidity levels, namely northern-type and southern-type products (Ravyts
67 et al., 2012). In the northern-type, the pH drops below 5.0 during fermentation and stays more or
68 less at that level throughout ripening. On the contrary, in the southern-type, the pH drops only

69 moderately during fermentation and increases during the drying phase, resulting in a final pH
70 between 5.5 and 6.0 (Demeyer et al., 2000).

71 Since the pH drop during meat fermentation not only influences flavour by creating an acid
72 taste but also through proteolysis, it is important to better understand the details of this mechanism.
73 Therefore, the aim of this study was to identify the peptides originated from actin degradation,
74 which is one of the most abundant proteins in meat, during ripening of dry fermented sausages at
75 two different pH values.

76

77 **2. MATERIAL AND METHODS**

78

79 *2.1. Dry fermented sausage preparation*

80 Dry fermented sausages were prepared by mixing lean pork (70.5 %), pork backfat (27.0 %),
81 sodium chloride (2.5 %), sodium nitrate (0.015 %, m/m), sodium ascorbate (0.05 %, m/m), and a
82 starter culture strain (*Lactobacillus sakei* CTC 494) originating from the culture collection of the
83 Research Group of Industrial Microbiology and Food Biotechnology (Vrije Universiteit Brussel,
84 Brussels, Belgium). Two pH treatments were installed by adding either 0.25 %, m/m (high-pH
85 treatment) or 0.7 %, m/m (low-pH treatment) of dextrose. The batter was stuffed into collagen
86 casings of 50 mm diameter (Naturin, Weinheim, Germany) and ripened for 28 days in a climate
87 chamber. During the first four days, fermentation was performed at a temperature of 24 °C and a
88 relative humidity of 94 %. For the drying process, the temperature was dropped to 12 °C and
89 relative humidity was set at 94 % for the first 14 days and at 80 % for the last 10 days. Samples
90 were taken at days 0, 4 (end of fermentation), and 28 (end of ripening). The manufacturing
91 processes and sampling were repeated in triplicate, resulting in three independent replicates.

92

93 *2.2 pH and weight loss*

94 In each manufacturing process, three randomly selected sausages per treatment were weighed and
95 the pH was recorded after their preparation and during ripening. The pH was measured directly in
96 the sausages [ISO 2917 (1999)] and the pH meter was calibrated in buffers of pH 4.0 and 7.0.
97 Weight loss was expressed as a percentage of the initial weight and the mean of the three records
98 was calculated.

99

100 *2.3 Peptide extraction*

101 The peptide extraction was carried out as described by Mora et al. (2015). Briefly, peptides were
102 extracted in 0.01 N HCl and proteins were precipitated by addition of EtOH. Finally, the peptide
103 extract was dried in a rotary evaporator and the peptides were dissolved in 25 ml 0.01 N HCl.

104

105 *2.4 Size-exclusion chromatography*

106 Size-exclusion chromatography was carried out to select peptides between 500 and 4000 Da.
107 Peptides were fractionated using an Akta Purifier (GE Healthcare Life Sciences, Uppsala, Sweden)
108 on a Sephadex G25 Fine column (2.6 × 60 cm) and 0.01 N HCl was used as mobile phase at a flow
109 rate of 1 mL/min. The fraction corresponding to elution volumes from 80 to 220 mL was collected
110 and aliquots of 100 µL were lyophilised.

111

112 *2.5 LCMS^E analysis*

113 Lyophilised peptides were suspended in 40 µl of 100 mM NH₄HCO₂ (pH 10). Samples were
114 filtered (0.22 µm) and centrifuged. The peptide identification was done according to Devos et al.
115 (2015) by liquid chromatography coupled to mass spectrometry in a data-independent positive
116 mode of acquisition (LCMS^E). Briefly, 2 µl sample was injected on a NanoAcquity UPLC[®] system
117 (Waters Corporation, Milford, MA, USA) for peptide separation. Solvent A1 and B1 were
118 composed of 20 mM ammonium formate in water and acetonitrile (pH 10), respectively. Solvent A2

119 and B2 were composed of 0.1 % formic acid in water and 0.1 % formic acid in acetonitrile,
120 respectively. The sample was loaded onto an Xbridge™ BEH130 C18 column (300 μm × 50 mm, 5
121 μm; Waters) at 50 % solvent B1 at 2 μL/min. Peptides were eluted and trapped on a Symmetry®
122 C18 trapping column (180 μm × 20 mm, 5 μm; Waters Corporation) and finally separated on a HSS
123 T3 C18 analytical column (75 μm × 250 mm, 1.8 μm; Waters Corporation) at 40°C at 250 nL/min
124 by increasing the acetonitrile concentration from 5 to 50 % B2 over 60 min.

125 The outlet of the column was directly connected to a PicoTip Emitter (uncoated SilicaTip™
126 10 ± 1 μm, New Objective, Woburn, MA, US) mounted on a Nanolockspray source of a SYNAPT™
127 G1 HDMS mass spectrometer (Waters Corporation). Accurate mass data were collected by
128 alternating between low (5 V) and high (ramping from 15 to 35 V) energy scan functions
129 (Geromanos et al., 2009). The selected m/z range was 125–2000 Da. The capillary voltage was set
130 to 3.0 kV, the sampling cone voltage was 26 V, and the extraction cone voltage was 2.65 V. The
131 source temperature was set at 80°C.

132 The LCMS^E data were processed using the ProteinLynx Global SERVER™ v2.5 (PLGS,
133 Waters Corporation) (Geromanos et al., 2009). The identification of actin peptides was done by
134 using a Uniprot database containing 753 protein entries from actin of the organism *Sus scrofa*
135 (downloaded from the Uniprot website, March 2016). The primary digest reagent was set as “none”.
136 The precursor and fragment ion tolerance were determined automatically. The default protein
137 identification criteria used included a detection of minimal three fragment ions per peptide, and
138 minimal three fragment ions per protein. Methionine oxidation was selected as a variable
139 modification. A false positive rate of 4 % was allowed.

140

141 2.6 Identification of peptides cleaved by cathepsins B and D

142 To identify the actin-derived peptides that were likely generated by the action of cathepsins B and D
143 during meat fermentation, the previously determined cleavage sites of these enzymes on bovine F-
144 actin, were used (Hughes et al., 1999, 2000).

145

146 **3. RESULTS**

147

148 *3.1 pH and weight loss*

149 The initial pH in the raw meat was 5.69 and 5.68 for the high-pH and low-pH treatments
150 respectively. During the fermentation phase, the pH decreased to 5.23 (\pm 0.02) and 4.89 (\pm 0.01) in
151 the high-pH and low-pH treatments, respectively. In the high-pH treatment, the pH increased
152 throughout the drying phase to a final pH of 5.98 (\pm 0.07) on day 28. On the contrary, in the low-pH
153 treatment, the pH decreased to a minimum of 4.76 (\pm 0.01) on day 15 and subsequently increased to
154 a final pH of 4.96 (\pm 0.01) at the end of ripening on day 28.

155 Weight loss showed similar trends in both treatments. Indeed, all samples lost about 2 % and 30 %
156 of the initial weight at the end of fermentation and ripening, respectively.

157

158 *3.2 Peptides identified from actin degradation*

159 The extracted peptides from the raw meat and the sausages at the end of fermentation and ripening
160 were subjected to identification by LCMS^E. In the raw meat, no peptides arising from actin were
161 found. At the end of fermentation, however, 52 and 42 peptides were identified in the high-pH and
162 low-pH sausages, respectively (Tables 1 and 2). At this stage, both sausage types showed 37
163 peptides in common (highlighted in bold in the tables). At the end of ripening, the number of
164 peptides identified increased to 66 in the high-pH sausages (Table 3). In the low-pH sausages, 144
165 peptides were identified, of which 47 peptides were also found in the high-pH variant (Table 4).
166 The peptides identified covered more than 50 % of the entire actin sequence and were generated
167 from all four actin subdomains, in which nine actin regions containing the majority of the identified
168 peptides are highlighted in Figure 1. Most peptides were identified in the regions 1 and 9, which are
169 near the N- and C-terminus and are located in the first subdomain together with the regions 3, 4 and
170 5 (region 1 is also partially lying in the second subdomain). Regions 2 and 7 were located in the

171 second and fourth subdomain, respectively. Finally, in the third subdomain, peptides were mainly
172 released from regions 6 and 8. On average, more peptides were identified at the end of ripening.
173 However, in the regions 6 and 9, similar amounts of peptides were identified at both days.

174 The identified peptides were mapped on to the 3D actin structure (Figure 2). Although after
175 the fermentation phase the number of peptides identified in the two pH treatments was similar,
176 some peptides (in red) were identified only at high pH (Figure 2A). On the contrary, at the end of
177 ripening, the higher number of peptides identified in the low-pH compared to the high-pH treatment
178 is reflected in the higher number of blue zones in the 3D structure (Figure 2B). However, the
179 majority of peptides were identified in both treatments and are highlighted in yellow.

180 Based on the known specific cleavage sites of cathepsins B and D (Figure 1), an analysis of
181 the proteolytic breakdown by the latter enzymes was done. At the end of fermentation, 19 and 17
182 peptides (on a total of 52 and 42) were cleaved off either from the N or C-terminus, at high and low
183 pH, respectively (Tables 1 and 2). Throughout ripening, cleavage site analysis indicated that
184 cathepsins B and D remained active leading to 29 and 67 identified peptides (on a total of 66 and
185 144) at high and low pH, respectively. The peptides Thr₁₀₈-Lys₁₂₀ and Ile₃₅₉-Lys₃₇₅ that were present
186 in the low-pH samples were likely generated by cathepsin D. In addition, cathepsin D was probably
187 responsible for the following cleavage sites: Cys₁₂-Asp₁₃, Thr₇₉-Asn₈₀, Thr₉₁-Phe₉₂, Ile₁₅₃-Val₁₅₄,
188 Leu₁₅₅-Asp₁₅₆, Lys₂₄₀-Ser₂₄₁, Trp₃₅₈-Ile₃₅₉, Thr₉₁-Phe₉₂, His₁₀₃-Pro₁₀₄, and Lys₁₂₀-Met₁₂₁. According
189 to the known cleavage sites for cathepsin B, this enzyme likely cleaved actin at the following sites:
190 Ala₂₄-Gly₂₅, Thr₆₈-Leu₆₉, Met₄₉-Gly₅₀, His₇₅-Gly₇₆, Gly₇₆-Ile₇₇, Lys₈₆-Ile₈₇, Thr₁₀₅-Leu₁₀₆, Thr₂₅₁-
191 Ile₂₅₂, Lys₃₃₀-Ile₃₃₁, and Glu₃₆₃-Tyr₃₆₄. Because of the occurrence of some common cleavage sites by
192 both cathepsins, it was not possible to denote which of both enzymes mainly cleaved at the sites
193 Gly₂₂-Phe₂₃, Arg₃₀-Ala₃₁, Phe₉₂-Tyr₉₃, Arg₉₇-Val₉₈, and Leu₁₀₇-Thr₁₀₈.

194 A lot of identified peptides were truncated differing only by a single amino acid. In
195 particular for the peptides Ile₁₅₃-Gly₁₇₀, Tyr₂₄₂-Arg₂₅₈, and Trp₃₅₈-His₃₇₃, single amino acids were
196 cleaved after already four days of fermentation and smaller peptides were generated in both high-

197 pH and low-pH sausages. In addition to the peptides reported above for the fermentation phase, the
198 peptides Met₄₉-Gly₆₅ and Ser₂₃₇-Arg₂₅₆ (at high and low pH) as well as Asp₁₃-Ala₃₁ and Phe₂₃-Gly₄₄
199 (at low pH only) were intensively hydrolysed at the end of the ripening phase. Similarly, several
200 peptides were truncated differing by two or three amino acids, indicating the release of dipeptides
201 and tripeptides. In the final products, the dipeptides Thr-Lys, Tyr-Ala, and Met-Gln were released
202 in the high-pH sausages. The dipeptides and tripeptides Gln-Gly, Arg-His, Val-Phe, Met-Gln, Gly-
203 Ser-Gly, Leu-Tyr-Arg, Ile-Val-Gly, Ile-Val-His, and Ile-Leu-Thr were released in the low-pH
204 sausages.

205

206 **4. DISCUSSION**

207

208 Since actin is one of the most abundant proteins in muscle (Dominguez and Holmes, 2011), its
209 degradation during ripening of dry fermented sausages is of interest for flavour formation.
210 Breakdown of actin during ripening has been demonstrated previously, usually via electrophoresis
211 techniques (Hughes et al., 2002). Yet, more detailed information on its degradation patterns requires
212 state-of-the-art analysis via proteomic analysis. In the present paper, the small peptides generated
213 from actin hydrolysis were identified by LCMS^E. The results showed that a high degree of
214 proteolysis took place since the identified peptides covered more than 50 % of the actin sequence
215 and peptides were identified from all four subdomains. In a recent study, Lopez et al. (2015)
216 identified some peptides arising from actin in fermented sausage models. Several peptides identified
217 from that group originated from the regions 1, 3, and 7 that were also highlighted in the present
218 study, confirming that those regions are susceptible to proteolysis.

219 It is evident that actin proteolysis started already during the fermentation phase: whereas no
220 actin peptides were found in the raw meat, a more or less comparable peptide generation became
221 already clear after four days of fermentation for both types of sausages. After 28 days of ripening,
222 however, the number of peptides identified was more than twofold higher in the low-pH sausages.

223 In contrast to the rather brief fermentation step, ripening lasted for a longer time and led to more
224 pronounced pH differences (pH 4.9 *versus* pH 6.0 at the end of ripening). Demeyer et al. (2000)
225 also reported lower actin degradation in sausages with higher pH and this might be due to the low
226 optimum pH of cathepsin D like muscle enzymes, which play a major role in actin hydrolysis
227 (Molly et al., 1997). In dry fermented sausages, proteolysis is generally divided in two steps: firstly,
228 endopeptidases break down intact proteins generating small peptides; secondly, the generated
229 peptides are further degraded by exopeptidases which release single amino acids, dipeptides, and
230 tripeptides. The lysosomal cathepsins B and D are believed to be the main endopeptidases
231 responsible for the first protein breakdown in dry fermented sausages (Molly et al., 1997). The
232 identification of the cleavage sites of cathepsins B and D on bovine actin, which has the same
233 sequence of pig actin, by Hughes et al. (1999, 2000) allowed to understand which peptides were
234 likely generated by these enzymes. Indeed, some of these cleavage sites were also the starting point
235 of several peptides identified at day 4 in the present study, supporting the finding that cathepsins are
236 already likely active during the first days of fermentation (Demeyer et al., 1992). The pH at the end
237 of fermentation was in the activity range of cathepsin B and D in both treatments. Indeed, Schwartz
238 and Bird (1977) reported that rabbit actin was degraded by cathepsins B and D at pH 5.0. These
239 enzymes remain stable and active for several months during processing of dry cured meat products
240 (Toldrà et al., 1993) and it is not surprising that the number of peptides likely generated by
241 cathepsin B and D increased at the end of ripening in the present study. This was particularly
242 evident in the low-pH treatment where the number of identified peptides greatly increased at the end
243 of ripening. In the high-pH sausages, the contribution of these enzymes was more limited during the
244 drying phase since their activity is very low at pH 6.0 (Schwartz and Bird, 1977). Although for
245 some peptides the contribution of cathepsins B and D seems indisputable, this might be doubtful for
246 other peptides. Indeed, in some cases the same peptide but with an additional residue after the
247 cathepsin cleavage site was also identified. At this point, it is impossible to know whether the

248 shorter peptide was the result of endopeptidase activity or if it was generated from the longer
249 peptide by hydrolysis of the additional residue made by an exopeptidase.

250 The further degradation of peptides exerted by exopeptidases releases amino acids,
251 dipeptides, and tripeptides from the N- and C-terminus. These enzymes, in dry fermented sausages,
252 are either of muscle or microbial origin (Mora et al., 2015). In the present study, the activity of
253 aminopeptidases and carboxypeptidases was evident since a lot of identified peptides were
254 truncated differing by a single amino acid. There are five aminopeptidases known to be active post-
255 mortem: leucyl, arginyl, alanyl, pyroglutamyl, and methionyl aminopeptidases (Toldrá, 2006). Ile,
256 Leu, Tyr, and Gly were probably released by alanyl aminopeptidase (Flores et al., 1996). Methionyl
257 aminopeptidase mainly cleaves Lys, Ala and Leu (Flores et al., 2000) and arginyl aminopeptidase
258 cleaves basic amino acids (Flores et al. 1993). These aminopeptidases have optimum activities at
259 neutral pH and only retain some activity at pH 5.0 (Toldrá, 2006). Nevertheless, the action of
260 exopeptidases generated more unique peptides at low pH. We suggest that some endopeptidases like
261 cathepsins, which have optimum activity at very acidic pH, provided more substrates to
262 exopeptidases at low pH. From the C-terminus, the activity of carboxypeptidases was also evident,
263 although knowledge about these enzymes in meat products is still limited. There are two lysosomal
264 carboxypeptidases (A and B) known to have optimal activity at acidic pH, with the former cleaving
265 hydrophobic amino acids and the latter having a wider activity (Toldrá, 2006). Several peptides
266 were also truncated differing by two or three amino acids, indicating that dipeptidyl, such as DPP I
267 and DPP II (Sentandreu and Toldrá, 2000, 2001), and tripeptidyl peptidases were active during
268 ripening.

269 In addition to the above, the contribution of microbial enzymes cannot be ignored. In the
270 present study, all sausages were inoculated with *L. sakei*. This species, which has often been
271 isolated from spontaneously fermented dry fermented sausages, is often used as starter culture for
272 its high competitiveness and ability to produce antimicrobial compounds with strong antilisterial
273 activity (Leroy and De Vuyst, 2005; Ravyts et al., 2012). Moreover, the background microbiota that

274 is usually constituted of coagulase-negative microbiota may also be relevant for proteolytic activity,
275 as well as for the further conversion of amino acids in aroma compounds (Sánchez Mainar et al.,
276 2016). It is generally believed that meat endogenous enzymes initiate proteolysis and the
277 contribution of microbial enzymes is relevant only in a later stage of ripening (Hughes et al., 2002;
278 Molly et al., 1997). The species *L. sakei* has been shown to possess some proteolytic activity
279 (Candogan and Acton, 2004), which was in particular reported to release Leu and Ala from peptides
280 (Sanz and Toldrá, 1997).

281 Peptides, especially the ones with a low molecular mass < 5000 Da, are potential flavour
282 compounds and flavour precursors. The taste of peptides depends on the conformational
283 characteristics and on the amino acid composition. Charges on the side chains or on the terminals
284 can give salty or sour taste, hydrophobic side chains can give bitter taste and umami taste was
285 reported for some small peptides in meat products (Temussi, 2011). Nevertheless, there are only
286 few studies that investigated the influence of peptides on the taste of meat products. Henriksen and
287 Stahnke (1997) evaluated the effect of fractions of small peptides and free amino acids extracted
288 from dry fermented sausages on taste. The results suggested that bitterness and sourness were
289 correlated with hydrophobic amino acids and glutamic acid, respectively, and bouillon taste was
290 provided by a mixture of different peptides and amino acids. The results of the present study
291 revealed that numerous peptides of low molecular weight are generated by actin degradation,
292 especially at low pH conditions. Therefore, actin might be a key protein in determining the final
293 taste characteristics in dry fermented sausages considering the high abundance of this protein.

294

295 **5. CONCLUSION**

296

297 In the present study, the generation of peptides from actin, which is one of the most abundant
298 proteins in muscle, was studied during ripening of dry fermented sausages through LCMS^E. The
299 understanding of proteolysis is of great importance since small peptides and amino acids influence

300 the sensory characteristics of these products. The results showed that actin is already being
301 hydrolysed during the fermentation phase, but proteolysis is most intense during the drying phase.
302 In addition, a more acid pH profile, characteristic of northern-type fermented sausages, generated
303 more unique peptides. The relevance of the findings of the present study will need to be confirmed
304 and contrasted with the analysis of proteolytic products of other major muscle proteins in future
305 research.

306

307 **Acknowledgements**

308 This work was financially supported by the Fund for Scientific Research – Flanders (FWO-
309 Vlaanderen; project G.0327.12) and by the Hercules Foundation (project AUG019). The authors are
310 grateful to Stijn De Waele, Msc and Dr. Gonzalez Vandriessche for support during peptide
311 purification and mass spectrometric analysis, respectively.

312

313 **Conflict of interest**

314 The authors declare that no competing interests exist.

315

316 **REFERENCES**

- 317 Candogan, K., & Acton, C. (2004). Proteolytic activity of bacterial starter cultures for meat
318 fermentation. *Journal of Muscle Foods* 15, 23–34.
- 319 Defernando, G.D.G., & Fox, P.F. (1991). Study of proteolysis during the processing of a dry
320 fermented pork sausage. *Meat Science*, 30(4), 367–383.
- 321 Demeyer, D., Claeys, E., Ötles, S., Caron, L., & Verplaetse, A. (1992). Effect of meat species on
322 proteolysis during dry sausage fermentation. *Proceedings of the 38th International Conference*
323 *on Meat Science and Technology, Clermont-Ferrand, France*, 775.
- 324 Demeyer, D., Raemaekers, M., Rizzo, A., Holck, A., De Smedt, A., ten Brink, B., Hagen, B.,
325 Montel, C., Zanardi, E., Murbrekk, E., Leroy, F., Vandendriessche, F., Lorentsen, K.,

326 Venema, K., Sunesen, L., Stahnke, L., De Vuyst, L., Talon, R., Chizzolini, R., & Eerola, S.
327 (2000). Control of bioflavour and safety in fermented sausages: first results of a European
328 project. *Food Research International*, 33(3-4), 171-180.

329 Devos, S., Van Oudenhove, L., Stremersch, S., Van Putte, W., De Rycke, R., Van Driessche, G.,
330 Vitse, J., Raemdonck, K., Devreese, B. (2015). The effect of imipenem and diffusible
331 signaling factors on the secretion of outer membrane vesicles and associated Ax21 proteins in
332 *Stenotrophomonas maltophilia*. *Frontiers in Microbiology*, 6.

333 Dominguez, R., & Holmes, K.C. (2011). Actin structure and function. *Annual Review of*
334 *Biophysics*, 40, 169–186.

335 Flores, M., Aristoy, M.C., & Toldrá, F. (1993). HPLC purification and characterization of porcine
336 muscle aminopeptidase B. *Biochimie*, 75, 861–867.

337 Flores, M., Aristoy, M.C., & Toldrá, F. (1996). HPLC purification and characterization of soluble
338 alanyl aminopeptidase from porcine skeletal muscle. *Journal of Agricultural and Food*
339 *Chemistry*. 44, 2578–2583.

340 Flores, M., Marina, M., & Toldrá, F. (2000). Purification and characterization of a soluble
341 methionyl aminopetidase from porcine skeletal muscle. *Meat Science*, 56(3), 247–254.

342 Geromanos, S.J., Vissers, J.P., Silva, J.C., Dorschel, C.A., Li G.Z., Gorenstein, M.V., Bateman, R.
343 H., Langridge, J.I. (2009). The detection, correlation, and comparison of peptide precursor
344 and product ions from data independent LC-MS with data dependant LC-MS/MS.
345 *Proteomics*, 9, 1683–1695.

346 Henriksen, A.P., & Stahnke, L.H. (1997). Sensory and chromatographic evaluations of water
347 soluble fractions from dried sausages. *Journal of Agricultural and Food Chemistry*, 45, 2679–
348 2684.

349 Hughes, M.C., O'Neill, E.E., McSweeney, P.L.H., & Healy, Á. (1999). Proteolysis of bovine F-
350 actin by cathepsin B. *Food Chemistry*, 64(4), 525–530.

351 Hughes, M.C., Healy, Á., McSweeney, P.L.H., & O'Neill, E.E. (2000). Proteolytic specificity of
352 cathepsin D on bovine F-actin. *Meat Science*, 56(2), 165–172.

353 Hughes, M.C., Kerry, J.P., Arendt, E.K., Kenneally, P.M., McSweeney, P.L.H., & O'Neill, E.E.
354 (2002). Characterization of proteolysis during the ripening of semi-dry fermented sausages.
355 *Meat Science*, 62(2), 205–216.

356 Leroy, F., & De Vuyst, L. (2005). Simulation of the effect of sausage ingredients and technology on
357 the functionality of the bacteriocin-producing *Lactobacillus sakei* CT 494 strain. *International*
358 *Journal of Food Microbiology*, 100, 141-152.

359 López, C.M., Sentandreu, M.A., Vignolo, G.M., & Fadda, S.G. (2015). Proteomic and peptidomic
360 insights on myofibrillar protein hydrolysis in a sausage model during fermentation with
361 autochthonous starter cultures. *Food Research International*, 78, 41–49.

362 Molly, K., Demeyer, D., Johansson, G., Raemaekers, M., Ghistelinck, M., & Geenen, I. (1997). The
363 importance of meat enzymes in ripening and flavour generation in dry fermented sausages.
364 First results of a European project. *Food Chemistry*, 59(4), 539–545.

365 Mora, L., Gallego, M., Escudero, E., Reig, M., Aristoy, M.-C., & Toldrá, F. (2015). Small peptides
366 hydrolysis in dry-cured meats. *International Journal of Food Microbiology*, 212, 9-15.

367 Ordóñez, J.A., Hierro, E.V., Bruna, J.M., & de la Hoz, L. (1999). Changes in the components of
368 dry-fermented sausages during ripening. *Critical Reviews in Food Science and Nutrition*, 39,
369 329–367.

370 Ravyts, F., De Vuyst, L., & Leroy, F. (2012). Bacterial diversity and functionalities in food
371 fermentations. *Engineering in Life Sciences*, 12(4), 356–367.

372 Sánchez Mainar, M., Stavropoulou, D.A., & Leroy, F. (2016). Exploring the metabolic
373 heterogeneity of coagulase-negative staphylococci to improve the quality and safety of
374 fermented meats: a review. *International Journal of Food Microbiology*, in press
375 (doi:10.1016/j.ijfoodmicro.2016.05.021).

376 Sanz, Y., & Toldrá, F. (1997). Purification and characterization of an aminopeptidase from
377 *Lactobacillus sake*. *Journal of Agricultural and Food Chemistry*, 45, 1552-1558.

378 Schwartz, W., & Bird, J.W.C. (1997). Degradation of myofibrillar proteins by cathepsins B and D.
379 *Biochemical Journal*, 167, 811–820.

380 Sentandreu, M.A., & Toldrá, F. (2000). Purification and biochemical properties of dipeptidyl
381 peptidase I from porcine skeletal muscle. *Journal of Agricultural and Food Chemistry*, 48,
382 5014–5022.

383 Sentandreu, M.A., & Toldrá, F., (2001). Partial purification and characterisation of dipeptidyl
384 peptidase II from porcine skeletal muscle. *Meat Science*, 57(1), 93–103.

385 Spaziani, M., Del Torre, M., & Stecchini, M.L. (2009). Changes of physicochemical,
386 microbiological, and textural properties during ripening of Italian low-acid sausages.
387 Proteolysis, sensory and volatile profiles. *Meat Science*, 81(1), 77–85.

388 Stavropoulou, D.A., Borremans, W., De Vuyst, L., De Smet, S., & Leroy, F. (2015). Amino acid
389 conversions by coagulase-negative staphylococci in a rich medium: assessment of inter-and
390 intraspecies heterogeneity. *International Journal of Food Microbiology*, 212, 34-40.

391 Temussi, P. A. (2012). The good taste of peptides. *Journal of Peptide Science*, 18, 73-82.

392 Toldrá F., Rico E., & Flores J. (1993). Cathepsin B, D, H and L activities in the processing of dry-
393 cured ham. *Journal of the Science of Food and Agriculture*, 62, 157-161.

394 Toldrá, F. (2006). *Biochemistry of Processing Meat and Poultry*. In Y. H. Hui (Ed), *Food*
395 *Biochemistry and Food Processing*, Blackwell Publishing, Ames, Iowa, USA.

396 Verplaetse, A., Demeyer, D., Gerard, S., & Buys, E. (1992). Endogenous and bacterial proteolysis
397 in dry sausage fermentation. *Proceedings of the 38th International Conference on Meat*
398 *Science and Technology*, Clermont-Ferrand, France, 851–854.

399

400 **Figure Captions**

401 **Figure 1.** Actin sequence and 3D structure. The majority of the identified peptides were lying in the
402 underlined regions.

403 **Figure 2.** Peptides were mapped on the 3D structure. Peptides identified for high-pH and low-pH
404 fermentations are shown in red and blue, respectively. Common peptides are shown in yellow.

1 **Table 1.** Peptides of actin identified by LCMS^E in high-pH dry fermented sausages after 4 days of
 2 ripening (fermentation phase)

N.	P _i ^a	Peptide sequence	P ₁ ^b	P. s. s. ^c	z ^d	m/z ^e	Info
1	E	DETTALVCDNGSGLVK	A	4	2	811.3924	
2	C	DNGSGLVKAGFAGDDAPR	A	12	3	582.9503	CD ^f (N-ter), CDB ^h (C-ter)
3	K	AGFAGDDAPR	A	20	2	488.7295	
4	G	FAGDDAPRAVFP SIVG	R	22	2	809.9159	CBD ^h (N-ter)
5	G	FAGDDAPRAVFP S	I	22	2	675.3301	CBD ^h (N-ter)
6	D	SYVGDEAQS ^h KRG	I	53	2	648.816	
7	Y	VGDEAQS ^h KRG	I	55	2	523.7644	
8	F	YNELRVAPEEHPTL	L	92	3	556.6154	CBD ^h (N-ter)
9	F	YNELRVAPEE	H	92	2	610.3024	CBD ^h (N-ter)
10	L	RVAPEEHPTL	L	96	2	574.8065	
11	R	VAPEEHPTL	L	97	2	496.7555	
12	A	PEEHPTL	L	99	2	411.7003	
13	T	GIVLDSGDGVTHNVPIYEG	Y	151	3	647.9892	
14	G	IVLDSGDGVTHNVPIYEG	Y	152	2	942.9739	
15	G	IVLDSGDGVTHNVPIYE	G	152	2	914.4625	
16	I	VLDSGDGVTHNVPIYEG	Y	153	2	886.4331	CD ^f (N-ter)
17	I	VLDSGDGVTHNVPIYE	G	153	2	857.9189	CD ^f (N-ter)
18	I	VLDSGDGVTHNVPIY	E	153	2	793.3958	CD ^f (N-ter)
19	I	VLDSGDGVTHNVPI	I	153	2	655.3221	CD ^f (N-ter)
20	V	LDSGDGVTHNVPIYEG	Y	154	2	836.8954	
21	L	DSGDGVTHNVPIYEG	Y	155	2	780.3534	CD ^f (N-ter)
22	L	DSGDGVTHNVPIYE	G	155	2	751.843	CD ^f (N-ter)
23	L	DSGDGVTHNVPIY	E	155	2	687.3205	CD ^f (N-ter)
24	D	SGDGVTHNVPIYEG	Y	156	2	722.8397	
25	S	DGVTHNVPIYEG	Y	158	2	650.8137	
26	L	EKSYELPDGQVITIGN	E	238	2	881.946	
27	S	YELPDGQVITIGNERFR	C	241	3	669.6759	
28	S	YELPDGQVITIGNERF	R	241	2	925.9726	
29	S	YELPDGQVITIGNER	F	241	2	852.4363	
30	S	YELPDGQVIT	I	241	1	1134.5723	CB ^g (C-ter)
31	Y	ELPDGQVITIGNERF	R	242	2	844.4377	
32	Y	ELPDGQVITIGNER	F	242	2	770.9025	
33	M	SGGTTMYPGIADRMQ	K	301	2	792.862	
34	M	SGGTTMYPGIADR	M	301	2	663.312	
35	S	TFQQMWITKQ ^h EY	D	352	3	534.9266	
36	M	WITKQ^hEYDEAGPSIVH	R	357	3	624.9725	
37	M	WITKQ^hEYDEAGPS	I	357	2	762.3543	
38	W	ITKQ^hEYDEAGPSIVH	R	358	3	562.9473	CD ^f (N-ter)
39	W	ITKQ^hEYDEAGPS	I	358	2	669.3152	CD ^f (N-ter)
40	I	TKQ^hEYDEAGPSIVH	R	359	2	787.3807	
41	K	Q^hEYDEAGPSIVH	R	361	2	672.8068	
42	Q	EYDEAGPSIVHRK	C	362	2	750.8767	
43	Q	EYDEAGPSIVHR	K	362	2	686.8291	
44	Q	EYDEAGPSIVH	R	362	2	608.781	
45	E	YDEAGPSIVHRK	C	363	3	457.9011	CB ^g (N-ter)
46	E	YDEAGPSIVHR	K	363	2	622.3057	CB ^g (N-ter)
47	E	YDEAGPSIVH	R	363	2	544.2557	CB ^g (N-ter)
48	Y	DEAGPSIVHRK	C	364	2	604.824	
49	Y	DEAGPSIVHR	K	364	2	540.7746	
50	Y	DEAGPSIVH	R	364	2	462.7249	
51	E	AGPSIVHR	K	366	2	418.7418	
52	E	AGPSIVH	R	366	1	680.3765	

3 ^a Position of the amino acid residue preceding the peptide sequence (N-terminus).

4 ^b Position of the amino acid residue following the peptide sequence (C-terminus).

5 ^c Peptide sequence start in actin.

6 ^d Charge (+).

7 ^e Mass to charge ratio.

8 ^f Cathepsin D cleavage site.

9 ^g Cathepsin B cleavage site.

10 ^h Cathepsin B and D common cleavage site.

11 Peptides that were also found in low pH dry fermented sausages are indicted in bold.

12

13 **Table 2.** Peptides of actin identified by LCMS^E in low-pH dry fermented sausages after 4 days of
 14 ripening (fermentation phase)

N.	P ₁ ^a	Peptide sequence	P ₁ ^b	P. s. s. ^c	z ^d	m/z ^e	Info
1	G	FAGDDAPRAVFP SIVG	R	22	2	809.9204	CBD ^h (N-ter)
2	G	FAGDDAPRAVFP S	I	22	2	675.3318	CBD ^h (N-ter)
3	H	GIITNWDDMEK	I	75	2	661.3103	CB ^g (N-ter), CB ^g (C-ter)
4	F	YNELRVAPEEHPTL	L	92	3	556.6168	CBD ^h (N-ter)
5	F	YNELRVAPEEHPT	L	92	3	518.9217	CBD ^h (N-ter)
6	F	YNELRVAPEE	H	92	2	610.3063	CBD ^h (N-ter)
7	L	RVAPEEHPTL	L	96	2	574.8041	
8	A	PEEHPTL	L	99	2	411.7017	
9	G	IVLDSGDGVTHNVPIYEG	Y	152	2	942.9772	
10	I	VLDSGDGVTHNVPIYEG	Y	153	2	886.4335	CD ^f (N-ter)
11	I	VLDSGDGVTHNVPIYE	G	153	2	857.9228	CD ^f (N-ter)
12	I	VLDSGDGVTHNVPIY	E	153	2	793.3984	CD ^f (N-ter)
13	V	LDSGDGVTHNVPIYEG	Y	154	2	836.8956	
14	L	DSGDGVTHNVPIYEG	Y	155	2	780.3545	CD ^f (N-ter)
15	L	DSGDGVTHNVPIYE	G	155	2	751.8453	CD ^f (N-ter)
16	L	DSGDGVTHNVPIY	E	155	2	687.3256	CD ^f (N-ter)
17	D	SGDGVTHNVPIYEG	Y	156	2	722.8389	
18	G	DGVTHNVPIYEG	Y	158	2	650.8106	
19	G	YALPHAIM	R	170	2	458.2395	
20	S	YELPDGQVITIGNERFR	C	241	3	669.6783	
21	S	YELPDGQVITIGNERF	R	241	2	925.9741	
22	S	YELPDGQVITIGNER	F	241	2	852.4399	
23	Y	ELPDGQVITIGNERF	R	242	2	844.4383	
24	Y	ELPDGQVITIGNER	F	242	2	770.9013	
25	M	WITKQ ^e YDEAGPSIVHRK	C	357	4	540.0309	
26	M	WITKQ^eYDEAGPSIVH	R	357	3	624.9791	
27	M	WITKQ^eYDEAGPS	I	357	2	762.3554	
28	W	ITKQ^eYDEAGPSIVH	R	358	3	562.9483	CD ^f (N-ter)
29	W	ITKQ^eYDEAGPS	I	358	2	669.3148	CD ^f (N-ter)
30	I	TKQ^eYDEAGPSIVH	R	359	2	787.3782	
31	K	Q^eYDEAGPSIVH	R	361	2	672.8107	
32	Q	EYDEAGPSIVHRK	C	362	2	750.8771	
33	Q	EYDEAGPSIVH	R	362	2	599.7724	
34	E	YDEAGPSIVHRK	C	363	3	457.9007	CB ^g (N-ter)
35	E	YDEAGPSIVHR	K	363	2	622.3088	CB ^g (N-ter)
36	E	YDEAGPSIVH	R	363	2	544.2598	CB ^g (N-ter)
37	Y	DEAGPSIVHRK	C	364	2	604.8242	
38	Y	DEAGPSIVHR	K	364	2	540.7753	
39	Y	DEAGPSIVH	R	364	2	462.724	
40	D	EAGPSIVHRK	C	365	2	547.3041	
41	E	AGPSIVHR	K	366	2	418.7422	
42	E	AGPSIVH	R	366	1	680.3748	

15 ^a Position of the amino acid residue preceding the peptide sequence (N-terminus).

16 ^b Position of the amino acid residue following the peptide sequence (C-terminus).

17 ^c Peptide sequence start in actin.

18 ^d Charge (+).

19 ^e Mass to charge ratio.

20 ^f Cathepsin D cleavage site.

21 ^g Cathepsin B cleavage site.

22 ^h Cathepsin B and D common cleavage site.

23 Peptides that were also found in high pH dry fermented sausages are indicated in bold.

24 **Table 3.** Peptides of actin protein identified by LCMS^E in high-pH dry fermented sausages after 28
 25 days of ripening

N.	P ₁ ^a	Peptide sequence	P ₁ ^b	P. s. s. ^c	z ^d	m/z ^e	Info
1	C	DNGSGLVKAGFAGDD	A	12	2	711.8276	CD ^f (N-ter)
2	K	AGFAGDDAPRAVFPISIVG	R	20	2	873.9467	
3	A	GFAGDDAPRAVFPISIVG	R	21	2	838.4287	
4	G	FAGDDAPRAVFPISIVG	R	22	2	809.9159	CBD ^h (N-ter)
5	F	AGDDAPRAVFPISIVGRP	R	23	3	575.6369	
6	F	AGDDAPRAVFPISIVGR	P	23	3	543.2865	
7	F	AGDDAPRAVFPISIVG	R	23	2	736.3813	
8	G	MGQKDSYVGDEAQSKRG	I	48	3	619.2906	
9	G	MGQKDSYVGDEAQ	S	48	2	714.3076	
10	M	GQKDSYVGDEAQSKRG	I	49	3	575.6074	CB ^g (N-ter)
11	M	GQKDSYVGDEAQSK	R	49	2	756.3522	CB ^g (N-ter)
12	M	GQKDSYVGDEAQS	K	49	2	692.3051	CB ^g (N-ter)
13	M	GQKDSYVGDEAQ	S	49	2	648.7893	CB ^g (N-ter)
14	Q	KDSYVGDEAQSKR	G	51	3	494.9124	
15	H	GIITNWDDMEK	I	75	2	661.3104	CB ^g (N-ter), CB ^g (C-ter)
16	G	IITNWDDMEK	I	76	2	632.7996	CB ^g (N-ter), CB ^g (C-ter)
17	I	ITNWDDMEK	I	77	2	576.2543	CB ^g (C-ter)
18	I	TNWDDMEKIWHHT	F	78	3	571.587	CD ^f (N-ter), CD ^f (C-ter)
19	W	DMEKIWHHT	F	81	2	656.292	CD ^f (C-ter)
20	F	YNELRVAPEEHPTL	L	92	3	556.614	CBD ^h (N-ter)
21	F	YNELRVAPEEHPT	L	92	3	518.9213	CBD ^h (N-ter), CB ^g (C-ter)
22	F	YNELRVAPEE	H	92	2	610.3028	
23	Y	NELRVAPEEHPT	L	93	2	696.3501	CB ^g (C-ter)
24	L	RVAPEEHPTL	L	96	2	574.8061	
25	R	VAPEEHPTLL	T	97	2	553.2978	CBD ^h (N-ter), CBD ^h (C-ter)
26	L	YASGRITGIVLDSGDGVTHNVPIYEG	Y	144	3	893.4371	CD ^f (N-ter)
27	T	GIVLDSGDGVTHNVPIYEG	Y	151	2	971.4831	
28	G	IVLDSGDGVTHNVPIYEG	Y	152	2	942.9721	
29	I	VLDSGDGVTHNVPIYEGYA	L	153	2	1003.4841	CD ^f (N-ter)
30	I	VLDSGDGVTHNVPIYEG	Y	153	2	886.4301	CD ^f (N-ter)
31	I	VLDSGDGVTHNVPIYE	G	153	2	857.9226	CD ^f (N-ter)
32	I	VLDSGDGVTHNVPI	I	153	2	655.323	CD ^f (N-ter)
33	V	LDSGDGVTHNVPIYEG	Y	154	2	836.8967	
34	L	DSGDGVTHNVPIYEG	Y	155	2	780.3542	CD ^f (N-ter)
35	L	DSGDGVTHNVPI	I	155	2	549.2497	CD ^f (N-ter)
36	D	SGDGVTHNVPIYEG	Y	156	2	722.843	
37	S	GDGVTHNVPIYEG	Y	157	2	679.3209	
38	G	DGVTHNVPIYEG	Y	158	2	650.8119	
39	S	SLEKSYELPDGQVITIGNER	F	236	3	750.0474	
40	S	SLEKSYELPDGQVIT	I	236	2	839.9317	CB ^g (C-ter)
41	L	EKSYELPDGQVITIGNER	F	238	3	683.345	
42	L	EKSYELPDGQVITIGN	E	238	2	881.951	
43	E	KSYELPDGQVITIGNER	F	239	2	959.9952	
44	E	KSYELPDGQVITIGN	E	239	2	817.4255	
45	K	SYELPDGQVITIGNERF	R	240	2	969.4918	CD ^f (N-ter)
46	K	SYELPDGQVITIGNER	F	240	2	895.9536	CD ^f (N-ter)
47	Y	ELPDGQVITIGNERF	R	242	2	844.4362	
48	Y	ELPDGQVITIGNER	F	242	2	770.905	
49	I	GMESAGIHETTYNS	I	269	2	748.8201	
50	R	KDLYANNVMSGGTTM	Y	292	2	801.3673	
51	M	SGGTTMYPGIADRMQ	K	301	2	792.8641	
52	M	SGGTTMYPGIADR	M	301	2	663.3132	
53	M	SGGTTMYPGIAD	R	301	1	1169.5151	
54	W	ITKQEYDEAGPSIVH	R	358	3	562.9504	CD ^f (N-ter)
55	W	ITKQEYDEAGPS	I	358	2	669.3137	CD ^f (N-ter)
56	I	TKQEYDEAGPSIVHRK	C	359	3	619.9816	
57	I	TKQEYDEAGPSIVHR	K	359	3	577.2846	
58	I	TKQEYDEAGPSIVH	R	359	2	787.3806	
59	I	TKQEYDEAGPS	I	359	2	612.7735	
60	K	QEYDEAGPSIVH	R	361	2	672.808	
61	Q	EYDEAGPSIVHR	K	362	3	452.213	
62	Q	EYDEAGPSIVH	R	362	2	608.7801	
63	E	YDEAGPSIVHR	K	363	2	622.3092	CB ^g (N-ter)
64	E	YDEAGPSIVH	R	363	2	544.2584	CB ^g (N-ter)

65	Y	DEAGPSIVHR	K	364	2	540.7759
66	E	AGPSIVHR	K	366	2	418.7395

26 ^a Position of the amino acid residue preceding the peptide sequence (N-terminus).

27 ^b Position of the amino acid residue following the peptide sequence (C-terminus).

28 ^c Peptide sequence start in actin.

29 ^d Charge (+).

30 ^e Mass to charge ratio.

31 ^f Cathepsin D cleavage site.

32 ^g Cathepsin B cleavage site.

33 Peptides that were also found in low pH dry fermented sausages after 28 days of ripening are indicted in bold.

34 **Table 4.** Peptides of actin protein identified by LCMS^E in low-pH dry fermented sausages after 28
 35 days of ripening

N.	P ₁ ^a	Peptide sequence	P ₁ ^b	P. s. s. ^c	z ^d	m/z ^e	Info
1	C	DNGSGLVKAGFAGDDAPRA	V	12	3	606.6272	CD ^f (N-ter)
2	C	DNGSGLVKAGFAGDDAPR	A	12	3	582.952	CD ^f (N-ter), CBD ^h (C-ter)
3	D	NGSGLVKAGFAGDDAPRAVFPISIVG	R	13	3	801.4185	
4	D	NGSGLVKAGFAGDDAPRA	V	13	3	568.287	
5	D	NGSGLVKAGFAGDDAPR	A	13	3	544.6087	CBD ^h (C-ter)
6	N	GSGLVKAGFAGDDAPRA	V	14	3	530.2726	
7	N	GSGLVKAGFAGDDAPR	A	14	3	500.5875	CBD ^h (C-ter)
8	G	LVKAGFAGDDAPRA	V	17	3	463.2476	
9	G	LVKAGFAGDDAPR	A	17	2	658.8565	CBD ^h (C-ter)
10	L	VKAGFAGDDAPR	A	18	2	602.3102	CBD ^h (C-ter)
11	V	KAGFAGDDAPRA	V	19	2	588.2925	
12	K	AGFAGDDAPRAVFPIS	I	20	2	739.3563	
13	K	AGFAGDDAPRA	V	20	2	524.2473	
14	K	AGFAGDDAPR	A	20	2	488.7296	CBD ^h (C-ter)
15	A	GFAGDDAPRAVFPIS	I	21	2	703.8424	
16	A	GFAGDDAPRA	V	21	2	488.7285	
17	G	FAGDDAPRAVFPISIVGRPRHQG	V	22	4	588.3061	CBD ^h (N-ter)
18	G	FAGDDAPRAVFPISIVGRPRH	Q	22	3	722.3819	CBD ^h (N-ter)
19	G	FAGDDAPRAVFPISIVGRP	R	22	3	624.6606	CBD ^h (N-ter)
20	G	FAGDDAPRAVFPISIVGR	P	22	3	592.3103	CBD ^h (N-ter)
21	G	FAGDDAPRAVFPISIVG	R	22	2	809.9169	CBD ^h (N-ter)
22	G	FAGDDAPRAVFPIS	I	22	2	675.34	CBD ^h (N-ter)
23	G	FAGDDAPRAVFP	S	22	2	631.8156	CBD ^h (N-ter)
24	G	FAGDDAPRAVFP	P	22	2	583.2892	CBD ^h (N-ter)
25	F	AGDDAPRAVFPISIVGRP	R	23	3	575.6375	
26	F	AGDDAPRAVFPISIVGR	P	23	3	543.2854	
27	F	AGDDAPRAVFPISIVG	R	23	2	736.3833	
28	A	GDDAPRA	V	24	1	683.3168	CB ^g (N-ter)
29	D	DAPRAVFPIS	I	26	2	480.251	
30	A	VFPISIVGRPRHQG	V	31	3	483.9377	
31	F	PSIVGRPRHQG	V	33	2	602.3381	
32	R	HQGVVMVGMGQK	D	41	2	586.286	
33	G	VMVGMGQKDS	Y	44	2	526.2501	
34	V	GMGQKDSYVGDEAQSQRG	I	47	3	638.2976	
35	G	MGQKDSYVGDEAQSQRGILT	L	48	3	728.3632	CB ^g (C-ter)
36	G	MGQKDSYVGDEAQSQRG	I	48	3	619.2863	
37	G	MGQKDSYVGDEAQSQR	G	48	3	600.2819	
38	G	MGQKDSYVGDEAQSQR	R	48	3	548.2491	
39	G	MGQKDSYVGDEAQ	S	48	2	714.3086	
40	M	GQKDSYVGDEAQSQRG	I	49	3	575.6082	CB ^g (N-ter)
41	M	GQKDSYVGDEAQSQR	G	49	3	556.5999	CB ^g (N-ter)
42	M	GQKDSYVGDEAQSQR	R	49	2	756.3502	CB ^g (N-ter)
43	D	SYVGDEAQSQR	G	53	2	620.3017	
44	S	YVGDEAQSQRGILT	L	54	2	768.9041	CB ^g (C-ter)
45	S	YVGDEAQSQRG	I	54	2	605.297	
46	S	YVGDEAQSQR	G	54	2	576.7844	
47	Y	VGDEAQSQRG	I	55	2	523.7643	
48	V	GDEAQSQRGILT	L	56	2	637.842	CB ^g (C-ter)
49	L	TLKYPIE	H	67	2	432.2482	
50	H	GIITNWDDMEK	I	75	2	661.3095	CB ^g (N-ter), CB ^g (C-ter)
51	G	IITNWDDMEKIWHHTF	Y	76	3	695.999	CB ^g (N-ter), CBD ^h (C-ter)
52	G	IITNWDDMEKIWHHT	F	76	3	646.9729	CB ^g (N-ter), CD ^f (C-ter)
53	G	IITNWDDMEK	I	76	2	632.8001	CB ^g (N-ter), CB ^g (C-ter)
54	I	ITNWDDMEKIWHHTF	Y	77	3	658.303	CBD ^h (C-ter), CD ^f (C-ter)
55	I	ITNWDDMEKIWHHT	T	77	3	575.5944	
56	I	ITNWDDMEK	I	77	2	576.2577	CB ^g (C-ter)
57	I	TNWDDMEKIWHHTF	Y	78	3	620.6089	CD ^f (N-ter), CBD ^h (C-ter)
58	I	TNWDDMEKIWHHT	F	78	3	571.5958	CD ^f (N-ter), CD ^f (C-ter)
59	I	TNWDDMEKIWHHT	T	78	3	537.9025	CD ^f (N-ter)
60	I	TNWDDMEK	I	78	2	519.7155	CD ^f (N-ter), CB ^g (C-ter)
61	T	NWDDMEKIWHHT	F	79	3	537.9053	CD ^f (C-ter)
62	N	WDDMEKIWHHTF	Y	80	3	548.9097	CBD ^h (C-ter)
63	N	WDDMEKIWHHT	T	80	3	466.203	

64	W	DDMEKIWHHT	F	81	2	656.2975	CD ^f (C-ter)
65	F	YNELRVAPEEHPTL	L	92	3	556.6167	CBD ^h (N-ter)
66	F	YNELRVAPEEHPT	L	92	3	518.9219	CBD ^h (N-ter)
67	F	YNELRVAPEE	H	92	2	610.3025	CBD ^h (N-ter)
68	E	LRVAPEEHPTL	L	95	2	631.3515	
69	E	LRVAPEEH	P	95	2	475.7531	CD ^f (C-ter)
70	L	RVAPEEHPTL	L	96	2	574.805	
71	L	RVAPEEHPT	L	96	2	518.2621	CB ^g (C-ter)
72	R	VAPEEHPTLL	T	97	2	553.2994	CBD ^h (N-ter), CBD ^h (C-ter)
73	R	VAPEEHPT	L	97	2	440.2114	CBD ^h (N-ter), CB ^g (C-ter)
74	A	PEEHPTL	L	99	2	411.7021	
75	L	LTEAPLNPKAN	R	106	2	584.3263	
76	L	TEAPLNPKANREKM	T	107	3	533.6141	CBD ^h (N-ter)
77	L	TEAPLNPKANREK	M	107	3	489.9309	CBD ^h (N-ter), CD ^f (C-ter)
78	L	TEAPLNPKANRE	K	107	2	670.3511	CBD ^h (N-ter)
79	L	TEAPLNPKAN	R	107	2	527.7844	CBD ^h (N-ter)
80	G	IVLDSGDGVTHNVPIYEG	Y	152	2	942.9775	
81	G	IVLDSGDGVTHNVPIYE	G	152	2	914.462	
82	I	VLDSGDGVTHNVPIYEG	Y	153	2	886.4333	CD ^f (N-ter)
83	I	VLDSGDGVTHNVPIYE	G	153	2	857.9201	CD ^f (N-ter)
84	I	VLDSGDGVTHNVPIY	E	153	2	793.3949	CD ^f (N-ter)
85	I	VLDSGDGVTHNVPIY	I	153	2	655.3256	CD ^f (N-ter)
86	L	DSGDGVTHNVPIYEG	Y	155	2	780.3532	CD ^f (N-ter)
87	L	DSGDGVTHNVPIYE	G	155	2	751.8449	CD ^f (N-ter)
88	L	DSGDGVTHNVPIY	E	155	2	687.3255	CD ^f (N-ter)
89	L	DSGDGVTHNVPIY	I	155	2	549.2509	CD ^f (N-ter)
90	D	SGDGVTHNVPIYEG	Y	156	2	722.8416	
91	D	SGDGVTHNVPIY	I	156	2	491.7312	
92	S	GDGVTHNVPIYEG	Y	157	2	679.3201	
93	G	DGVTHNVPIYEG	Y	158	2	650.814	
94	Y	ALPHAIMRL	D	171	2	511.2992	
95	T	AASSSSLEKSYELPDGQVITIGN	E	231	2	1183.5982	
96	A	AASSSSLEKSYELPDGQVIT	I	232	2	1005.9979	CB ^g (C-ter)
97	S	SLEKSYELPDGQVITIGNERF	R	236	3	799.0719	
98	S	SLEKSYELPDGQVITIGN	E	236	2	982.0108	
99	S	SLEKSYELPDGQVIT	I	236	2	839.9343	CB ^g (C-ter)
100	S	SLEKSYELPDGQVI	T	236	2	789.4068	
101	S	LEKSYELPDGQVITIGNER	F	237	3	721.0399	
102	S	LEKSYELPDGQVIT	I	237	2	796.4186	CB ^g (C-ter)
103	L	EKSYELPDGQVITIGN	E	238	2	881.9505	
104	L	EKSYELPDGQVIT	I	238	2	739.8722	
105	K	SYELPDGQVITIGNERF	R	240	2	969.4866	CD ^f (N-ter)
106	K	SYELPDGQVITIGNER	F	240	2	895.9547	CD ^f (N-ter)
107	S	YELPDGQVITIGNERFR	C	241	3	669.6787	
108	S	YELPDGQVITIGNERF	R	241	2	925.9733	
109	S	YELPDGQVITIGNER	F	241	2	852.4358	
110	S	YELPDGQVIT	I	241	1	1134.5734	
111	Y	ELPDGQVITIGNERF	R	242	2	844.4392	
112	Y	ELPDGQVITIGNER	F	242	2	770.9043	
113	Y	ELPDGQVI	T	242	1	870.4615	
114	E	TLFQPSF	I	261	1	839.3847	
115	Y	NSIMKCDIDIRK	D	281	2	718.3576	
116	D	LYANNVMSSGGTTMYPGIADRMQKE	I	294	3	883.0785	
117	A	NNVMSSGGTTMYPGIADRMQKE	I	297	3	767.3495	
118	A	NNVMSSGGTTMYPGIADR	M	297	2	892.4089	
119	M	SGGTTMYPGIADRMQKE	I	301	3	614.62	
120	M	SGGTTMYPGIADRMQK	E	301	3	571.6075	
121	M	SGGTTMYPGIADRMQ	K	301	2	792.8625	
122	M	SGGTTMYPGIADR	M	301	2	663.3142	
123	M	SGGTTMYPGIAD	R	301	1	1169.5235	
124	K	IIAPPERKYS	V	330	2	587.3361	CB ^g (N-ter)
125	M	WITKQEYDEAGPSIVHRK	C	357	4	540.0306	
126	M	WITKQEYDEAGPSIVHR	K	357	3	677.0105	
127	M	WITKQEYDEAGPS	I	357	2	762.3605	
128	W	ITKQEYDEAGPSIVHRK	C	358	4	493.5112	CD ^f (N-ter)
129	W	ITKQEYDEAGPSIVHR	K	358	3	614.984	CD ^f (N-ter)
130	W	ITKQEYDEAGPSIVH	R	358	3	562.9486	CD ^f (N-ter)
131	W	ITKQEYDEAGPS	I	358	2	669.3159	CD ^f (N-ter)
132	I	TKQEYDEAGPSIVHRK	C	359	3	619.9871	

133	I	TKQEYDEAGPSIVHR	K	359	3	577.2883	
134	I	TKQEYDEAGPSIVH	R	359	2	787.38	
135	I	TKQEYDEAGPS	I	359	2	612.7782	
136	Q	EYDEAGPSIVHRK	C	362	2	750.8781	
137	Q	EYDEAGPSIVHR	K	362	2	686.8298	
138	Q	EYDEAGPSIVH	R	362	2	608.7845	
139	E	YDEAGPSIVHRK	C	363	2	686.3556	CB ^g (N-ter)
140	E	YDEAGPSIVHR	K	363	2	622.3098	CB ^g (N-ter)
141	E	YDEAGPSIVH	R	363	2	544.2598	CB ^g (N-ter)
142	Y	DEAGPSIVHRK	C	364	2	604.8217	
143	Y	DEAGPSIVHR	K	364	2	540.7767	
144	E	AGPSIVH	R	366	1	680.3776	

36 ^a Position of the amino acid residue preceding the peptide sequence (N-terminus).

37 ^b Position of the amino acid residue following the peptide sequence (C-terminus).

38 ^c Peptide sequence start in actin.

39 ^d Charge (+).

40 ^e Mass to charge ratio.


41 ^f Cathepsin D cleavage site.

42 ^g Cathepsin B cleavage site.

43 Peptides that were also found in high pH dry fermented sausages after 28 days of ripening are indicated in bold.

1 **Figure 1.**

2




3

4 * Cleavage sites of cathepsin D according to Hughes et al. (2000)

5 # Cleavage sites of cathepsin B according to Hughes et al. (1999)

6

7 **Figure 2.**



8