



**Lancaster University Management School**  
**Working Paper**  
**2006/049**

**Empirical pricing kernels obtained from the UK index  
options market**

Xiaoquan Liu, Mark Shackleton, Stephen Taylor and  
Xinzhong Xu

The Department of Accounting and Finance  
Lancaster University Management School  
Lancaster LA1 4YX  
UK

© Xiaoquan Liu, Mark Shackleton, Stephen Taylor and Xinzhong Xu  
All rights reserved. Short sections of text, not to exceed  
two paragraphs, may be quoted without explicit permission,  
provided that full acknowledgement is given.

The LUMS Working Papers series can be accessed at <http://www.lums.lancs.ac.uk/publications/>  
LUMS home page: <http://www.lums.lancs.ac.uk/>

# Empirical Pricing Kernels Obtained from the UK Index Options Market

Xiaoquan Liu\*  
Mark B Shackleton†  
Stephen J Taylor‡  
Xinzhong Xu§

October 16, 2006

---

\*Department of Accounting, Finance, and Management, University of Essex, Colchester CO4 3SQ, UK. Email: liux@essex.ac.uk.

†Department of Accounting and Finance, Lancaster University, Lancaster LA1 4YX, UK. Email: m.shackleton@lancaster.ac.uk.

‡Department of Accounting and Finance, Lancaster University, Lancaster LA1 4YX, UK. Email: s.taylor@lancaster.ac.uk.

§Guanghua School of Management, Peking University, Peking and Cardiff Business School, Cardiff University, UK. Email: xux@gsm.pku.edu.cn.

## **Abstract**

Empirical pricing kernels for the UK equity market are derived as the ratio between risk-neutral densities, inferred from FTSE 100 index options, and historical real-world densities, estimated from time series of the index. The kernels thus obtained are almost compatible with a risk averse representative agent, unlike similar estimates for the US market.

*JEL classifications:* G12; G13

*Keywords:* Pricing kernels; Risk-neutral densities; Index options; Risk aversion

# 1 Introduction

The pricing kernel assumes a central role in asset pricing literature, as it succinctly summarizes investors' risk and time preferences. With a correctly identified pricing kernel, asset pricing becomes a straightforward discounting of future payoffs by the kernel.

In this article, we estimate empirical pricing kernels from the options market, as option prices have been shown to contain incremental information in forecasting future volatilities and price distributions compared to the time series of asset prices<sup>1</sup>. We express the empirical pricing kernels as a ratio between risk-neutral densities (RND), inferred from FTSE 100 index options, and historical densities obtained from time series of the index, averaged across time to minimize the impact of measurement errors.

In particular, we assume that risk-neutral densities follow certain distributions. They can be either a mixture of two lognormal densities (MLN), a generalized beta distribution of the second kind (GB2), or a flexible spline function, all of which are easy to estimate and able to capture the stylized facts of negative skewness and excess kurtosis that are associated with index distributions.

Using a sample of 126 months data from July 1993 to December 2003, we find that the average empirical pricing kernel for the UK equity market is generally downward sloping and does not exhibit the puzzling hump shape documented by Jackwerth (2000), Rosenberg and Engle (2002), and Brown and Jackwerth (2002). This work is also related to Brennan et al (2006), which adopts the traditional asset pricing approach by identifying state variables that fully describe the investment opportunities and specifying flexible functional forms for the pricing kernels.

The rest of the paper is organized as follows. Section 2 introduces the risk-

---

<sup>1</sup>A vast literature has documented that option-implied volatility provides better forecasts of future volatilities than realized volatility. Poon and Granger (2003) and Taylor (2005) provide recent survey evidence. In terms of density forecasts, Liu et al (2006) demonstrates that distributions from option prices are better forecasts than those obtained from asset price histories.

neutral and historical densities and the estimation procedures. Section 3 discusses data. Section 4 presents empirical results and describes the empirical pricing kernels for the UK. Finally, Section 5 concludes.

## 2 Risk-neutral and historical densities

### 2.1 Mixture of lognormal densities (MLN)

Following Ritchey (1990) and Melick and Thomas (1997), the risk-neutral density of the asset price when options expiry can be defined as a mixture of lognormal densities. The MLN densities are flexible and easy to estimate, with the possibility of attaching an economic interpretation to the parameters when the component densities are determined by specific states of the world when the options expire. The MLN density function in this study is the following weighted average of two lognormal densities  $g_{LN}$ ,

$$g_{MLN}(x|\theta) = wg_{LN}(x|F_1, \sigma_1, T) + (1 - w)g_{LN}(x|F_2, \sigma_2, T) \quad (1)$$

with

$$g_{LN}(x|F, \sigma, T) = \frac{1}{x\sigma\sqrt{2\pi T}} \exp \left[ -\frac{1}{2} \left( \frac{\log x - (\log F - \sigma^2 T/2)}{\sigma\sqrt{T}} \right)^2 \right]. \quad (2)$$

The parameter vector is  $\theta = (F_1, F_2, \sigma_1, \sigma_2, w)$ , with  $0 \leq w \leq 1$  and  $F_1, F_2, \sigma_1, \sigma_2 > 0$ . The parameters  $F_1, \sigma_1$ , and  $w$  denote the mean, volatility, and weight of the first lognormal density, while  $F_2, \sigma_2$ , and  $1 - w$  are the mean, volatility, and weight of the second lognormal density.

The density is risk-neutral when its expectation equals the current futures price  $F$ , i.e. when  $wF_1 + (1 - w)F_2 = F$ . The theoretical European option pricing formula is then simply the weighted average of two option prices given by the Black (1976) formula, denoted by  $c_B(\cdot)$ ,

$$c(X|\theta, r, T) = wc_B(F_1, T, X, r, \sigma_1) + (1 - w)c_B(F_2, T, X, r, \sigma_2). \quad (3)$$

## 2.2 Generalized beta distribution (GB2)

The generalized beta distribution of the second kind (GB2) was first proposed by Bookstaber and McDonald (1987) and utilized by Anagnou-Basioudis et al (2005). The GB2 density incorporates four positive parameters  $\theta = (a, b, p, q)$  that permits general combinations of the mean, variance, skewness, and kurtosis of a positive random variables. The GB2 density function is defined as,

$$g_{GB2}(x|a, b, p, q) = \frac{ax^{ap-1}}{b^{ap}B(p, q)[1 + (x/b)^a]^{p+q}} \quad (4)$$

with  $B(j, k) = \Gamma(j)\Gamma(k)/\Gamma(j + k)$ . The density is risk-neutral when

$$F = \frac{bB(p + \frac{1}{a}, q - \frac{1}{a})}{B(p, q)}. \quad (5)$$

The parameter  $b$  is seen to be a scale parameter, while the product of  $a$  and  $q$  determines the maximum number of moments and hence the asymptotic shape of the right tail.

The theoretical option pricing formula depends on the cumulative distribution (c.d.f.) function of the GB2 density, denoted  $G_{GB2}$ , which is a function of the c.d.f. of the beta distribution, denoted  $G_\beta$ ,

$$G_{GB2}(x|a, b, p, q) = G_{GB2}((x/b)^a|1, 1, p, q) = G_\beta(y(x, a, b)|p, q) \quad (6)$$

with  $y(x, a, b) = (x/b)^a/(1 + (x/b)^a)$ . If the density is risk-neutral, so that (5) applies, the European call option prices are given by

$$c(X|\theta) = e^{-rT} \int_X^\infty (x - X)g_{GB2}(x|a, b, p, q)dx \quad (7)$$

$$= Fe^{-rT} \left[ 1 - G_\beta(y(X, a, b) | p + \frac{1}{a}, q - \frac{1}{a}) \right] - Xe^{-rT} [1 - G_\beta(y(X, a, b) | p, q)].$$

## 2.3 Flexible densities

To make sure that the parametric densities discussed above are not inferior to more flexible density curves, we also infer risk-neutral densities defined by spline functions, as estimated by Bliss and Panigirtzoglou (2004). We apply their methodology to obtain implied volatilities, denoted by  $\sigma(\Delta|\theta)$ , which are a function of option delta,  $\Delta$ , and a parameter vector  $\theta$ . Numerical methods then give call prices  $c$  as functions of strikes  $X$  and hence define the risk-neutral densities as  $g(X) = e^{rT} \partial^2 c / \partial X^2$ . For options on futures,  $\Delta$  is defined as a function of the Black-Scholes at-the-money volatility  $\sigma_A$

$$\begin{aligned} \Delta(X) &= e^{-rT} N(d_1(X)) & (8) \\ d_1(X) &= \frac{\log(F/X) + \sigma_A^2 T / 2}{\sigma_A \sqrt{T}}. \end{aligned}$$

The spline function  $\sigma(\Delta|\theta)$  is defined over  $0 \leq \Delta \leq \exp(-rT)$ . It is composed of linear pieces and cubic polynomials, defined on intervals determined by knot points  $\Delta_1 < \Delta_2 < \dots < \Delta_N$ . Each cubic is defined over an interval from  $\Delta_i$  to  $\Delta_{i+1}$ , while the function is linear for  $\Delta \leq \Delta_1$  and  $\Delta \geq \Delta_N$ . The coefficients of the lines and cubics are constrained by the requirement that  $\sigma(\Delta|\theta)$  and its first two derivatives are continuous functions. The spline function has  $N$  free parameters [Lange (1998, page 104)] and there is a unique spline with the required properties that passes through a given set of  $N$  points  $(\Delta_i, \sigma_i)$ . Taking the knot points as given, the parameter vector is the corresponding set of implied volatilities  $\theta = (\sigma_1, \dots, \sigma_N)$ .

## 2.4 Estimation of the RND parameters

The RND parameter vector  $\theta$  is estimated once a month with 4-weeks to maturity so that the densities are non-overlapping for each of the three density

functions. For the MLN and GB2 densities,  $\theta$  is obtained by minimizing the following average squared difference between observed market call prices and theoretical option prices:

$$\frac{1}{N} \sum_{i=1}^N (c_{\text{market}}(X_i) - c(X_i|\theta))^2 \quad (9)$$

with

$$c(X_i|\theta) = e^{-rT} \int_{X_i}^{\infty} (x - X_i)g(x|\theta)dx, 1 \leq i \leq N.$$

In these equations,  $N$  is the number of European option prices used for a particular day,  $g(x|\theta)$  is a parametric density function, and  $c(X|\theta)$  is the associated theoretical option pricing formula, given by either equation (3) or equation (7).

The estimates of  $\theta$  for the flexible densities are obtained by minimizing a function that combines two criteria, namely the accuracy and the smoothness of the fitted spline function  $\sigma(\Delta|\theta)$ . From  $N$  market prices, we derive implied volatilities and hence co-ordinates  $(\Delta_i, \sigma_{\text{market}}(X_i))$ . Then for a set of weights  $w_i$ , we select  $\sigma_1, \dots, \sigma_n$  to minimize

$$\eta \sum_{i=1}^m w_i (\sigma_{\text{market}}(X_i) - \sigma_i)^2 + (1 - \eta) \int_{\Delta_1}^{\Delta_M} \sigma''(\Delta|\sigma_1, \dots, \sigma_m)^2 d\Delta. \quad (10)$$

There is a straightforward solution to this optimization problem [Lange (1998, page 111)]. Implementation requires making a subjective choice for the parameter  $\eta$ , that controls the trade-off between accuracy and smoothness. Following Bliss and Panigirtzoglou (2002, 2004), appropriate weights are proportional to option vega, so we use  $w_i = \exp(-d_1(X_i)^2/2)/\sqrt{(2\pi)}$  with  $d_1(X_i)$  given by equation (8).

## 2.5 Historical densities

ARCH models for daily index returns are estimated and simulated to provide historical real-world densities. The simulated ARCH models must accommo-



date the stylized facts documented in the literature, including a time-varying conditional mean, a persistent conditional volatility, and an asymmetric response of volatility to positive and negative returns. We choose the GJR-GARCH(1,1)-MA(1)-M specification, following Glosten, Jagannathan and Runkle (1993) and Engle and Ng (1993). The conditional mean  $\mu_t$  and the conditional variance  $h_t$  of the daily index return  $r_t$  are as follows,

$$\begin{aligned} h_t &= \omega + \beta h_{t-1} + (\alpha_1 + \alpha_2 D_{t-1})(r_{t-1} - \mu_{t-1})^2, \\ \mu_t &= \xi + \lambda(h_{t-1})^{1/2} + \Theta(r_{t-1} - \mu_{t-1}), \\ D_t &= 1 \quad \text{if } r_t \leq \mu_t, \\ D_t &= 0 \quad \text{otherwise.} \end{aligned} \tag{11}$$

Ten years of daily index returns prior to each estimation date  $t_i$  are used to estimate the ARCH parameters  $\theta = (\omega, \beta, \alpha_1, \alpha_2, \xi, \lambda, \Theta)$ , by maximizing the quasi-log-likelihood function which assumes the conditional distributions are normal. These estimates are consistent even when the normality assumption is false [Bollerslev and Wooldridge (1992)].

The parameters obtained from the returns information up to selected times  $t_i$  are used to simulate the ARCH equations for 4-week periods that end on option expiry dates. A large number,  $M$ , of simulations of the final asset level  $S_{T,i}$  are obtained for each month  $i$ . The historical real-world density  $\tilde{g}_i$  is then the smooth function obtained by using the Gaussian kernel with bandwidth  $H = 0.9\psi/\sqrt[5]{M}$  and with  $\psi$  the standard deviation of the simulated final levels. We have set  $M$  equal to 100,000. Our formula for the bandwidth  $H$  is recommended by Silverman (1986, page 48) and used by Rosenberg and Engle (2002).

### 3 Data

The futures and options contracts are written on the FTSE 100 index and they are traded at the London International Financial Futures and Options

Exchange (LIFFE). Futures and options have the same expiry month and share a common expiry time, 10:30 on the third Friday. European options can then be valued by assuming that they are written on the futures contract, and hence spot levels of the index are not needed.

Daily settlement values for futures and options prices with 4-week to maturity are obtained from LIFFE, for 126 consecutive expiry months from July 1993 to December 2003 inclusive. The call and put implieds for the same contract are almost identical, as should be expected from put-call parity. We average the call and put implieds and use them to calculate European call prices for each contract.

On average, 37 exercise prices are available for each month. The exercise prices are always separated by 50 index points. Table 1 provides summary information about the option prices.

There are twelve expiry dates per annum for the options but the futures contracts are traded for only one expiration date each quarter. Synthetic futures prices must be calculated for the remaining eight months. Fair futures prices,  $F$ , are the future value of the current spot prices  $S$  minus the present value of dividends expected during the life of the futures contract

$$F = e^{rT}(S - PV(\text{dividends})). \quad (12)$$

We have obtained actual dividend payments for the 100 component companies of the index from DataStream, and computed the present value of dividends by assuming that future expected dividends can be approximated by realized dividend payments.

Risk-free interest rates are collected from DataStream. We prefer the London Eurocurrency rate to the UK treasury bill rate, because the Eurocurrency rate is a market rate accessible to AA corporate borrowers.

Figure 1 shows typical estimates of the risk-neutral densities, which exhibit marked negative skewness that has already been extensively documented. The MLN and GB2 densities are almost identical, while the spline density

differs in the left tail. The historical real-world density is very different from the risk-neutral ones, with much less skewness and kurtosis.

## 4 Empirical pricing kernels

Empirical pricing kernels are estimated by using RNDs obtained from the options market and historical real-world densities from index returns. Three empirical pricing kernels  $M(x) = e^{-rT}g(x)/\tilde{g}(x)$  are constructed for each option expiry date, with  $g(x)$  either the MLN, GB2 or spline RND and  $\tilde{g}(x)$  the historical density from GARCH simulations. The geometric mean of each set of kernels is computed, across expiry dates, to reduce the impact of the noise created firstly by fitting the RNDs and secondly by using different sources of information to find  $g$  and  $\tilde{g}$ . We plot the geometric means of the three sets of ratios  $g(yF)/\tilde{g}(yF)$  against the moneyness variable  $y = x/F$  in Figure 2. All three graphed kernels are generally decreasing functions of  $x/F$ , although they are almost flat between 0.98 and 1.05.

None of our empirical pricing kernels for the UK equity market resembles those of Ait-Sahalia and Lo (2000), Jackwerth (2000), Brown and Jackwerth (2002) and Rosenberg and Engle (2002) for the US market. These researchers estimate very clear hump-shaped kernels, using S&P 500 data that ends in 1995, which challenges economic theory and indicates that the representative agent has a risk-seeking utility function in some wealth region. The risk-seeking range obtained by Jackwerth (2000) is  $0.96 \leq x/F \leq 1.01$  while Rosenberg and Engle (2002) obtain  $0.96 \leq x/F \leq 1.02$ . Brennan et al (2006) also use FTSE 100 index options data to estimate the option pricing kernel, approximated by a three-term Chebyshev polynomial whose state variables are the real interest rate, the maximum Sharpe ratio, and volatility. The pricing kernel is upward sloping in the region  $1.03 \leq x/F \leq 1.05$ .

## 5 Conclusion

Using a dataset from July 1993 to December 2003, we derive three series of risk-neutral densities, namely the mixture of two lognormal densities, the generalized beta distribution, and a flexible spline distribution from FTSE 100 index options data. We also fit GARCH models to the time series of FTSE 100 index returns and simulate historical real-world densities. The empirical pricing kernel obtained from the two sets of densities is broadly downward sloping and therefore consistent with economic theories.

## References

- Ait-Sahalia, Y. and A. W. Lo, 2000, Nonparametric risk management and implied risk aversion, *Journal of Econometrics*, 94, 9-51.
- Anagnou-Basioudis, I., M. Bedendo, S. D. Hodges, and R. Tompkins, 2005, Forecasting accuracy of implied and GARCH-based probability density functions, *Review of Futures Markets*, 11, 41-66.
- Black, F., 1976, The pricing of commodity contracts, *Journal of Financial Economics*, 3, 167-179.
- Bliss, R. and N. Panigirtzoglou, 2002, Testing the stability of implied probability density functions, *Journal of Banking and Finance*, 26, 381-422.
- Bliss, R. and N. Panigirtzoglou, 2004, Option-implied risk aversion estimates, *Journal of Finance*, 59, 407-446.
- Bollerslev, T., and J. M. Wooldridge, 1992, Quasi-maximum likelihood estimation and inference in dynamic models with time varying covariances, *Econometric Reviews*, 11, 143-172.
- Bookstaber, R., and J. MacDonald, 1987, A general distribution for describing security price returns, *Journal of Business*, 60, 401-424.
- Brennan, M. J., X. Liu, and Y. Xia, 2006, Option pricing kernels and ICAPM, Unpublished manuscript, UCLA and University of Essex.
- Brown, P., and J. Jackwerth, 2002, The pricing kernel puzzle: reconciling index option data and economic theory, Unpublished manuscript, University of Wisconsin.
- Engle, R. F., and V. K. Ng, 1993, Measuring and testing the impact of news on volatility, *Journal of Finance*, 48, 1749-1778.
- Glosten, L. R., R. Jagannathan, and D. Runkle, 1993, Relationship between the expected value and the volatility of the nominal excess return on stocks, *Journal of Finance*, 48, 1779-1801.
- Jackwerth, J., 2000, Recovering risk aversion from option prices and realized returns, *Review of Financial Studies*, 13, 433-451.

- Lange, K., 1998, *Numerical Analysis for Statisticians*, Springer Verlag.
- Liu, X., M. B. Shackleton, S. J. Taylor, and X. Xu, 2006, Closed-form transformations from risk-neutral to real-world distributions, *Journal of Banking and Finance*, forthcoming.
- Melick, W. R., and C. P. Thomas, 1997, Recovering an asset's implied PDF from option prices: an application to crude oil during the Gulf crisis, *Journal of Financial and Quantitative Analysis*, 32, 91-115.
- Poon, S.-H., and C. W. J. Granger, 2003, Forecasting volatility in financial markets, *Journal of Economic Literature*, 41, 478-539.
- Ritchey, R. J., 1990, Call option valuation for discrete normal mixtures, *Journal of Financial Research*, 13, 285-296.
- Rosenberg, J. and R. F. Engle, 2002, Empirical pricing kernels, *Journal of Financial Economics*, 64, 341-372.
- Silverman, B. W., 1992, *Density Estimation for Statistics and Data Analysis*, Chapman & Hall.
- Taylor, S. J., 2005, *Asset Price Dynamics, Volatility, and Prediction*, Princeton University Press.

Table 1. Summary statistics for the dataset of FTSE 100 index option prices. The average number of option prices is 37 per month for the 126 expiry months from July 1993 to December 2003 inclusive. The moneyness of a call option is defined by  $X/F - 1$ , with  $X$  the exercise and  $F$  the futures price.

Moneyness	No. of options	Percentage (%)
$\leq -0.20$	19	0.4
$(-0.20, -0.10]$	392	8.3
$(-0.10, -0.03]$	849	18.0
$(-0.03, 0.03]$	702	14.9
$(0.03, 0.10]$	731	15.5
$(0.10, 0.20]$	818	17.3
$(0.20, 0.30]$	572	12.1
$(0.30, 0.40]$	303	6.4
$> 0.40$	335	7.1
Totals	4721	100.0

Figure 1. Three risk-neutral densities and the historical density on March 21, 1997






Figure 2. Empirical pricing kernels, as geometric averages across all expiry months

