

# Riemann-Hilbert problem associated with Angelesco systems

A. Branquinho<sup>a,1,\*</sup>, U. Fidalgo<sup>b,2</sup>, and A. Foulquié Moreno<sup>c,3</sup>

<sup>a</sup>*CMUC, Department of Mathematics, University of Coimbra, Largo D. Dinis, 3001-454 Coimbra, Portugal.*

<sup>b</sup>*Departamento de Matemáticas, Universidad Carlos III de Madrid, c/ Universidad 30, 28911 Leganés, Spain.*

<sup>c</sup>*Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal.*

---

## Abstract

Angelesco systems of measures with Jacobi type weights are considered. For such systems, strong asymptotics for the related multiple orthogonal polynomials are found as well as the Szegő-type functions. In the procedure, an approach from Riemann-Hilbert problem plays a fundamental role.

*Key words:* approximation by rational function, rate of convergence, simultaneous approximation,  
*1991 MSC:* 41A20,41A25,41A28

---

## 1 The statement of the Riemann-Hilbert problem

In this work the problem considered is a particular case of the general situation analyzed in [2]. However, due to the simplicity of the case considered, we are able to compute the Szegő-type functions in great detail (cf. (11)).

---

\* Corresponding author: A. Branquinho.

*Email addresses:* `ajplb@mat.uc.pt` (A. Branquinho), `ufidalgo@math.uc3m.es` (U. Fidalgo), `foulquie@ua.pt` (A. Foulquié Moreno).

<sup>1</sup> Research supported by CMUC/FCT.

<sup>2</sup> Research supported by grants MTM 2006-13000-C03-02 from Ministerio de Ciencia y Tecnología and CCG 06-UC3M/ESP-0690 of Universidad Carlos III de Madrid-Comunidad de Madrid and by grant SFRH/BPD/31724/2006 from Fundação para a Ciência e a Tecnologia.

<sup>3</sup> Research supported by UI Matemática e Aplicações from University of Aveiro.

Let  $\Delta_1 = [-\lambda, -1]$  and  $\Delta_2 = [1, \lambda]$  be two intervals on the real line  $\mathbb{R}$ . For each  $j = 1, 2$ , take a holomorphic function  $h_j$ , on a neighborhood  $\mathcal{V}_{h_j}$  of  $\Delta_j$ , i.e.  $h_j \in H(\mathcal{V}_{h_j})$ . We also require that such function  $h_j$  does not vanishes on  $\mathcal{V}_{h_j}$ , acquiring only positive values on  $\Delta_j$ . Observe that  $1/h_j \in H(\mathcal{V}_{h_j})$ ,  $j = 1, 2$ . Let us define the system of measures  $(\sigma_1, \sigma_2)$  where  $\sigma_1$  and  $\sigma_2$  have the differential form

$$d\sigma_j(x) = \frac{h_j(x)dx}{\sqrt{(\lambda - |x|)(|x| - 1)}}, \quad x \in \Delta_j, \quad j = 1, 2.$$

This system  $(\sigma_1, \sigma_2)$  belongs to the class of Angelesco systems introduced by Angelesco in [1]. Fix a multi-index  $\mathbf{n} = (n_1, n_2)$ , we say that a polynomial  $Q_{\mathbf{n}} \not\equiv 0$  is a type II multiple-orthogonal polynomial corresponding to a system  $(\sigma_1, \sigma_2)$ , if  $\deg Q_{\mathbf{n}} \leq |\mathbf{n}| = n_1 + n_2$  and  $Q_{\mathbf{n}}$  satisfies the following orthogonality conditions

$$\int_{\Delta_j} x^\nu Q_{\mathbf{n}}(x) d\sigma_j(x) = 0, \quad \nu = 0, \dots, n_j - 1, \quad j = 1, 2. \quad (1)$$

It is well known (see [1]) that for any multi-index  $\mathbf{n} = (n_1, n_2)$ , the polynomial  $Q_{\mathbf{n}}$  has for each  $j = 1, 2$ , exactly  $n_j$  simple zeros lying in the interior set of  $\Delta_j$ , which we represent by  $\mathring{\Delta}_j$ . We will denote the function of the second kind

$$R_{\mathbf{n}}^j(z) = \frac{1}{2\pi i} \int_{\Delta_j} Q_{\mathbf{n}}(x) \frac{d\sigma_j(x)}{x - z}. \quad (2)$$

Let us take a subset of multi-indices  $\Lambda = \{\mathbf{n} = (n, n) : n \in \mathbb{Z}\}$ . In the present article we obtain results about the strong asymptotics of the sequence of multiple-orthogonal polynomials  $\{Q_{\mathbf{n}} : \mathbf{n} \in \Lambda\}$ . An effective method for such study with this kind of so “very nice” measures, is analyzing the Riemann-Hilbert problem for multiple-orthogonal polynomials, which was introduced in [12]. Let us consider a  $3 \times 3$  matrix,  $Y$ , whose entries are complex functions  $Y_{s,k} : \mathbb{C} \setminus (\Delta_1 \cup \Delta_2) \rightarrow \mathbb{C}$ ,  $s, k = 1, 2, 3$ . Given a point  $x \in \mathring{\Delta}_1 \cup \mathring{\Delta}_2$ , the following matricial limits, where  $z \in \mathbb{C} \setminus (\Delta_1 \cup \Delta_2)$  tending to  $x$ , represent the formal pointwise non tangential limits of all entries of  $Y$  at the same time:

$$\lim_{z \rightarrow x} Y(z) = Y_+(x), \quad \Im m(z) > 0 \quad \text{and} \quad \lim_{z \rightarrow x} Y(z) = Y_-(x), \quad \Im m(z) < 0.$$

Let  $\delta_{s,k} : \mathbb{N}^2 \rightarrow \{0, 1\}$  denote the Kronecker delta function, i.e.  $\delta_{s,k} = 0$  when  $s \neq k$ , and  $\delta_{s,s} = 1$ ,  $s, k \in \mathbb{N}$ . Let us look for a matrix function  $Y$ , which satisfies the following conditions:

- (1) The entries of  $Y$ ,  $Y_{s,k}$ , belongs to  $H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2))$ , which we write as  $Y \in H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2))$ ;

(2) For each  $\Delta_j$ ,  $j = 1, 2$ , the so called jump condition takes place

$$Y_+(x) = Y_-(x) \begin{pmatrix} 1 & \frac{\delta_{1,j}h_1(x)}{\sqrt{(\lambda-|x|)(1-|x|)}} & \frac{\delta_{2,j}h_2(x)}{\sqrt{(\lambda-|x|)(1-|x|)}} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad x \in \overset{\circ}{\Delta}_j;$$

(3) Given a multi-index  $\mathbf{n} = (n, n) \in \Lambda$ , we require the following asymptotic condition at infinity,

$$Y(z) \begin{pmatrix} z^{-2n} & 0 & 0 \\ 0 & z^n & 0 \\ 0 & 0 & z^n \end{pmatrix} = \mathbb{I} + \mathcal{O}(1/z) \quad \text{as } z \rightarrow \infty,$$

where  $\mathbb{I}$  is the identity matrix of size  $3 \times 3$ ;

(4) For each  $i, j = 1, 2$ , we set the following behavior around the endpoints  $c_{1,1} = -\lambda$ ,  $c_{2,1} = -1$ ,  $c_{1,2} = 1$  and  $c_{2,2} = \lambda$ ,

$$Y(z) = \mathcal{O} \begin{pmatrix} 1 & \delta_{2,j} + \frac{\delta_{1,j}}{\sqrt{|z-c_{i,j}|}} & \delta_{1,j} + \frac{\delta_{2,j}}{\sqrt{|z-c_{i,j}|}} \\ 1 & \delta_{2,j} + \frac{\delta_{1,j}}{\sqrt{|z-c_{i,j}|}} & \delta_{1,j} + \frac{\delta_{2,j}}{\sqrt{|z-c_{i,j}|}} \\ 1 & \delta_{2,j} + \frac{\delta_{1,j}}{\sqrt{|z-c_{i,j}|}} & \delta_{1,j} + \frac{\delta_{2,j}}{\sqrt{|z-c_{i,j}|}} \end{pmatrix}, \quad \text{as } z \rightarrow c_{i,j}.$$

This problem, which consists in finding the matrix function  $Y$ , was called in [12] a Riemann-Hilbert problem for type II multiple orthogonal polynomials, and for the system of measures  $(\sigma_1, \sigma_2)$ , RHP in short. The solution  $Y$  is unique and has the form

$$Y(z) = \begin{pmatrix} Q_{\mathbf{n}}(z) & R_{\mathbf{n}}^1(z) & R_{\mathbf{n}}^2(z) \\ d_1 Q_{\mathbf{n}_-^1}(z) & d_1 R_{\mathbf{n}_-^1}^1(z) & d_1 R_{\mathbf{n}_-^1}^2(z) \\ d_2 Q_{\mathbf{n}_-^2}(z) & d_2 R_{\mathbf{n}_-^2}^1(z) & d_2 R_{\mathbf{n}_-^2}^2(z) \end{pmatrix}, \quad (3)$$

with

$$d_j^{-1} = -\frac{1}{2\pi i} \int_{\Delta_j} x^{n-1} Q_{\mathbf{n}_-^j}(x) d\sigma_j(x),$$

where  $\mathbf{n}_-^1 = (n-1, n)$  and  $\mathbf{n}_-^2 = (n, n-1)$ .

The key of our procedure is inspired in the works [2,3,9,10] and it is based in finding the relationship between  $Y$  and a matrix function  $R$  which is the solution of the following RHP:

(1)  $R : \mathbb{C} \rightarrow \mathbb{C}^{3 \times 3}$  belongs to  $H(\mathbb{C} \setminus \gamma)$ ;

- (2)  $R_+(\xi) = R_-(\xi)V_n(\xi)$ ,  $\xi \in \gamma$ ;
- (3)  $R(z) \rightarrow \mathbb{I}$  as  $z \rightarrow \infty$ ;

where  $V_n \in H(\mathcal{A})$ , with  $\mathcal{A} \subset \mathbb{C}$  a certain domain,  $V_n = \mathbb{I} + \epsilon_n$ , such that  $\epsilon_n \rightarrow 0$  uniformly on compact subsets of  $\mathcal{A}$  as  $n \rightarrow \infty$ , and  $\gamma$  is a contour or system of contours, that is contained in  $\mathcal{A}$ . In this case we can assure that

$$R = \mathbb{I} + \mathcal{O}(\epsilon_n) .$$

The RHP for  $Y$  is not normalized in the sense that the conditions (3) at infinity for  $Y$  and  $R$  are different. In order to normalize the RHP, we are going to modify  $Y$  in such a way that we set another RHP with the same contours (possibly different jump conditions), for which the solution tends to the identity matrix as  $z \rightarrow \infty$ . For normalizing we need to take into account the behavior of  $Y(z)$  for large  $z$ . This behavior depends on the distribution of the zeros of the multiple-orthogonal polynomials. The zero distribution of the orthogonal polynomials is usually given by an extremal problem in logarithmic potential theory. In section 2 we introduce some concepts and results which we will need about this theory and we will normalize the Riemann-Hilbert problem at infinity. In section 3 such a Riemann-Hilbert problem with oscillatory and exponentially decreasing jumps can be analyzed by using the steepest descent method introduced by Deift and Zhou (see [5,6]). The first work such that the orthogonal polynomials appear as solution of a Riemann-Hilbert problem is [7], and in [4] these ideas were for the first time applied to get strong asymptotics for orthogonal polynomials.

## 2 The equilibrium problem and the normalization at infinity

Let us fix  $j \in \{1, 2\}$ .  $\mathcal{M}_{1/2}(\Delta_j)$  denotes the set of all finite Borel measures whose supports, i.e.  $\text{supp}(\cdot)$ , are contained in  $\Delta_j$  with total variation  $1/2$ . Take  $\mu_j \in \mathcal{M}_{1/2}(\Delta_j)$  and define its logarithmic potential as follows

$$V^{\mu_j}(z) = \int \log \frac{1}{|z-x|} d\mu_j(x), \quad z \in \mathbb{C} .$$

For each pair of measures  $(\mu_1, \mu_2)$ , where  $\mu_j \in \mathcal{M}_{1/2}(\Delta_j)$ ,  $j = 1, 2$ , we define the quantities

$$m_j(\mu_1, \mu_2) = \min_{x \in \Delta_j} (2V^{\mu_j}(x) + V^{\mu_k}(x)) , \quad j, k = 1, 2, \quad j \neq k .$$

The following Proposition is deduced immediately from the results of [8].

**Proposition 1** *There exists a unique pair  $(\bar{\mu}_1, \bar{\mu}_2) \in \mathcal{M}_{1/2}(\Delta_1) \times \mathcal{M}_{1/2}(\Delta_2)$ ,*

which satisfies for  $j, k = 1, 2$

$$2V^{\bar{\mu}_j}(x) + V^{\bar{\mu}_k}(x) = m_j(\bar{\mu}_1, \bar{\mu}_2) = m_j, \quad x \in \text{supp}(\bar{\mu}_j) = \Delta_j, \quad j \neq k.$$

For each  $j = 1, 2$  the measure  $\bar{\mu}_j$  is absolutely continuous and has the following differential form

$$d\bar{\mu}_1(x) = \frac{\rho_1(x)dx}{\sqrt{(\lambda - |x|)(|x| - 1)}}, \quad d\bar{\mu}_2(x) = \frac{\rho_2(x)dx}{\sqrt{(\lambda - |x|)(|x| - 1)}},$$

where  $\rho_j$  is a function which has an analytic continuation to a neighborhood  $\mathcal{V}_{\rho_j}$  of the interval  $\Delta_j$ .

In what follows we consider  $\mathcal{V}_j = \mathcal{V}_{h_j} \cap \mathcal{V}_{\rho_j}$ . The pair  $(\bar{\mu}_1, \bar{\mu}_2)$  is called extremal or equilibrium pair of measures with respect to  $(\Delta_1, \Delta_2)$ . Let us denote for each  $j = 1, 2$  the analytic potentials

$$g_j(z) = \int_{\Delta_j} \log(z - x) d\bar{\mu}_j(x) = -V^{\bar{\mu}_j}(z) + i \int_{\Delta_j} \arg(z - x) d\bar{\mu}_j(x),$$

where  $\arg$  denotes the principal argument function.

Substituting the logarithmic potential in Proposition 1 we obtain for each  $j, k = 1, 2$  with  $j \neq k$  that

$$-(g_{j+}(x) + g_{j-}(x)) - g_{k-}(x) = m_j, \quad x \in \Delta_j.$$

Observe that if  $c_{1,1} = -\lambda$ ,  $c_{2,1} = -1$ ,  $c_{1,2} = 1$  and  $c_{2,2} = \lambda$ , then

$$g_{j+}(x) - g_{j-}(x) = \begin{cases} 0 & \text{if } c_{2,j} \leq x \\ i\pi & \text{if } c_{1,j} \geq x \\ 2i\pi \int_x^{c_{2,j}} d\bar{\mu}_j(t) & \text{if } x \in \Delta_j \end{cases}.$$

Let us introduce the matrices

$$G(z) = \begin{pmatrix} e^{-n(g_1(z)+g_2(z))} & 0 & 0 \\ 0 & e^{ng_1(z)} & 0 \\ 0 & 0 & e^{ng_2(z)} \end{pmatrix} \quad \text{and} \quad L = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-nm_1} & 0 \\ 0 & 0 & e^{-nm_2} \end{pmatrix}. \quad (4)$$

We define the matrix function  $T = LYGL^{-1}$ , where  $L, G$  are as in (4) and  $Y$  is given by (3). Hence  $T$  is the unique solution of the RHP:

- (1)  $T \in H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2))$ ;
- (2)  $T_+(x) = T_-(x)M(x)$ ,  $x \in \overset{\circ}{\Delta}_1 \cup \overset{\circ}{\Delta}_2$ ;
- (3)  $T(z) = \mathbb{I} + \mathcal{O}(1/z)$  as  $z \rightarrow \infty$ ;
- (4)  $T$  and  $Y$  have the same behavior on the endpoints of the intervals  $\Delta_j$ , for  $j = 1, 2$ ;

where the jump matrix  $M$  has the form

$$M(x) = \begin{pmatrix} e^{-2ni\pi \int_x^{c_{2,j}} d\bar{\mu}_j(t)} & \frac{\delta_{j,1} h_1(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} & \frac{\delta_{j,2} h_2(x) dx}{\sqrt{(\lambda-|x|)(|x|-1)}} \\ 0 & e^{2n\delta_{j,1} i\pi \int_x^{c_{2,1}} d\bar{\mu}_1(t)} & 0 \\ 0 & 0 & e^{2n\delta_{j,2} i\pi \int_x^{c_{2,2}} d\bar{\mu}_2(t)} \end{pmatrix}, \quad (5)$$

with  $x \in \mathring{\Delta}_j$ .

### 3 The opening of the lens

Let us consider

$$\phi_1(z) = -\pi \int_z^{-1} \frac{\rho_1(\zeta) d\zeta}{\sqrt{(\zeta + \lambda)(\zeta + 1)}}, \quad z \in \mathcal{V}_1$$

and

$$\phi_2(z) = -\pi \int_z^\lambda \frac{\rho_2(\zeta) d\zeta}{\sqrt{(\zeta - \lambda)(\zeta - 1)}}, \quad z \in \mathcal{V}_2.$$

We have considered  $\sqrt{(\zeta + \lambda)(\zeta + 1)}$  and  $\sqrt{(\zeta - \lambda)(\zeta - 1)}$  as analytic functions on  $\mathbb{C} \setminus \Delta_1$  and  $\mathbb{C} \setminus \Delta_2$ , respectively, where we have taken the branches which are positive for real  $\zeta > -1$  and  $\zeta > \lambda$ , respectively. Observe that for each  $j = 1, 2$ , the function  $\phi_j \in H(\mathcal{V}_j \setminus \Delta_j)$ , the real part of the functions  $\phi_{j\pm}$  vanish on  $\Delta_j$ ,  $\Re(\phi_{j\pm})(x) = 0$ ,  $x \in \Delta_j$ , and their derivatives

$$\phi'_{j\pm}(x) = \mp i\pi \frac{\rho_j(x)}{\sqrt{(\lambda - |x|)(|x| - 1)}}.$$

By the Cauchy-Riemann conditions we have that

$$\pm \frac{\partial \Re \phi_{\pm}}{\partial y}(x) > 0, \quad x \in \Delta_j.$$

Since  $\Re \phi_j$  is a harmonic function on  $\mathcal{V}_j \setminus \Delta_j$  we can assure that  $\Re \phi_j(z) > 0$ ,  $z \in \mathcal{V}_j \setminus \Delta_j$ .

Factorize the jump matrix function  $M$  in (5) as follows

$$M(x) = \begin{pmatrix} 1 & 0 & 0 \\ \frac{\delta_{j,1} e^{-2n\phi_{1-}(x)} \sqrt{(\lambda-|x|)(|x|-1)}}{h_1(x)} & 1 & 0 \\ \frac{\delta_{j,2} e^{-2n\phi_{2-}(x)} \sqrt{(\lambda-|x|)(|x|-1)}}{h_2(x)} & 0 & 1 \end{pmatrix}$$

$$\times \begin{pmatrix} 0 & \frac{\delta_{j,1}h_1(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} & \frac{\delta_{j,2}h_2(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} \\ -\frac{\delta_{1,j}\sqrt{(\lambda-|x|)(|x|-1)}}{h_1(x)} & \delta_{j,2} & 0 \\ -\frac{\delta_{2,j}\sqrt{(\lambda-|x|)(|x|-1)}}{h_2(x)} & 0 & \delta_{j,1} \end{pmatrix} \times \begin{pmatrix} 1 & 0 & 0 \\ \frac{\delta_{j,1}e^{-2n\phi_{1+}(x)}\sqrt{(\lambda-|x|)(|x|-1)}}{h_1(x)} & 1 & 0 \\ \frac{\delta_{j,2}e^{-2n\phi_{2+}(x)}\sqrt{(\lambda-|x|)(|x|-1)}}{h_2(x)} & 0 & 1 \end{pmatrix}.$$

Now we are going to follow a procedure analogous to the one in [3]. For each  $j = 1, 2$  let us fix a closed curve  $\gamma_j$  contained in  $\mathcal{V}_j$ , with the clockwise orientation. Set  $\Gamma_j$  the bounded connected component of  $\mathbb{C} \setminus \gamma_j$ . Let us introduce the matrix function  $S$ , defined by

$$S(z) = T(z) \begin{pmatrix} 1 & 0 & 0 \\ \frac{i\delta_{1,j}e^{-2n\phi_1(z)}\sqrt{(z+\lambda)(z+1)}}{h_1(z)} & 1 & 0 \\ \frac{i\delta_{2,j}e^{-2n\phi_2(z)}\sqrt{(z-\lambda)(z-1)}}{h_2(z)} & 0 & 1 \end{pmatrix}, \quad z \in \Gamma_j,$$

and  $S(z) = T(z)$ ,  $z \in \mathbb{C} \setminus \bar{\Gamma}_j$ .

The matrix function  $S$  satisfies the RHP:

- (1)  $S \in H(\mathbb{C} \setminus \cup_{j=1,2}(\Delta_j \cup \gamma_j))$ ;
- (2) The jump conditions  $j = 1, 2$  are,

$$S_+(x) = S_-(x) \begin{pmatrix} 0 & \frac{\delta_{1,j}h_1(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} & \frac{\delta_{2,j}h_2(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} \\ -\frac{\delta_{1,j}\sqrt{(\lambda-|x|)(|x|-1)}}{h_1(x)} & \delta_{2,j} & 0 \\ -\frac{\delta_{2,j}\sqrt{(\lambda-|x|)(|x|-1)}}{h_2(x)} & 0 & \delta_{1,j} \end{pmatrix},$$

when  $x \in \overset{\circ}{\Delta}_j$ , and if  $z \in \gamma_j$ ,

$$S_+(z) = S_-(z) \begin{pmatrix} 1 & 0 & 0 \\ \frac{i\delta_{1,j}e^{-2n\phi_1(z)}\sqrt{(z+\lambda)(z+1)}}{h_1(z)} & 1 & 0 \\ \frac{i\delta_{2,j}e^{-2n\phi_2(z)}\sqrt{(z-\lambda)(z-1)}}{h_2(z)} & 0 & 1 \end{pmatrix};$$

- (3)  $S(z) = \mathbb{I} + \mathcal{O}(1/z)$  as  $z \rightarrow \infty$ ;
- (4) The conditions for the endpoints are the same as for  $T$ .

Now, we consider the limiting problem, because for the matrix  $S$  the jump matrix function on each  $\gamma_j$  for  $j = 1, 2$  tends to the identity matrix when

$n \rightarrow \infty$ . We look for the matrix function  $N$  which satisfies the following RHP:

- (1)  $N \in H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2))$ ;
- (2) The jump conditions in  $\mathring{\Delta}_j$  for  $j = 1, 2$  are,

$$N_+(x) = N_-(x) \begin{pmatrix} 0 & \frac{\delta_{1,j} h_1(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} & \frac{\delta_{2,j} h_2(x)}{\sqrt{(\lambda-|x|)(|x|-1)}} \\ -\frac{\delta_{1,j} \sqrt{(\lambda-|x|)(|x|-1)}}{h_1(x)} & \delta_{2,j} & 0 \\ -\frac{\delta_{2,j} \sqrt{(\lambda-|x|)(|x|-1)}}{h_2(x)} & 0 & \delta_{1,j} \end{pmatrix}; \quad (6)$$

- (3)  $N(z) = \mathbb{I} + \mathcal{O}(1/z)$  as  $z \rightarrow \infty$ ;
- (4)  $N$  satisfies the same conditions for the endpoints as  $S$ .

Let us consider the matrix function  $K = [K_{k,l}]$ ,  $k, l = 1, 2, 3$  that is the solution of the RHP:

- (1)  $K \in H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2))$ ;
- (2) The jump conditions in  $\mathring{\Delta}_j$  for  $j = 1, 2$  are, because of (6),

$$K_+(x) = K_-(x) \begin{pmatrix} 0 & \frac{\delta_{1,j}}{\sqrt{(\lambda-|x|)(|x|-1)}} & \frac{\delta_{2,j}}{\sqrt{(\lambda-|x|)(|x|-1)}} \\ -\delta_{1,j} \sqrt{(\lambda-|x|)(|x|-1)} & \delta_{2,j} & 0 \\ -\delta_{2,j} \sqrt{(\lambda-|x|)(|x|-1)} & 0 & \delta_{1,j} \end{pmatrix} \quad (7)$$

- (3)  $K(z) = \mathbb{I} + \mathcal{O}(1/z)$  as  $z \rightarrow \infty$ ;
- (4) The conditions for the endpoints are the same as for  $N$ .

Notice that when  $h_j = 1$ ,  $j = 1, 2$ ,  $K$  and  $N$  have the same RHP. Analogously to the ideas in [3], let us again consider  $\sqrt{(z+\lambda)(z+1)}$  and  $\sqrt{(z-\lambda)(z-1)}$  as analytic functions on  $\mathbb{C} \setminus \Delta_1$  and  $\mathbb{C} \setminus \Delta_2$ , respectively, where we have taken the branches which are positive for real  $z > -1$  and  $z > \lambda$ , respectively,

$$\left(\frac{1}{i} \sqrt{(z+\lambda)(z+1)}\right)_{\pm}(x) = \pm \sqrt{(\lambda+x)(-x-1)}, \quad x \in \mathring{\Delta}_1$$

and

$$\left(\frac{1}{i} \sqrt{(z-\lambda)(z-1)}\right)_{\pm}(x) = \pm \sqrt{(\lambda-x)(x-1)}, \quad x \in \mathring{\Delta}_2.$$

For each  $k = 1, 2, 3$ , we rewrite (7) as

$$\begin{cases} \left(\frac{1}{i} \sqrt{(z+\lambda)(z+1)} K_{k,2}\right)_{\pm}(x) = (K_{k,1})_{\mp}(x) \\ (K_{k,3})_+(x) = (K_{k,3})_-(x) \end{cases}, \quad x \in \mathring{\Delta}_1$$



$$\begin{cases} \left(\frac{1}{i}\sqrt{(z-\lambda)(z-1)}K_{k,3}\right)_{\pm}(x) = (K_{k,1})_{\mp}(x) \\ (K_{k,2})_{+}(x) = (K_{k,2})_{-}(x) \end{cases}, \quad x \in \mathring{\Delta}_2$$

and we denote

$$\begin{aligned} \psi_0^k(z) &= K_{k,1}(z), \quad \psi_1^k(z) = \frac{1}{i}\sqrt{(z+\lambda)(z+1)}K_{k,2}(z) \\ \text{and } \psi_2^k(z) &= \frac{1}{i}\sqrt{(z-\lambda)(z-1)}K_{k,3}(z). \end{aligned}$$

Then from the relations (7), we may interpret each row  $k = 1, 2, 3$  of such matrix  $K$  as a function defined on a Riemann surface. Let  $\mathcal{R}$  define the Riemann surface which has two cuts. One of them connects the two branch points  $-\lambda$  and  $-1$  with the cut in the interval  $\Delta_1$ . The other cut is made in the interval  $\Delta_2$ , to connect the two other branch points  $1$  and  $\lambda$ . The sheet  $\mathcal{R}_0$  is glued to another sheet  $\mathcal{R}_1$  along the cut  $\Delta_1$ , and  $\mathcal{R}_0$  is also glued to  $\mathcal{R}_2$  along the interval  $\Delta_2$ . Let us denote by  $\psi^k$ ,  $k = 1, 2, 3$ , three multi-valued functions  $\psi^k = (\psi_0^k, \psi_1^k, \psi_2^k)$ , such that for each  $k = 1, 2, 3$  its components  $\psi_l^k$ ,  $l = 0, 1, 2$ ,  $k = 1, 2, 3$ , map the corresponding sheet  $\mathcal{R}_l$  onto  $\mathbb{C}$ , and satisfy:

- i)  $\psi_0^k \in H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2))$ ,  $\psi_j^k \in H(\mathbb{C} \setminus \Delta_j)$ ,  $j = 1, 2$ ;
- ii)  $\psi_{0\pm}^k = \psi_{j\mp}^k$ ,  $j = 1, 2$ ;
- iii)  $\psi_0^k = \delta_{k,1} + \mathcal{O}(1/z)$ , and  $\psi_l^k(z) = -iz\delta_{k,l+1} + \mathcal{O}(1)$ ,  $l = 1, 2$ , as  $z \rightarrow \infty$ ;
- iv)  $\psi_l^k(z) = \mathcal{O}(1)$ , at the endpoints.

Observe that  $\psi^1 : \mathcal{R} \rightarrow \mathbb{C}$  is a bounded holomorphic function on  $\mathcal{R}$ , where  $\lim_{z \rightarrow \infty} \psi_0^1(z) := \psi_0^1(\infty) = 1$ . This implies that  $\psi^1$  is the constant function identically equal to 1, i.e.  $\psi^1 \equiv 1$ . For the cases when  $k = 2, 3$ , G. López Lagomasino et al., [11], proved that up to complex constants  $c_1, c_2$

$$\psi^2(z) = \frac{c_1}{\varphi(z)} \quad \text{and} \quad \psi^3(z) = c_2 \frac{\varphi^1(z)}{\varphi(z)},$$

where

$$\varphi(z) = \left(\frac{1+a^2}{(1-a^2)^2}\right)^{1/3} (1+G^{-1}(z)), \quad \varphi^1(z) = \frac{1+G^{-1}(z)}{1-G^{-1}(z)}, \quad (8)$$

$$G(w) = \frac{H(w)}{H(a)}, \quad H(w) = w - \frac{(1-a^2)^2 w}{(1+a^2)(1-w^2)},$$

and  $a$  is the unique solution on the interval  $]0, 1[$  of the biquartic equation

$$a^8 + (16\lambda^2 - 8)a^6 + 18a^4 - 27 = 0. \quad (9)$$

In this case,  $H^{-1}(z)$  is the solution of the cubic equation

$$w^3 - zw^2 + \frac{a^4 - 3a^2}{1+a^2}w + z = 0. \quad (10)$$

Notice that given a value  $\lambda > 1$ , the equation (9) as well as (10) can be solved by elementary methods.

Let us find the diagonal  $3 \times 3$  matrix function  $D = \text{diag}(D_0, D_1, D_2)$ , such that  $N(z) = D^{-1}(\infty)K(z)D(z)$ . The conditions (7) imply that the entries of  $D$  must satisfy the following conditions

$$h_j(x)D_{0\pm}(x) = D_{j\mp}(x), \quad D_{k+}(x) = D_{k-}(x) \quad \text{when } x \in \overset{\circ}{\Delta}_j, \quad j, k = 1, 2, \quad k \neq j,$$

i.e.  $D_l$ ,  $l = 0, 1, 2$  are the *Szegő-type functions*.

Analogously to the function  $\psi_j^i$ , we obtain the following problem for the entries of  $D$ :

- i)  $D_0 \in H(\overline{\mathbb{C}} \setminus (\Delta_1 \cup \Delta_2)), D_j \in H(\overline{\mathbb{C}} \setminus \Delta_j), j = 1, 2;$
- ii)  $h_j(x)D_{0\pm}(x) = D_{j\mp}(x), j = 1, 2;$
- iii)  $D_l(z) = \mathcal{O}(1), l = 0, 1, 2,$  at the endpoints.

In order to find this matrix function  $D$ , we consider the function  $\varphi$  given by (8), such that its components  $\varphi_l$ ,  $l = 0, 1, 2$ , map the corresponding sheet  $\mathcal{R}_l$  on  $\mathbb{C}$ , and satisfy:

- i)  $\varphi_0 \in H(\mathbb{C} \setminus (\Delta_1 \cup \Delta_2)), \varphi_j \in H(\mathbb{C} \setminus \Delta_j), j = 1, 2;$
- ii)  $\varphi_{0\pm} = \varphi_{j\mp}, j = 1, 2;$
- iii)  $\varphi_0(z) = \mathcal{O}(z), \varphi_1(z) = \mathcal{O}(1/z),$  and  $\varphi_2(z) = \mathcal{O}(1),$  as  $z \rightarrow \infty;$
- iv)  $\varphi_0\varphi_1\varphi_2(\infty) = 1;$
- v)  $\varphi_l(z) = \mathcal{O}(1),$  at the endpoints.

We denote by  $\Sigma_j = \varphi_{0-}(\Delta_j) \cup \varphi_{0+}(\Delta_j)$ , for  $j = 1, 2$  the closed curves in the complex plane, with the clockwise orientation, and we denote by  $\Omega_j$  the interior set of  $\Sigma_j$  for  $j = 0, 1, 2$  and by  $\Omega_0$  the exterior set of  $\Sigma_1 \cup \Sigma_2$ . Taking into account the behavior of the functions  $\varphi_l$  at infinity,  $\Omega_l = \varphi_l(\mathcal{R}), l = 0, 1, 2$ . Using (10) we get that  $\varphi(z)$  is the solution of the cubic algebraic equation

$$w^3 - \left( \frac{1+a^2}{(1-a^2)^2} \right)^{1/3} (3+z)w^2 + \left( \frac{1+a^2}{(1-a^2)^2} \right)^{2/3} \left( 2z + \frac{3+a^4}{1+a^2} \right) w - 1 = 0,$$

that is equivalent to

$$z = \frac{w^3 - 3 \left( \frac{1+a^2}{(1-a^2)^2} \right)^{1/3} w^2 + \left( \frac{1+a^2}{(1-a^2)^2} \right)^{2/3} \frac{3+a^4}{1+a^2} w - 1}{\left( \frac{1+a^2}{(1-a^2)^2} \right)^{1/3} w^2 - 2 \left( \frac{1+a^2}{(1-a^2)^2} \right)^{2/3} w} =: r(w).$$

Using this rational function  $r$  we consider the complex function  $\tilde{D}$ , defined as

$$\tilde{D}(w) = \begin{cases} D_0(r(w)), & w \in \Omega_0 \\ D_1(r(w)), & w \in \Omega_1 \\ D_2(r(w)), & w \in \Omega_2 \end{cases}.$$

This function  $\tilde{D}$  verifies the multiplicative scalar Riemann-Hilbert problem

$$h_j(r(\xi))\tilde{D}(\xi)_- = \tilde{D}(\xi)_+, \quad \xi \in \Sigma_j, \quad j = 1, 2.$$

Taking into account that  $D_0D_1D_2$  is an entire function, and using the behavior at  $z = \infty$ , it follows that  $D_0D_1D_2 \equiv c$ , where  $c$  is a complex constant. We can choose a single valued branch of the complex logarithm, and we have the additive scalar Riemann-Hilbert problem

$$\log h_j(r(\xi)) + \log \tilde{D}(\xi)_- = \log \tilde{D}(\xi)_+, \quad \xi \in \Sigma_j, \quad j = 1, 2.$$

Using the Sokhotsky-Plemelj formula we obtain that

$$\log \tilde{D}(w) = \frac{1}{2\pi i} \sum_{j=1,2} \int_{\Sigma_j} \frac{\log h_j(r(\xi))}{\xi - w} d\xi,$$

and so, the Szegő-type functions, are given explicitly by,

$$D_l(z) = \exp \left\{ \frac{1}{2\pi i} \sum_{j=1,2} \varepsilon_j \int_{\Delta_j} \log h_j(x) \left( \frac{-\varphi'_{0+}(x)}{\varphi_{0+}(x) - \varphi_l(z)} + \frac{\varphi'_{0-}(x)}{\varphi_{0-}(x) - \varphi_l(z)} \right) dx \right\}, \quad (11)$$

for  $l = 0, 1, 2$ , where  $\varepsilon_j = 1$  if orientation of  $-\varphi_{0+}(\Delta_j) \cup \varphi_{0-}(\Delta_j)$  is in the clockwise direction, where we are considering that the intervals  $\Delta_j$ ,  $j = 1, 2$  are oriented from left to right, and  $\varepsilon_j = -1$  if this not happen.

For this functions  $D_l$  the behavior at the end points of the intervals  $\Delta_j$ , for  $j = 1, 2$  is  $\mathcal{O}(1)$  if we take into account the quadratic ramifications at these points suggested by the Riemann surface,  $\mathcal{R}$ .

Finally the matrix function  $N$  has the form

$$N(z) = \begin{pmatrix} \frac{D_0(z)}{D_0(\infty)} & \frac{i D_1(z)}{D_0(\infty)\sqrt{(z+\lambda)(z+1)}} & \frac{i D_2(z)}{D_0(\infty)\sqrt{(z-\lambda)(z-1)}} \\ \frac{D_0(z)\psi_0^2(z)}{D_1(\infty)} & \frac{i D_1(z)\psi_1^2(z)}{D_1(\infty)\sqrt{(z+\lambda)(z+1)}} & \frac{i D_2(z)\psi_2^2(z)}{D_1(\infty)\sqrt{(z-\lambda)(z-1)}} \\ \frac{D_0(z)\psi_0^3(z)}{D_2(\infty)} & \frac{i D_1(z)\psi_1^3(z)}{D_2(\infty)\sqrt{(z+\lambda)(z+1)}} & \frac{i D_2(z)\psi_2^3(z)}{D_2(\infty)\sqrt{(z-\lambda)(z-1)}} \end{pmatrix}.$$

Set  $R(z) = S(z)N^{-1}(z)$ . Since  $S$  and  $N$  have the same jump across  $\overset{\circ}{\Delta}_j$ ,  $j = 1, 2$ , we have that  $R_+(x) = R_-(x)$  for  $x \in \overset{\circ}{\Delta}_j$ ,  $j = 1, 2$ . From the

definition of  $R$ , and the endpoint conditions for  $N$ , we can also deduce that these endpoints are removable singularities. Hence  $R$  is an analytic function across the full intervals  $\Delta_1$  and  $\Delta_2$ , and it has jumps on the curves  $\gamma_j$ ,  $j = 1, 2$ . Then we have the following RHP for  $R$ :

- (1)  $R \in H(\mathbb{C} \setminus (\gamma_1 \cup \gamma_2))$ ;
- (2) The jump conditions are for  $j = 1, 2$

$$R_+(z) = R_-(z) N(z) \begin{pmatrix} 1 & 0 & 0 \\ \frac{i \delta_{1,j} e^{-2n\phi_1(z)} \sqrt{(z+\lambda)(z+1)}}{h_1(z)} & 1 & 0 \\ \frac{i \delta_{2,j} e^{-2n\phi_2(z)} \sqrt{(z-\lambda)(z-1)}}{h_2(z)} & 0 & 1 \end{pmatrix} N^{-1}(z) \quad \text{if } z \in \gamma_j;$$

- (3)  $R(z) = \mathbb{I} + \mathcal{O}(1/z)$ .

Then in each compact  $\mathcal{K} \subset \mathbb{C} \setminus (\gamma_1 \cup \gamma_2)$ , using the same argument as in [10], we have that  $R = \mathbb{I} + \mathcal{O}(e^{-cn})$ , with  $c(\mathcal{K}) > 0$  uniformly as  $n \rightarrow \infty$ , so it holds uniformly in compact sets of the indicated region that

$$Y(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{nm_1} & 0 \\ 0 & 0 & e^{nm_2} \end{pmatrix} \left( \mathbb{I} + \mathcal{O}(e^{-cn}) \right) N(z) \times \begin{pmatrix} e^{n(g_1(z)+g_2(z))} & 0 & 0 \\ 0 & e^{-n(m_1+g_1(z))} & 0 \\ 0 & 0 & e^{-n(m_2+g_2(z))} \end{pmatrix},$$

$z \in \mathbb{C} \setminus (\bar{\Gamma}_1 \cup \bar{\Gamma}_2)$ , and

$$Y(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{nm_1} & 0 \\ 0 & 0 & e^{nm_2} \end{pmatrix} \left( \mathbb{I} + \mathcal{O}(e^{-cn}) \right) N(z) \times \begin{pmatrix} 1 & 0 & 0 \\ \frac{-i \delta_{1,j} e^{-2n\phi_1(z)} \sqrt{(z+\lambda)(z+1)}}{h_1(z)} & 1 & 0 \\ \frac{-i \delta_{2,j} e^{-2n\phi_2(z)} \sqrt{(z-\lambda)(z-1)}}{h_2(z)} & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{n(g_1(z)+g_2(z))} & 0 & 0 \\ 0 & e^{-n(m_1+g_1(z))} & 0 \\ 0 & 0 & e^{-n(m_2+g_2(z))} \end{pmatrix},$$

$z \in \Gamma_j$ , where  $N$  is given by (3).

Finally, we state the main result of this paper.

**Theorem 1** *The type II multiple orthogonal polynomial given by (1), has on any compact  $\mathcal{K} \subset \mathbb{C} \setminus (\Delta_1 \cup \Delta_2)$ , uniformly as  $n \rightarrow \infty$ , the following strong asymptotic behavior,*

$$\begin{aligned} Q_{\mathbf{n}}(z) &= \frac{D_0(z)}{D_0(\infty)} e^{n(g_1(z)+g_2(z))} \left(1 + \mathcal{O}(e^{-cn})\right), \\ d_1 Q_{\mathbf{n}_-^1}(z) &= \frac{D_0(z)}{D_0(\infty)} \psi_0^2(z) e^{n(m_1+g_1(z)+g_2(z))} \left(1 + \mathcal{O}(e^{-cn})\right), \\ d_2 Q_{\mathbf{n}_-^2}(z) &= \frac{D_0(z)}{D_0(\infty)} \psi_0^3(z) e^{n(m_2+g_1(z)+g_2(z))} \left(1 + \mathcal{O}(e^{-cn})\right), \end{aligned}$$

and also holds on any compact  $\mathcal{K} \subset \Delta_j$ ,  $j, k = 1, 2$ ,  $j \neq k$ ,

$$\begin{aligned} Q_{\mathbf{n}}(x) &= \left\{ \frac{D_{0+}(x)}{D_0(\infty)} e^{ng_{j+}(x)} + \frac{D_{0-}(x)}{D_0(\infty)} e^{ng_{j-}(x)} \right\} e^{ng_k(x)} \left(1 + \mathcal{O}(e^{-cn})\right), \\ d_1 Q_{\mathbf{n}_-^1}(x) &= \left\{ \frac{D_{0+}(x)}{D_1(\infty)} e^{ng_{j+}(x)} \psi_{0+}^2(x) + \frac{D_{0-}(x)}{D_1(\infty)} e^{ng_{j-}(x)} \psi_{0-}^2(x) \right\} \\ &\quad \times e^{n(g_k(x)+m_1)} \left(1 + \mathcal{O}(e^{-cn})\right), \\ d_2 Q_{\mathbf{n}_-^2}(x) &= \left\{ \frac{D_{0+}(x)}{D_2(\infty)} e^{ng_{j+}(x)} \psi_{0+}^3(x) + \frac{D_{0-}(x)}{D_2(\infty)} e^{ng_{j-}(x)} \psi_{0-}^3(x) \right\} \\ &\quad \times e^{n(g_k(x)+m_2)} \left(1 + \mathcal{O}(e^{-cn})\right). \end{aligned}$$

We can also state:

**Theorem 2** *The second kind function given by (2), has on any compact  $\mathcal{K}$  of the indicated region, uniformly as  $n \rightarrow \infty$ , the following strong asymptotic behavior,*

$$\begin{aligned} R_{\mathbf{n}}^1(z) &= \frac{i D_1(z) e^{-n(m_1+g_1(z))}}{D_0(\infty) \sqrt{(z+\lambda)(z+1)}} \left(1 + \mathcal{O}(e^{-cn})\right), \quad z \in \mathbb{C} \setminus \Delta_1, \\ R_{\mathbf{n}}^2(z) &= \frac{i D_2(z) e^{-n(m_2+g_2(z))}}{D_0(\infty) \sqrt{(z-\lambda)(z-1)}} \left(1 + \mathcal{O}(e^{-cn})\right), \quad z \in \mathbb{C} \setminus \Delta_2, \\ d_1 R_{\mathbf{n}_-^1}(z) &= \frac{i D_1(z) \psi_1^2(z) e^{-ng_1(z)}}{D_1(\infty) \sqrt{(z+\lambda)(z+1)}} \left(1 + \mathcal{O}(e^{-cn})\right), \quad z \in \mathbb{C} \setminus \Delta_1, \\ d_1 R_{\mathbf{n}_-^2}(z) &= \frac{i D_2(z) \psi_2^2(z) e^{-n(m_2-m_1+g_2(z))}}{D_1(\infty) \sqrt{(z-\lambda)(z-1)}} \left(1 + \mathcal{O}(e^{-cn})\right), \quad z \in \mathbb{C} \setminus \Delta_2, \\ d_2 R_{\mathbf{n}_-^1}(z) &= \frac{i D_1(z) \psi_1^3(z) e^{-n(m_1-m_2+g_1(z))}}{D_2(\infty) \sqrt{(z+\lambda)(z+1)}} \left(1 + \mathcal{O}(e^{-cn})\right), \quad z \in \mathbb{C} \setminus \Delta_1, \\ d_2 R_{\mathbf{n}_-^2}(z) &= \frac{i D_2(z) \psi_2^3(z) e^{-ng_2(z)}}{D_2(\infty) \sqrt{(z-\lambda)(z-1)}} \left(1 + \mathcal{O}(e^{-cn})\right), \quad z \in \mathbb{C} \setminus \Delta_2. \end{aligned}$$

## References

- [1] M.A. ANGELESCO, *Sur deux extensions des fractions continues algébriques*, C.R. Acad. Sci. Paris, **18** (1919), 262-263.
- [2] A.I. APTEKAREV, A.B.J. KUIJLAARS, AND W. VAN ASSCHE, *Asymptotics of Hermite-Padé rational approximants for two analytic functions with separated pairs of branch points (case of genus 0)*, Internat. Math. Research Papers, Vol. 2008, Article ID rpm007, 128 pages.
- [3] A.I. APTEKAREV AND W. VAN ASSCHE, *Scalar and matrix Riemann-Hilbert approach to the strong asymptotics of Padé approximants and complex orthogonal polynomials with varying weight*, J. of Approx. Theory, **129** (2) (2004), 129-166.
- [4] P. DEIFT, T. KRIECHERBAUER, K. T.-R. MCLAUGHLIN, S. VENAKIDES, AND X. ZHOU, *Strong asymptotics of orthogonal polynomials with respect to exponential weights*, Comm. Pure Appl. Math., **52** (12) (1999), 1491-1552.
- [5] P. DEIFT, X. ZHOU, *A steepest descent method for oscillatory Riemann-Hilbert problems*, Bull. Amer. Math. Soc., **26** (1) (1992), 119-123.
- [6] P. DEIFT, X. ZHOU, *A steepest descent method for oscillatory Riemann-Hilbert problems, Asymptotic for the MKdV equation*, Ann. of Math., Second Series, **137** (2) (1993), 295-368.
- [7] A.S. FOKAS, A.R. ITS, AND A.V. KITAEV, *The isomonodromy approach to matrix models in 2D quantum gravity*, Comm. Math. Phys., **147** (2) (1992), 395-430.
- [8] A.A. GONCHAR AND E.A. RAKHMANOV, *On Convergence of Simoultaneous Padé Approximants for Systems of Functions of Markov Type*, Proceedings of the Steklov Institute of Mathematics, **3** (1983), 31-50.
- [9] A.B.J. KUIJLAARS, *Riemann-Hilbert analysis for orthogonal polynomials*, In Orthogonal Polynomials and Special Functions (E. Koelink and W. Van Asshe, eds.), Lect. Notes Math., **1817**, Springer-Verlag, Berlin, 2003, 167-210.
- [10] A.B.J. KUIJLAARS, W. VAN ASSCHE, F. WIELONSKY, *Quadratic Hermite-Padé approximation to the exponential function: a Riemann-Hilbert approach*, Constr. Approx., **21**(3) (2005), 351-412.
- [11] G. LÓPEZ LAGOMASINO, D. PESTANA, J.M. RODRÍGUEZ, AND D. YAKUBOVICH, Personal communication.
- [12] W. VAN ASSCHE, J.S. GERONIMO AND A.B.J. KUIJLAARS, *Riemann-Hilbert Problems for Multiple Orthogonal Polynomials, Special Functions 2000: Current Perspectives and Futire Directions*, (J. Bustoz et al., eds.), Kluwer, Dordrecht, (2001), 23-59.