

Symbiosis (2012) 56:129–137
DOI 10.1007/s13199-012-0168-0

The smaller vesicomimid bivalves in the genus *Isorropodon* (Bivalvia, Vesicomiyidae, Pliocardiinae) also harbour chemoautotrophic symbionts

Clara F. Rodrigues · Marina R. Cunha · Karine Olu · Sébastien Duperron

Received: 9 March 2012 / Accepted: 11 May 2012 / Published online: 25 May 2012
© Springer Science+Business Media B.V. 2012

Abstract Species of *Isorropodon* are vesicomimid bivalves for which little information is available regarding host phylogeny and bacterial symbioses. In this study we investigated the symbioses in three *Isorropodon* species from three cold seep areas: *Isorropodon bigoti* (Gulf of Guinea), *Isorropodon megadesmus* (Gulf of Cadiz) and *Isorropodon perplexum* (Eastern Mediterranean). Analysis of bacterial 16S ribosomal RNA gene sequences demonstrated that each vesicomimid species harbours a single symbiont phylotype, that symbionts from the three species cluster together, and that they are closely related to other known vesicomimid symbionts. These results are confirmed by other marker genes (encoding 23S rRNA and APS reductase) and by fluorescence in situ hybridization. Due to their extended depth range and transoceanic distribution *Isorropodon* species are interesting examples to further study evolutionary processes in bivalve hosts and their associated symbionts.

Keywords Vesicomiyids · *Isorropodon* · Symbioses · Reduced environments · Atlantic · Mediterranean

C. F. Rodrigues · S. Duperron
Université Pierre et Marie Curie, UMR 7138 (UPMC CNRS IRD MNHN), Systématique, Adaptation, Evolution,
7, quai St. Bernard, bâtiment A,
75005 Paris, France

C. F. Rodrigues (✉) · M. R. Cunha
Departamento de Biologia & CESAM, Universidade de Aveiro,
Campus Universitário de Santiago,
3810-193 Aveiro, Portugal
e-mail: clara.rodrigues@ua.pt

K. Olu
Laboratoire Environnement Profond, Département Etudes des
Ecosystèmes Profonds, Centre Ifremer de Brest,
BP 71, 29280 Plouzané, France

1 Introduction

Vesicomiyidae clams are among the dominant metazoan fauna in deep-sea sulphide-rich habitats (e.g. hydrothermal vents, cold seeps, whale carcasses) and occur worldwide from 77°N to 70°S at depths from 100 to 9,050 m (Krylova and Sahling 2010). Genera within the subfamily Pliocardiinae (considered large-sized vesicomiyids, with shell lengths in the range of 1 to 30 cm) display distinctive shared features including reduced gut and feeding groove, indicating a large dependence upon (intracellular) chemoautotrophic bacteria for their nutrition (Krylova and Sahling 2010). Indeed all large-sized vesicomiyids studied so far live in obligate symbiosis with sulphur-oxidising *Gammaproteobacteria* located within their gill epithelial cells, and data regarding the diversity, structure and functioning of symbioses have been published by several authors (see Kim et al. 1995; Imhoff et al. 2003; Hurtado et al. 2003; Stewart et al. 2008; Stewart and Cavanaugh 2009; Stewart et al. 2009). The advent of symbiosis in this group is probably linked to its radiation that started in Eocene (from about 55 MYA) (Amano and Kiel 2007). Unlike most chemosynthetic symbioses found in metazoan, the transmission of bacterial symbionts from one generation to the next in vesicomiyids is predominantly vertical, although occasional acquisition of alternative lineages from the environment is also documented (Cary and Giovannoni 1993; Dubilier et al. 2008; Stewart et al. 2008). Vertical transmission in vesicomiyids leads to host/symbiont co-speciation patterns which are not observed for example in mytilids, despite the close relatedness of sulphur-oxidising symbionts in these two bivalve families (Peek et al. 1998). Genome sequencing of symbionts has emphasized several traits typical of maternally-transmitted bacteria, including genome size reduction coupled with the loss of certain genes (Kim et al. 1995; Harada

et al. 2009; Kuwahara et al. 2008; Stewart et al. 2008). The systematics and evolution of Pliocardiinae hosts have also been studied but their taxonomy is still under debate as genera are typically found non-monophyletic in phylogenetic reconstructions (Kojima et al. 1995; Baco et al. 1999; Osborn 2001; Goffredi et al. 2003; Kojima et al. 2004, 2006; Decker et al. 2012). Yet, apart from large-sized Pliocardiinae vesicomyids, there are in this subfamily smaller-sized species which have been overlooked, and for which very little data is available, both for hosts and associated symbionts. Among these, the genus *Isorropodon* includes nine accepted species reported from the Eastern Atlantic Ocean along the west coast of Africa from Namibia to Mauritania (*Isorropodon bigoti* Cosel and Salas 2001, *I. curtum* Cosel and Salas 2001, *I. megadesmus* Oliver et al. 2011, *I. striatum* (Thiele & Jaeckel, 1931) and *I. tenina* (Allen, 2001)), the Southern and North Western Atlantic basins (*I. elongatum* (Allen, 2001)), the Eastern Mediterranean Sea (*I. perplexum*, Sturany, 1986 the type species), the Norwegian Sea (*I. nyeggaensis* Krylova et al. 2011) and the Japan and Aleutian Trenches (*I. fossajaponicum* Okutani et al. 2000) (see for review Krylova et al. 2011; Oliver et al. 2011). The vertical range of *Isorropodon* is very broad, extending from 150 m to 6,809 m and the genus has been found in several habitats including oil drilling platform areas, soft mud, vegetal debris and mud volcanoes (Krylova and Sahling 2010; Krylova et al. 2011). Transmission electron micrographs of *Isorropodon perplexum* gill tissue showed the presence of bacteriocytes enclosing sulphur-oxidising bacteria (Olu-Le Roy et al. 2004). The thick and large gills of *I. bigoti* and *I. megadesmus* suggest that these species also harbour bacteria (Cosel and Salas 2001; Oliver et al. 2011) although these were not yet identified. In addition, carbon and sulphur stable isotope signatures (−31 and −1‰, respectively) in *Isorropodon perplexum* tissues indicate that thiotrophy may significantly contribute to their nutrition (Olu-Le Roy et al. 2004; Carlier et al. 2010). The only available molecular data for symbiotic bacteria are from the abundant endosymbionts occurring in the gills of *I. fossajaponicum* (initially ascribed to the genus *Calyptogena*) from the Japan Trench (Okutani et al. 2000; Fujiwara et al. 2000). Recently, we recovered a few specimens of three *Isorropodon* morphospecies from the Eastern Mediterranean, Gulf of Cadiz and Gulf of Guinea cold seeps. In order to verify the monophyly of *Isorropodon*, we sequenced cytochrome oxidase I (COI), a molecular marker previously used in vesicomyid studies (Baco et al. 1999; Peek et al. 1997; Goffredi et al. 2003; Decker et al. 2012). In order to characterize potential bacterial partners associated with *Isorropodon* species, we used comparative 16S rRNA gene sequence analysis and fluorescence in situ hybridization (FISH). Because 16S rRNA displays notably low variation among vesicomyid symbionts (Stewart et al. 2008, 2009), we also sequenced fragments of the 23S rRNA- and adenosine 5'-phosphosulfate (APS) reductase-encoding genes, to better

assess the level of relatedness to other vesicomyid symbionts. This work presents the first molecular characterization of Atlantic *Isorropodon* vesicomyids and their associated symbionts. Because of their transoceanic and vertical distribution, *Isorropodon* clams may provide good models for further studies of host/symbiont biogeography and evolution.

2 Material and methods

2.1 Specimens


The specimens analysed belong to three *Isorropodon* morphospecies from three distinct sites: *Isorropodon bigoti* (Gulf of Guinea), *Isorropodon megadesmus* (Gulf of Cadiz) and *Isorropodon perplexum* (Eastern Mediterranean) (Fig. 1a). Three specimens of *Isorropodon bigoti* were collected at cold seeps on a pockmark at the Guinness Site (Gulf of Guinea, cruise WACS, ROV dive 433, 1°34.65'S, 8°32.91'E, 582 m depth). Morphologically similar to *I. perplexum*, *I. bigoti* is distinguished by its larger size, the much less visible and more spaced growth lines and the more conspicuous wavy pattern of growth (Cosel and Salas 2001). This species was initially collected in an oil field at 150 m off Congo and deeper near Mauritania at 900–1,200 m. It was further sampled in a methane hydrate area in the Congo basin (500–800 m depth), (Cosel and Olu 2009). A single specimen of *I. megadesmus* was collected at the Darwin Mud volcano (Gulf of Cadiz, cruise TTR17, station AT664Gr, 35°23.52'N, 07°11.48'W, 1,128 m depth). This species is characterized by its large ligament that rises well above the dorsal margin of the shell and extends posteriorly beyond the nymph to fill the escutcheon and was previously reported only for Captain Arutyunov mud volcano (Oliver et al. 2011). Three specimens of *I. perplexum* were collected at the Amsterdam mud volcano (Eastern Mediterranean, Cruise MSM13/4, Station MSM13/994-1, 35°20.06'N, 30°16.17'W, 2,024 m depth). Shells from this species are thin-walled, ovate to elliptical in outline, with prosogyrate beaks. The escutcheon is present and lunular incision indistinct or missing (Cosel and Salas 2001).

The specimens were dissected immediately following collection and their gills were preserved for symbiont characterization using molecular and fluorescent in situ hybridization. Shells were kept for morphological comparison with documented species.

2.2 DNA extraction, polymerase chain reaction, and sequencing

Total DNA was extracted from ethanol-preserved gill tissue of all specimens using the DNeasy Blood and Tissue Kit (QIAGEN, CA, USA). A fragment of mtCOI and of three bacterial genes (16S ribosomal RNA (rRNA), 23S rRNA

Fig. 1 *Isorropodon* species of this study. Map with sampling locations (a). ML phylogenetic tree based on the mitochondrial COI gene of vesicomids (515 bp) (b). Scale bar represents estimated 5 % sequence substitution. Bootstrap support values over 50 % are shown. Taxonomy according to the WORMS database as of 01/03/2012 (GENBANK name also included; in grey). Species studied herein are represented in red, EM: Eastern Mediterranean; GoC: Gulf of Cadiz and GoG: Gulf of Guinea



and APS reductase) were amplified from all specimens (see Table 1 for PCR details).

For mtCOI, 23S rRNA and APS reductase amplification, products were purified using a gel extraction kit (QIAGEN) and were sequenced at GATC Biotech (Conzanz, Germany). For each species, a 16S rRNA gene library was

constructed from three independent PCR products of each individual clam, that were pooled, cleaned using a QIAquick Kit (QIAGEN) and cloned (TA™ cloning kit with pCR® TOPO 2.1. vector, Invitrogen, CA, USA). Between 14 and 29 positive clones were selected from each specimen and full-length inserts sequenced. Sequences were checked

Table 1 PCR conditions and primers used in this study DS: amplification products were purified and sequenced directly. GL: a gene library was constructed

Gene	Gene-length	Primers (5' > 3')	PCR conditions	Reference	
COI	Mitochondrial gene~560 nt	VesLCO -TTAATAGGAA CTGCTTTTAG VesHCO -TCACCAAACC AGCAGGATC	35 cycles of 30 s@94 °C, 60 s@45 °C, and 75 s@ 72 °C	Peek et al. 1997	DS
16S	Small subunit rRNA~1500nt	27 F - AGAGTTTGATC (AC)TGGCTAG 1492R - ACGG(CT)ACCTT GTTACGACTT	24 cycles of 45 s@94 °C, 45 s@45 °C, and 105 s@ 72 °C	Lane 1991	GL
23S	Large subunit rRNA~900 nt	3505 F - GACCGTCAGCTAA GGTCCCAA 4761R -CCAGTCAAAC TACCCAC CATG	35 cycles of 45 s@94 °C, 45 s@53 °C, and 2 min@ 72 °C	Stewart et al. 2009	DS
APS	adenosine 5'-phosphosulfate reductase - ~390nt	AprA-1-FW - TGGCAGATCATGA TYMAYGG Apar1-5-RV - GCGCCAACY GGRCRCA'	32 cycles of 1 min@92 °C, 1 min@58 °C, and 1 min@ 72 °C	Meyer and Kuever 2007	DS

for chimaeras with Chimera Check Ribosomal Database Project II (<http://rdp.cme.msu.edu/>, Cole et al. 2003), and most similar sequences were identified using Blast (<http://www.ncbi.nlm.nih.gov>) (Altschul et al. 1997)

2.3 Phylogenetic analysis

Sequence alignments including Blast hits and representative sequences were generated for the mtCOI-, 16S rRNA-, the 23S rRNA- and the APS reductase-encoding genes using ClustalX and edited manually (Thompson et al. 1997). Phylogenetic relationships were reconstructed using Maximum Likelihood (ML) with the PHYLIP package (Felsenstein 1993). ML used a transition/transversion ratio of two, empirical base frequencies, no rate variation among sites, and randomized input of sequences. Robustness of clades was evaluated by 100 ML bootstrap replicates. For proteins (APS) ML were analysed using the PROML function with the JTT model for probabilities of change between aminoacids.

16S rRNA was further analysed using Bayesian analysis with Mr. Bayes under a GTR model and Gamma rates run for $4 \cdot 10^6$ generations (discarding 10 % samples as burn-in) (Ronquist and Huelsenbeck 2003). Robustness of clades of bacterial 16S rRNA gene sequences was evaluated by 1,000 ML bootstrap replicates and computation of Bayesian posterior probabilities.

2.4 Statistical analysis

Correlation between hosts and symbiont genetic distances, and geographical distances were tested using the Mantel test (PAST, Hammer et al. 2001). Two geographical scenarios were investigated: (1) Great Arc distances between pairs of locations were estimated from the geographical coordinates; and (2) Minimum oceanic distances around continents and islands were estimated using GOOGLE EARTH v.5. The shortest paths between all pairs of locations were traced around present-day continental shelf margins that intervened between locations (Zielinski et al. 2009).

2.5 Fluorescence in situ hybridization

Gill tissue was fixed on board using 4 % formaldehyde in filtered seawater for 2 h at 4 °C, two rinses in filtered seawater and dehydration in increasing ethanol series (50, 80 and 100 %, 10 min each). Gills were embedded in PEG distearate: 1-hexadecanol (9:1) wax and cut into 7- μ m-thick sections using a microtome. Sections were hybridized as previously described using Cy3- and Cy5 labelled probes and a negative control (Duperron et al. 2007). Probe Gam42 (5'- GCCTCCCACATCGTTT-3') that targets the 23S

rRNA (large subunit) was used to highlight the presence of *Gammaproteobacteria*, and probe ImedT2-193 (5'-TAGAGGCCTCCTTTA-3') that targets the 16S rRNA (small subunit), originally designed for mussel symbionts, was also used here since mussel symbionts are closely related with vesicomid symbionts. All probes were tested at 30 % formamide. Hybridized sections were observed under an Olympus BX61 epifluorescence microscope (Olympus, Japan).

3 Results and discussion

3.1 Host characterization

The 560 bp fragments of mtCOI amplified from all specimens of the three *Isorropodon* morphospecies displayed 90 % similarity with the mtCOI sequence from *Calyptogena nautilei* (AB110759). COI from *I. bigoti* (Gulf of Guinea, sequences identical for the three specimens) and *I. megadesmus* (Gulf of Cadiz) presented 3.4 % sequence divergence whereas the sequences of *I. bigoti* and *I. perplexum* (Eastern Mediterranean, sequences identical for the three specimens) presented 7 % divergence. The same level of divergence (7.5 %) occurred between COI from *I. megadesmus* and *I. perplexum*. Phylogenetic analysis indicated that these three *Isorropodon* morphospecies formed a monophyletic group (Fig. 1b) related to *Calyptogena nautilei*, but with a low bootstrap support. COI genetic distances did not correlate with geographic distances (Mantel Test, $p > 0.6$). The two Atlantic species presented higher sequence similarity compared with *I. perplexum* suggesting the occurrence of a biogeographical barrier. Previous studies on fishes and crustaceans have shown the effect of three putative barriers to gene flow between the Atlantic and the Mediterranean: Strait of Gibraltar, Almeria–Oran Front and Ibiza Channel (Garcia-Merchan et al. 2012). The level of divergence among COI sequences from the three *Isorropodon* species is within the range of values documented between distinct populations of the species *Archivesica gigas* which may, in fact, be a species complex (Osborn 2001). Goffredi et al. 2003 study on the “*pacifica/lepta*” species documented divergence levels between 0.2 and 2.8 % (intraspecific) and between 3.9 and 10.3 % (interspecific), levels that were consistent with the ones reported for other described species of vesicomids (Peek et al. 1997, 2000). Baco et al. (1999) and Kojima et al. (2006) documented much lower intra-specific levels of COI variations (from 0.1 to 1.8 %) in *Calyptogena kilmeri*, *Calyptogena elongata* and *Calyptogena solidissima*. The level of divergence reported here is congruent with inter-specific, rather than intra-specific, variations. Moreover, the three *Isorropodon* morphospecies displayed distinctive morphological features and we can

reasonably assume at this stage that they actually represent distinct species.

Overall the Vesicomidae phylogenetic tree based on COI data displayed low resolution between the various lineages, and genera did not appear monophyletic. For example, *Isorropodon fossajaponicum* (initially described as *Calypptogena fossajaponica*) (Okutani et al. 2000), did not branch within the *Isorropodon* clade (Fig. 1). Since *I. perplexum* is the type species of the genus, the name should be restricted to the group documented here, but this emphasizes the need for a revision of the whole family, including the use of additional molecular markers. Indeed, because genera were initially poorly defined in Vesicomidae, recent work has attempted to clarify the taxonomy of the family (see Krylova and Sahling 2010 for review). However the taxonomic and phylogenetic study of this group is still in progress, and the incongruence between clades and named genera is fairly evident from our, as well as from other authors' phylogenies (Peek et al. 1997; Goffredi et al. 2003; Kojima et al. 2004; Decker et al. 2012).

3.2 Bacterial symbioses in *Isorropodon*

A total of 130 clones were analysed from 16S rRNA-based libraries: 24 from *I. megadesmus*, 48 from the three specimens of *I. perplexum*, and 58 from the three specimens of *I. bigoti*. A single phylotype was recovered within each of the three species. Sequences for the three species differed from each other by 1.4 to 3 % (Table 2), and displayed over 98 % sequence similarity with other vesicomid symbionts. Similar levels of divergence (1.7 to 2.8 %) were also observed among the 23S rRNA-encoding gene sequences. APS reductase (encoding a key enzyme of sulphur metabolism) nucleic acid sequences were identical for *I. megadesmus* and *I. perplexum*, and both differed by 3.6 % from that of *I. bigoti* (Table 2).

Table 2 Similarity matrix between symbiont sequences from the three *Isorropodon* species. Genes encoding 16S rRNA, 23S rRNA and APS are presented

	Host name	<i>I. bigoti</i>	<i>I. megadesmus</i>	<i>I. perplexum</i>
16S	<i>I. bigoti</i>	1	0.980	0.970
	<i>I. megadesmus</i>	–	1	0.986
	<i>I. perplexum</i>	–	–	1
23S	<i>I. bigoti</i>	1	0.981	0.972
	<i>I. megadesmus</i>	–	1	0.983
	<i>I. perplexum</i>	–	–	1
APS	<i>I. bigoti</i>	1	0.964	0.964
	<i>I. megadesmus</i>	–	1	1
	<i>I. perplexum</i>	–	–	1


In the 16S rRNA- (Fig. 2a), the 23S rRNA- (Fig. 3a) and the APS-reductase-based trees (Fig. 3b), *Isorropodon* sequences clustered with other vesicomid associated symbionts. Trees based on each of the three genes indicated monophyly of the three *Isorropodon*-associated bacterial sequences, although with low to moderate bootstrap support (54, 88 and 67, respectively). For all genes, genetic distances between symbionts were not correlated both with Great Arc and Minimum oceanic distances (Mantel tests, $p > 0.6$).

Gill tissue of all specimens of the three species hybridized successfully with the *Gammaproteobacteria*-specific probe Gam42, and with the probe Imed-T2-193, initially designed for mussel-associated thiotrophs but also targeting vesicomid symbionts due to closeness between these phylotypes (Duperron et al. 2008). Positive signals were located within a single epithelial cell type occurring in the lateral zone of gill filaments (known as bacteriocytes, Fisher 1990). Signals occupied most of the volume within bacteriocytes indicating very dense symbiont populations in all specimens (Fig. 2b and c). Overall, the smaller *Isorropodon* species display symbioses resembling those documented in larger vesicomid species, with dense bacterial populations in the gills. The identification of a sulphur metabolism (APS reductase), the phylogenetic clustering of all vesicomid symbionts, and the presence of sulphide in the clam's habitat all suggest these symbionts oxidize sulphide like those of larger vesicomids. Furthermore, the tissue $\delta^{13}\text{C}$ values of -31.1‰ reported for *Isorropodon perplexum* (Olu-Le Roy et al. 2004) and between -33.1 and -32.7‰ for *Isorropodon bigoti* suggest that thioautotrophy contributes significantly to the nutrition of the hosts.

3.3 Host/symbiont relationships and the evolution of *Isorropodon*

The monophyly of symbiont sequences mirrors that of their *Isorropodon* hosts. In the host tree, *I. megadesmus* and *I. bigoti* are more closely related, but with low bootstrap support, suggesting that COI does not resolve well within-group relationships. Altogether our results do not contradict the hypothesis of maternal inheritance and host/symbiont co-speciation postulated for other Vesicomidae (Peek et al. 1998; Stewart et al. 2008). However, vesicomid symbionts display highly similar 16S rRNA sequences (range: 91.9 to 99.9 %; mean: 97.0 ± 1.4 %; $n=26$ sequences (23 retrieved from Genbank and three from this study)), which does not allow to resolve symbiont phylogeny properly. Although 23S rRNA is often considered a more variable marker (Stewart et al. 2009), it is only slightly more variable among the 10 available vesicomid symbiont sequences (range: 91.1 to 99.6 %; mean: 94.9 ± 2.4 %; $n=10$ sequences (7 retrieved from Genbank and three from this study)), and additional markers will be necessary to settle the issue of


Fig. 2 Phylogenetic tree based on 16S rRNA sequences (1,267 positions in the alignment). Scale bar represents estimated 2 % sequence substitution. Values at nodes indicate posterior probabilities (Bayesian/bootstrap values (ML) over 50 % are shown). Sequences in *red* are from this study. ML and Bayesian analysis yielded identical tree topologies. Host taxonomy according to the WORMS database (GENBANK name also included; in *grey*). FISH hybridization on *Isorropodon bigoti* (**b**) and *Isorropodon megadesmus* gill transverse sections (**c**) with the Gam-42 (stained in *green*) and ImedT2-193 (stained in *red*) with an overlay result (*yellow*). No signal was observed from the ciliated zones, whereas all bacteriocytes from the lateral zones displayed dense FISH signal (in *yellow*), indicating bacteriocytes filled with bacteria. Nuclei of vesicomid cells, which are counterstained with DAPI, appear *blue*



co-speciation. Other vesicomid species are found in the Eastern Atlantic, but *I. perplexum* is to our knowledge the only species documented in the Mediterranean. Larger vesicomids tentatively attributed to the genus *Calyptogena*

were sampled in the Miocene fossil record, together with large *Bathymodiolus* type mytilids and lucinids (Taviani 1994). Because of episodes such as the Messinian Salinity crisis (5.96–5.33 MYA, Duggen et al. 2003), it can be

Fig. 3 Phylogenetic trees based on a fragment of the 23S rRNA encoding gene (796 positions analysed) (**a**) and on a fragment of the gene-encoding alpha subunit of APS reductase (125 amino acid positions analysed) (**b**). Scale bars represent estimated 5 % sequence substitution. For both trees, robustness of clades was evaluated by 100 ML bootstrap replicates. Bootstrap support values over 50 % are shown. Sequences in red are from this study. Host taxonomy according to the WORMS database (GENBANK name also included; in grey)



hypothesized that *I. perplexum* colonized the Mediterranean from Eastern Atlantic ancestors recently. The Strait of Gibraltar may act as an important limitation to connectivity between these two biogeographic regions which may explain the higher genetic divergence of Atlantic vs. Mediterranean species. A different trend is found in the symbionts, with genetic difference between *I. perplexum* and *I. megadesmus* symbionts than between Atlantic species. This suggests that symbiont transmission may not be strictly vertical in vesicomysids, and symbiont movement between hosts or

acquisition from the environment may contribute to incongruence between host and symbiont phylogenies. Most *Isorropodon* species were reported from the Atlantic Ocean (7 out of 9 species). This geographical distribution, and the wide vertical range (from 150 to 6,809 m) (Krylova et al. 2011), in combination with the high level of adaptation to reducing environments, makes *Isorropodon* a prospective model to study biogeographical, ecological and depth-related evolutionary processes in hosts and symbionts.

Acknowledgments Thanks are due to the chief-scientists and crew of the several cruises where specimens were collected: TTR17 Leg 2 (Gulf of Cadiz, RV *Logatchev*): Luis Pinheiro (Departamento de Geociências, Universidade de Aveiro) and Michael Ivanov (Moscow State University); MSM13/4 (Eastern Mediterranean; RV *MS Merian*, ROV *Quest*): Antje Boetius and Frank Wenzhofer (Max Planck Institute); WACS (Gulf of Guinea; RV *Pourquoi Pas?*, ROV *Victor 6000*). The authors are indebted to Nelly Leger of the team AMEX (UPMC) for her help in the lab. Finally we are grateful to the two anonymous reviewers and the editor Dr. Charles Fisher for their constructive and helpful comments.

This research was supported by the European Commission Seventh Framework Programme (FP7/2007-2013) under the HERMIONE project contract 226354, French ANR DeepOases, GDRE—DIWOOD and CHEMOCO ESF EURODEEP (FCT; EURODEEP/0001/2007). CFR was also supported by a post-doctoral fellowship (SFRH/BPD/64154/2009) from the Fundação para a Ciência e a Tecnologia (FCT, Portugal).

References

- Allen JA (2001) The family Kelliellidae (Bivalvia: Heterodonta) from the deep Atlantic and its relationship with the family Vesicomidae. *Zool J Linn Soc-Lond* 131:199–226
- Altschul SF, Madden TL, Schaeffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25(17):3389–3402
- Amano K, Kiel S (2007) Fossil vesicomid bivalves from the North Pacific region. *Veliger* 49:270–293
- Baco AR, Smith CR, Peek AS, Roderick GK, Vrijenhoek RC (1999) The phylogenetic relationships of whale-fall vesicomid clams based on mitochondrial COI DNA sequences. *Mar Ecol Prog Ser* 182:137–147
- Carlier A, Ritt B, Rodrigues CF, Sarrazin J, Olu K, Grall J, Clavier J (2010) Heterogeneous energetic pathways and carbon sources on deep eastern Mediterranean cold seep communities. *Mar Biol* 157(11):2545–2565
- Cary SC, Giovannoni SJ (1993) Transovarial inheritance of endosymbiotic bacteria in clams inhabiting deep-sea hydrothermal vents and cold seeps. *Proc Natl Acad Sci* 90:5695–5699
- Cole JR, Chai B, Marsh TL, Farris RJ, Wang Q, Kulam SA, Chandra S, McGarrell DM, Schmiedl TM, Garrity GM, Tiedje JM (2003) The Ribosomal Database Project (RDP-II): previewing a new autoaligner that allows regular updates and the new prokaryotic taxonomy. *Nucleic Acids Res* 31(1):442–443
- Cosel RV, Olu K (2009) Large Vesicomidae (Mollusca: Bivalvia) from cold seeps in the gulf of Guinea off the coasts of Gabon, Congo and northern Angola. *Deep-Sea Res II* 56:2350–2379
- Cosel RV, Salas C (2001) Vesicomidae (Mollusca: Bivalvia) of the genera *Vesicomya*, *Waisiuconcha*, *Isorropodon* and *Callogonia* in the eastern Atlantic and the Mediterranean. *Sarsia* 86(4–5):333–366
- Decker C, Olu K, Cunha RL, Arnaud-Haond S (2012) Phylogenetic relationships among cold-seep vesicomid bivalves from the Gulf of Guinea and their western Atlantic and Pacific counterparts. *PLoS One* 7(4):e33359. doi:10.1371/journal.pone.0033359
- Dubilier N, Bergin C, Lott C (2008) Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nat Rev Microbiol* 5:725–740
- Duggen S, Hoernie K, van den Bogaard P, Rupke L, Phipps Morgan J (2003) Deep roots of the Messinian salinity crisis. *Nature* 422:602–606
- Duperron S, Fiala-Medioni A, Caprais JC, Olu K, Sibuet M (2007) Evidence for chemoautotrophic symbiosis in a Mediterranean cold seep clam (Bivalvia: Lucinidae): comparative sequence analysis of bacterial 16S rRNA, APS reductase and RubisCO genes. *FEMS Microbiol Ecol* 59:64–70
- Duperron S, Halary S, Lorion J, Sibuet M, Gail F (2008) Unexpected co-occurrence of six bacterial symbionts in the gills of the cold seep mussel *Idas* sp. (Bivalvia: Mytilidae). *Environ Microbiol* 10(2):433–445
- Felsenstein J (1993) PHYLIP (phylogeny inference package) 3.57c edn. Department of Genetics, University of Washington, Seattle
- Fisher CR (1990) Chemoautotrophic and methanotrophic symbioses in marine invertebrates. *Rev Aquat Sci* 2(3,4):399–436
- Fujiwara Y, Kojima S, Mizota C, Maki Y, Fukikura K (2000) Phylogenetic characterization of the endosymbionts of the deepest-living vesicomid clam, *Calyptogena fossajaponica*, from the Japan Trench. *Venus Jap Jour Mal* 59(4):307–316
- Garcia-Merchan VH, Robainas-Barcia A, Abelló P, Macpherson E, Palero F, Garcia-Rodriguez M, Gil de Sola L, Pascual M (2012) Phylogeographic patterns of decapod crustaceans at the Atlantic-Mediterranean transition. *Mol Phylogenet Evol* 62:664–672
- Goffredi SK, Hurtado LA, Hallam S, Vrijenhoek RC (2003) Evolutionary relationships of deep-sea vent and cold seep clams (Mollusca: Vesicomidae) of the “pacifica/lepta” species complex. *Mar Biol* 142:311–320
- Hammer T, Harper DAT, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontol Electron* 4:1–9
- Harada M, Yoshida T, Kuwahara H, Shimamura S, Takaki Y, Kato C, Miwa T, Miyake H, Maruyama T (2009) Expression of genes for sulfur oxidation in the intracellular chemoautotrophic symbiont of the deep-sea bivalve *Calyptogena okutanii*. *Extremophiles* 13(6):895–903
- Hurtado LA, Mateos M, Lutz RA, Vrijenhoek RC (2003) Coupling of bacterial endosymbiont and host mitochondrial genomes in the hydrothermal vent clam *Calyptogena magnifica*. *Appl Environ Microbiol* 69(4):2058–2064
- Imhoff JF, Sahling H, Suling J, Kath T (2003) 16S rDNA-based phylogeny of sulphur-oxidising bacterial endosymbionts in marine bivalves from cold-seeps environments. *Mar Ecol Prog Ser* 249:39–51
- Kim Y, Yasuda M, Yamagishi A, Oshima T, Ohta S (1995) Characterization of the endosymbiont of a deep-sea bivalve *Calyptogena soyoae*. *Appl Environ Microbiol* 61(2):823–827
- Kojima S, Kobayashi T, Hashimoto J, Ohta S (1995) RFLP analysis of a mitochondrial gene, Cytochrome Oxidase I (COI) of three species of the genus *Calyptogena* around Japan. *J Oceanogr* 51:257–259
- Kojima S, Fujikura K, Okutani T (2004) Multiple trans-Pacific migrations of deep-sea vent/seep endemic bivalves in the family Vesicomidae. *Mol Phylogenet Evol* 32:396–406
- Kojima S, Tsuchida E, Numanami H, Fujikura K, Okutani T (2006) Synonymy of *Calyptogena solidissima* with *Calyptogena kawamurai* (Bivalvia: Vesicomidae) and its population structure revealed by mitochondrial DNA sequences. *Zool Sci* 23:835–842
- Krylova EM, Sahling H (2010) Vesicomidae (Bivalvia): current taxonomy and distribution. *PLoS One* 5(4):e9957
- Krylova EM, Gebruk AY, Portnova DA, Todt C, Haflidason H (2011) New species of the genus *Isorropodon* (Bivalvia: Vesicomidae: Pliocardiinae) from cold methane seeps at Nyegga (Norwegian Sea, Voring Plateau, Storrega slide). *J Mar Biol Assoc UK* 91(5):1135–1144
- Kuwahara H, Takaki Y, Yoshida T, Shimamura S, Takishita K, Reimer J, Kato C, Maruyama T (2008) Reductive genome evolution in chemoautotrophic intracellular symbionts of deep-sea *Calyptogena* clams. *Extremophiles* 12(3):365–374

- Lane DJ (1991) 16S/(23S rRNA sequencing. In: Stakebrandt E, Goodfellow M (eds) Nucleic acid techniques in bacterial systematics. Wiley, New York, pp 115–175
- Meyer B, Kuever J (2007) Molecular analysis of the distribution and phylogeny of dissimilatory adenosine-59-phosphosulfate reductase-encoding genes (*aprBA*) among sulfuroxidizing prokaryotes. *Microbiol* 153:3478–3498
- Okutani T, Fujikura K, Kojima S (2000) New taxa and review of vesicomimid bivalves collected from the northwest Pacific by deep-sea research systems of Japan Marine Science & Technology Center. *Venus Jap Jour Mal* 59:83–101
- Oliver PG, Rodrigues CF, Cunha MR (2011) Chemosymbiotic bivalves from the mud volcanoes of the Gulf of Cadiz, NE Atlantic, with descriptions of new species of Solemyidae, Lucinidae and Vesicomimidae. *Zookeys* 113:1–38
- Olu-Le Roy K, Sibuet M, Fiala-Médioni A, Gofas S, Salas C, Mariotti A, Foucher JP, Woodside J (2004) Cold seep communities in the deep eastern Mediterranean Sea: composition, symbiosis and spatial distribution on mud volcanoes. *Deep-Sea Res I* 51:1915–1936
- Osborn M (2001) Molecular systematics of the vesicomimid clams: *Calyptogena kilmeri* and *Vesicomima gigas*. MBARI report. pp 7
- Peek AS, Gustafson RG, Lutz RA, Vrijenhoek RC (1997) Evolutionary relationships of deep-sea hydrothermal vent and cold-water clams (Bivalvia: Vesicomimidae) results from the mitochondrial cytochrome oxidase subunit I. *Mar Biol* 130:151–161
- Peek AS, Feldman RA, Lutz RA, Vrijenhoek RC (1998) Cospeciation of chemoautotrophic bacteria and deep-sea clams. *Proc Natl Acad Sci* 95:9962–9966
- Peek AS, Gaut BS, Feldman RA, Barry JP, Kochevar RE, Lutz RA, Vrijenhoek RC (2000) Neutral and Nonneutral Mitochondrial Genetic Variation in Deep-Sea Clams from the Family Vesicomimidae. *J Mol Evol* 50 (2):141–153
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(2):1572–1574
- Stewart FJ, Cavanaugh CM (2009) Pyrosequencing analysis of endosymbiont population structure: co-occurrence of divergent symbiont lineages in a single vesicomimid host clam. *Environ Microbiol* 11(8):2136–2147
- Stewart FJ, Young CR, Cavanaugh CM (2008) Lateral symbiont acquisition in a maternally transmitted chemosynthetic clam endosymbiosis. *Mol Biol Evol* 25(4):673–687
- Stewart FJ, Young CR, Cavanaugh CM (2009) Evidence for homologous recombination in intracellular chemosynthetic clam. *Mol Biol Evol* 26(6):1391–1404
- Sturany R (1896). Zoologische Ergebnisse VII. Mollusken I (Prosobranchier und Opisthobranchier; Scaphopoden; Lamellibranchier) gesammelt von S.M. Schiff “Pola” 1890-1894. Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftlichen Classe 63:1–36, pl.1–2
- Taviani M (1994) The “calcarei a Lucina” macrofauna reconsidered: deep-sea faunal oases from Miocene-age cold vents in the Romagna Apennine, Italy. *Geo-Mar Lett* 14:185–191
- Thiele J, Jaekel S (1931) Muscheln der Deutschen Tiefsee Expedition. *Wissenschaftliche Ergebnisse der Deutschen Tiefsee Expedition 1898–1899* 21:159–268
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The Clustal X-Windows interface: flexible strategies for multiple sequence alignments aided by quality analysis tools. *Nucleic Acids Res* 25(24):4876–4882
- Zielinski FU, Pernthaler A, Duperron S, Raggi L, Giere O, Borowski C, Dubilier N (2009) Widespread occurrence of an intranuclear bacterial parasite in vent and seep bathymodiolin mussels. *Environ Microbiol* 11(5):1150–1167