

**Microwave Synthesis of
cisDichlorotetrakis(dimethylsulfoxide)ruthenium(II)**

HARVEY, Amanda <<http://orcid.org/0000-0003-3720-2602>>, DRAGANJAC, Mark, CHIU, Sheng, SNELL, R and BENJAMIN, Ellis

Available from Sheffield Hallam University Research Archive (SHURA) at:
<http://shura.shu.ac.uk/19160/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

HARVEY, Amanda, DRAGANJAC, Mark, CHIU, Sheng, SNELL, R and BENJAMIN, Ellis (2009). Microwave Synthesis of cisDichlorotetrakis(dimethylsulfoxide)ruthenium(II). *Journal of the Arkansas Academy of Science*, 63 (26), p. 185.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

2009

Microwave Synthesis of cis-Dichlorotetrakis(dimethylsulfoxide)ruthenium(II)

A. Harvey

Arkansas State University

Mark Draganjac

Arkansas State University, mdraganj@astate.edu

S. Chui

Arkansas State University

R. Snell

Arkansas State University

E. Benjamin

Arkansas State University

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

 Part of the [Physical Chemistry Commons](#)

Recommended Citation

Harvey, A.; Draganjac, Mark; Chui, S.; Snell, R.; and Benjamin, E. (2009) "Microwave Synthesis of cis-Dichlorotetrakis(dimethylsulfoxide)ruthenium(II)," *Journal of the Arkansas Academy of Science*: Vol. 63 , Article 26. Available at: <http://scholarworks.uark.edu/jaas/vol63/iss1/26>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This General Note is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Microwave Synthesis of *cis*-Dichlorotetrakis(dimethylsulfoxide)ruthenium(II)

A. Harvey, M. Draganjac¹, S. Chui, R. Snell, and E. Benjamin

Department of Chemistry and Physics, Arkansas State University, State University, AR 72467

¹ Correspondence: mdraganj@astate.edu

The *cis*-RuCl₂(dms_o)₄ compound has been shown to be a versatile starting reagent for the preparation of numerous Ru complexes (Evans et al. 1973, Alessio et al. 1991, Landgrafe and Sheldrick 1994). *cis*-RuCl₂(dms_o)₄ is prepared by refluxing RuCl₃·xH₂O in DMSO for 5 min followed by concentration to half volume and subsequent precipitation with acetone (Evans et al. 1973). Due to the high boiling point of DMSO (189 °C), actual reaction time approaches 3 hr. The reduction in volume is also critical, as too little reduction leads to reduced yields. The difficulties in the synthesis of *cis*-RuCl₂(dms_o)₄ and the length of reaction time led us to explore the use of microwave technologies for improving the preparation of the title compound.

The starting materials, RuCl₃·xH₂O and dimethyl sulfoxide, were purchased from Aldrich Chemical Company. All reagents were used as purchased, without further purification. An Agilent 8453 UV-visible Spectroscopy system and a Midac IR spectrophotometer were used for the analysis of the compounds. The microwave synthesis was performed using a CEM Discover microwave reactor.

For the microwave reaction, 0.10 g (0.38 mmol) RuCl₃·xH₂O was added to 2 mL DMSO in a microwave reaction tube under N₂ or Ar and heated to 135 °C for 10 min. Upon standing, a yellow precipitate formed and was isolated by filtration. The product was washed with acetone and air-dried, yield = 49%, melting point = 193 °C. The visible and infrared spectra for our product matched those reported in the literature (Evans et al. 1973, Alessio et al. 1988). Reaction times can be decreased to three min. by increasing the surface area inside the microwave reaction tube.

Though the yield from the microwave reaction is lower than the reported literature yield (Evans et al. 1973), decrease in reaction times, lower energy expenditures and reduction in waste due to the use of smaller solvent volumes makes the microwave synthesis a viable alternative to the conventional preparative method for *cis*-RuCl₂(dms_o)₄.

Acknowledgments

The authors thank Dr. William Burns for aid in obtaining IR spectra and the Department of Chemistry and Physics for their support of this project.

Literature Cited

- Alessio E, G Mestroni, G Nardin, WM Attia, M Calligaris, G Sava, and S Zorzet.** 1988. *cis*- and *trans*-dihalotetrakis(dimethylsulfoxide) ruthenium(II) complexes (RuX₂(DMSO)₄; X = Cl, Br): synthesis, structure, and antitumor activity. *Inorganic Chemistry* 27(23):4099-106.
- Alessio E, G Balducci, M Calligaris, G Costa, and WM Attia.** 1991. Synthesis, molecular structure, and chemical behavior of hydrogen *trans*-bis(dimethyl sulfoxide)-tetrachlororuthenate(III) and *mer*-trichloro-tris(dimethyl sulfoxide) ruthenium(III): the first fully characterized chloride-dimethyl sulfoxide-ruthenium(III) complexes. *Inorganic Chemistry* 30(4):609-18.
- Evans IP, A Spencer, and G Wilkinson.** 1973. Dichlorotetrakis(dimethylsulphoxide)-ruthenium(II) and its use as a source material for some new ruthenium(II) complexes. *J.C.S. Dalton Trans* 1973:204-9.
- Landgrafe C and WS Sheldrick.** 1994. Structure and reactions of the thioether half-sandwich ruthenium(II) complexes [Ru(MeCN)₃([9]aneS₃)] [CF₃SO₃]₂ and [Ru(MeCN)₂(PPh₃)([9]aneS₃)] [CF₃SO₃]₂ ([9]aneS₃ = 1,4,7-trithiacyclononane). *J.C.S. Dalton Trans* 1994:1885-93.