



The University of Bradford Institutional Repository

<http://bradscholars.brad.ac.uk>

This work is made available online in accordance with publisher policies. Please refer to the repository record for this item and our Policy Document available from the repository home page for further information.

To see the final version of this work please visit the publisher's website. Access to the published online version may require a subscription.

Link to publisher's version: <https://doi.org/10.1177/1533317515603500>

Citation: Barbosa A, Nolan M, Sousa L et al (2016) Effects of psychoeducational intervention for direct care workers caring for people with dementia: results from a 6-month follow-up study. *American Journal of Alzheimer's Disease & Other Dementias*® 31(2): 144-155.

Copyright statement: The final, definitive version of this paper has been published in *American Journal of Alzheimer's Disease & Other Dementias*®, vol 31/ issue 2 by SAGE Publications Ltd, All rights reserved. © 2015 The Authors.

Effects of a psycho-educational intervention for direct care workers caring for people with dementia: Results from a 6-month follow-up study

Ana Barbosa

MSc, Department of Health Sciences, University of Aveiro, Portugal. E-mail: anabarbosa@ua.pt

Mike Nolan

PhD, School of Nursing and Midwifery, University of Sheffield, Northern General Hospital, Sheffield, United Kingdom. E-mail: m.r.nolan@sheffield.ac.uk

Liliana Sousa

PhD, Department of Health Sciences, University of Aveiro; Center for Health Technology and Services Research (CINTESIS), University of Aveiro, Portugal. E-mail: lilianax@ua.pt

Alda Marques

PhD, School of Health Sciences, University of Aveiro; Center for Health Technology and Services Research (CINTESIS), University of Aveiro, Portugal. E-mail: amarques@ua.pt

Daniela Figueiredo

PhD, School of Health Sciences, University of Aveiro; Center for Health Technology and Services Research (CINTESIS), University of Aveiro, Portugal. E-mail: daniela.figueiredo@ua.pt

Word counts (abstract and text): 3924

Funding support: This work was supported by grants (grant numbers SFRH/BD/72460/2010 and RIPD/CIF/109464/2009) from the Foundation for Science and Technology (FCT).

Disclosure: The authors have reported no conflicts of interest

Correspondence address: Ana Barbosa, Department of Health Sciences, University of Aveiro, Campus Universitário de Santiago, Agra do Crasto - edifício 30, Aveiro, Portugal. E-mail: anabarbosa@ua.pt. Tel: +351 234 370 213. Fax: +351 234 401 597

Abstract

This study aimed to assess the effects of a psycho-educational intervention, designed to improve direct care workers' stress, burnout and job satisfaction and person-centered communicative behavior with people with dementia. A pretest-posttest control group design was conducted in four aged-care facilities. Two experimental facilities received a psycho-educational intervention; two control facilities received an education-only. Data were gathered from fifty three care workers at baseline, immediately and six months after the intervention, through self-administrated instruments and video-recorded morning care sessions.

The experimental group showed a significant decrease in care workers' burnout and a significant improvement in several communicative behaviors (e.g., involvement). Stress levels deteriorated at six months and no intervention effects were found for job satisfaction. The findings highlight the importance of providing care workers with both technical competences and tools for stress management as this might be associated with a reduction of their levels of exhaustion and improved communicative behaviors.

Keywords

Aged-care facilities, dementia, person-centered care, direct care workers, psycho-educational intervention

Introduction

People with dementia are one of the fastest growing groups of people living with long-term conditions. The number of people living with dementia worldwide in 2013 was estimated at 44.35 million, reaching 75.62 million in 2030 and 135.46 million in 2050 ¹. Along with these projections, there will be an associated increase in demand for long-term care mainly provided by aged-care facilities ¹.

In an aged-care facility, the bulk of care that residents receive is provided by direct care workers (DCWs) ². These workers are responsible for helping frail and disabled older adults carry out the most basic activities of daily life, such as bathing, dressing, toileting, and eating, during which they also provide the personal interaction that is essential to residents' quality of life and care ². However, stressors resulting from inadequate education and training in dementia care, high workload, interpersonal conflicts or lack of management support have been shown to be prevalent in DCWs' work ^{3,4}. Such stressors place DCWs at high risk of experiencing stress, burnout and job dissatisfaction which are known to create a disruption in the worker-resident relationship and hinder the delivery of quality care ^{3,5}.

The provision of education to DCWs has been long regarded as an essential component for improving the quality of dementia care ^{6,7}. This is often designed to improve DCWs' skills based on person-centered care (PCC) approaches, which have become synonymous with "best practice" ^{8,9}.

Person-centered dementia care has its roots in the work of Tom Kitwood ¹⁰, who was inspired by Carl Rogers and his client-centered counseling. Kitwood (1997), soon followed by Nolan et al. (2004) and their *relationship-centered care*, stressed the influence of interpersonal relationships as an essential aspect for understanding the dementia experience, theorizing that some of the deterioration seen in people with dementia was caused not only by the disease process itself, but also by how the person is treated. These authors emphasized the relational nature of PCC and the need to provide workers with the skills they need to enhance positive interactions (e.g., consult or validate) (labeled *positive person work* - PPW) and reduce negative

communicative behaviors (e.g., ignore or infantilization) that depersonalize experiences of the person with dementia (labeled *malignant social psychology* - MSP)¹⁰, thus creating an enriched environment of care⁸.

In notable randomized controlled studies, DCWs were provided with PCC based education-only interventions designed to reduce the residents' levels of agitation and aggression¹¹⁻¹³ and to enhance residents' engagement in daily activities¹⁴. Researchers have demonstrated the potential for these interventions to improve DCWs' knowledge and enhance their willingness to encourage residents' autonomy, independence and communication¹⁵; however their effects on stress, burnout or job satisfaction are modest, and often neither long-lasting or significant^{16,17}. This suggests that interventions are mainly focused on improving DCWs' knowledge and instrumental skills and are less concerned with their emotional and relational skills, which, despite the rhetoric of PCC are still undervalued. Providing DCWs with both technical competences and tools for stress and emotional management holds promise as a means of driving forward benefits for DCWs and care provision. This approach may better prepare DCWs to deal with their multifaceted and emotionally demanding job, potentially improving person-centered interactions, job satisfaction and wellbeing. However, to the authors' knowledge, no studies assessing the efficacy of psycho-educational interventions in the context of formal care have been conducted.

This study aimed to examine the immediate and the 6-month effects of a Person Centered Care based psycho-educational intervention (PCC-based PE intervention) targeted at DCWs caring for people with dementia in aged-care facilities. It was hypothesized that, compared with an education-only intervention (control group), an intervention offering both educational and emotional support would reduce DCWs' perceived stress, burnout and job dissatisfaction. It was also expected that, compared to the control group, the PCC-based PE intervention would decrease the frequency and duration of DCWs' MSP behaviors and enhance the frequency and duration of PPW behaviors.

Design

This experimental study used a pre-posttest control group design and was conducted in four aged-care residential facilities. The study was approved by an ethics committee [names deleted to maintain the integrity of the review process].

Data were collected at i) baseline assessment (T1), in all facilities 3 weeks prior to the intervention; ii) posttest (T2), 2 weeks after the end of the intervention and iii) follow-up (T3), 6 months after the intervention.

Procedure

The facilities of the local area where the study was conducted were stratified into groups by the staff/resident ratio and residents with dementia/total of residents' ratio. Then, two pairs of facilities of the same created group were approached by the research team and were given the opportunity to participate in the study. All four facilities agreed to participate and were randomly allocated to the experimental group - PE intervention - or control group - education-only intervention – using a random number generator. Randomization could not occur at the individual level due to possible treatment effects if the same facility functioned as both experimental and control sites. Study facilities were private, non-profit institutions of collective accommodation with more than 30 licensed beds and with a staff/resident ratio between 1:2 or 1:3.

Sample

The study sample includes DCWs (may be called under different names in different countries, for example, nursing aides or care assistants), who represent the largest component of the long-term care workforce and are responsible for helping frail and disabled older adults carry out the most intimate and basic activities of daily living. To be included in the study, DCWs had to be employed for at least two months (so adjustments to the residents and facility had been achieved) and provide morning personal care (i.e., period of time between 07am and 12am that

involved activities related to bathing, grooming, dressing and toileting) to people with a diagnosis of moderate to severe dementia. Temporary DCWs and trainees were excluded as it was not possible to ensure their participation until the end of the study. The identification of the eligible DCWs was supported by facility managers. Three DCWs were excluded from the study due to being temporary.

A meeting with the fifty-eight identified DCWs was scheduled to provide detailed information about the study and invite them to participate. They were informed about the voluntary nature of their participation and their anonymity and confidentiality were assured. All DCWs agreed to participate and their informed signed consent was obtained at the end of the meeting.

Twenty-seven DCWs received a psycho-educational intervention and 31 DCWs participated in the control group.

Of these, 53 DCWs completed all three rounds of data collection. Two dropouts occurred in the control group and 3 in the experimental group. The dropouts were due to DCWs' absence from work during the assessment periods, as a result of sick leave (n=2), vacation (n=2) or dismissal (n=1) (Diagram 1).

[Diagram 1]

The legal guardians of the identified residents were also contacted, informed about the study and asked to sign a written informed consent. From 51 residents with moderate-to-severe dementia, 47 participated (one legal guardian refused participation, one resident refused permanently to be assessed by video and two residents died before collecting any data).

Intervention

PCC-based PE intervention

The experimental facilities received a PCC-based PE intervention informed by: (1) relevant literature on PE approaches, PCC and dementia ^{11,18}; (2) findings from a previous pilot study conducted by the authors' research team [names deleted to maintain the integrity of the review process]; and (3) interviews with DCWs and managers about instrumental and emotional needs [names deleted to maintain the integrity of the review process]. The intervention included 8 weekly 90 minute sessions, coordinated by a gerontologist and a physical therapist with training and experience in PCC approaches and psycho-educational groups. Each session followed a similar format, organized into two components: education and support.

The education component aimed to provide DCWs with: i) principles to integrate PCC within the care routines (e.g. incorporate biographical knowledge in personal care); ii) basic knowledge about dementia; and iii) PCC-based interaction strategies, including motor stimulation (e.g. encourage the person to perform one task or a part of it) and multisensory stimulation (e.g. provide a gentle massage while washing resident's hair). In the 3 days following each PE session, the gerontologist and the physical therapist assisted DCWs individually during morning care, clarifying doubts and making suggestions to help them implement a more PCC approach.

The supportive component aimed to provide DCWs with coping strategies to manage work-related stress and prevent burnout (e.g., time-management, assertiveness and problem-solving). At the end of each supportive component, relaxation techniques, stretching and strengthening exercises were practiced. Detailed information about the intervention can be found elsewhere [names deleted to maintain the integrity of the review process] and is summarized in Table 1.

PCC-based education-only intervention

The control facilities received an education-only intervention. The coordination, length, order and content of the sessions were the same as the educational component of the PE intervention. It was the absence of the supportive component that distinguished both interventions. Each

participant was assisted during morning care by the same professionals, who helped DCWs to deliver a more PCC and clarified doubts that emerged from sessions.

[Table 1]

Measures

DCWs' perceived stress

DCWs' perceived stress was measured using the Portuguese version of the Perceived Stress Scale (PSS)¹⁹. The PSS is a 14-item self-report questionnaire rated on a 5-point Likert-type scale ranging from "never" (0) to "very often" (4). The items evaluate the degree to which individuals believe their life has been unpredictable, uncontrollable, and overloaded during the previous month. Higher scores correspond to higher degrees of perceived stress.

The Cronbach's alpha coefficient of the scale showed a score of $\alpha=0,88$. Scores for the criterion validity ranged between 0,4 and 0,8 and the examination of the factorial validity with the one-factor structure accounted for 43.96% of variance. Overall, the acceptable psychometric properties of the Portuguese version of the PSS are similar to those obtained in other versions²⁰.

DCWs' burnout

The 22-item Maslach Burnout Inventory (MBI) - Human Services Survey was used to assess the DCWs' experience of burnout²¹. The MBI is the most widely used instrument to assess burnout and is divided into three subscales: 8 items assess emotional exhaustion (EE), i.e. feelings of being emotionally exhausted by one's work; 5 items measure depersonalization (DP), i.e. the negative attitudes toward recipients' care or treatment; and 8 items assess personal accomplishment (PA), i.e., feelings of competence and successful achievement in work. The respondents are asked to report the frequency with which such feelings are experienced on a 7-point Likert-type scale, ranging from "never" (0) to "every day" (6). A combination of high scores on EE and DP, and a low score on PA, correspond to a high level of burnout. The Cronbach's alpha coefficient for the Portuguese version showed a score of $\alpha=0.75$ and

reliability coefficients of 0.80 for EE, 0.71 for DP and 0.70 for PA. The validity of the three-factor structure of the MBI was found to provide a reasonable fit to the data, explaining 43.4% of the total variance. The psychometric properties of the used version are consistent with the ones of previous studies²².

DCWs' job satisfaction

The short-form Minnesota Satisfaction Questionnaire (MSQ)²³ was used to assess DCWs' job satisfaction. It includes 20 items rated on a 5-point Likert scale ranging from "extremely dissatisfied" (1) to "extremely satisfied" (5). Item responses are summed or averaged to create a total score – the lower the score, the lower the level of job satisfaction. Besides a total score, the MSQ can also be scored for intrinsic and extrinsic satisfaction. The intrinsic subscale includes 6 items with scores ranging between 1 and 30 and refers to how people feel about the nature of the job tasks themselves. The extrinsic satisfaction subscale contains 8 items ranging from 8 to 40 and refers to how people feel about aspects of the work situation that are external to the job tasks. The psychometric properties of the used version of the MSQ are acceptable. High internal consistency was found for the global scale ($\alpha=0.93$), and for the intrinsic ($\alpha=0.88$) and extrinsic subscales ($\alpha=0.82$). These values are higher than the ones found for the original scale²⁴. Strong test-retest correlations were found (>0.80), which denotes adequate stability coefficients and corroborates the findings obtained in previous studies²⁴. The factorial analysis confirmed the adequacy of the two-factor structure of the MSQ, which explained 62.69% of the total variance.

DCWs' person-centered communication

In order to capture both DCWs' verbal and non-verbal communicative behaviors, video-recordings of morning care routines were used. Video-recordings were performed in the resident's bedroom; the moment DCWs entered the room was defined as the starting point and when they left the room as the ending point. Bathing was not recorded to assure privacy to the person with dementia. Some procedures were followed to minimize participants' or residents' reactivity (i.e., response during data collection that affects the natural course of behavior as a

result of being observed): i) prior to data collection, several video-recordings were performed in order to familiarize participants with the methodology; ii) DCWs were instructed to stop or remove the video camera if they noticed any resident's negative reaction caused by the device presence; and iii) once the cameras were placed on a tripod and adequately positioned, the researcher left the room so that a further source of disruption could be avoided.

To prevent random DCWs' communicative behaviors, DCWs were video-recorded thrice at each assessment point. In total, 474 morning care sessions were video-recorded. At baseline, 3 participants were only recorded twice as they were absent from work.

DCWs' communicative behavior was studied by analyzing the frequency and duration of a list of mutually exclusive behaviors (ethogram). The categories described in the Kitwood's dialectical framework¹⁰, relevant literature on staff's verbal and non-verbal communication^{25,26} and preliminary observations of the video recordings informed the construction of the ethogram. The final list comprised 18 verbal communicative behaviors (Table 2) and 8 non-verbal communicative behaviors (Table 3). One coder (1st author) rated the DCWs' communicative behaviors according to the ethogram using specialized software, Noldus Observer XT (version 11.0) (Noldus International Technology, Wageningen, Netherlands). The coder was previously trained to use the software.

[Table 2]

[Table 3]

Data Analysis

Repeated measures analysis of variance (ANOVA) were used to determine the existence of significant differences on DCWs' perceived stress, burnout, job satisfaction and person-centered interactions at three points in time. Specifically, a series of one between-subjects variable (experimental vs control) and one within-subjects variable (pretest, posttest, follow-up) repeated measures ANOVA was performed. This statistical technique was used to test intervention, time, and intervention by time interaction effects. *Partial eta squared* (η^2), which corresponds to the

Effect Size, was interpreted as small (≥ 0.05), medium (0.05-0.25), large (0.25-0.50) and very large (≥ 0.50)²⁷.

All variables were previously tested for normality. The level of significance was set at 0.05. All analyses were conducted using the SPSS v20.0 (IBM Corp., Armonk, NY).

Inter-observer reliability

Inter-observer reliability with two independent coders was performed for 30% of the videos. This value is similar to those of previous studies²⁸. The frequency and duration of each category in each moment were considered, using the intra-class correlation coefficient (ICC) equation (2.1) and the Bland and Altman method. The ICC(2,1) values were interpreted as follows: >0.75 was excellent, 0.40–0.75 was moderate and <0.40 was poor²⁹. The results ranged between 0.45 and 1.0, indicating a moderate to excellent reliability.

Bland and Altman 95% limits of agreement were measured and the scatter plots were analyzed for all categories. A good agreement between the coders was found and no evidence of systematic bias was observed.

Comparison of Sample at Baseline

The experimental and control groups were compared on the basis of demographic variables and measured outcomes. Significance was set at the 0.05 level. The t-test for independent samples was used to compare the two groups on the basis of age and length of service and DCWs' outcomes; χ^2 tests were used to compare the two groups on the basis of dichotomous variables, including gender, education and marital status. Participants were all female with a mean age of 44.72 ± 9.02 years. The majority were married (67.2%), 46.4% had primary and middle school education and 41.4% high school. The average length of service was 9.61 ± 3.72 years. None of the differences were statistically significant at baseline (Table 4).

[Table 4]

Results

DCWs' perceived stress, burnout and job satisfaction

Data concerning DCWs' outcomes is displayed in Table 5.

A negative significant time interaction effect was obtained on perceived stress, with both groups reporting higher scores at 6 month follow-up than at baseline and immediately after the intervention ($p > 0.001$). Effect sizes were large ($\eta^2 = 0.36$).

Analyses showed a significant time interaction effect on the MBI subscale 'personal accomplishment'. After a decrease in personal accomplishment scores immediately after the PE intervention, at 6 months, scores had improved in the experimental group and the control group showed a decline over time in this variable ($p > 0.05$). Effect sizes were moderate ($\eta^2 = 0.08$).

Although no significant differences were found for the remaining MBI subscales, DCWs from the PE intervention showed improved levels of emotional exhaustion and depersonalization at 6 months follow-up. In the control group the values of all the MBI subscales deteriorated at 6 months follow-up.

No significant differences were obtained for total, intrinsic or extrinsic job satisfaction. A modest but positive change on total job satisfaction from pre to post-test, followed by a deterioration at 6 months follow-up, was found for both groups.

[Table 5]

DCWs' person-centered communication

Data concerning DCWs' person-centered communication can be found in Table 6.

Within the PE group, the frequency of the majority (24 out of 26 behaviors) of person-centered behaviors improved immediately after the intervention (T2). However, 22 out of 26 behaviors dropped at 6 months follow-up. For 10 of these behaviors, the frequencies were lower than those found at baseline. Among these, significant time and group interaction effects were found for the frequency of 'inform' ($p > 0.01$, $\eta^2_{\text{partial}} = 0.18$) and 'laugh' ($p > 0.01$,

$\eta^2_{\text{partial}}=0.10$). In the PE group, their frequency improved from T1 to T2 and dropped at follow-up. In the control group, the frequency of 'inform' reduced at T2 and values were sustained at follow-up, whereas 'laugh' improved at follow-up. Also, significant time effects were obtained for the frequency of 'validation' ($p>0.01$, $\eta^2_{\text{partial}}=0.10$) and 'play' ($p>0.05$, $\eta^2_{\text{partial}}=0.07$), with both groups experiencing a decline over time in these variables. Improvements were noticed in the frequency of 'involvement', 'withholding' and 'orientation'. Positive significant time effects were found for the frequency of the first two behaviors. The frequency of 'involvement' ($p>0.05$, $\eta^2_{\text{partial}}=0.07$) and 'withholding' ($p>0.05$, $\eta^2_{\text{partial}}=0.11$) improved over time in the experimental group and reduced at 6 months follow-up in the control group.

Concerning the duration of person-centered interactions, an improvement in 8 out of 11 variables was noticed. Among these, a significant time effect was found for the duration of 'social conversation' ($p>0.01$, $\eta^2_{\text{partial}}=0.09$), with both groups reporting an improvement over time in this variable. Declines over time were found for 'conversation about the person', 'instrumental conversation' and 'resident-directed eye gaze'. Significant time effects were obtained for the duration of 'instrumental conversation' ($p>0.05$, $\eta^2_{\text{partial}}=0.08$) and 'resident-directed eye gaze' ($p>0.05$, $\eta^2_{\text{partial}}=0.07$), with the experimental and control groups deteriorating over time.

[Table 6]

Discussion

This study sought to examine the effects of a PCC-based PE intervention on DCWs' stress, burnout, job satisfaction and person-centered communication.

Findings suggest that a PE intervention can positively impact DCWs' burnout. Compared to DCWs in the control group, those who received the PE intervention reported a decrease in their levels of burnout (with significant findings found for the MBI subscale 'personal accomplishment') at both post-intervention and 6 months follow-up. This promising result

suggests that, over time, adding a supportive component to PCC-training might enable DCWs to feel pleased about themselves and satisfied with their accomplishments on the job.

The findings did not support the hypothesis that, compared to education-only intervention, a PE intervention would improve DCWs' job satisfaction. A modest but positive change on total job satisfaction from pre to post-test, followed by a deterioration at 6 months follow-up, was found for both groups. One explanation for these results might be related to the measure used to assess job satisfaction. The MSQ is based on the conceptualization of job satisfaction as a multidimensional construct, considering several aspects that were not covered by the intervention (e.g., managers' support and organization conditions). Also, despite the guarantee of confidentiality, participants might have been reluctant to answer questions related to leadership's role or policies of the organization. The individual assistance during morning care in both groups can also offer some light about the lack of significant differences between groups. By allowing workers to have immediate guidance and support to handle challenging situations, it can have an independent effect on DCWs' job satisfaction. The extent to which individual assistance may impact DCWs' outcomes requires further consideration.

Strongest effects were found immediately after the intervention, with diminishing strength at the 6 month follow-up, on perceived stress and on most DCWs' person-centered communicative behaviors. Several reasons may explain these findings. First, the intervention ran for a short period of time (8 weeks). As it is important to keep DCWs under intervention long enough so they can experience lasting positive changes, maintenance strategies, such as 'booster sessions' (i.e., brief, periodic contacts intended to remind participants of intervention goals or encourage them to continue using the learned skills) are clearly needed. Moreover, it is possible that the results have been influenced by uncontrolled factors, such as the facility organizational culture. Contextual factors, in particular a supportive leadership and a reward culture of openness and accuracy, have been repeatedly cited in the literature as critically important to the success of interventions in terms of outcomes for DCWs³⁰. According to the literature, DCWs are often not acknowledged by their managers and feel that their work is

unappreciated, which negatively affects their state of well-being and care provision ³¹.

Therefore, efforts in improving leadership and management skills can be determinant to engender more positive impacts on DCWs. This might involve, for example, training to prepare for the challenges of leading change and support to provide adequate staff supervision and positive feedback systems ³⁰. However, more research is needed to determine what leadership skills can indeed be helpful. Another factor may be related to the progressive nature of dementia, which means that the symptoms gradually worsen over time increasing DCWs' stress and reducing job satisfaction. Also, the increased perceived stress at 6 month follow-up can be the result of an enhanced DCWs' awareness of stress on the job; the repeatedly completion of a measure on perceived stress may contribute to increase DCWs' understanding and consciousness of stress over time. At last, the possible pressure to provide better care after the interventions could have triggered higher stress levels. Nevertheless, some person-centered behaviors were positively affected by the PE intervention and should be highlighted. DCWs from the experimental group experienced a significant improvement in the frequency of involvement and withholding and in the duration of social conversation. Also, positive but no significant improvements were found in the duration of multisensory stimulation and several non-verbal communicative behaviors, including smile, laugh and affective touch. These are promising results as it is becoming increasingly acknowledged that good dementia care is underpinned by interpersonal relationships between people with dementia and DCWs that rely more on emotional, sensitive, and empathetic interactions rather than on verbal expressiveness ⁹. The fact that at 6 months-follow up results were more positive for the duration of behaviors suggest that DCWs spend more time communicating with fewer interruptions. Besides, these findings indicate that DCWs might have selected over time those communicative behaviors that showed to be more effective, making more use of them.

Results must be, however, interpreted with caution as DCWs were not always recorded with the same resident and this may have influenced the results. Future studies should try to create and follow the same dyads. Besides, DCWs' performance, particularly immediately after

the intervention, may have been influenced by the “Hawthorne effect”, which means that DCWs being aware of video-recording possibly behaved differently. Though, DCWs’ familiarization with the methodology may have minimized this effect at 6 months follow-up³².

Other methodological limitations have also to be considered. The sample size was relatively small and might have reduced the statistical power to detect more significant changes. Hence when cluster designs are used, there are two sources of variance in the observations: the variability of patients within a cluster; and the variability between clusters. These two sources combine to produce an increase in the variance, and both must be considered in the analysis. The effect of the increased variance due to a cluster design is to increase the size of the standard errors and thus to widen confidence intervals and increase p-values, compared with a study of the same size using individual randomization³³. In effect, the sample size is reduced and power is lost, and thus, sample sizes have to be inflated. Moreover, it was not possible to blind researchers to the experimental or control groups or assessments. Future studies with a double-blinded design should be conducted to clarify the findings.

Nevertheless, our findings are of interest as they provide evidence that a PE intervention may be an effective approach to reduce DCWs’ burnout levels and improve person-centered behaviors. Further research is warranted to determine the extent of the benefits of this approach on residents with dementia and on other DCWs’ outcomes such as depression, anxiety and perceived mastery. One of the strengths of the study is the consideration of a 6-months follow up evaluation. This is important as most studies tend to rely only on pre and immediately posttest assessments¹⁷. Also, the use of video-recording provided a suitable method to assess interactions. Video-recording enables replaying and reviewing the data, to control the observer’s fatigue and to achieve deeper levels of observation and analysis that are not possible to achieve by means of real-time observations³². The high inter-observer reliability obtained for the ethogram further supports its reliability and validity to measure PCC interactions. Yet, conducting future evaluations of the ethogram in order to further develop its acceptability, utility and validity is strongly recommended.

Overall, the results suggested that providing DCWs with training and emotional support is more effective in reducing burnout and improve adequate communicative behaviors than an education-only intervention. These findings highlight the importance of interventions in dementia care settings to go beyond DCWs' knowledge and instrumental skills to also address emotional skills. The addition of booster follow-up sessions to help maintain and extend the positive long-term effects of the intervention is highly encouraged. Also, DCWs' outcomes are largely associated to factors within the organization, thus culture-change initiatives (e.g., breaking down hierarchies, leadership commitment and DCWs empowerment) are further encouraged, as this is determinant to achieve and sustain practice changes.

Acknowledgements

The authors gratefully acknowledge all institutions and DCWs for their participation in this study.

References

1. Alzheimer's Disease International. *World Alzheimer Report 2013: An analysis of long-term care for dementia*. London: Alzheimer's Disease International;2013.
2. Stone R. The Long-Term Care Workforce: From Accidental to Valued Profession. In: Wolf D, Folbre N, eds. *Universal Coverage of Long-Term Care in the United States: Can We Get There from Here?* United States: Russell Sage Foundation; 2012.
3. Gray-Stanley JA, Muramatsu N. Work stress, burnout, and social and personal resources among direct care workers. *Research in Developmental Disabilities*. May-Jun 2011;32(3):1065-1074.
4. Edberg A-K, Bird M, Richards DA, Woods R, Keeley P, Davis-Quarrell V. Strain in nursing care of people with dementia: Nurses' experience in Australia, Sweden and United Kingdom. *Aging & Mental Health*. 2008;12(2):236-243.
5. Edvardsson D, Sandman P, Nay R, Karlsson S. Predictors of job strain in residential dementia care nursing staff. *J Nurs Manag*. 2009;17(1):59-65.
6. Menne HL, Ejaz FK, Noelker LS, Jones JA. Direct Care Workers' Recommendations for Training and Continuing Education. *Gerontology & Geriatrics Education*. 2007;28(2):91-108.
7. Nolan M, Davies S, Brown J, et al. The role of education and training in achieving change in care homes: a literature review. *Journal of Research in Nursing*. 2008;13(5):411-433.
8. Nolan M, Davies S, Brown J, Keady J, Nolan J. Beyond person-centred care: a new vision for gerontological nursing. *J Clin Nurs*. Mar 2004;13(3a):45-53.
9. Brooker D. *Person-centred dementia care: making services better*. London: Jessica Kingsley Publishers; 2007.
10. Kitwood T. *Dementia Reconsidered: The Person Comes First* Buckingham: Open University Press; 1997.
11. Chenoweth L, King MT, Jeon YH, et al. Caring for Aged Dementia Care Resident Study (CADRES) of person-centred care, dementia-care mapping, and usual care in dementia: a cluster-randomised trial. *Lancet Neurol*. Apr 2009;8(4):317-325.
12. Fossey J. Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial. *BMJ*. 2006;332(7544):756-761.
13. Sloane PD, Hoefler B, Mitchell CM, et al. Effect of person-centered showering and the towel bath on bathing-associated aggression, agitation, and discomfort in nursing home residents with dementia: a randomized, controlled trial. *J Am Geriatr Soc*. 2004;52(11):1795-1804.
14. Sidani S, Streiner D, Leclerc C. Evaluating the effectiveness of the abilities-focused approach to morning care of people with dementia. *Int J Older People Nurs*. Mar 2012;7(1):37-45.
15. Edvardsson D, Winblad B, Sandman P. Person-centred care of people with severe Alzheimer's disease: current status and ways forward. *The Lancet Neurology*. 2008;7(4):362-367.
16. van den Pol-Grevelink A, Jukema JS, Smits CH. Person-centred care and job satisfaction of caregivers in nursing homes: a systematic review of the impact of different forms of person-centred care on various dimensions of job satisfaction. *Int J Geriatr Psychiatry*. Mar 2012;27(3):219-229.
17. Barbosa A, Sousa L, Nolan M, Figueiredo D. Effects of person-centered care approaches to dementia care on staff: A systematic review. *American Journal of Alzheimer's Disease and Other Dementias*. January 20, 2014 2014.
18. Van Weert J, Vandulmen A, Spreeuwenberg P, Ribbe M, Bensing J. Effects of snoezelen, integrated in 24h dementia care, on nurse-patient communication during morning care. *Patient Education and Counseling*. 2005;58(3):312-326.

19. Pais Ribeiro J, Marques T. A avaliação do stresse: a propósito de um estudo de adaptação da escala de percepção de stresse. *Psicologia, Saúde & Doenças*. 2009;10(2):237-248.
20. Lee E-H. Review of the psychometric evidence of the perceived stress scale. *Asian Nursing Research*. 2012;6(4):121-127.
21. Melo B, Gomes A, Cruz J. Desenvolvimento e adaptação de um instrumento de avaliação psicológica do “burnout” para os profissionais de Psicologia. In: Associação dos Psicólogos Portugueses, ed. *Avaliação psicológica: formas e contextos*. Braga: Lusografe; 1999:596-603.
22. Poghosyan L, Aiken LH, Sloane DM. Factor structure of the Maslach Burnout Inventory: An analysis of data from large scale cross-sectional surveys of nurses from eight countries. *International Journal of Nursing Studies*. 2009;46(7):894-902.
23. Ferreira J, Fernandes R, Haase RF, Santos E. Minnesota Satisfaction Questionnaire - Short Form: estudo de adaptação e validação para a língua portuguesa. *Psicologica*. 2009;51:251-281.
24. Weiss DJ, Dawis RV, England GW. *Manual for the Minnesota Satisfaction Questionnaire*. Minneapolis: University of Minneapolis;1967.
25. Caris-Verhallen W, Kerkstra A, Bensing J. Non-verbal behaviour in nurse-elderly patient communication. *Journal of Advanced Nursing*. 1999;29(4):808-818.
26. Williams K, Harris B, Lueger A, Ward K, Wassmer R, Weber A. Visual cues for person-centered communication. *Clinical Nursing Research*. 2011;20(4):448-461.
27. Cohen J. *Statistical Power Analysis for the Behavioral Sciences* 2nd ed. New Jersey: Lawrence Erlbaum Associates; 1988.
28. Bourgeois MS, Dijkstra K, Burgio LD, Allen RS. Communication skills training for nursing aides of residents with dementia: The impact of measuring performance. *Clin Gerontologist*. 2004;27(1-2):119-138.
29. Fleiss J. *The Design and Analysis of Clinical Experiments*: Taylor & Francis; 1986.
30. Jeon YH, Merlyn T, Chenoweth L. Leadership and management in the aged care sector: a narrative synthesis. *Australasian Journal on Ageing*. 2010;29(2):54-60.
31. Barbosa A, Nolan M, Sousa L, Figueiredo D. Supporting direct care workers in dementia care: effects of a psychoeducational intervention. *Am J Alzheimers Dis Other Demen*. 2015;30(2):130-138.
32. Haidet KK, Tate J, Divirgilio-Thomas D, Kolanowski A, Happ MB. Methods to improve reliability of video-recorded behavioral data. *Res Nurs Health*. Aug 2009;32(4):465-474.
33. Campbell MK, Elbourne DR, Altman DG. CONSORT statement: extension to cluster randomised trials. *BMJ*. 2004;328(7441):702-708.
34. Bower S, Bower G. *Asserting yourself: A practical guide for positive change*. Boston, MA: Da Carpo; 2004.
35. Bono E. *Six thinking hats*. New York, NY: Little, Brown and Company.; 1985.

Table 1. Content of the interventions

Session	Component	Experimental Group	Control Group
1	Educative	<i>Information about PCC and dementia:</i> Information about the concept and principles of PCC. Basic information on dementia, its causes, symptoms and evolution.	
	Supportive	<i>Emotional impact of caregiving:</i> The positive and negative impacts of the caregiving experience on personal and professional life; Abdominal breathing.	-
2	Educative	<i>Communication in dementia:</i> Communicative behavioral strategies to interact with residents with dementia. (e.g. give simple choices; use validation; allows time to respond; use individual's name and eye contact).	
	Supportive	<i>Conflict management:</i> Improving assertiveness through the DESC technique (Describe; Explain; Specify; Conclude) technique ³⁴ . Stretching and strengthening exercises.	-
3	Educative	<i>Challenging behaviors:</i> Information about challenging behaviors and strategies to deal with them.	
	Supportive	<i>Teamwork:</i> The importance, benefits and constraints to teamwork; strategies to enhance cooperation between DCWs (e.g. active listen, positive feedback). Cognitive relaxation technique.	-
4	Educative	<i>The environment and dementia:</i> Strategies to enhance the physical and social environment for the person with dementia (e.g. decrease background noise; post signs as reminders); information about the risk factors and strategies to prevent falls.	
	Supportive	<i>Dealing with emotions:</i> Improving emotion-management strategies through the activity "six colors to think" ³⁵ ; Stretching and strengthening exercises.	-
5	Educative	<i>Motor stimulation:</i> Information about motor stimulation; strategies to enhance residents' involvement in daily care (e.g., break the small steps of an activity); and techniques for the moving and handling of residents.	
	Supportive	<i>Time management:</i> The impact of poor time management on personal and professional life and tools for better time management (e.g. set priorities; use a planning tool). Mental body-scan.	-
6	Educative	<i>Multisensory stimulation - olfaction:</i> Information about multisensory stimulation; dementia-related olfactory changes and strategies to stimulate the olfaction during the daily care (e.g., use shower gel of different fragrances; place aroma diffusers in the bedroom)	
	Supportive	<i>Problem-solving:</i> Using the problem-solving technique: (a) identify the problem; (b) explain the problem; (c) create solutions; (d) choose one solution; (e) plan the implementation of the solution; (f) evaluate the efficacy. Stretching and strengthening exercises	-
7	Educative	<i>Multi-sensory stimulation – vision and tactile stimulation:</i> The importance of vision and touch for people with dementia, dementia-related visual and tactile changes; strategies to stimulate the vision (e.g. reality orientation) and touch (e.g. hand massage during bath)	
	Supportive	<i>Relaxation:</i> Yoga	-
8	Educative	<i>Multi-sensory stimulation – audition and taste:</i> The importance of audition and taste for people with dementia; dementia-related audition and taste changes; strategies to stimulate the audition (e.g., listen to residents' favorite song) and taste (e.g. brush the person's teeth with toothpastes of different flavors).	
		Celebration and finalization	

Table 2. Verbal communicative behaviors

Categories	Description
Consult	Consulting the person with dementia about his or her preferences, desires and needs. Includes questions that invite resident's judgment. Examples include: <ul style="list-style-type: none"> • Would you like your shoes on or off? • Do you want to wear a skirt or pants?
Inform	Guiding the resident in terms of what to expect and providing information about what is going to happen during the task. Examples include: <ul style="list-style-type: none"> • Now I'm going to comb your hair. • Today you will take a bath.
Involve	Giving the resident the opportunity to take care for him/herself as much as possible and just 'completing' the care task when necessary. Examples include: <ul style="list-style-type: none"> • Could you help me with this? • Hold the toothbrush with your hand.
Reward	Rewarding the person and his/her behavior, giving compliments and using expressions of encouragement. Examples include: <ul style="list-style-type: none"> • Well done, Sr. John. • You can do it, Sr. John.
Validate	Acknowledging the subjective reality of a person's emotions and feelings, and giving a response on the feeling level, without correcting the residents' reality or frame of reference, even if it is chaotic. Using statements to interpret or recognize the emotional state of the resident during the interaction. Examples include: <ul style="list-style-type: none"> • This is distressing for you, I understand. • How do you feel about it?
Assess comfort	Conveying interest and concern for the welfare and comfort of the person with dementia. Examples include: <ul style="list-style-type: none"> • How are you feeling today? • Does your leg hurt?
Distract	Amusing the person through humorous commentaries or distracting him/her in a positive way by guiding the conversation away from something unpleasant.
Sensory stimulation	Providing sensory information, without the intervention of concepts and intellectual understanding; for example through music, touch or aromas. Examples include: <ul style="list-style-type: none"> • Feel how nice and soft this towel is. • This cream smells good!
Conversation about the person	Showing interest in the resident's life or background. Examples include: <ul style="list-style-type: none"> • You were a teacher, weren't you? • You used to like gardening, didn't you?
Social conversation	Friendly conversation that conveys an interest in the resident and is not related to instrumental care. Includes statements that acknowledge that the resident said something. Examples include: <ul style="list-style-type: none"> • You have a very nice dress. Where did you get it? • Thank you!
Task-oriented conversation	Communication that is related to task accomplishment or focused on nursing or therapeutic topics. Examples include: <ul style="list-style-type: none"> • Where are your glasses? • The doctor said not to eat bread.
Conversation with a third person	Communication to a third person. Examples include: <ul style="list-style-type: none"> • Can you please give me a towel? (to another DCW)

Ignore	Ignoring residents' statements by responding with an unrelated statement or question, interrupting or changing the topic of conversation. Carrying on a conversation in the presence of a person as if he/she is not present. Examples include: <ul style="list-style-type: none"> • Today she [<i>the resident</i>] is very friendly.
Infantilize	Patronizing or treating and talking to the person with dementia as if he/she was a child. Examples include: <ul style="list-style-type: none"> • Good girl, you behaved so well.
Invalidate	Failing to acknowledge the subjective reality of a person's experience and especially what he or she is feeling. Correcting the resident on cognitive facts. Examples include: <ul style="list-style-type: none"> • Your husband is dead. • It's Wednesday today, not Monday.
Mockery	Disdain, pointing out or making fun of residents' behavior or actions. Placing the person towards his/her difficulties. Examples include: <ul style="list-style-type: none"> • What's my name? Have you forgotten?
Criticise	Showing disapproval or criticize residents' performance or behavior. Examples include: <ul style="list-style-type: none"> • That's wrong. You are hopeless.
Impose	Forcing a person to do something, overriding desire or denying the possibility of choice on his or her part. Statements can be considered dominating or controlling. Examples include: <ul style="list-style-type: none"> • You will dress this sweater because it is the freshest you have. • Be quiet.

Table 3. Non-verbal communicative behaviors

Affirmative Nodding	Nodding head as a sign of approval, encouragement, or interest in the resident.
Resident-Directed Eye Gaze	Looking at the face of the resident.
Smile	Expression in which the corners of the mouth are directed upwards, denoting affability towards the resident.
Laugh	Opening the mouth (totally or partially), making a sound commonly associated with the act of laughing.
Withholding	Refusing a residents' request or question. Includes statements from the resident that the DCW does not acknowledge (e.g. resident asks if she can return to her room and the DCW does not respond).
Affective touch	Spontaneous and affective touch that is not necessary for the completion of a task (e.g. a pat on the back, a hug).
Guiding touch	Using touch to draw the person's attention or guide him/her for a task.
Instrumental touch	Deliberate physical contact, which is necessary for the completion of a task.

Table 4. DCWs' socio-demographic characteristics

Outcome	Total (n=58) n (%)	Experimental group (n=27) n (%)	Control Group (n=31) n (%)	p-value
Gender				
Female	58 (100.0)	27 (100.0)	31 (100.0)	-
Age in years				
M (SD)	44.72 (9.02)	43.37 (10.00)	45.90 (8.04)	0.290 ^a
Marital Status				
Married	39 (67.2)	17 (63.0)	22 (71.0)	
Widowed	3 (5.2)	1 (3.7)	2 (6.5)	
Single	4 (6.9)	2 (7.4)	2 (6.5)	0.887 ^b
Divorced/separated	9 (15.5)	5 (15.5)	4 (12.9)	
Other	3 (5.2)	2 (7.4)	1 (3.2)	
Education				
Primary school	15 (25.9)	4 (14.8)	11 (35.5)	
Middle school	12 (20.7)	6 (22.2)	6 (19.4)	
High school	24 (41.4)	11 (40.7)	13 (41.9)	0.144 ^b
College degree	1 (1.7)	1 (3.7)	0 (0.0)	
Other	6 (10.3)	5 (18.5)	1 (3.2)	
Length of service (years)				
M (SD)	9.61 (3.72)	9.84 (4.86)	9.42 (2.51)	0.678 ^a

Abbreviations: M, mean; SD, standard deviation

^a t-test^b χ^2

Table 5. Changes in DCWs' stress, burnout and job satisfaction

Outcome	Experimental group (n=24)			Control group (n=29)			Time effect	ES	Group x Time effect	ES
	T1	T2	T3	T1	T2	T3				
	Mean (SD)			Mean (SD)						
PSS	19.42 (5.90)	18.79 (6.48)	27.25 (4.59)	20.55 (6.31)	20.10 (4.79)	25.55 (6.99)	F=28.255, <i>df</i> (2) p=0.000	0.36	F=1.346, <i>df</i> (2) p=0.265	0.03
MBI										
Emotional Exhaustion	17.0 (11.41)	15.8 (8.60)	13.82 (9.76)	12.67 (10.59)	15.42 (9.72)	16.25 (10.70)	F=0.132, <i>df</i> (2) p=0.876	0.00	F=2.145, <i>df</i> (2) p=0.123	0.05
Depersonalization	5.25 (5.05)	6.88 (6.40)	3.20 (3.85)	6.07 (5.71)	5.52 (4.01)	6.54 (5.89)	F=0.848; <i>df</i> (2) p=0.431	0.02	F=3.048, <i>df</i> (2) p=0.052	0.06
Personal accomplishment	40.42 (4.94)	38.54 (8.42)	39.50 (5.80)	40.69 (6.20)	37.31 (8.02)	34.17 (8.48)	F=4.604, <i>df</i> (2) p=0.012	0.08	F=2.649, <i>df</i> (2) p=0.076	0.05
MSQ										
MSQ intrinsic	24.29 (2.31)	24.00 (2.16)	24.33 (2.16)	22.55(4.19)	21.55(3.53)	20.62 (5.19)	F=1.619, <i>df</i> (2) p= 0.203	0.03	F=1.733; <i>df</i> (2) p=0.182	0.01
MSQ extrinsic	26.58 (4.67)	26.37 (3.56)	26.04 (4.42)	24.62(4.84)	25.59(4.15)	26.55 (6.41)	F=0.635, <i>df</i> (2); p=0.532	0.04	F=2.007; <i>df</i> (2); p=0.140	0.04
Total	72.96 (5.99)	73.83 (8.48)	72.29 (7.92)	68.14 (9.05)	68.55 (10.13)	68.48 (15.38)	F=0.209, <i>df</i> (2) p=0.811	0.00	F=0.163, <i>df</i> (2) p=0.849	0.00

Notes: M= mean; SD= standard deviation; ES= effect sizes; *df*= degrees of freedom

Table 6. Changes in DCWs' verbal and non-verbal communicative behaviour

Categories	Type	Experimental group (n=24)			Control group (n=29)			time p-value	Partial eta squared	group×time p-value	Partial eta squared
		T1	T2	T3	T1	T2	T3				
		Mean (SD)			Mean (SD)						
<i>Verbal communicative behaviour</i>											
Consult	Freq	1.02 (1.02)	1.05 (1.13)	0.82 (1.05)	1.60 (1.83)	1.24 (1.28)	0.80 (1.18)	0.097	0.05	0.412	0.02
Inform	Freq	8.69 (4.40)	9.45 (5.60)	4.70 (3.68)	8.23 (4.41)	6.99 (3.91)	6.98 (5.20)	0.013*	0.16	0.008*	0.18
Involve	Freq	3.46 (3.15)	3.55 (3.20)	4.85 (3.03)	3.50 (3.28)	6.13 (2.72)	5.00 (2.64)	0.027*	0.07	0.100	0.05
	Dur	13.06 (14.92)	17.58 (18.04)	24.00 (14.88)	19.14 (21.89)	28.68 (18.94)	21.36 (12.61)	0.053	0.06	0.099	0.05
Reward	Freq	1.37 (1.49)	1.81 (1.72)	1.49 (1.70)	0.75 (0.70)	1.34 (1.17)	1.38 (1.25)	0.048*	0.06	0.454	0.02
Validate	Freq	0.32 (0.70)	0.35 (0.83)	0.02 (0.10)	0.13 (0.34)	0.17 (0.34)	0.03 (0.10)	0.006*	0.37	0.101	0.02
Assess comfort	Freq	0.35 (0.48)	0.35 (0.48)	0.19 (0.32)	0.38 (0.58)	0.43 (0.79)	0.49 (0.74)	0.964	0.00	0.531	0.01
Distract	Freq	0.41 (0.84)	0.55(1.13)	0.09 (0.21)	0.25 (0.60)	0.15 (0.34)	0.08 (0.17)	0.486	0.00	0.130	0.04
Sensory stimulation	Freq	0.12 (0.23)	0.33 (0.46)	0.32 (0.42)	0.11 (0.33)	0.32 (0.82)	0.44 (1.45)	0.023*	0.09	0.904	0.00
	Dur	0.60 (1.74)	1.71 (2.78)	2.06 (3.11)	0.34 (0.98)	3.08 (9.59)	1.06 (3.55)	0.129	0.44	0.399	0.01
Social conversation	Freq	4.69 (2.64)	5.28 (2.84)	5.19 (3.00)	6.44 (4.51)	6.95 (4.18)	6.13 (3.09)	0.306	0.02	0.993	0.00
	Dur	33.39 (30.55)	43.79 (23.04)	46.75 (40.92)	37.30 (25.44)	49.74 (33.38)	63.40 (45.99)	0.009* *	0.55	0.732	0.00
Conversation about the person	Freq	0.02 (0.11)	0.03 (0.14)	0.02 (0.08)	0.12 (0.25)	0.06 (0.16)	0.07 (0.17)	0.509	0.00	0.210	0.03
	Dur	0.09 (0.35)	0.43 (2.09)	0.05 (0.23)	1.17 (2.76)	0.80 (2.08)	1.99 (8.08)	0.907	0.55	0.09	0.01
Instrumental conversation	Freq	2.79 (2.10)	2.50 (2.37)	3.18 (2.62)	2.10 (1.66)	1.94 (1.15)	2.81 (2.34)	0.608	0.00	0.972	0.00
	Dur	13.41 (11.25)	10.68 (9.81)	19.43 (19.03)	10.73 (9.85)	10.73 (9.96)	15.01 (13.96)	0.040*	0.62	0.082	0.01
Conversation with others	Freq	5.14 (4.02)	4.86 (2.93)	3.89 (2.84)	5.55 (4.06)	4.02 (2.89)	3.89 (2.84)	0.006*	0.09	0.462	0.01
	Dur	21.33 (17.63)	25.32 (14.81)	20.52 (13.83)	14.68 (17.90)	20.77 (12.02)	19.14 (12.52)	0.182	0.04	0.628	0.01
Ignore	Freq	1.82 (1.47)	1.70 (1.73)	2.14 (2.07)	2.12 (2.10)	1.63 (1.56)	1.65 (1.75)	0.613	0.00	0.457	0.02
Infantilize	Freq	0.02 (0.09)	0.00	0.00	0.28 (0.51)	0.12 (0.28)	0.21 (0.42)	0.334	0.02	0.409	0.02

Invalidate	Freq	0.07 (0.27)	0.00	0.02 (0.07)	0.13 (0.25)	0.17 (0.39)	0.20 (0.10)	0.909	0.00	0.418	0.02
Mockery	Freq	0.11 (0.22)	0.09 (0.22)	0.34 (0.70)	0.54 (0.69)	0.25 (0.44)	0.57 (1.12)	0.074	0.05	0.543	0.01
Criticise	Freq	0.47 (0.73)	0.07 (0.25)	0.28 (0.49)	0.46 (0.72)	0.57 (0.75)	0.24 (0.53)	0.386	0.02	0.051	0.06
Impose	Freq	0.49 (0.89)	0.16 (0.37)	0.37 (0.60)	0.56 (0.58)	0.83 (1.57)	0.22 (0.47)	0.228	0.059	0.104	0.09
<i>Non-verbal communicative behaviour</i>											
Affirmative Nodding	Freq	0.45 (0.85)	0.74 (1.16)	0.22 (0.65)	0.55 (0.84)	0.51 (0.99)	0.72 (1.19)	0.630	0.02	0.170	0.07
Resident-directed eye gaze	Freq	1.90 (1.99)	2.08 (1.93)	0.85 (0.98)	1.91 (2.36)	2.03 (2.26)	1.91 (2.34)	0.106	0.05	0.169	0.04
	Dur	19.18 (32.24)	26.56 (40.08)	4.17 (6.03)	25.43 (56.24)	19.68 (33.73)	11.69 (17.92)	0.032*	0.07	0.461	0.02
Smile	Freq	0.32 (0.62)	0.35 (0.62)	0.33 (0.43)	0.31 (0.86)	0.23 (0.54)	0.85 (1.21)	0.120	0.08	0.109	0.09
	Dur	0.61 (1.25)	0.71 (1.76)	1.29 (3.12)	2.65 (11.90)	0.89 (2.11)	5.99 (13.81)	0.199	0.03	0.373	0.02
Laugh	Freq	1.15 (1.05)	1.78 (1.71)	0.94 (1.12)	1.01 (1.01)	0.62 (0.82)	1.03 (1.14)	0.526	0.01	0.004*	0.106
	Dur	9.24 (17.38)	7.39 (12.30)	8.61 (19.40)	10.28 (29.11)	2.10 (3.03)	3.75 (5.09)	0.550	0.01	0.934	0.001
Withholding	Freq	0.41 (0.76)	0.06 (0.16)	0.00	0.08 (0.23)	0.01 (0.06)	0.07 (0.29)	0.024*	0.114	0.06	0.11
Affective touch	Freq	0.69 (0.57)	0.68 (0.62)	0.48 (0.56)	1.18 (1.33)	0.75 (1.00)	0.94 (1.28)	0.324	0.02	0.393	0.02
	Dur	2.22 (3.28)	1.69 (1.96)	1.86 (3.61)	3.53 (4.42)	4.05 (9.03)	3.68 (7.08)	0.992	0.00	0.854	0.00
Instrumental touch	Freq	10.33 (2.61)	10.21 (2.90)	10.82 (2.25)	9.82 (2.79)	8.99 (2.34)	10.44 (2.92)	0.085	0.05	0.616	0.01
	Dur	281.06 (169.8)	311.31 (52.67)	296.29 (43.72)	326.37 (251.40)	268.9 (69.58)	284.08 (51.45)	0.849	0.00	0.27	0.03
Awareness touch	Freq	0.16 (0.49)	0.56 (1.39)	0.23 (0.38)	0.44 (0.71)	0.44 (0.61)	0.43 (0.63)	0.343	0.02	0.34	0.02
