

# ANALYSIS & PDE

Volume 7

No. 2

2014

BEN ANDREWS, MAT LANGFORD AND JAMES MCCOY

**CONVEXITY ESTIMATES FOR HYPERSURFACES MOVING  
BY CONVEX CURVATURE FUNCTIONS**

# CONVEXITY ESTIMATES FOR HYPERSURFACES MOVING BY CONVEX CURVATURE FUNCTIONS

BEN ANDREWS, MAT LANGFORD AND JAMES MCCOY

We consider the evolution of compact hypersurfaces by fully nonlinear, parabolic curvature flows for which the normal speed is given by a smooth, convex, degree-one homogeneous function of the principal curvatures. We prove that solution hypersurfaces on which the speed is initially positive become weakly convex at a singularity of the flow. The result extends the convexity estimate of Huisken and Sinestrari [*Acta Math.* **183**:1 (1999), 45–70] for the mean curvature flow to a large class of speeds, and leads to an analogous description of “type-II” singularities. We remark that many of the speeds considered are positive on larger cones than the positive mean half-space, so that the result in those cases also applies to non-mean-convex initial data.

## 1. Introduction

Given a smooth, compact immersion  $X_0 : M^n \rightarrow \mathbb{R}^{n+1}$ ,  $n > 1$ , we consider smooth families  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  of smooth immersions  $X(\cdot, t)$  solving the curvature flow

$$\frac{\partial X}{\partial t}(x, t) = -s(x, t)\nu(x, t), \quad X(\cdot, 0) = X_0, \quad (1-1)$$

where  $\nu$  is the outer unit normal field of the solution, and the speed  $s$  is determined by a function of the principal curvatures  $\kappa_i$  (with respect to  $\nu$ ). That is,

$$s(x, t) = f(\kappa_1(x, t), \dots, \kappa_n(x, t)). \quad (1-2)$$

We require that the speed function  $f$  satisfies the following conditions:

### Conditions.

- (i)  $f \in C^\infty(\Gamma)$  for some connected, open, symmetric cone  $\Gamma \subset \mathbb{R}^n$ .
- (ii)  $f$  is monotone increasing in each argument.
- (iii)  $f$  is homogeneous of degree one.
- (iv)  $f > 0$ .
- (v)  $\Gamma$  is preserved by the flow (1-1).

---

Research partly supported by ARC Discovery Projects grants DP0556211, DP120100097. Langford acknowledges the support and hospitality of the Mathematical Sciences Center at Tsinghua University, and the Institute for Mathematics and its Applications at the University of Wollongong, where part of this work was completed.

*MSC2010*: primary 53C44; secondary 35K55.

*Keywords*: convexity estimates, curvature flows, fully nonlinear.



Condition (v) is intended as follows: Let  $X$  be a solution of (1-1)–(1-2) such that the initial hypersurface satisfies  $(\kappa_1(x, 0), \dots, \kappa_n(x, 0)) \in \Gamma$  for all  $x \in M$ . Then there is a connected, open, symmetric subcone  $\Gamma_0$  of  $\Gamma$  satisfying  $\overline{\Gamma_0} \setminus \{0\} \subset \Gamma$  such that the principal curvatures of the solution satisfy  $(\kappa_1(x, t), \dots, \kappa_n(x, t)) \in \Gamma_0$  for all  $(x, t) \in M \times [0, T)$ . We refer to  $\Gamma_0$  as a *preserved cone* of the flow. This is discussed further below.

Observe that, since the normal points outwards and  $f$  is homogeneous, we lose no generality in assuming further that  $(1, \dots, 1) \in \Gamma$ , and that  $f$  is normalised such that  $f(1, \dots, 1) = 1$ . Furthermore, since  $f$  is symmetric, we may at each point reorder the principal curvatures such that  $\kappa_n \geq \dots \geq \kappa_1$ .

For most of the paper, we will also require that  $f$  satisfies the following two conditions, which are somewhat distinct from conditions (i)–(v):

**Conditions.**

- (vi)  $f$  is locally convex.
- (vii)  $(\partial f / \partial z_p - \partial f / \partial z_q)|_z \geq 0$  whenever  $z \in \Gamma$  is such that  $z_p \geq z_q$ .

We will say that  $s$  is an *admissible speed* for the flow (1-1) if  $s$  is given by (1-2) such that  $f$  satisfies conditions (i)–(vii).

Some discussion of conditions (i)–(vii) is in order: The symmetry of  $f$  is a geometric condition — it allows us to write  $s$  as a smooth function of the Weingarten map of the solution, which ensures geometric invariance of the flow. The monotonicity of  $f$  then ensures that the flow is parabolic, which guarantees short time existence of a solution if the principal curvatures of the initial immersion lie in  $\Gamma$ . Condition (v) is then a requirement that the principal curvatures do not “move out of”  $\Gamma$  during the flow. In general, some such condition is necessary (see [Andrews et al. 2013b, Theorem 3]), although, in particular, it automatically holds in each of the following situations (Lemma 2.4):

**Ancillary conditions.**

- (viii) Conditions (i)–(iv) and (vi) hold, and  $\Gamma$  is convex.
- (ix) Conditions (i)–(iv) and (vi) hold, and  $f|_{\partial\Gamma} = 0$ .
- (x) Conditions (i)–(iv) hold, and  $n = 2$ .

For the purposes of Theorem 1.1, however, we need only assume that the weaker condition (v) holds. We remark that ancillary condition (ix) makes sense because any function satisfying conditions (i)–(iv) has a continuous extension to  $\partial\Gamma$ . This is proved for  $\Gamma = \Gamma_+$  in [Andrews et al. 2013b], but the proof is easily modified for the present situation.

In the presence of condition (i), conditions (vi)–(vii) are equivalent to requiring that the speed is a smooth, convex function of the Weingarten map (Lemma 2.1). We note that condition (vii) is automatically true in each of the following situations:

**Ancillary conditions.**

- (xi) Conditions (i)–(iii) and (vi) hold, and  $\Gamma$  is convex.
- (xii) Conditions (i)–(iii) and (vi) hold, and  $f$  extends as a convex function to  $\mathbb{R}^n$  (for example, if  $f|_{\partial\Gamma} = 0$ ).
- (xiii) Conditions (i)–(iv) and (vi) hold, and  $n = 2$ .

The above assertions are discussed in greater detail in [Section 2](#).

We now list some examples of admissible speeds.

**Examples 1.1.** The following functions define admissible speeds for the flow (1-1):

- (1) *The arithmetic mean:*  $f(z_1, \dots, z_n) = z_1 + \dots + z_n$  on the half-space  $\Gamma = \{z \in \mathbb{R}^n : z_1 + \dots + z_n > 0\}$ . The corresponding flow is the (mean convex) mean curvature flow.
- (2) *The power means:*  $f_p(z_1, \dots, z_n) = (z_1^p + \dots + z_n^p)^{1/p}$ , for  $p \geq 1$ , on the positive cone  $\Gamma_+^n = \{z \in \mathbb{R}^n : z_i > 0 \text{ for all } i\}$ . The case  $p = 2$  corresponds to the flow by the norm of the Weingarten map.
- (3) *Positive linear combinations:* If  $f_1, \dots, f_k$  are admissible on  $\Gamma$ , then, for all  $(s_1, \dots, s_k) \in \Gamma_+^k$ , the function  $f = s_1 f_1 + \dots + s_k f_k$  is admissible on  $\Gamma$ . For example, the function

$$f(z_1, \dots, z_n) = z_1 + \dots + z_n + \sqrt{z_1^2 + \dots + z_n^2}$$

on the cone  $\Gamma_+$  defines an admissible speed. In fact, the functions

$$f_\alpha(z_1, \dots, z_n) = z_1 + \dots + z_n + \alpha \sqrt{z_1^2 + \dots + z_n^2}$$

for  $\alpha \in [0, 1]$  on the larger cones  $\Gamma_\alpha = \{z \in \mathbb{R}^n : z_1 + \dots + z_n + \alpha \sqrt{z_1^2 + \dots + z_n^2} > 0\}$  define admissible speeds. We remark that the cones  $\Gamma_\alpha$  contain the half-space  $\{z \in \mathbb{R}^n : z_1 + \dots + z_n > 0\}$  when  $\alpha > 0$ .

- (4) *Concave functions:* If  $g \in C^\infty(\Gamma)$  is symmetric, homogeneous degree one and concave, then an admissible speed is defined by the function  $f = H - \varepsilon g$  on the subcone of  $\Gamma$  for which  $H > \varepsilon g$  and  $\dot{g}^i < 1/\varepsilon$  for all  $i$ . The class of concave functions discussed in [\[Andrews 2007\]](#) then provide an interesting class of admissible speeds.
- (5) *Convex homogeneous combinations:* Let  $\phi$  satisfy conditions (i)–(iv) and (vi)–(vii) on a cone  $\tilde{\Gamma} \subset \mathbb{R}^k$ , and suppose that the functions  $f_1, \dots, f_k$  define admissible speeds on a cone  $\Gamma_k \subset \mathbb{R}^n$ . Then the function

$$f(z_1, \dots, z_n) := \phi(f_1(z_1, \dots, z_n), \dots, f_k(z_1, \dots, z_n))$$

on the cone  $\{z \in \Gamma : (f_1(z), \dots, f_k(z)) \in \tilde{\Gamma}\}$  defines an admissible speed. For example, the function  $f_\varepsilon(z_1, \dots, z_n) = H_p(z_1 + \varepsilon H, \dots, z_n + \varepsilon H)$  on the cone  $\Gamma_\varepsilon := \{z \in \mathbb{R}^n : z_i + \varepsilon H > 0 \text{ for all } i\}$  defines an admissible speed.

Curvature problems of the form (1-1)–(1-2) have been studied extensively, although mostly under the assumption that the initial hypersurface is locally convex, that is, having Weingarten map everywhere positive definite. The most well-known result in this case is Huisken’s theorem [\[1984\]](#), which states that, when the speed is given by the mean curvature, uniformly locally convex initial hypersurfaces remain uniformly locally convex and shrink to round points, “round” meaning that the solution approaches total umbilicity at the final point. Chow showed that this behaviour is true also for the flows by the  $n$ -th root of the Gauss curvature [\[1985\]](#), and, if an initial curvature pinching condition is assumed, the square root of the scalar curvature [\[1987\]](#). Each of these flows satisfy conditions (i)–(iv) on the positive cone  $\Gamma = \Gamma_+ := \{x \in \mathbb{R}^n : x_i > 0, i = 1, \dots, n\}$ . More general degree-one homogeneous speeds were treated

in [Andrews 1994a; 2007; 2010], where it was shown that uniformly convex hypersurfaces will contract to round points under the flow (1-1)–(1-2), so long as the speed satisfies conditions (i)–(iv) and, in addition, either

- (1)  $n = 2$ , or
- (2)  $f$  is convex, or
- (3)  $f$  is concave, and *inverse concave*, that is, the function

$$f_*(z_1, \dots, z_n) = f(z_1^{-1}, \dots, z_n^{-1})^{-1}$$

is concave.

These conditions were weakened in [Andrews et al. 2013b], and their necessity demonstrated by the construction, in dimensions  $n > 2$ , of concave speed functions satisfying conditions (i)–(iv) for which convex initial hypersurfaces do not remain convex under the corresponding flow [ibid., Theorem 3].

In the case of nonconvex initial hypersurfaces, much less is known about the behaviour of solutions of (1-1), although in many cases the analogy with the mean curvature flow continues. For example, a simple calculation shows that spheres shrink to points in finite time under flows (1-1)–(1-2) satisfying conditions (i)–(iv). The avoidance principle (see<sup>1</sup> [Andrews et al. 2013a, Theorem 5]) then implies that any compact solution of (1-1) must become singular in finite time. If, in addition, the flow admits second derivative Hölder estimates (for example, if the speed function is a concave or convex function of the principal curvatures [Evans 1982; Krylov 1982], or if  $n = 2$  [Andrews 2004]), one can deduce, by standard methods, that a singularity is characterised by a curvature blow-up [Andrews et al. 2012].

For the mean curvature flow, a crucial part of the current understanding of singularities is the asymptotic convexity estimate of Huisken and Sinestrari [1999a], which states that any mean convex initial hypersurface flowing by mean curvature becomes weakly convex at a singularity. This, together with the monotonicity formula of Huisken [1990] and the Harnack inequality of Hamilton [1995b] allows a rather complete description of singularities in the positive mean curvature case. We note that asymptotic convexity is necessary for the application of the Harnack inequality to deduce that “fast-forming” or “type-II” singularities are asymptotic to convex translation solutions of the flow.

For other flows, the understanding of singularities is far less developed, except in some specific settings such as axial symmetry (see [McCoy et al. 2014], for example). There are several reasons for this: First, there is no analogue available for the monotonicity formula, which is used to show that “slowly forming” or “type-I” singularities of the mean curvature flow are asymptotically self-similar. Second, there is in general no Harnack inequality available sufficient to classify type-II singularities, although the latter is known for quite a wide subclass of flows [Andrews 1994b]. And finally, there is so far no analogue of the Huisken–Sinestrari asymptotic convexity estimate for most other flows, with the notable exception of the recent result of Alessandrini and Sinestrari, which applies to a class of flows by functions of the mean

---

<sup>1</sup>We remark that the avoidance principle proved in [Andrews et al. 2013a, Theorem 5] is not in general true when the cone of definition of the speed is nonconvex. However, a slight modification reveals that it is still possible to compare compact solutions with spheres.

curvature having a certain asymptotic behaviour [Alessandroni and Sinestrari 2010]. In a companion paper [Andrews et al. 2012], we were able to exploit the simplified structure of the evolution equation for the second fundamental form in two dimensions (see also [Schulze 2006; Andrews 2007; McCoy 2011]) to prove that an asymptotic convexity estimate holds in surprising generality for flows of surfaces, namely for any surface flow (1-1)–(1-2) satisfying conditions (i)–(iv). On the other hand, one would expect this result should fail in higher dimensions in such generality, due to the aforementioned examples of “nice” speeds which fail to preserve local convexity of initial data. In this paper, we show that an asymptotic convexity estimate is possible in higher dimensions in the presence of the additional convexity conditions (vi)–(vii).

**Theorem 1.1.** *Let  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a solution of (1-1) with  $s$  an admissible speed. Then for all  $\varepsilon > 0$  there is a constant  $C_\varepsilon > 0$  such that*

$$-\kappa_1(x, t) \leq \varepsilon s(x, t) + C_\varepsilon$$

for all  $(x, t) \in M \times [0, T)$ .

The proof of Theorem 1.1 utilises a Stampacchia–De Giorgi iteration procedure analogous to those of [Huisken 1984; Huisken and Sinestrari 1999b; 1999a; Chow 1985; 1987] (see also [Andrews et al. 2012]), in contrast to the result of [Alessandroni and Sinestrari 2010] (see also [Schulze 2006]), which is proved using the maximum principle. We remark that, by carefully constructing our curvature pinching function, we are able to avoid the rather technical induction on the elementary symmetric functions of curvature that is necessary in [Huisken and Sinestrari 1999a].

Combining Theorem 1.1 with the Harnack estimate of [Andrews 1994b] (see also [Hamilton 1995b]) as in [Huisken and Sinestrari 1999b; 1999a], we are led to the following classification of type-II blow-up limits about type-II singularities.

**Corollary 1.2.** *If  $s$  is an admissible speed, then any type-II blow-up limit of a solution of the corresponding flow (1-1) about a type-II singularity decomposes as a product  $X : \Sigma^k \times \mathbb{R}^{n-k} \rightarrow \mathbb{R}^{n+1}$ , such that  $X|_{\Sigma^k} : \Sigma^k \times \mathbb{R} \rightarrow \mathbb{R}^{k+1} \subset \mathbb{R}^{n+1}$  is a strictly convex ( $k$ -dimensional) translation solution of the flow (1-1).*

Corollary 1.2 is proved in Section 6.

## 2. Notation and preliminary results

We now describe some important background results necessary for the subsequent sections. We begin with flow-independent results to do with symmetric functions, and prove, in Lemma 2.2, that each of the ancillary conditions (xi)–(xiii) implies condition (vii). We then discuss flow-dependent results, and prove, in Lemma 2.4, that each of the ancillary conditions (viii)–(x) implies condition (v). We follow the conventions of [Andrews et al. 2013b; Andrews 2007; 2010; McCoy 2005], where proofs or references for much of this section may be found. Many of the results can also be found in the book [Gerhardt 2006].

The curvature function  $f$  is a smooth, symmetric function defined on an open, convex, symmetric cone  $\Gamma$ . Denote by  $\mathcal{S}_\Gamma$  the cone of symmetric  $n \times n$  matrices with  $n$ -tuple of eigenvalues,  $\lambda := (\lambda_1, \dots, \lambda_n)$ , lying in  $\Gamma$ . A result of Glaeser [1963] implies that there is a smooth,  $\text{GL}(n)$ -invariant function  $F : \mathcal{S}_\Gamma \rightarrow \mathbb{R}$

such that  $f(\lambda(A)) = F(A)$ . The invariance of  $F$  under similarity transformations implies that the speed  $s(x, t) = f(\kappa_1(x, t), \dots, \kappa_n(x, t))$  is a well-defined, smooth function of the Weingarten map  ${}^{\circ}\mathcal{W}$ , that is,  $s(x, t) = F({}^{\circ}\mathcal{W}(x, t)) := F(W(x, t))$ , where  $W(x, t)$  is the component matrix of  ${}^{\circ}\mathcal{W}(x, t)$  with respect to some basis for  $T_x^*M \otimes T_xM$ . If we restrict attention to orthonormal bases, then  $W_i^j = h_{ij}$ , where the  $h_{ij}$  are the components of the second fundamental form.

We shall use dots to indicate derivatives of  $f$  and  $F$  as follows:

$$\begin{aligned} \dot{f}^i(\lambda)v_i &:= \left. \frac{d}{ds} \right|_{s=0} f(\lambda + sv), & \ddot{f}^{ij}(\lambda)v_iv_j &:= \left. \frac{d^2}{ds^2} \right|_{s=0} f(\lambda + sv), \\ \dot{F}^{ij}(A)B_{ij} &:= \left. \frac{d}{ds} \right|_{s=0} F(A + sB), & \ddot{F}^{pq,rs}(A)B_{pq}B_{rs} &:= \left. \frac{d^2}{ds^2} \right|_{s=0} F(A + sB). \end{aligned} \tag{2-1}$$

The derivatives of  $f$  and  $F$  are related in the following way:

**Lemma 2.1** [Gerhardt 1990; Andrews 1994a; 2007]. *Suppose that the function  $f$  satisfies condition (i). Define the function  $F : \mathcal{S}_\Gamma \rightarrow \mathbb{R}$  by  $F(A) := f(\lambda(A))$  as above. Then for any diagonal  $A \in \mathcal{S}_\Gamma$  we have*

$$\dot{F}^{kl}(A) = \dot{f}^k(\lambda(A))\delta^{kl}, \tag{2-2}$$

and for any diagonal  $A \in \mathcal{S}_\Gamma$  and symmetric  $B \in \text{GL}(n)$ , we have

$$\ddot{F}^{pq,rs}(A)B_{pq}B_{rs} = \ddot{f}^{pq}(\lambda(A))B_{pp}B_{qq} + 2 \sum_{p>q} \frac{\dot{f}^p(\lambda(A)) - \dot{f}^q(\lambda(A))}{\lambda_p(A) - \lambda_q(A)} (B_{pq})^2. \tag{2-3}$$

Note that (2-3) holds (as a limit) even if  $A$  has eigenvalues of multiplicity greater than one.

In particular, in an orthonormal frame of eigenvectors of  ${}^{\circ}\mathcal{W}$ , we have

$$\begin{aligned} \dot{F}^{kl}({}^{\circ}\mathcal{W}) &= \dot{f}^k(\kappa)\delta^{kl} \\ \ddot{F}^{pq,rs}({}^{\circ}\mathcal{W})B_{pq}B_{rs} &= \ddot{f}^{pq}(\kappa)B_{pp}B_{qq} + 2 \sum_{p>q} \frac{\dot{f}^p(\kappa) - \dot{f}^q(\kappa)}{\kappa_p - \kappa_q} (B_{pq})^2. \end{aligned}$$

Observe that, by (2-2), conditions (i)–(ii) imply that (1-1)–(1-2) is parabolic. The methods of [Gerhardt 2006, Section 2.5] (see also [Giga and Goto 1992] and [Baker 2010]) then imply short time existence of solutions, so long as the principal curvatures of the initial immersion lie in  $\Gamma$ .

It follows from (2-3) that the function  $F$  is convex if and only if the function  $f$  is convex and satisfies  $(\dot{f}^p - \dot{f}^q)(z_p - z_q) \geq 0$ . We now show that in most cases of interest the second condition is automatic.

**Lemma 2.2.** *Suppose that  $f$  satisfies one of the ancillary conditions (xi), (xii) or (xiii). Then  $f$  satisfies condition (vii).*

*Proof.* Suppose first that condition (xi) is satisfied, so that  $\Gamma$  is convex. If  $\Gamma = \Gamma_+$  then the claim is proved in [Andrews 1994a, Lemma 2.2] (see also [Ecker and Huisken 1989]). However, the proof applies to any convex cone: Consider an arbitrary point  $z \in \Gamma$ . Since  $f$  is smooth and convex, for any  $v \in \mathbb{R}^n$  and any

$s \in \mathbb{R}$  such that  $z + sv \in \Gamma$  we have

$$0 \leq \frac{d^2}{ds^2} f(z + sv) = \frac{d}{ds} \dot{f}^i(z + sv)v_i.$$

Therefore, if  $s > 0$ ,

$$\dot{f}^i(z + sv)v_i \geq \dot{f}^i(z)v_i.$$

Setting  $v = -(e_p - e_q)$ , where  $e_i$  is the basis vector in the direction of  $z_i$ , we obtain

$$(\dot{f}^p - \dot{f}^q)|_z \geq (\dot{f}^p - \dot{f}^q)|_{z-s(e_p-e_q)}.$$

If  $z_p \geq z_q$  then there is some  $s_0 > 0$  such that  $(z - s_0(e_p - e_q))_p = (z - s_0(e_p - e_q))_q$ . By the symmetry and convexity of  $\Gamma$ , this point is in  $\Gamma$ . Since  $f$  is symmetric,  $\dot{f}^p = \dot{f}^q$  at this point and the claim follows.

Now suppose that (xii) is satisfied, so that  $f$  extends to a convex, symmetric function on  $\mathbb{R}^n$ . If the extension is smooth, then the claim follows as above. If not, then we need to be more careful; we make use of the fact that the difference quotient  $(f(\gamma(s)) - f(\gamma(t)))/(s - t)$  is nondecreasing in both  $s$  and  $t$  along all lines  $\gamma(s) = z + sv$ .

Consider a point  $z \in \Gamma$  and a direction  $v \in \mathbb{R}^n$ . Then, for any  $s \in \mathbb{R}$  and any  $s_0 > 0$ , we have

$$\frac{f(z + sv) - f(z + s_0v)}{s - s_0} \geq \frac{f(z + sv) - f(z)}{s} \geq \lim_{s \searrow 0} \frac{f(z + sv) - f(z)}{s} = \dot{f}^i|_z v_i.$$

Setting  $v = -(e_p - e_q)$ , it follows that

$$-(\dot{f}^p - \dot{f}^q)|_z = \dot{f}^i|_z v_i \leq \frac{f(z + sv) - f(z + s_0v)}{s - s_0} \leq \lim_{s \nearrow s_0} \frac{f(z + sv) - f(z + s_0v)}{s - s_0} = \psi'_-(0),$$

where we have defined  $\psi(\sigma) := f(z + (\sigma + s_0)v)$ . We note that the left derivative  $\psi'_-(0)$  exists, and is no greater than the right derivative  $\psi'_+$ , by convexity of  $\psi$ . Supposing without loss of generality that  $z_p \geq z_q$ , we may choose  $s_0$  such that  $z_p - s_0 = z_q + s_0$ . With this choice, it is easily checked that  $\psi$  is an even function. Since  $\psi$  is convex, we have

$$\begin{aligned} \psi'_-(0) &\leq \psi'_+(0) = \lim_{s \searrow 0} \frac{\psi(s) - \psi(0)}{s} \\ &= -\lim_{s \nearrow 0} \frac{\psi(-s) - \psi(0)}{s} = -\lim_{s \nearrow 0} \frac{\psi(s) - \psi(0)}{s} = -\psi'_-(0). \end{aligned}$$

It follows that  $\psi'_-(0) \leq 0$  and we obtain  $(\dot{f}^p - \dot{f}^q)|_z \geq 0$  as required.

Finally, suppose that (xiii) is satisfied, so that  $\Gamma \subset \mathbb{R}^2$ . Consider some point  $z \in \Gamma$  and suppose  $p \neq q$  are such that  $z_p \geq z_q$ . Since  $f$  is homogeneous of degree one, we have  $f = \dot{f}^1 z_1 + \dot{f}^2 z_2$ . Then, since  $f$ ,  $\dot{f}^1$  and  $\dot{f}^2$  are positive on  $\Gamma$ , we must have  $z_p > 0$ . Now,

$$2f = 2(\dot{f}^p z_p + \dot{f}^q z_q) = (\dot{f}^p - \dot{f}^q)(z_p - z_q) + (\dot{f}^p + \dot{f}^q)(z_p + z_q),$$

so that

$$(\dot{f}^p - \dot{f}^q)(z_p - z_q) = 2f - (\dot{f}^p + \dot{f}^q)(z_p + z_q).$$



If  $z_p + z_q \leq 0$ , then we are done (since  $f, f^1$  and  $f^2$  are positive). Otherwise,  $z$  lies in the open, symmetric, convex cone  $\{z \in \mathbb{R}^2 : z_1 + z_2 > 0\}$ . But we have just proved that the claim already holds in this case. This completes the proof.  $\square$

In the following, we are interested in the behaviour of solutions of the flow equation (1-1)–(1-2). We consider speeds  $s = f(\kappa)$  such that  $f$  satisfies condition (i), and denote the corresponding function of  $W$  by  $F$ . We will use the following convention in order to simplify notation: If  $g$  satisfies condition (i), and  $G(A) = g(\lambda(A))$  is the corresponding function on  $\mathcal{S}_\Gamma$ , then we write  $g(x, t) \equiv g(\kappa(x, t))$  and  $G(x, t) \equiv G(W(x, t))$ . Similarly,  $\dot{G}(x, t) \equiv \dot{G}(W(x, t))$  and  $\ddot{G}(x, t) \equiv \ddot{G}(W(x, t))$ . This convention makes the notation  $s$  for the speed unnecessary, and from here on the speed will be denoted by  $F$ .

We recall the following evolution equations:

**Lemma 2.3** [Andrews 1994a; 2007; Andrews et al. 2013b; Gerhardt 2006; McCoy 2005]. *Let*

$$X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$$

*be a solution of the flow (1-1)–(1-2) such that  $f$  satisfies conditions (i)–(iii). Then the following evolution equations hold along  $X$ :*

- (1)  $(\partial_t - \mathcal{L})h_i^j = (\nabla_i dF)^j + F h_i^k h_k^j = \ddot{F}^{pq,rs} \nabla_i h_{pq} \nabla^j h_{rs} + \dot{F}^{kl} h_{kl}^2 h_i^j.$
- (2)  $(\partial_t - \mathcal{L})F = F \dot{F}^{kl} h_{kl}^2.$
- (3)  $\partial_t d\mu = -HF d\mu.$
- (4)  $(\partial_t - \mathcal{L})G = (\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} + \dot{G}^{pq} h_{pq} \dot{F}^{kl} h_{kl}^2.$

*Here  $\mathcal{L}$  is the elliptic operator  $\dot{F}^{ij} \nabla_i \nabla_j$ ,  $h_{ij}^2 = h_i^k h_{kj}$ ,  $\mu(t)$  is the measure induced on  $M$  by the immersion  $X(\cdot, t)$ , and  $G$  is any function given by  $G(x, t) := g(\kappa_1(x, t), \dots, \kappa_n(x, t))$  for some smooth, symmetric  $g : \Gamma \rightarrow \mathbb{R}$ .*

Applying the maximum principle to Lemma 2.3(2), we see that  $F$  remains positive for all  $t \in [0, T)$  whenever it is initially positive. It then follows from Euler’s theorem and the monotonicity of  $f$  that the largest principal curvature also remains positive.

In the case that  $g$  is homogeneous of degree one, Euler’s theorem simplifies Lemma 2.3(4) to

$$(\partial_t - \mathcal{L})G = (\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} + \dot{F}^{kl} h_{kl}^2 G. \tag{2-4}$$

It follows that

$$(\partial_t - \mathcal{L})\left(\frac{G}{F}\right) = \frac{1}{F} (\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} - \frac{2}{F} \dot{F}^{kl} \nabla_k F \nabla_l \left(\frac{G}{F}\right). \tag{2-5}$$

Therefore  $\max_{M \times \{t\}} (G/F)$  will be nonincreasing in  $t$  whenever  $G$  satisfies the condition

$$(\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \leq 0. \tag{2-6}$$

These observations help us to find preserved cones for the flow: Suppose that  $f$  satisfies conditions (i)–(iii). If there is a smooth, nonnegative, symmetric, homogeneous degree-one function  $g : \Gamma \rightarrow \mathbb{R}$  such

that

$$(\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) T_{kpq} T_{lrs} \leq 0$$

for any totally symmetric  $T \in \mathbb{R}^n \otimes \mathbb{R}^n \otimes \mathbb{R}^n$ , where  $G$  is the corresponding function on  $\mathcal{S}_\Gamma$ , then any solution of the corresponding flow admits a preserved cone. Namely, the cone

$$\Gamma_0 := \left\{ z \in \mathbb{R}^n : g(z) < \max_{M \times \{0\}} \left( \frac{G}{F} \right) f(z) \right\}$$

is preserved.

In general, finding such a function  $g$  will be highly specific to the choice of flow speed  $f$ , however, in many cases we can be sure preserved cones exists:

**Lemma 2.4.** *Suppose  $f$  satisfies one of the ancillary conditions (viii), (ix), or (x). Then  $f$  satisfies condition (v).*

*Proof.* Suppose that condition (viii) holds, so that the cone  $\Gamma$  is convex. It follows from Lemma 2.2 that condition (vii) holds, so that  $\ddot{F} \geq 0$  by Lemma 2.1. Let  $X$  be a solution of (1-1)–(1-2). Then the Weingarten map of  $X$  satisfies

$$(\partial_t - \mathcal{L})h_i^j \geq \dot{F}^{kl} h_{kl}^2 h_i^j. \tag{2-7}$$

Let  $\Gamma_0$  be the interior of the symmetrised convex conic hull in  $\mathbb{R}^n$  of the principal curvatures of  $X_0$ . Then  $\overline{\Gamma_0} \setminus \{0\} \subset \Gamma$ . The preservation of  $\Gamma_0$  by the flow follows by applying a slight modification of Hamilton’s tensor maximum principle [1986, Section 3] to (2-7) (for details, see [Andrews 2007, Theorem 3.2] and [Andrews and Hopper 2011, Chapter 6]).

Now suppose that (ix) is satisfied, so that  $f$  vanishes on  $\partial\Gamma$ . If  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  is a solution of the corresponding flow, then  $F$  is initially positive, and the maximum principle implies that it remains so. Then we may consider the function  $G_1(x, t) := g_1(\kappa_1(x, t), \dots, \kappa_n(x, t))$ , where  $g_1$  is the function defined by (3-1) of the following section. Observe that  $f$  extends to a convex function on  $\mathbb{R}^n$  by setting  $f = 0$  outside  $\Gamma$ , so that, by Lemma 2.2, condition (vii) holds. Then we may proceed as in Lemma 3.2 to obtain

$$Z := (\dot{G}_1^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}_1^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \leq 0, \tag{2-8}$$

and it follows that  $G_1/F \leq c_0 := \max_{M \times \{0\}} G_1/F$ . So consider  $\Gamma_0 := \{z \in \mathbb{R}^n : g_1(z) < c_0 f(z)\}$ . Since  $g_1(z) = 0$  if and only if  $z \in \overline{\Gamma}_+ \cap \Gamma$  and, by convexity of the extension of  $f$ ,  $\{z \in \mathbb{R}^n : z_1 + \dots + z_n > 0\} \subset \Gamma$ , we have  $(\partial\Gamma \cap \partial\Gamma_0) \setminus \{0\} = \emptyset$ . It follows that  $\Gamma_0$  is a preserved cone.

Finally, consider the case that condition (x) holds, so that  $\Gamma \subset \mathbb{R}^2$ . Observe that, in this case, it is sufficient to obtain an estimate on the pinching ratio of the solution (which in this case follows from an estimate on  $G_1/F$ ), since any open, connected, symmetric cone  $\Gamma$  in  $\mathbb{R}^2$  that contains the positive ray is of the form  $\{z \in \mathbb{R}^2 : z_{\min} > \varepsilon z_{\max}\}$ . However, we can no longer use any convexity properties of  $f$  to control  $G_1/F$ , and the above proof that  $Z \leq 0$  no longer applies. On the other hand, by carefully analysing each of the terms in the expression for  $Z$ , it is possible to write the terms involving second derivatives of the speed as gradient terms, and the remaining terms turn out to be automatically favourable

for obtaining the desired estimate on  $Z$ . We refer the reader to the papers [Andrews 2007; Andrews et al. 2012] for the proof of this assertion.  $\square$

The existence of a preserved cone ensures that the flow is uniformly parabolic:

**Lemma 2.5.** *Let  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a solution of (1-1), with an admissible speed  $F$ . Then there is a constant  $c_1 > 0$  such that for all  $(x, t) \in M \times [0, T)$  it holds that*

$$\frac{1}{c_1}|v|^2 \leq \dot{F}^{kl}(x, t)v_kv_l \leq c_1|v|^2$$

for all  $v \in T_xM$ , where  $|\cdot|$  is the norm induced on  $TM$  by the immersion  $X(\cdot, t)$ .

*Proof.* In an orthonormal frame of eigenvectors of the Weingarten map, we have, by (2-2), that  $\dot{F}^{kl} = \dot{f}^k\delta^{kl}$ . Let  $\Gamma_0$  be a preserved cone for the flow. Since  $\overline{\Gamma_0} \setminus \{0\} \subset \Gamma$ , and  $\dot{f}^k > 0$  on  $\Gamma$  for all  $k$ , we see that the derivatives  $\dot{f}^k$  are bounded by positive constants on the compact set  $K := \{z \in \overline{\Gamma_{c_0}} : |z| = 1\}$ . Since the derivatives  $\dot{f}^k$  are homogeneous of degree zero, these bounds extend to the cone  $\overline{\Gamma_{c_0}} \setminus \{0\}$ , which completes the proof.  $\square$

The following long time existence result then follows using standard methods.

**Proposition 2.6** [Andrews et al. 2012]. *Let  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a maximally extended solution of (1-1), with an admissible speed. Then  $T < \infty$ , and  $\max_{M \times \{t\}} |\mathcal{W}| \rightarrow \infty$  as  $t \rightarrow T$ .*

We now focus on the proof of Theorem 1.1 and Corollary 1.2, so for the rest of the paper we will assume that  $f$  defines an admissible speed, and  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  is a maximally extended solution of the corresponding flow (1-1).

### 3. The pinching function

In this section, we carefully construct an appropriate curvature pinching function to be used in the proof of Theorem 1.1. That is, we construct a smooth, symmetric, homogeneous (degree-one, say) function  $G(x, t) = g(\kappa_1(x, t), \dots, \kappa_n(x, t))$  of the principal curvatures that vanishes only if the hypersurface is weakly convex. Our goal is to show that the ratio  $G/F$  vanishes asymptotically along the flow. In particular, this ratio should be nonincreasing. In view of (2-5) we would therefore like  $G$  to satisfy

$$(\dot{G}^{kl}\ddot{F}^{pq,rs} - \dot{F}^{kl}\ddot{G}^{pq,rs})\nabla_k h_{pq}\nabla_l h_{rs} \leq 0.$$

In fact, as we shall see, the following two estimates will be essential.

**Properties.** (1) *For all  $\varepsilon > 0$ , there is a constant  $c_\varepsilon > 0$  such that*

$$(\dot{G}^{kl}\ddot{F}^{pq,rs} - \dot{F}^{kl}\ddot{G}^{pq,rs})\nabla_k h_{pq}\nabla_l h_{rs} \leq -c_\varepsilon \frac{|\nabla^q \mathcal{W}|^2}{F}$$

whenever  $G > \varepsilon F$ .

(2) *For all  $\varepsilon > 0$ , there is a constant  $\gamma_\varepsilon > 0$  such that*

$$(F\dot{G}^{kl} - G\dot{F}^{kl})h_{kl}^2 \leq -\gamma_\varepsilon F|\mathcal{W}|^2$$

whenever  $G > \varepsilon F$ .

These estimates are needed to show that the positive part of the function  $G_{\varepsilon,\sigma} := (G/F - \varepsilon)F^\sigma$  is bounded in  $L^p(M \times [0, T])$  for any  $\varepsilon > 0$ , so long as  $\sigma$  is sufficiently small. This is done in Section 4. The proof of Theorem 1.1 then follows from standard arguments, which we recall in Section 5. But first, we construct our pinching function. We first try a smoothed out version of the natural choice,  $\max\{-\kappa_1, 0\}$ . The function we obtain possesses the second of the above properties, but the first property only weakly (that is, with  $c_\varepsilon = 0$ ). By making this function slightly more convex (namely, strictly convex in nonradial directions) we are able to obtain a function satisfying both estimates uniformly (without harming the other properties).

We begin with a smooth function  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  which is strictly convex and positive, except on  $\mathbb{R}_+$ , where it vanishes identically. Such a function is easily constructed; for example, we could use

$$\phi(r) = \begin{cases} r^4 e^{-1/r^2} & \text{if } r < 0, \\ 0 & \text{if } r \geq 0. \end{cases}$$

Now consider the following function, defined on  $\Gamma$ :

$$g_1(z) := f(z) \sum_{i=1}^n \phi\left(\frac{z_i}{f(z)}\right). \tag{3-1}$$

Observe that  $g_1$  is nonnegative and vanishes on (and only on)  $\bar{\Gamma}_+ \cap \Gamma$ . Furthermore,  $g_1$  is clearly smooth, symmetric, and homogeneous of degree one. We now calculate

$$\begin{aligned} \dot{g}_1^k &= \dot{f}^k \sum_{i=1}^n \phi\left(\frac{z_i}{f}\right) + \sum_{i=1}^n \dot{\phi}\left(\frac{z_i}{f}\right) \left(\delta_i^k - \frac{z_i}{f} \dot{f}^k\right) \\ &= \dot{\phi}\left(\frac{z_k}{f}\right) + \dot{f}^k \sum_{i=1}^n \left[ \phi\left(\frac{z_i}{f}\right) - \frac{z_i}{f} \dot{\phi}\left(\frac{z_i}{f}\right) \right]. \end{aligned}$$

It follows easily from the convexity of  $\phi$  that  $\phi(r) - r\dot{\phi}(r) \leq \phi(0) = 0$ . Since  $\phi$  is positive and  $\dot{\phi}$  vanishes on  $\mathbb{R}_+$ , we must also have  $\dot{\phi}(r) \leq 0$  for all  $r \in \mathbb{R}$ . Moreover, equality holds in the above inequalities only if  $r \geq 0$ . Therefore  $\dot{g}_1^k(z) \leq 0$  for each  $k$ , with equality if and only if  $z \in \bar{\Gamma}_+ \cap \Gamma$ .

Now compute

$$\ddot{g}_1^{pq} = \ddot{f}^{pq} \sum_{i=1}^n \left[ \phi\left(\frac{z_i}{f}\right) - \frac{z_i}{f} \dot{\phi}\left(\frac{z_i}{f}\right) \right] + \frac{1}{f} \sum_{i=1}^n \ddot{\phi}\left(\frac{z_i}{f}\right) \left(\delta_i^p - \frac{z_i}{f} \dot{f}^p\right) \left(\delta_i^q - \frac{z_i}{f} \dot{f}^q\right).$$

and

$$\dot{g}_1^k \ddot{f}^{pq} - \dot{f}^k \ddot{g}_1^{pq} = \dot{\phi}\left(\frac{z_k}{f}\right) \ddot{f}^{pq} - \frac{\dot{f}^k}{f} \sum_{i=1}^n \ddot{\phi}\left(\frac{z_i}{f}\right) \left(\delta_i^p - \frac{z_i}{f} \dot{f}^p\right) \left(\delta_i^q - \frac{z_i}{f} \dot{f}^q\right). \tag{3-2}$$

This forms a nonpositive definite matrix for each  $k$ . Finally, consider

$$\dot{g}_1^k \frac{\dot{f}^p - \dot{f}^q}{z_p - z_q} - \dot{f}^k \frac{\dot{g}_1^p - \dot{g}_1^q}{z_p - z_q} = \dot{\phi}\left(\frac{z_k}{f}\right) \frac{\dot{f}^p - \dot{f}^q}{z_p - z_q} - \dot{f}^k \frac{\dot{\phi}(z_p/f) - \dot{\phi}(z_q/f)}{z_p - z_q}. \tag{3-3}$$



This is also nonpositive for each  $k$ , since convexity of  $\phi$  implies  $\frac{\dot{\phi}(r) - \dot{\phi}(s)}{r - s} \geq 0$ . Putting (3-2) and (3-3) together using Lemma 2.1, we see that

$$(\dot{G}_1^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}_1^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \leq 0.$$

To obtain the uniform estimate, we modify the function  $g_1$  to introduce a slightly stronger convexity property. We use the good convexity properties of the Euclidean norm: Consider the function  $g$  defined by

$$g := K(g_1, g_2) := \frac{g_1^2}{g_2}, \tag{3-4}$$

where  $g_2$  is a positive, monotone, degree-one homogeneous function of the principal curvatures which is strictly convex in nonradial directions. The function defined by

$$g_2(z) := Rf(z) + \sum_{i=1}^n z_i - |z|$$

has the properties we require, so long as the constant  $R > 0$  may be chosen such that  $g_2 > 0$  (at least along the flow). Let's first show that such a choice is possible.

**Lemma 3.1.** *There exists a constant  $R > 0$  such that*

$$RF(x, t) + H(x, t) - |^q W(x, t)| > 0$$

for all  $(x, t) \in M \times [0, T)$ .

*Proof.* Define  $G_2(x, t) := g_2(\kappa_1(x, t), \dots, \kappa_n(x, t))$ . Since  $F(\cdot, 0) > 0$  and  $M$  is compact, we may choose  $R > 0$  such that  $G_2(\cdot, 0) > 0$ . By (2-4), it suffices to show that

$$(\dot{G}_2^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}_2^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \geq 0.$$

First calculate

$$\dot{g}_2^k = R \dot{f}^k + 1 - \frac{z_k}{|z|}$$

and

$$\ddot{g}_2^{pq} = R \ddot{f}^{pq} - \frac{1}{|z|^3} (|z|^2 \delta_{pq} - z_p z_q).$$

It follows that

$$\dot{g}_2^k \ddot{f}^{pq} - \dot{f}^k \ddot{g}_2^{pq} = \left(1 - \frac{z_k}{|z|}\right) \ddot{f}^{pq} + \frac{\dot{f}^k}{|z|^3} (|z|^2 \delta_{pq} - z_p z_q), \tag{3-5}$$

which, by the Cauchy–Schwarz inequality, is nonnegative definite for each  $k$ .

Finally,

$$\dot{g}_2^k \frac{\dot{f}^p - \dot{f}^q}{z_p - z_q} - \dot{f}^k \frac{\dot{g}_2^p - \dot{g}_2^q}{z_p - z_q} = \left(1 - \frac{z_k}{|z|}\right) \frac{\dot{f}^p - \dot{f}^q}{z_p - z_q} + \frac{1}{|z|} \dot{f}^k,$$

which is also nonnegative definite for each  $k$ . It now follows from (2-2) and (2-3) that

$$(\dot{G}_2^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}_2^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \geq 0$$

as required. □

So the function  $G$  is well defined. We show that it also satisfies property (1) (page 416) weakly:

**Lemma 3.2.** *There is a constant  $c_0 > 0$  such that*

$$G(x, t) \leq c_0 F(x, t)$$

for all  $(x, t) \in M \times [0, T)$ .

*Proof.* By a straightforward calculation, we find

$$(\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) = \dot{K}^1 (\dot{G}_1^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}_2^{pq,rs}) + \dot{K}^2 (\dot{G}_2^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}_2^{pq,rs}) - \dot{F}^{kl} \ddot{K}^{\alpha\beta} \dot{g}_\alpha^p \dot{g}_\beta^q$$

at any diagonal matrix. Noting that  $\dot{K}^1(x, y) > 0$ ,  $\dot{K}^2(x, y) < 0$  and  $\ddot{K}(x, y) \geq 0$  whenever  $x$  and  $y$  are positive, we see that

$$(\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \leq 0. \tag{3-6}$$

In view of (2-5), the claim now follows from the maximum principle. □

We now show that  $G$  satisfies the required properties (1) and (2) (page 416) uniformly:

**Lemma 3.3.** *For all  $\varepsilon > 0$  there exist constants  $c_2 > 0$  and  $\gamma > 0$  such that*

$$-c_2 \frac{|\nabla^q W|^2}{F} \leq (\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \leq -\frac{1}{c_2} \frac{|\nabla^q W|^2}{F} \tag{3-7}$$

and

$$(F \dot{G}^{kl} - G \dot{F}^{kl}) h_{kl}^2 \leq -\gamma F |\nabla^q W|^2 \tag{3-8}$$

whenever  $G > \varepsilon F$ .

*Proof.* Let  $A \in GL(n)$  be a diagonal matrix and  $T \in \mathbb{R}^n \otimes \mathbb{R}^n \otimes \mathbb{R}^n$  be a totally symmetric tensor. Define

$$Q(A, T) := -(\dot{G}^{kl} \ddot{F}^{pq,rs} - \dot{F}^{kl} \ddot{G}^{pq,rs})|_A T_{kpq} T_{lrs} \geq 0. \tag{3-9}$$

Recalling the application of the Cauchy–Schwarz inequality to (3-5) reveals that equality occurs in (3-9) only if  $T$  is radial, that is, if for each  $k$  we have  $T_{kpq} = \mu_k A_{pq}$  for some constant  $\mu_k$ .

Define the set  $\Gamma_\varepsilon := \{x \in \Gamma : \varepsilon f(z) \leq g(z) \leq c_0 f(z)\}$ . Then, to prove (3-7), we need to demonstrate uniform positive bounds for  $FQ(A, T)$  whenever  $A$  has eigenvalues in  $\Gamma_\varepsilon$  and  $|T| \neq 0$ . Since  $Q$  is homogeneous of degree two with respect to  $T$ , we may assume without loss of generality that  $|T| = 1$ . Moreover, since  $Q$  is homogeneous of degree  $-1$  with respect to  $A$ , it suffices to obtain the required bounds on the compact slice  $K := \{A \in \mathcal{S}_\Gamma : \varepsilon F(A) \leq G(A) \leq c_0 F(A), |A| = 1\}$ . The upper bound now follows immediately from the continuity of  $Q$ .

To prove the lower bound, it suffices to show that  $Q(A, T) = 0$  for  $A \in K$  only if  $|T| = 0$ . We have seen that  $Q(A, T)$  can only vanish if  $T$  is radial. Then, since  $A$  is diagonal, it follows that  $T$  is also diagonal:  $T_{klm} \neq 0$  only if  $k = l = m$ . Since  $A \neq 0$ , there is some  $p$  for which  $\lambda_p(A) \neq 0$ . But, since  $T_{klm} = \mu_k A_{lm} = \mu_k \lambda_l(A) \delta_{lm}$ , we have for any  $k$

$$T_{kkk} = \frac{\lambda_k(A)}{\lambda_p(A)} T_{kpp}.$$

But  $T_{kpp}$  vanishes unless  $k = p$ . Thus  $T$  has at most one nonzero component:  $T_{ppp}$ . It follows that  $A$  has at most one nonzero eigenvalue: If instead we had  $\lambda_q > 0$  for some  $q \neq p$ , then we could obtain the contradiction  $T_{ppp} = (\lambda_p/\lambda_q)T_{qpp} = 0$ . Since  $A \in \mathcal{S}_{\Gamma_\varepsilon} \subset \mathcal{S}_\Gamma$ , we must have  $\lambda_p(A) > 0$ . But this implies that  $G(A) = 0$ , so that  $A \notin K$ , a contradiction. Therefore  $Q$  can only vanish if  $T$  vanishes. This completes the proof of (3-7).

For the second estimate, we observe that, in an orthonormal basis of eigenvectors of  ${}^{\circ}W$ ,

$$(F\dot{G}^{kl} - G\dot{F}^{kl}) \leq F\dot{G}^{kl} = F\dot{g}^k\delta^{kl} \leq 2F\frac{g_1}{g_2}\dot{g}_1^{kl}\delta^{kl}.$$

Now  $g_1/g_2$  is positive on  $\Gamma_\varepsilon$  and therefore has a strictly positive lower bound on the compact slice  $\Gamma_\varepsilon \cap \{|z| = 1\}$ . Similarly,  $\dot{g}_1^k < 0$  on  $\Gamma_\varepsilon$ , and therefore has a strictly negative upper bound on  $\Gamma_\varepsilon \cap \{|z| = 1\}$ . Since both terms are homogeneous of degree zero, these bounds extend unharmed to  $\Gamma_\varepsilon$ , and the claim follows.  $\square$

Now consider, for some positive constants  $\varepsilon$  and  $\sigma$ , the function

$$G_{\varepsilon,\sigma} := \left(\frac{G}{F} - \varepsilon\right)F^\sigma.$$

Observe that the upper bound  $G/F < c_0$  implies

$$G_{\varepsilon,\sigma} < c_0F^\sigma. \quad (3-10)$$

**Lemma 3.4.** *The function  $G_{\varepsilon,\sigma}$  satisfies the evolution equation*

$$\begin{aligned} (\partial_t - \mathcal{L})G_{\varepsilon,\sigma} &= F^{\sigma-1}(\dot{G}^{kl}\ddot{F}^{pq,rs} - \dot{F}^{kl}\ddot{G}^{pq,rs})\nabla_k h_{pq}\nabla_l h_{rs} \\ &\quad + \frac{2(1-\sigma)}{F}\langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F - \frac{\sigma(1-\sigma)}{F^2}|\nabla F|_F^2 + \sigma G_{\varepsilon,\sigma}|{}^{\circ}W|_F^2, \end{aligned} \quad (3-11)$$

where we have introduced the notation  $\langle u, v \rangle_F := \dot{F}^{kl}u_k v_l$  and  $|{}^{\circ}W|_F^2 := \dot{F}^{kl}h_{kl}^2$ .

*Proof.* We first compute

$$\nabla G_{\varepsilon,\sigma} = F^{\sigma-1}\left(\nabla G - \frac{G}{F}\nabla F\right) + \frac{\sigma}{F}G_{\varepsilon,\sigma}\nabla F.$$

It follows that

$$\mathcal{L}G_{\varepsilon,\sigma} = F^{\sigma-1}\left(\mathcal{L}G - \frac{G}{F}\mathcal{L}F\right) + \frac{\sigma}{F}G_{\varepsilon,\sigma}\mathcal{L}F - 2\frac{\sigma-1}{F}\langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F - \frac{\sigma(1-\sigma)}{F^2}G_{\varepsilon,\sigma}|\nabla F|_F^2. \quad (3-12)$$

Therefore,

$$\begin{aligned} (\partial_t - \mathcal{L})G_{\varepsilon,\sigma} &= F^{\sigma-1}\left((\partial_t - \mathcal{L})G - \frac{G}{F}(\partial_t - \mathcal{L})F\right) + \frac{\sigma}{F}G_{\varepsilon,\sigma}(\partial_t - \mathcal{L})F \\ &\quad + 2\frac{1-\sigma}{F}\langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F - \frac{\sigma(1-\sigma)}{F^2}G_{\varepsilon,\sigma}|\nabla F|_F^2 \\ &= F^{\sigma-1}(\dot{G}^{kl}\ddot{F}^{pq,rs} - \dot{F}^{kl}\ddot{G}^{pq,rs})\nabla_k h_{pq}\nabla_l h_{rs} + \sigma G_{\varepsilon,\sigma}|h|_F^2 \\ &\quad + 2\frac{1-\sigma}{F}\langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F + \frac{\sigma(1-\sigma)}{F^2}G_{\varepsilon,\sigma}|\nabla F|_F^2 \end{aligned}$$

as required.  $\square$

Just as for the mean curvature flow, it is the final two terms of the evolution equation (3-11) that obstruct the application of the maximum principle. We will proceed by the Stampacchia–De Giorgi iteration method as applied in [Huisken 1984; Huisken and Sinestrari 1999b]. The first step is to show that the spatial  $L^p$  norms of the positive part,  $(G_{\varepsilon,\sigma})_+ := \max\{G_{\varepsilon,\sigma}, 0\}$ , of  $G_{\varepsilon,\sigma}$  are nonincreasing in  $t$ , so long as  $\sigma$  is sufficiently small. As in [Huisken 1984; Huisken and Sinestrari 1999b; 1999a], this leads to a uniform upper bound on  $G_{\varepsilon,\sigma}$  for small, nonzero  $\sigma$ .

#### 4. The integral estimates

**Proposition 4.1.** *For all  $\varepsilon > 0$  there exist constants  $\ell, L > 0$  such that for all  $p > L$  and  $0 < \sigma < \ell p^{-\frac{1}{2}}$ , the  $L^p(M, \mu(t))$  norm of  $(G_{\varepsilon,\sigma})_+$  is nonincreasing in  $t$ .*

To simplify notation somewhat, we fix  $\varepsilon > 0$  and denote  $E := (G_{\varepsilon,\sigma})_+$ . Then  $E^p$  is  $C^1$  in  $t$  for  $p > 1$ , with  $\partial_t E^p = p E^{p-1} \partial_t G_{\varepsilon,\sigma}$ . The evolution equation (3-11) for  $G_{\varepsilon,\sigma}$  then implies

$$\begin{aligned} \frac{d}{dt} \int E^p d\mu &= p \int E^{p-1} \mathcal{L}G_{\varepsilon,\sigma} d\mu - p \int E^{p-1} F^{\sigma-1} Q d\mu \\ &\quad + 2(1-\sigma)p \int E^{p-1} \frac{\langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F}{F} d\mu - \sigma(1-\sigma)p \int E^p \frac{|\nabla F|_F^2}{F^2} d\mu \\ &\quad + \sigma p \int E^p |\mathcal{W}|_F^2 d\mu - \int E^p HF d\mu, \end{aligned} \tag{4-1}$$

where we have defined  $Q = (\dot{F}^{kl} \ddot{G}^{pq,rs} - \dot{G}^{kl} \ddot{F}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs}$ . It will be useful to estimate  $|\nabla F|_F$  in terms of  $|\nabla \mathcal{W}|$ :

**Lemma 4.2.** *There is a constant  $c_3 > 0$  for which  $|\nabla F|_F^2 \leq c_3 |\nabla \mathcal{W}|^2$ .*

*Proof.* Since  $\nabla_k F = \dot{f}^p \nabla_k h_{pp}$  in an orthonormal basis of eigenvectors of  $\mathcal{W}$ , the claim follows from the uniform positive bounds on  $\dot{f}^i$  along the flow. □

For  $p > 2$ , we can integrate the first term of (4-1) by parts:

$$\int E^{p-1} \mathcal{L}G_{\varepsilon,\sigma} d\mu = -(p-1) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|_F^2 d\mu - \int E^{p-1} \ddot{F}^{kl,rs} \nabla_k h_{rs} \nabla_l G_{\varepsilon,\sigma} d\mu.$$

The first term on the right will be useful. We estimate the second term (when  $G_{\varepsilon,\sigma} > 0$ ) using Young’s inequality as follows:

$$\begin{aligned} -\ddot{F}^{kl,rs} \nabla_k h_{rs} \nabla_l G_{\varepsilon,\sigma} &\leq \frac{2c_4}{F} \sum_{k,l,r,s} |\nabla_k h_{rs} \nabla_l G_{\varepsilon,\sigma}| \\ &\leq c_4 E \sum_{k,l,r,s} \left( \frac{(\nabla_k h_{rs})^2}{p^{\frac{1}{2}} F^2} + \frac{p^{\frac{1}{2}} (\nabla_l G_{\varepsilon,\sigma})^2}{E^2} \right) \\ &= c_4 E \left( p^{-\frac{1}{2}} \frac{|\nabla \mathcal{W}|^2}{F^2} + p^{\frac{1}{2}} \frac{|\nabla G_{\varepsilon,\sigma}|^2}{E^2} \right), \end{aligned} \tag{4-2}$$

where we have estimated each of the homogeneous terms  $\ddot{F}^{kl,rs}$  above by  $2c_4/F$ .



A useful term is obtained from the second term of (4-1) using the first estimate of Lemma 3.3. We estimate the third term using Young’s inequality as follows:

$$\int E^p \left\langle \frac{\nabla G_{\varepsilon,\sigma}}{E}, \frac{\nabla F}{F} \right\rangle_F d\mu \leq \frac{p^{\frac{1}{2}}}{2} \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|_F^2 d\mu + \frac{p^{-\frac{1}{2}}}{2} \int E^p \frac{|\nabla F|_F^2}{F} d\mu. \tag{4-3}$$

Putting this back together, we obtain:

**Lemma 4.3.** *For all  $\sigma \in (0, 1)$  it holds that*

$$\begin{aligned} \frac{d}{dt} \int E^p d\mu \leq & ((c_1 + c_4)p^{\frac{3}{2}} - c_1^{-1}p(p-1)) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|^2 d\mu \\ & + \left( (c_3 + c_4)p^{\frac{1}{2}} - \frac{1}{c_0 c_2} p \right) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu + c_5(\sigma p + 1) \int E^p |{}^q W|^2 d\mu. \end{aligned} \tag{4-4}$$

*Proof.* Since  $-HF/|{}^q W|^2$  is homogeneous of degree zero in the principal curvatures, it may be estimated above by some constant, which allows us to estimate the final term in (4-1). Now apply the estimates of Lemmata 2.5, 4.2 and 3.3, and the inequalities (3-10), (4-2) and (4-3) to the remaining terms.  $\square$

Notice that, for any fixed large  $p$ , the first two terms of (4-4) become nonpositive for sufficiently small  $\sigma$  (of order  $p^{-\frac{1}{2}}$ ). We now estimate the final term in a similar fashion.

**Proposition 4.4.** *There are positive constants  $A_1, A_2, A_3, B_1, B_2$ , independent of  $p$  and  $\sigma$ , such that*

$$\int E^p |{}^q W|^2 \leq (A_1 p^{\frac{3}{2}} + A_2 p^{\frac{1}{2}} + A_3) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|^2 d\mu + (B_1 p^{\frac{1}{2}} + B_2) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu. \tag{4-5}$$

*Proof.* We begin with the commutation formula (see [Andrews and Baker 2010, Proposition 5])

$$\nabla_k \nabla_l h_{pq} = \nabla_p \nabla_q h_{kl} + h_{kl} h_{pq}^2 - h_{pq} h_{kl}^2 + h_{kq} h_{pl}^2 - h_{pl} h_{kq}^2,$$

which holds on a general hypersurface of  $\mathbb{R}^{n+1}$ . This contracts to the Simons-type identity

$$\mathcal{L}h_{pq} = \dot{F}^{kl} \nabla_p \nabla_q h_{kl} + F h_{pq}^2 - \dot{F}^{kl} h_{pq} h_{kl}^2 + \dot{F}^{kl} h_{kq} h_{pl}^2 - \dot{F}^{kl} h_{pl} h_{kq}^2.$$

Contracting further with  $\dot{G}$  yields

$$\dot{G}^{pq} \mathcal{L}h_{pq} = \dot{G}^{pq} \dot{F}^{kl} \nabla_p \nabla_q h_{kl} + (F \dot{G}^{kl} - G \dot{F}^{kl}) h_{kl}^2.$$

On the other hand, we have

$$\dot{F}^{kl} \nabla_p \nabla_q h_{kl} = \nabla_p \nabla_q F - \ddot{F}^{kl,rs} \nabla_p h_{rs} \nabla_q h_{kl},$$

so that

$$\dot{G}^{pq} \mathcal{L}h_{pq} = \dot{G}^{pq} \nabla_p \nabla_q F - \dot{G}^{pq} \ddot{F}^{kl,rs} \nabla_p h_{rs} \nabla_q h_{kl} + (F \dot{G}^{kl} - G \dot{F}^{kl}) h_{kl}^2. \tag{4-6}$$

We now recall (3-12):

$$\begin{aligned} \mathcal{L}G_{\varepsilon,\sigma} &= F^{\sigma-1} \left( \mathcal{L}G - \frac{G}{F} \mathcal{L}F \right) + \frac{\sigma}{F} G_{\varepsilon,\sigma} \mathcal{L}F - 2 \frac{1-\sigma}{F} \langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F + \frac{\sigma(1-\sigma)}{F^2} G_{\varepsilon,\sigma} |\nabla F|_F^2 \\ &= F^{\sigma-1} \left( \dot{F}^{kl} \dot{G}^{pq} \nabla_k \nabla_l h_{pq} + \dot{F}^{kl} \ddot{G}^{pq,rs} \nabla_k h_{pq} \nabla_l h_{rs} - \frac{G}{F} \mathcal{L}F \right) \\ &\quad + \frac{\sigma}{F} G_{\varepsilon,\sigma} \mathcal{L}F - 2 \frac{1-\sigma}{F} \langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F + \frac{\sigma(1-\sigma)}{F^2} G_{\varepsilon,\sigma} |\nabla F|_F^2. \end{aligned} \quad (4-7)$$

Putting (4-6) and (4-7) together, we obtain

$$\begin{aligned} \mathcal{L}G_{\varepsilon,\sigma} &= F^{\sigma-1} (\dot{F}^{kl} \ddot{G}^{pq,rs} - \dot{G}^{kl} \ddot{F}^{pq,rs}) \nabla_k h_{pq} \nabla_l h_{rs} \\ &\quad + F^{\sigma-2} (F \dot{G}^{kl} - G \dot{F}^{kl}) \nabla_k \nabla_l F + F^{\sigma-1} (F \dot{G}^{kl} - G \dot{F}^{kl}) h_{kl}^2 \\ &\quad + \frac{\sigma}{F} G_{\varepsilon,\sigma} \mathcal{L}F - 2 \frac{(1-\sigma)}{F} \langle \nabla F, \nabla G_{\varepsilon,\sigma} \rangle_F + \frac{\sigma(1-\sigma)}{F^2} G_{\varepsilon,\sigma} |\nabla F|_F^2. \end{aligned} \quad (4-8)$$

The first and third terms on the right may be estimated from below using Lemma 3.3.

Applying Young's inequality to the term involving the inner product, we obtain

$$-2 \frac{(1-\sigma)}{F} \langle \nabla F, \nabla G_{\varepsilon,\sigma} \rangle_F \leq (1-\sigma) E \left( \frac{|\nabla F|_F^2}{F^2} + \frac{|\nabla G_{\varepsilon,\sigma}|_F^2}{E^2} \right)$$

wherever  $G_{\varepsilon,\sigma} > 0$ . Recalling the estimates of Lemmata 2.5, 3.3 and 4.2, and Equation (3-10), we obtain

$$\begin{aligned} \mathcal{L}G_{\varepsilon,\sigma} &\leq (c_0 c_1 + c_2 + c_0 c_3) F^\sigma \frac{|\nabla^q W|^2}{F^2} + F^{\sigma-2} (F \dot{G}^{kl} - G \dot{F}^{kl}) \nabla_k \nabla_l F \\ &\quad - \gamma F^\sigma |W|^2 + \frac{\sigma}{F} G_{\varepsilon,\sigma} \mathcal{L}F + c_0 c_1 F^\sigma \frac{|\nabla G_{\varepsilon,\sigma}|_F^2}{E^2}. \end{aligned}$$

Now put the  $\gamma F^\sigma |W|^2$  term on the left, multiply the equation by  $E^p F^{-\sigma}$ , and integrate over  $M$  to obtain

$$\begin{aligned} \gamma \int E^p |W|^2 d\mu &\leq - \int E^p F^{-\sigma} \mathcal{L}G_{\varepsilon,\sigma} d\mu + (c_0 c_1 + c_2 + c_0 c_3) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu \\ &\quad + \int E^p F^{-2} (F \dot{G}^{kl} - G \dot{F}^{kl}) \nabla_k \nabla_l F d\mu \\ &\quad + \sigma \int E^{p+1} F^{-1-\sigma} \mathcal{L}F d\mu + c_0 c_1 \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|_F^2 d\mu. \end{aligned} \quad (4-9)$$

Integrating the first term on the right by parts, we obtain the following estimate:

**Lemma 4.5.** *If  $\sigma \in (0, 1)$  and  $p > 2$ , there are constants  $C_1, C_2, D_1 > 0$ , independent of  $\sigma$  and  $p$ , such that*

$$- \int E^p F^{-\sigma} \mathcal{L}G_{\varepsilon,\sigma} d\mu \leq (C_1 p + C_2) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|_F^2 d\mu + D_1 \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu.$$

*Proof.* Integrating by parts, we find

$$\begin{aligned} &- \int E^p F^{-\sigma} \mathcal{L}G_{\varepsilon,\sigma} d\mu \\ &= p \int E^{p-1} F^{-\sigma} |\nabla G_{\varepsilon,\sigma}|_F^2 d\mu - \sigma \int E^p F^{-\sigma-1} \langle \nabla G_{\varepsilon,\sigma}, \nabla F \rangle_F d\mu + \int E^p F^{-\sigma} \ddot{F}^{kl,rs} \nabla_k h_{rs} \nabla_l G_{\varepsilon,\sigma} d\mu. \end{aligned}$$

Estimating each of the coefficients of  $\ddot{F}$  above by  $2c_4/F$  and applying Young's inequality to the second and third terms, we obtain

$$\begin{aligned}
 - \int E^p F^{-\sigma} \mathcal{L}G_{\varepsilon,\sigma} d\mu &\leq c_0 p \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|_F^2 d\mu + \frac{c_0 \sigma}{2} \int E^p \left( \frac{|\nabla G_{\varepsilon,\sigma}|_F^2}{E^2} + \frac{|\nabla F|_F^2}{F^2} \right) d\mu \\
 &\quad + \frac{c_0 c_4}{2} \int E^p \left( \frac{|\nabla^q W|^2}{F^2} + \frac{|\nabla G_{\varepsilon,\sigma}|^2}{E^2} \right) d\mu.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 - \int E^p F^{-\sigma} \mathcal{L}G_{\varepsilon,\sigma} d\mu &\leq \left( c_0 c_1 p + \frac{c_0 c_1 \sigma}{2} + \frac{c_0 c_4}{2} \right) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|^2 d\mu + \left( \frac{c_0 c_1 c_2 \sigma}{2} + \frac{c_0 c_4}{2} \right) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu. \quad \square
 \end{aligned}$$

In the same way, we obtain the following estimate on the third term of (4-9):

**Lemma 4.6.** *There are constants  $C_3, C_4, D_3, D_4 > 0$ , independent of  $p > 2$  and  $\sigma \in (0, 1)$ , such that*

$$\begin{aligned}
 \int E^p F^{-2} (F \dot{G}^{kl} - G \dot{F}^{kl}) \nabla_k \nabla_l F d\mu &\leq (C_3 p^{\frac{3}{2}} + C_4) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|^2 d\mu + (D_3 p^{\frac{1}{2}} + D_4) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu.
 \end{aligned}$$

And the fourth term:

**Lemma 4.7.** *There are constants  $C_5, C_6, D_5, D_6 > 0$ , independent of  $p$  and  $\sigma$ , such that*

$$\int E^{p+1} F^{-1-\sigma} \mathcal{L}F d\mu \leq (C_5 p^{\frac{3}{2}} + C_6) \int E^{p-2} |\nabla G_{\varepsilon,\sigma}|^2 d\mu + (D_5 p^{\frac{1}{2}} + D_6) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu.$$

This completes the proof of Proposition 4.4.

Combining Proposition 4.4 with Lemma 4.3, we obtain

$$\begin{aligned}
 \frac{d}{dt} \int E^p d\mu &\leq -(c_1 p^2 - \alpha_1 \sigma p^{\frac{5}{2}} - \alpha_2 \sigma p^2 - \alpha_3 p^{\frac{3}{2}} - \alpha_4 p) \int E^{p-2} |G_{\varepsilon,\sigma}|^2 d\mu \\
 &\quad - (\beta_1 p - \beta_2 \sigma p - \beta_3 p^{\frac{1}{2}} - \beta_4) \int E^p \frac{|\nabla^q W|^2}{F^2} d\mu.
 \end{aligned}$$

for some constants  $\alpha_i, \beta_i > 0$ , which are independent of  $\sigma$  and  $p$ . Proposition 4.1 follows easily.

### 5. Proof of Theorem 1.1

We are now able to proceed just as in [Huisken 1984, Section 5] and [Huisken and Sinestrari 1999b, Section 3], using Proposition 4.1 and the following lemma to derive the desired bound on  $G_{\varepsilon,\sigma}$ .

**Lemma 5.1** [Stampacchia 1966]. *Let  $\varphi : [k_0, \infty) \rightarrow \mathbb{R}$  be a nonnegative, nonincreasing function satisfying*

$$\varphi(h) \leq \frac{C}{(h-k)^\alpha} \varphi(k)^\beta, \quad h > k > k_0, \tag{5-1}$$

for some constants  $C > 0$ ,  $\alpha > 0$  and  $\beta > 1$ . Then

$$\varphi(k_0 + d) = 0,$$

where  $d^\alpha = C\varphi(k_0)^{\beta-1}2^{\alpha\beta/(\beta-1)}$ .

Now, given any  $k \geq k_0$ , where  $k_0 := \sup_{\sigma \in (0,1)} \sup_M G_{\varepsilon,\sigma}(\cdot, 0)$ , set

$$v_k(x, t) := (G_{\varepsilon,\sigma}(x, t) - k)_+^{p/2} \quad \text{and} \quad A_k(t) := \{x \in M : v_k(x, t) > 0\}.$$

We will show that  $\varphi(k) = |A_k| := \int_0^T \int_{A_k(t)} d\mu(\cdot, t) dt$  satisfies the conditions of Stampacchia's lemma for some  $k_1 \geq k_0$ . This provides us with a constant  $d$  for which  $|A_{k_1+d}|$  vanishes. [Theorem 1.1](#) then follows. Observe that  $|A_k|$  is nonnegative and nonincreasing with respect to  $k$ . Then we only need to demonstrate that an inequality of the form [\(5-1\)](#) holds.

**Lemma 5.2.** *There are constants  $L_1 \geq L$  and  $c_6 > 0$  such that for all  $p > L_1$  we have*

$$\frac{d}{dt} \int v_k^2 d\mu + c_1^{-1} \int |\nabla v_k|^2 d\mu \leq c_6(\sigma p + 1) \int_{A_k} F^2 G_{\varepsilon,\sigma}^p d\mu. \tag{5-2}$$

*Proof.* Observe that

$$\frac{d}{dt} \int v_k^2 d\mu = \int_{A_k} p(G_{\varepsilon,\sigma} - k)_+^{p-1} \partial_t G_{\varepsilon,\sigma} d\mu - \int v_k^2 H F d\mu.$$

The result is then obtained by proceeding as in [Lemma 4.3](#), applying

$$|\nabla v_k|^2 = \frac{p^2}{4} (G_{\varepsilon,\sigma} - k)_+^{p-2} |\nabla G_{\varepsilon,\sigma}|^2,$$

and estimating  $|\mathcal{W}|^2 \leq C F^2$  using the degree-zero homogeneity of  $|\mathcal{W}|^2/F^2$ . □

Now set  $\sigma' = \sigma + n/p$ . Then

$$\int_{A_k} F^n d\mu \leq \int_{A_k} F^n \frac{(G_{\varepsilon,\sigma})_+^p}{k^p} d\mu = k^{-p} \int_{A_k} (G_{\varepsilon,\sigma'})_+^p d\mu \leq k^{-p} \int (G_{\varepsilon,\sigma'})_+^p d\mu. \tag{5-3}$$

If  $p \geq \max\{L_1, 4n^2/\ell^2\}$  and  $\sigma \leq (\ell/2)p^{-\frac{1}{2}}$ , then  $p \geq L_1$  and  $\sigma' \leq \ell p^{-\frac{1}{2}}$ , so that, by [Proposition 4.1](#),

$$\int_{A_k} F^n d\mu \leq k^{-p} \int (G_{\varepsilon,\sigma'}(\cdot, 0))_+^p d\mu_0 \leq \mu_0(M) \left(\frac{k_0}{k}\right)^p. \tag{5-4}$$

Choosing  $k$  sufficiently large, the right hand side of this inequality can be made arbitrarily small. We will use this fact in conjunction with the following Sobolev inequality to exploit the good gradient term in [\(5-2\)](#).

**Lemma 5.3 [Huisken 1984].** *There is a constant  $c_S$  (independent of  $\sigma$ ,  $p$ , and  $\varepsilon$ ) such that*

$$\left(\int v_k^{2q} d\mu\right)^{\frac{1}{q}} \leq c_S \left(\int |\nabla v_k|^2 d\mu + \left(\int_{A_k} F^n d\mu\right)^{\frac{2}{n}} \left(\int v_k^{2q} d\mu\right)^{\frac{1}{q}}\right), \tag{5-5}$$

where  $q$  is equal to  $n/(n-2)$  if  $n > 2$ , or any positive number if  $n = 2$ .



*Proof.* Since we have the estimate  $H^2 < CF^2$  (by degree-zero homogeneity of the quantity  $H^2/F^2$ ) this follows from the Michael–Simon–Sobolev inequality [1973] just as in [Huisken 1984].  $\square$

It follows from (5-4) and (5-5) that there is some  $k_1 > k_0$  such that for all  $k > k_1$  we have

$$\left( \int v_k^{2q} d\mu \right)^{\frac{1}{q}} \leq 2c_S \int |\nabla v_k|^2 d\mu.$$

Therefore, from (5-2), we have for all  $k > k_1$

$$\frac{d}{dt} \int v_k^2 d\mu + \frac{1}{2c_1c_S} \left( \int v^{2q} d\mu \right)^{\frac{1}{q}} \leq c_6(\sigma p + 1) \int_{A_k} F^2 G_{\varepsilon,\sigma}^p d\mu.$$

Integrating this over time, and noting that  $A_k(0) = \emptyset$ , we find (since we may assume  $2c_1c_S \geq 1$ ) that

$$\sup_{[0,T)} \left( \int_{A_k} v_k^2 d\mu \right) + \int_0^T \left( \int v^{2q} d\mu \right)^{\frac{1}{q}} dt \leq 4c_1c_Sc_6(\sigma p + 1) \int_0^T \int_{A_k} F^2 G_{\varepsilon,\sigma}^p d\mu dt. \tag{5-6}$$

We now exploit the interpolation inequality for  $L^p$  spaces:

$$|f|_{q_0} \leq |f|_r^{1-\theta} |f|_q^\theta, \tag{5-7}$$

where  $\theta \in (0, 1)$  and  $1/q_0 = \theta/q + (1 - \theta)/r$ . Setting  $r = 1$  and  $\theta = 1/q_0$ , we may assume  $1 < q_0 < q$ . Then applying (5-7) we find

$$\int_{A_k} v_k^{2q_0} d\mu \leq \left( \int_{A_k} v_k^2 d\mu \right)^{q_0-1} \left( \int_{A_k} v^{2q} d\mu \right)^{\frac{1}{q}}.$$

Now, applying the Hölder inequality, we find

$$\left( \int_0^T \int_{A_k} v_k^{2q_0} d\mu dt \right)^{\frac{1}{q_0}} \leq \left( \sup_{[0,T)} \int_{A_k} v_k^2 d\mu \right)^{\frac{q_0-1}{q_0}} \left( \int_0^T \left( \int_{A_k} v^{2q} d\mu \right)^{\frac{1}{q}} dt \right)^{\frac{1}{q_0}}.$$

Using Young’s inequality,  $ab \leq (1 - 1/q_0)a^{q_0/(q_0-1)} + (1/q_0)b^{q_0}$ , on the right hand side, we obtain

$$\begin{aligned} \left( \int_0^T \int_{A_k} v_k^{2q_0} d\mu dt \right)^{\frac{1}{q_0}} &\leq \left( 1 - \frac{1}{q_0} \right) \sup_{[0,T)} \int_{A_k} v_k^2 d\mu + \frac{1}{q_0} \int_0^T \left( \int_{A_k} v^{2q} d\mu \right)^{\frac{1}{q}} dt \\ &\leq \sup_{[0,T)} \int_{A_k} v_k^2 d\mu + \int_0^T \left( \int_{A_k} v^{2q} d\mu \right)^{\frac{1}{q}} dt. \end{aligned}$$

Recalling (5-6), we arrive at

$$\left( \int_0^T \int_{A_k} v_k^{2q_0} d\mu dt \right)^{\frac{1}{q_0}} \leq 4c_1c_Sc_6(\sigma p + 1) \int_0^T \int_{A_k} F^2 G_{\varepsilon,\sigma}^p d\mu dt. \tag{5-8}$$

Application of the Hölder inequality yields the inequalities

$$\int_0^T \int_{A_k} F^2 G_{\varepsilon,\sigma}^p d\mu dt \leq |A_k|^{1-\frac{1}{r}} \left( \int_0^T \int_{A_k} F^{2r} G_{\varepsilon,\sigma}^{pr} d\mu dt \right)^{\frac{1}{r}} \leq c_7 |A_k|^{1-\frac{1}{r}} \tag{5-9}$$

and

$$\int_0^T \int_{A_k} v_k^2 d\mu dt \leq |A_k|^{1-\frac{1}{q_0}} \left( \int_0^T \int_{A_k} v_k^{2q_0} d\mu dt \right)^{\frac{1}{q_0}}, \tag{5-10}$$

where the integral on the right hand side of (5-9) was estimated in a similar manner to (5-4), with  $c_7 := k_0^2(T\mu_0(M))^{1/r}$  (so long as  $\sigma \leq (\ell/4)p^{-1/2}$ , and  $2r > L_2 := \max\{L_1, 4n^2/\ell^2, 64/\ell^2\}$ , say). Finally, for  $h > k \geq k_1$  we may estimate

$$|A_h| := \int_0^T \int_{A_h} d\mu dt = \int_0^T \int_{A_h} \frac{(G_{\varepsilon,\sigma} - k)_+^p}{(G_{\varepsilon,\sigma} - k)_+^p} d\mu dt \leq \int_0^T \int_{A_h} \frac{(G_{\varepsilon,\sigma} - k)_+^p}{(h - k)^p} d\mu dt.$$

Since  $A_h(t) \subset A_k(t)$  for all  $t \in [0, T)$ , and  $v_k^2 := (G_{\varepsilon,\sigma} - k)_+^p$ , we obtain

$$(h - k)^p |A_h| \leq \int_0^T \int_{A_k} v_k^2 d\mu dt. \tag{5-11}$$

Putting together estimates (5-8), (5-9), (5-10) and (5-11), we arrive at

$$|A_h| \leq \frac{4c_1 c_5 c_6 c_7 (\sigma p + 1)}{(h - k)^p} |A_k|^\gamma$$

for all  $h > k \geq k_1$ , where  $\gamma := 2 - 1/q_0 - 1/r$ . Now fix  $p := 2L_2$  and choose  $\sigma < (\ell/4)p^{-\frac{1}{2}}$  sufficiently small that  $\sigma p < 1$ . Then, choosing  $r > \max\{q_0/(q_0 - 1), L_2\}$ , so that  $\gamma > 1$ , we may apply Stampacchia’s lemma. We conclude that  $|A_k| = 0$  for all  $k > k_1 + d$ , where  $d^p = c_1 c_5 c_6 c_7 2^{3+\gamma p/(\gamma-1)} |A_{k_1}|^{\gamma-1}$ . We note that  $d$  is finite, since  $T$  is finite and

$$\int_{A_{k_1}} d\mu \leq \int_{A_{k_1}} \frac{(G_{\varepsilon,\sigma})_+^p}{k_1^p} d\mu \leq k_1^{-p} \int (G_{\varepsilon,\sigma})_+^p d\mu \leq k_1^{-p} \int (G_{\varepsilon,\sigma}(\cdot, 0))_+^p d\mu_0,$$

where the final estimate follows from Proposition 4.1.

It follows that

$$G \leq \varepsilon F + (k_1 + d)F^{1-\sigma} \leq 2\varepsilon F + C_\varepsilon$$

for some suitably large constant  $C_\varepsilon > 0$ . Theorem 1.1 follows.

### 6. Rescaling about type-II singularities

We now analyse the structure of fast forming singularities. Let  $X : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a smooth, compact solution of (1-1) satisfying the following ansatz: For all  $C > 0$  there is a time  $t_C \in [0, T)$  such that

$$\max_{x \in M} |{}^q W(x, t)|^2 \geq \frac{C}{T - t} \tag{6-1}$$

for all  $t \in [t_C, T)$ . We say that the flow undergoes a *type-II singularity*. To analyse the shape of type-II singularities, we consider, following Hamilton [1995a] and Huisken and Sinestrari [1999b], the following sequence of parabolic rescalings: For each  $k \in \mathbb{N}$ , choose a sequence  $(t_k)$  of times  $t_k \in [0, T - 1/k)$  and a

sequence  $(x_k)$  of points  $x_k \in M$  such that

$$|\mathcal{W}(x_k, t_k)|^2 \left(T - \frac{1}{k} - t_k\right) = \max_{(x,t) \in M \times [0, T-1/k]} |\mathcal{W}(x, t)|^2 \left(T - \frac{1}{k} - t\right).$$

Now set

$$L_k := |\mathcal{W}(x_k, t_k)|^2, \quad \alpha_k := -L_k t_k, \quad \sigma_k := L_k \left(T - \frac{1}{k} - t_k\right).$$

**Lemma 6.1.** *As  $k \rightarrow \infty$ , we have*

$$t_k \rightarrow T, \quad L_k \rightarrow \infty, \quad \alpha_k \rightarrow -\infty, \quad \sigma_k \rightarrow \infty.$$

*Proof.* By the ansatz (6-1), for all  $R > 0$  there exists  $t_R \in [0, T)$  and  $x_R \in M$  such that

$$|\mathcal{W}(x_R, t_R)|^2 (T - t_R) > 2R.$$

On the other hand, there is some sufficiently large  $k_R \in \mathbb{N}$  such that

$$t_R < T - \frac{1}{k}, \quad |\mathcal{W}(x_R, t_R)|^2 \left(T - \frac{1}{k} - t_R\right) > R$$

for all  $k > k_R$ . Therefore, by definition,

$$\sigma_k \geq |\mathcal{W}(x_R, t_R)|^2 \left(T - \frac{1}{k} - t_R\right) > R$$

for all  $k > k_R$ . Since  $R$  was arbitrary, we find  $\sigma_k \rightarrow \infty$  as  $k \rightarrow \infty$ .

Since  $(T - 1/k - t_k)$  is bounded, it follows from the definition of  $\sigma_k$  that  $L_k \rightarrow \infty$  as  $k \rightarrow \infty$ . Therefore, since  $|\mathcal{W}|$  remains bounded whilst  $t < T$ , we must have  $t_k \rightarrow T$ . It follows that  $\alpha_k \rightarrow -\infty$ .  $\square$

Now consider the rescalings

$$X_k(x, t) = \sqrt{L_k} \left( X \left( x, \frac{t}{L_k} + t_k \right) - X(x_k, t_k) \right) \quad \text{for } t \in [\alpha_k, \sigma_k].$$

It is straightforward to compute

$$\begin{aligned} \frac{\partial X_k}{\partial t}(x, t) &= -L_k^{-\frac{1}{2}} F \left( x, \frac{t}{L_k} + t_k \right) v \left( x, \frac{t}{L_k} + t_k \right); \\ \frac{\partial X_k}{\partial x^i}(x, t) &= \sqrt{L_k} \frac{\partial X}{\partial x^i} \left( x, \frac{t}{L_k} + t_k \right) \Rightarrow (g_k)_{ij}(x, t) = L_k g_{ij} \left( x, \frac{t}{L_k} + t_k \right) \\ &\Rightarrow (g_k)^{ij}(x, t) = \frac{1}{L_k} g^{ij} \left( x, \frac{t}{L_k} + t_k \right); \end{aligned}$$

and

$$\begin{aligned} v_k(x, t) &= v \left( x, \frac{t}{L_k} + t_k \right) \Rightarrow {}^k D_i v_k(x, t) = {}^k D_i v \left( x, \frac{t}{L_k} + t_k \right) \\ &\Rightarrow \mathcal{W}_k(x, t) = L_k^{-\frac{1}{2}} \mathcal{W} \left( x, \frac{t}{L_k} + t_k \right) \\ &\Rightarrow F_k(x, t) = L_k^{-\frac{1}{2}} F \left( x, \frac{t}{L_k} + t_k \right), \end{aligned}$$

where we used the script  $k$  to distinguish quantities related to the rescaling  $X_k$  (in particular,  ${}^k D$  is the pullback of the Euclidean connection along  $X_k$ ). We refer to the sequence  $(X_k)$  as a *blow-up sequence*.

Observe that the rescalings satisfy the flow equation (1-1). We also note the following properties (compare [Huisken and Sinestrari 1999b, Lemma 4.4]):

**Lemma 6.2.** (1) For each  $k \in \mathbb{N}$ ,  $X_k(x_k, 0) = 0$  and  $|\mathring{W}(x_k, 0)| = 1$ .

(2) For any  $\varepsilon > 0$  and  $\Sigma > 0$  there exists  $k_0 \in \mathbb{N}$  such that  $\sigma_k > \Sigma$  and

$$\max_{M \times [\alpha_{k_0}, \Sigma]} |\mathring{W}_k|^2 \leq 1 + \varepsilon \tag{6-2}$$

for all  $k \geq k_0$ .

(3) For any  $\varepsilon > 0$  there exists  $C_\varepsilon$  such that

$$-\kappa_1^{(k)}(x, t) \leq \varepsilon F_k(x, t) + \frac{C_\varepsilon}{\sqrt{L_k}} \tag{6-3}$$

for all  $(x, t) \in M \times [\alpha_k, \sigma_k]$ , where  $\kappa_1^{(k)}$  is the smallest principal curvature of  $X_k$ .

*Proof.* Part (1) is immediate from the definitions and our calculation of  $\mathring{W}_k$ .

To prove part (2), first note that

$$|\mathring{W}_k(x, t)|^2 = L_k^{-1} |\mathring{W}(x, L_k^{-1}t + t_k)|^2.$$

By the definition of  $L_k$  and the choice of  $(x_k, t_k)$  we also have

$$|\mathring{W}(x, L_k^{-1}t + t_k)|^2 \left(T - \frac{1}{k} - (L_k^{-1}t + t_k)\right) \leq L_k \left(T - \frac{1}{k} - t_k\right).$$

Therefore

$$|\mathring{W}_k(x, t)|^2 \leq \frac{T - \frac{1}{k} - t_k}{T - \frac{1}{k} - t_k - L_k^{-1}t} = \frac{\sigma_k}{\sigma_k - t} = 1 + \frac{t}{\sigma_k - t}.$$

Since  $\sigma_k \rightarrow \infty$ , the claim follows.

For part (3), we have

$$\kappa_1^k(x, t) = \frac{1}{\sqrt{L_k}} \kappa_1(x, L_k^{-1}t + t_k).$$

Therefore, by Theorem 1.1, for all  $\varepsilon > 0$  there exists  $C_\varepsilon$  such that

$$-\kappa_1^k(x, t) \leq \frac{1}{\sqrt{L_k}} (\varepsilon F(x, L_k^{-1}t + t_k) + C_\varepsilon) = \varepsilon F_k(x, t) + \frac{C_\varepsilon}{\sqrt{L_k}}$$

for all  $(x, t) \in M \times [-\alpha_k, \sigma_k]$ . □

*Proof of Corollary 1.2.* Since the flow speed is a convex function of the Weingarten map, the flow admits second derivative Hölder estimates, and we may proceed as in [Baker 2011, Section 3], using Lemma 6.2, to obtain a sublimit  $X_\infty : M_\infty \times I_\infty \rightarrow \mathbb{R}^{n+1}$  of the blow-up sequence. Since for each  $k$  the rescaled immersion  $X_k$  is a solution of the flow on the time interval  $[\alpha_k, \sigma_k]$ , we deduce from Lemma 6.1 that  $X_\infty$  is an eternal solution of the flow (1-1) (that is,  $I_\infty = \mathbb{R}$ ). Part (3) of Lemma 6.2 implies that  $X_\infty$

is weakly convex. Applying the strong tensor maximum principle [Hamilton 1982] (see also [Andrews 2007, Theorem 3.1]) to the evolution equation for the Weingarten map

$$\partial_t h_i^j = \mathcal{L}h_i^j + \ddot{F}^{pq,rs} \nabla_i h_{pq} \nabla^j h_{rs} + \dot{F}^{kl} h_{kl}^2 h_i^j,$$

we deduce, just as in [Huisken and Sinestrari 1999a, Theorem 4.1], that the rank of  $\mathcal{W}$  is constant and its null-space is invariant under parallel transport. The same use of Frobenius' theorem as in [Huisken 1993, Theorem 5.1] (compare [Hamilton 1986]) then implies that  $M_\infty$  splits isometrically as a product  $\mathbb{R}^{n-k} \times \Sigma_\infty^k$  for some  $1 \leq k \leq n$ , where  $\Sigma_\infty^k$  is strictly convex. Moreover,  $X_\infty|_{\Sigma_\infty^k}$  solves the flow (1-1) in  $\mathbb{R}^{k+1}$ .

Now observe that, by Lemma 6.2(i) and (ii), the maximum value of  $|\mathcal{W}_\infty|$  is 1, and occurs at  $(x_\infty, 0)$ ; it follows that the maximum value of  $F$  is also attained here. We complete the proof by applying the differential Harnack inequality of [Andrews 1994b] to deduce that  $X_\infty|_{\Sigma_\infty^k}(\Sigma_\infty^k)$  moves by translation (compare [Hamilton 1995b]).

**Proposition 6.3.** *Let  $X : \Sigma^k \times \mathbb{R} \rightarrow \mathbb{R}^{k+1}$  be a strictly convex, eternal solution of (1-1) with admissible speed  $F$  such that  $\sup_{\Sigma \times \mathbb{R}} F$  is attained. Then  $X$  moves by translation.*

*Proof.* Consider the function  $\Phi(A) = -F(A^{-1})$ , where  $F : \mathcal{S}_+ \rightarrow \mathbb{R}$  gives the flow speed as a function of the Weingarten map (here,  $\mathcal{S}_+$  is the cone of symmetric, positive definite matrices). For any  $A \in \mathcal{S}_+$ ,  $B \in GL(n)$ , we have

$$\dot{\Phi}|_A(B) = \frac{d}{ds} \Big|_{s=0} \Phi(A + sB) = -\frac{d}{ds} \Big|_{s=0} F([A + sB]^{-1}) = \dot{F}|_A(A^{-1}BA^{-1})$$

and

$$\ddot{\Phi}|_A(B, B) = \frac{d^2}{ds^2} \Big|_{s=0} \Phi(A + sB) = -\ddot{F}|_A(A^{-1}BA^{-1}, A^{-1}BA^{-1}) - 2\dot{F}|_A(A^{-1}BA^{-1}BA^{-1}).$$

Since  $\ddot{F} \geq 0$ ,  $\dot{F} > 0$ , and  $F > 0$ , it follows that

$$\ddot{\Phi} + \frac{1 - \alpha}{\alpha} \frac{\dot{\Phi} \otimes \dot{\Phi}}{\Phi} \leq 0$$

for all  $\alpha \in (0, 1)$ . That is,  $\Phi$  is  $\alpha$ -concave for all  $\alpha \in (0, 1)$ . Thus Corollary 5.11 of [Andrews 1994b] may be applied. We deduce that any strictly convex solution of (1-1) satisfies

$$\partial_t F - g(\mathcal{W}^{-1}(\text{grad } F), \text{grad } F) + \frac{(\alpha - 1)F}{\alpha(t - t_0)} \geq 0 \tag{6-4}$$

for all  $t > t_0$ , where  $t_0$  is the initial time, and  $\text{grad}$  is the gradient operator on  $M$ . It follows that any strictly convex, eternal solution of (1-1) satisfies

$$P := \partial_t F - g(\mathcal{W}^{-1}(\text{grad } F), \text{grad } F) \geq 0.$$

Moreover, (6-4) is deduced from the maximum principle applied to the time evolution of  $P$ , such that equality is attained at a space-time point only if equality holds identically. Since by assumption  $\sup_{\Sigma \times \mathbb{R}} F$  is attained,  $P$  vanishes identically.

We now recall from [Andrews 1994b, Equation 5.2] that, in the Gauss map parametrisation, the Harnack quantity  $P$  satisfies:

$$(\partial_t - \bar{\mathcal{L}})P = \dot{\Phi}(\text{Id})P + \ddot{\Phi}(\bar{Q}, \bar{Q}),$$

where  $\bar{Q}$  is the time derivative of the inverse of the Weingarten map in the Gauss map parametrisation, and  $\bar{\mathcal{L}}$  is the contraction of the covariant Hessian on  $S^n$  by  $\dot{\Phi}$ . Since  $P$  is identically zero, this simply says  $\ddot{\Phi}(\bar{Q}, \bar{Q}) = 0$ . Recalling the equation for  $\ddot{\Phi}$ , positive definiteness of  $\dot{F}$  and strict convexity of  $\Sigma$  imply that  $\bar{Q}$  must vanish. Returning to the standard parametrisation (for example, using [Andrews 1994b, Lemma 3.10]), we find  $0 = Q = -\mathcal{W}^{-1} \circ (\partial_t \mathcal{W} + \nabla_V \mathcal{W}) \circ \mathcal{W}^{-1}$ , where we have defined the vector field  $V := -\mathcal{W}^{-1}(\text{grad } F)$ . Substituting  $\partial_t \mathcal{W} = \nabla \text{grad } F + F \mathcal{W}^2$ , we have, for all  $u \in T\Sigma$ ,

$$\begin{aligned} 0 &= \nabla_u \text{grad } F + F \mathcal{W}^2(u) + \nabla_u \mathcal{W}(V) \\ &= \nabla_u(\text{grad } F + \mathcal{W}(V)) + \mathcal{W}(F \mathcal{W}(u) - \nabla_u V). \end{aligned}$$

It follows that  $\nabla V - F \mathcal{W} = 0$ .

Now define the vector field  $T := X_* V - F\nu$ . Then, for any  $u \in T\Sigma$ ,

$${}^X D_u T = (\nabla_u V - F \mathcal{W}(u)) - g(\mathcal{W}(V) + \text{grad } F, u)\nu = 0.$$

Furthermore,

$${}^X D_t T = {}^X D_t X_* V - \partial_t F \nu - F \text{grad } F,$$

where  ${}^X D$  is the pullback of the Euclidean connection  $D$  by  $X$ . Since  $P \equiv 0$ , this becomes

$${}^X D_t T = {}^X D_t X_* V - g(\mathcal{W}^{-1}(\text{grad } F), \text{grad } F)\nu - F \text{grad } F = {}^X D_t X_* V + g(V, \text{grad } F)\nu - F \text{grad } F.$$

Since  $V$  is tangential, we have

$$\langle {}^X D_t X_* V, \nu \rangle = -\langle X_* V, {}^X D_t \nu \rangle = -g(V, \text{grad } F).$$

Thus the normal component of  ${}^X D_t T$  is zero. The tangential part of  ${}^X D_t X_* V$  is  $({}^X D_t X_* V)^\top = -F \mathcal{W}(V) = F \text{grad } F$ ; so the tangential component of  ${}^X D_t T$  also vanishes. We have proved that  $T$  is parallel. Now set  $\tilde{X}(x, t) := X(\phi(x, t), t)$ , where  $\phi$  is the solution of  $d\phi^i/dt = V^i$  with initial condition  $\phi(x, 0) = x$ . Then

$$\frac{\partial \tilde{X}}{\partial t} = \frac{\partial X}{\partial x^i} \frac{d\phi^i}{dt} + \frac{\partial X}{\partial t} = T. \quad \square$$

This completes the proof of Corollary 1.2. □

### References

[Alessandrini and Sinestrari 2010] R. Alessandrini and C. Sinestrari, “Convexity estimates for a nonhomogeneous mean curvature flow”, *Math. Z.* **266**:1 (2010), 65–82. MR 2011m:53115 Zbl 1197.53080

[Andrews 1994a] B. Andrews, “Contraction of convex hypersurfaces in Euclidean space”, *Calc. Var. Partial Differential Equations* **2**:2 (1994), 151–171. MR 97b:53012 Zbl 0805.35048

[Andrews 1994b] B. Andrews, “Harnack inequalities for evolving hypersurfaces”, *Math. Zeitschrift* **217**:2 (1994), 179–197. MR 95j:58178 Zbl 0807.53044

[Andrews 2004] B. Andrews, “Fully nonlinear parabolic equations in two space variables”, preprint, 2004. arXiv math/0402235



- [Andrews 2007] B. Andrews, “Pinching estimates and motion of hypersurfaces by curvature functions”, *J. Reine Angew. Math.* **608** (2007), 17–33. [MR 2008i:53087](#) [Zbl 1129.53044](#)
- [Andrews 2010] B. Andrews, “Moving surfaces by non-concave curvature functions”, *Calc. Var. Partial Differential Equations* **39**:3-4 (2010), 649–657. [MR 2011k:53082](#) [Zbl 1203.53062](#)
- [Andrews and Baker 2010] B. Andrews and C. Baker, “Mean curvature flow of pinched submanifolds to spheres”, *J. Differential Geom.* **85**:3 (2010), 357–395. [MR 2012a:53122](#) [Zbl 1241.53054](#)
- [Andrews and Hopper 2011] B. Andrews and C. Hopper, *The Ricci flow in Riemannian geometry: a complete proof of the differentiable 1/4-pinching sphere theorem*, Lecture Notes in Math. **2011**, Springer, Heidelberg, 2011. [MR 2012d:53208](#) [Zbl 1214.53002](#)
- [Andrews et al. 2012] B. Andrews, M. Langford, and J. A. McCoy, “Convexity estimates for surfaces moving by curvature functions”, preprint, 2012. To appear in *J. Differential Geom.*
- [Andrews et al. 2013a] B. Andrews, M. Langford, and J. A. McCoy, “Non-collapsing in fully non-linear curvature flows”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **30**:1 (2013), 23–32. [MR 3011290](#) [Zbl 1263.53059](#)
- [Andrews et al. 2013b] B. Andrews, J. A. McCoy, and Y. Zheng, “Contracting convex hypersurfaces by curvature”, *Calc. Var. Partial Differential Equations* **47**:3-4 (2013), 611–665. [MR 3070558](#) [Zbl 06187283](#)
- [Baker 2010] R. C. Baker, *The mean curvature flow of submanifolds of high codimension*, thesis, Australian National University, Canberra, 2010. [arXiv 1104.4409](#)
- [Baker 2011] R. C. Baker, “A partial classification of type I singularities of the mean curvature flow in high codimension”, preprint, 2011. [arXiv 1104.4592v1](#)
- [Chow 1985] B. Chow, “Deforming convex hypersurfaces by the  $n$ th root of the Gaussian curvature”, *J. Differential Geom.* **22**:1 (1985), 117–138. [MR 87f:58155](#) [Zbl 0589.53005](#)
- [Chow 1987] B. Chow, “Deforming convex hypersurfaces by the square root of the scalar curvature”, *Invent. Math.* **87**:1 (1987), 63–82. [MR 88a:58204](#) [Zbl 0608.53005](#)
- [Ecker and Huisken 1989] K. Ecker and G. Huisken, “Immersed hypersurfaces with constant Weingarten curvature”, *Math. Ann.* **283**:2 (1989), 329–332. [MR 90c:53150](#) [Zbl 0643.53043](#)
- [Evans 1982] L. C. Evans, “Classical solutions of fully nonlinear, convex, second-order elliptic equations”, *Comm. Pure Appl. Math.* **35**:3 (1982), 333–363. [MR 83g:35038](#) [Zbl 0469.35022](#)
- [Gerhardt 1990] C. Gerhardt, “Flow of nonconvex hypersurfaces into spheres”, *J. Differential Geom.* **32**:1 (1990), 299–314. [MR 91k:53016](#) [Zbl 0708.53045](#)
- [Gerhardt 2006] C. Gerhardt, *Curvature problems*, Series in Geometry and Topology **39**, International Press, Somerville, MA, 2006. [MR 2007j:53001](#) [Zbl 1131.53001](#)
- [Giga and Goto 1992] Y. Giga and S. Goto, “Geometric evolution of phase-boundaries”, pp. 51–65 in *On the evolution of phase boundaries* (Minneapolis, MN, 1990–1991), edited by M. E. Gurtin and G. B. McFadden, IMA Vol. Math. Appl. **43**, Springer, New York, 1992. [MR 94g:35226](#) [Zbl 0771.35027](#)
- [Glaeser 1963] G. Glaeser, “Fonctions composées différentiables”, *Ann. of Math. (2)* **77** (1963), 193–209. [MR 26 #624](#) [Zbl 0106.31302](#)
- [Hamilton 1982] R. S. Hamilton, “Three-manifolds with positive Ricci curvature”, *J. Differential Geom.* **17**:2 (1982), 255–306. [MR 84a:53050](#) [Zbl 0504.53034](#)
- [Hamilton 1986] R. S. Hamilton, “Four-manifolds with positive curvature operator”, *J. Differential Geom.* **24**:2 (1986), 153–179. [MR 87m:53055](#) [Zbl 0628.53042](#)
- [Hamilton 1995a] R. S. Hamilton, “The formation of singularities in the Ricci flow”, pp. 7–136 in *Surveys in differential geometry* (Cambridge, MA, 1993), vol. II, edited by C. C. Hsiung and S.-T. Yau, International Press, Cambridge, MA, 1995. [MR 97e:53075](#) [Zbl 0867.53030](#)
- [Hamilton 1995b] R. S. Hamilton, “Harnack estimate for the mean curvature flow”, *J. Differential Geom.* **41**:1 (1995), 215–226. [MR 95m:53055](#) [Zbl 0827.53006](#)
- [Huisken 1984] G. Huisken, “Flow by mean curvature of convex surfaces into spheres”, *J. Differential Geom.* **20**:1 (1984), 237–266. [MR 86j:53097](#) [Zbl 0556.53001](#)

- [Huisken 1990] G. Huisken, “Asymptotic behavior for singularities of the mean curvature flow”, *J. Differential Geom.* **31**:1 (1990), 285–299. [MR 90m:53016](#) [Zbl 0694.53005](#)
- [Huisken 1993] G. Huisken, “Local and global behaviour of hypersurfaces moving by mean curvature”, pp. 175–191 in *Differential geometry, I: Partial differential equations on manifolds* (Los Angeles, CA, 1990), edited by R. Greene and S.-T. Yau, Proc. Sympos. Pure Math. **54**, Amer. Math. Soc., Providence, RI, 1993. [MR 94c:58037](#) [Zbl 0791.58090](#)
- [Huisken and Sinestrari 1999a] G. Huisken and C. Sinestrari, “Convexity estimates for mean curvature flow and singularities of mean convex surfaces”, *Acta Math.* **183**:1 (1999), 45–70. [MR 2001c:53094](#) [Zbl 0992.53051](#)
- [Huisken and Sinestrari 1999b] G. Huisken and C. Sinestrari, “Mean curvature flow singularities for mean convex surfaces”, *Calc. Var. Partial Differential Equations* **8**:1 (1999), 1–14. [MR 99m:58057](#) [Zbl 0992.53052](#)
- [Krylov 1982] N. V. Krylov, “Ограниченно неоднородные эллиптические и параболические уравнения”, *Izv. Akad. Nauk SSSR Ser. Mat.* **46**:3 (1982), 487–523. Translated as “Boundedly inhomogeneous elliptic and parabolic equations” in *Math. USSR-Izv.* **20**:3 (1983), 459–492. [MR 84a:35091](#) [Zbl 03806019](#)
- [McCoy 2005] J. A. McCoy, “Mixed volume preserving curvature flows”, *Calc. Var. Partial Differential Equations* **24**:2 (2005), 131–154. [MR 2006g:53098](#) [Zbl 1079.53099](#)
- [McCoy 2011] J. A. McCoy, “Self-similar solutions of fully nonlinear curvature flows”, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **10**:2 (2011), 317–333. [MR 2012g:53139](#) [Zbl 1234.53018](#)
- [McCoy et al. 2014] J. McCoy, F. Mofarreh, and G. Williams, “Fully nonlinear curvature flow of axially symmetric hypersurfaces with boundary conditions”, *Ann. Mat. Pura Appl.* (2014). To appear; posted online March 2013.
- [Michael and Simon 1973] J. H. Michael and L. M. Simon, “Sobolev and mean-value inequalities on generalized submanifolds of  $\mathbb{R}^n$ ”, *Comm. Pure Appl. Math.* **26** (1973), 361–379. [MR 49 #9717](#) [Zbl 0256.53006](#)
- [Schulze 2006] F. Schulze, “Convexity estimates for flows by powers of the mean curvature”, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **5**:2 (2006), 261–277. [MR 2007b:53138](#) [Zbl 1150.53024](#)
- [Stampacchia 1966] G. Stampacchia, *Équations elliptiques du second ordre à coefficients discontinus*, Séminaire de Mathématiques Supérieures **16**, Les Presses de l’Université de Montréal, Montréal, QC, 1966. [MR 40 #4603](#) [Zbl 0151.15501](#)

Received 21 Dec 2012. Accepted 23 Jul 2013.

BEN ANDREWS: [ben.andrews@anu.edu.au](mailto:ben.andrews@anu.edu.au)

*Current address:* Mathematical Sciences Institute, Australian National University, Building 27, Acton ACT 0200, Australia  
Mathematical Sciences Center, Tsinghua University, Beijing 100084, China

MAT LANGFORD: [mathew.langford@anu.edu.au](mailto:mathew.langford@anu.edu.au)

*Mathematical Sciences Institute, Australian National University, Building 27, Acton ACT 0200, Australia*

JAMES MCCOY: [jamesm@uow.edu.au](mailto:jamesm@uow.edu.au)

*Institute for Mathematics and its Applications, School of Mathematics and Applied Statistics, University of Wollongong, Wollongong NSW 2522, Australia*

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)

University of California  
Berkeley, USA

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>	Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Vaughan Jones	U.C. Berkeley & Vanderbilt University <a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a>	Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>
Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 7 No. 2 2014

---

Two-phase problems with distributed sources: regularity of the free boundary DANIELA DE SILVA, FAUSTO FERRARI and SANDRO SALSA	267
Miura maps and inverse scattering for the Novikov–Veselov equation PETER A. PERRY	311
Convexity of average operators for subsolutions to subelliptic equations ANDREA BONFIGLIOLI, ERMANNO LANCONELLI and ANDREA TOMMASOLI	345
Global uniqueness for an IBVP for the time-harmonic Maxwell equations PEDRO CARO and TING ZHOU	375
Convexity estimates for hypersurfaces moving by convex curvature functions BEN ANDREWS, MAT LANGFORD and JAMES MCCOY	407
Spectral estimates on the sphere JEAN DOLBEAULT, MARIA J. ESTEBAN and ARI LAPTEV	435
Nondispersive decay for the cubic wave equation ROLAND DONNINGER and ANIL ZENGINOĞLU	461
A non-self-adjoint Lebesgue decomposition MATTHEW KENNEDY and DILIAN YANG	497
Bohr’s absolute convergence problem for $\mathcal{H}_p$ -Dirichlet series in Banach spaces DANIEL CARANDO, ANDREAS DEFANT and PABLO SEVILLA-PERIS	513