

Accepted Manuscript

The *microRNA ame-miR-279a* regulates sucrose responsiveness of forager honey bees (*Apis mellifera*)

Fang Liu, Tengfei Shi, Wei Yin, Xin Su, Lei Qi, Zachary Y. Huang, Shaowu Zhang, Linsheng Yu



PII: S0965-1748(17)30140-6

DOI: [10.1016/j.ibmb.2017.09.008](https://doi.org/10.1016/j.ibmb.2017.09.008)

Reference: IB 2994

To appear in: *Insect Biochemistry and Molecular Biology*

Received Date: 29 March 2016

Revised Date: 20 August 2017

Accepted Date: 14 September 2017

Please cite this article as: Liu, F., Shi, T., Yin, W., Su, X., Qi, L., Huang, Z.Y., Zhang, S., Yu, L., The *microRNA ame-miR-279a* regulates sucrose responsiveness of forager honey bees (*Apis mellifera*), *Insect Biochemistry and Molecular Biology* (2017), doi: 10.1016/j.ibmb.2017.09.008.


This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Abundance of *ame-miR-279a*

High



Low

Overexpression/inhibition of
ame-miR-279aRegulation of the sucrose
responsiveness

1

2 **The *microRNA ame-miR-279a* regulates sucrose responsiveness of**
3 **forager honey bees (*Apis mellifera*)**

4

5 **Fang Liu ^{*1a}, Tengfei Shi^{1a}, Wei Yin^b, Xin Su^a, Lei Qi^a, Zachary Y. Huang^{*c}, Shaowu Zhang^d,**
6 **Linsheng Yu^a**

7

8 ^a College of Animal Science and Technology, Anhui Agricultural University, 230000, Hefei, Anhui,
9 China; ^b The Core Facility, Zhejiang University School of Medicine, Zhejiang University,
10 Hangzhou 310058, China; ^c Department of Entomology, Michigan State University, East Lansing,
11 Michigan, United States of America; ^d Research School of Biology, College of Medicine, Biology
12 and Environment, The Australian National University, Australia.

13

14

15 *Correspondence: Zachary Y. Huang, Department of Entomology, Michigan State University, East
16 Lansing, Michigan, United States of America e-mail: bees@msu.edu; Fang Liu, College of Animal
17 Science and Technology, Apiculture Research Institute, Anhui Agricultural University, 230000,
18 Hefei, Anhui, China. Tel.: +86 18297912164; e-mail: lfxiaomifeng@ahau.edu.cn;

19 ¹These authors contributed equally to this work.

20

21 **Abstract**

22 Increasing evidence demonstrates that microRNAs (miRNA) play an important role in the
23 regulation of animal behaviours. Honey bees (*Apis mellifera*) are eusocial insects, with honey bee
24 workers displaying age-dependent behavioural maturation. Many different miRNAs have been
25 implicated in the change of behaviours in honey bees and *ame-miR-279a* was previously shown to
26 be more highly expressed in nurse bee heads than in those of foragers. However, it was not clear
27 whether this difference in expression was associated with age or task performance. Here we show
28 that *ame-miR-279a* shows significantly higher expression in the brains of nurse bees relative to
29 forager bees regardless of their ages, and that *ame-miR-279a* is primarily localized in the Kenyon
30 cells of the mushroom body in both foragers and nurses. Overexpression of *ame-miR-279a*
31 attenuates the sucrose responsiveness of foragers, while its absence enhances their sucrose
32 responsiveness. Lastly, we determined that *ame-miR-279a* directly target the mRNA of *Mblk-1*.
33 These findings suggest that *ame-miR-279a* plays important roles in regulating honey bee division
34 of labour.

35

36 **Key words:** microRNA, Honey bee, proboscis extension reflex, sucrose responsiveness, division
37 of labour.

38

39 1. Introduction

40 The honey bee (*Apis mellifera*. L) is a eusocial insect and a good model organism
41 to study the mechanisms and evolution of social behaviours (Robinson et al., 2005).
42 The workers in the colony exhibit age-related division of labour: young honey bees
43 usually engage in within-nest tasks such as brood care (“nursing”), while the old
44 honey bees forage outside for different resources (pollen, nectar, water and propolis)
45 (Winston, 1987; Robinson, 1992). However, the division of labour is very flexible:
46 bees can accelerate or reverse their behavioural development according to the colony
47 needs (Robinson, 1992; Huang and Robinson, 1996).

48 Numerous studies have focused on the molecular mechanisms underpinning
49 division of labour. Behavioural changes are associated with gene expression changes
50 in the honey bee brain (Whitfield et al., 2003). A number of genes, such as *period*
51 (Toma et al., 2000), *acetylcholinesterase* (Shapira et al., 2001), *foraging* (Ben-shahar
52 et al., 2002, 2005) and *malvolio* (Ben-shahar et al., 2004) are reported to be involved
53 in the behavioural transition from nurse to forager. MicroRNAs (miRNAs) are
54 endogenous small non-coding RNAs (18~24nt) which downregulate gene expression
55 by mRNA cleavage or translation repression (Bartel, 2004). One single miRNA may
56 target many mRNAs, and a single mRNA may contain binding sites for many
57 different miRNAs. This leads to a complex regulatory system for biological processes,
58 such as cell proliferation, differentiation and apoptosis, embryonic development,
59 neurogenesis, immunity response and disease resistance (Ambros, 2004; Pillai, 2005;
60 Vasudevan, 2007; Legeai et al., 2010).

61 Several miRNAs were reported to be involved in the honey bee behavioural
62 maturation process. Behura and Whitfield (2010) found that *miR-276* was upregulated
63 in young nurses, and had obviously higher expression in young and old nurses than in
64 young and old foragers, suggesting its involvement in the behavioural maturation
65 from nurses to foragers. Hori et al. (2010) found that *ame-miR-276* and
66 *ame-miR-1000* are enriched in the optic lobes and in small type Kenyon cells of honey
67 bees and that their targets may encode neural function related genes. Greenburg et al.
68 (2012) found that *miR-2796* is highly expressed in bee brain, and binds to the coding

69 region of phospholipase C (PLC)-epsilon gene, which was implicated in neuronal
70 development and differentiation in mammals (Wing et al., 2003), and reported to be
71 transcriptionally regulated in association with division of labour in honey bees
72 (Tsuchimoto et al., 2004). Nunes et al. (2013) identified more than 70 miRNAs that
73 were regulated by the gene *vitellogenin*, and one of these was *ame-miR-279*, which
74 may be associated with foraging behavior. Still, the precise mechanism of how
75 miRNAs regulate the division of labour in honey bees is poorly understood.

76 Nine miRNAs were previously found to be significantly differentially expressed
77 between nurses and foragers. One of these was *ame-miR-279a*, which was
78 up-regulated in nurses, and *Mblk-1* was predicted as a candidate target of
79 *ame-miR-279a* through bioinformatics (Liu et al., 2012). In the present study, we
80 further investigate the role of *ame-miR-279a* in honey bee behavioural development.
81 We show that *ame-miR-279a* is mainly localized in the Kenyon cells of the honey bee
82 mushroom body, and overexpression of *ame-miR-279a* attenuates the sucrose
83 responsiveness of foragers, while its inhibition enhances their sucrose responsiveness.
84 Furthermore, we found that *ame-miR-279a* directly targets the mRNA of *Mblk-1*.

85

86 **2. Materials and methods**

87 *2.1 Honey bees collections*

88 European honey bees, *Apis mellifera*, were maintained according to standard
89 beekeeping practices at Anhui Agriculture University, Hefei, China. Nurses were
90 caught when they had their heads inside cells feeding the larvae. Foragers with
91 pollens on their corbiculae were captured at the entrance of the hive. One-day-old
92 honey bees were obtained by removing honeycombs with capped pupae from a typical
93 colony to an incubator (33 °C) until adults emerged. Each one-day-old honey bee was
94 marked with a paint dot on the thorax, and kept in the incubator for an hour before
95 being put back into the original colony. A total of 200-300 one-day-old honey bees
96 were marked from each typical colony, and three independent typical colonies were
97 used in this study. Three single-cohort colonies were also made, each with about 1000
98 one-day-old honey bees obtained as described, an unrelated mated queen, an empty

99 comb for queen to lay eggs, a comb containing some honey and pollen, all placed in
100 small hive boxes (Whitfield et al., 2003).

101 Twenty 12-day-old nurses (12N) and 30-day-old foragers (30F) were captured
102 respectively from each of the three typical colonies, while another twenty of
103 12-day-old nurses (12N) and 12-day-old (“precocious”) foragers (12PF), and
104 30-day-old (“overaged”) nurses (30ON) and 30-day-old foragers (30F) were captured
105 from the three single-cohort colonies. The collected honey bees were kept in an
106 incubator (33 °C) before their heads were removed for brain dissection to extract RNA
107 for real-time quantitative polymerase chain reaction (RT-qPCR) and northern blot
108 analysis. The honey bees for behavioural experiments were collected from typical
109 colonies. More details are provided later in Section 2.6.

111 2.2 Oversupply/inhibition of *ame-miR-279a* in honey bees

112 A mimic of *ame-miR-279a* with the sense strand (5’ugacuagauccacacucauaaa3’) and the antisense strand 5’aagaguguggaucuagucauu3’) including a 2 nt-3’overhang (UU) and 2 nt-5’trim was synthesized by GenPharma (Shanghai, China). An inhibitor (5’uuuagaguguggaucuaguca3’), a single stranded RNA exactly complementary to *ame-miR-279a* sequence was also synthesized. A mimic control by using nonsense sequence (sense: 5’uucuccgaacgugucacgutt3’, antisense: 5’acgugacacguucggagaatt3’) and an inhibitor control using nonsense sequence (5’caguacuuuuguguaguacaa3’) were also synthesized.

120 Twenty foragers from a typical colony were used in each treatment and feeding
121 treatments were carried out in three independent experiments. The bees were
122 cold-anaesthetized, secured in 0.5-ml Eppendorf tubes with a strip of insulating tape
123 (Supplementary Fig. S1), and kept in an incubator (28 °C, 70% relative humidity) for
124 at least an hour to recover. There were four groups of foragers in the experiment,
125 namely groups fed with the mimic of *ame-miR-279a* (M), the mimic control of
126 nonsense sequences (NS), the inhibitor of *ame-miR-279a* (I) and the inhibitor control
127 of nonsense sequences (INS) respectively. Each forager was fed with 10 µl 50%
128 sucrose solution containing 6.6 µg of each synthetic reagent. All the foragers were

129 fed to satiety with 50% sucrose solution after treatments (Fig. S2), and kept in the
130 incubator in darkness (28°C, 70% relative humidity). The *ame-miR-279a* expression
131 in the brains of the foragers was measured 24h after feeding.

132

133 2.3 RT-PCR and qRT-PCR analysis

134 Bee brains were dissected according to Whitfield et al (2003), then processed for
135 total RNA extraction using a miRNeasy Mini Kit (Qiagen, Germany). The sample
136 quality and quantity were confirmed using a NanoDrop (Thermo Fisher Scientific,
137 Wilmington, DE, USA), and the samples were stored at -80 °C.

138 Total RNA (0.5µg per sample) was reverse transcribed with a universal adaptor
139 primer and primeScript RTase. PCR was performed at the same time with specific
140 forward primer (Table 1) and Uni-miR qPCR primer according to the instructions of
141 the SYBR PrimeScript miRNA RT-PCR Kit (TakaRa). The reactions were performed
142 in a TC PCR Thermocycle Instrument (BIOER) under the following conditions: 50 °C
143 for 60 min, 85 °C for 5 s. The qRT-PCR assays were performed in the ABI
144 StepOnePlus™ Real-Time PCR system. Amplification was carried out in 25-µl
145 reaction volume, containing 10 µl SYBR premix Ex Taq II, 2 µl first strand cDNA, 6
146 µl RNase free water, 0.8 µl of 10 µM of each of F and R of the specific primer (Table
147 1). PCR conditions were 95 °C for 30 s, followed by 40 cycles of 95 °C for 5 s and 60
148 °C for 30 s, followed by the melting curve (60 °C—95 °C). *β-actin* was used as the
149 reference gene. For each gene, test reactions were amplified in quadruplicate along
150 with a no-template and a no-enzyme control. Relative gene expression was calculated
151 using the $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen, 2001).

152

153 2.4 Northern blot

154 Total RNA (15 µg per sample) from 20 honey bees brains was separated through a
155 15% denaturing polyacrylamide gel, then transferred to Hybond-N nylon membranes
156 by Mini Tans-Blot (Liuyi, Beijing, China) and cross-linked by exposing to ultraviolet
157 light. DNA oligonucleotides with reverse complementarity to specific sequences were
158 incorporated with a single digoxigenin-labeled dideoxyuridine-triphosphate

159 (DIG-ddUTP) (Schmitz et al., 1991) by terminal transferase. The sequence of
160 *ame-miR-279a* probe was 5'uuuauugaguguggaucuaguca3'. The probe hybridizations
161 and washes were performed at 65 °C according to the instructions of DIG Northern
162 Starter Kit (Roche, Shanghai, China). Finally, the blots were exposed to Kodak film
163 according to the method established by Ramkissoon et al. (2006).

164

165 *2.5 In Situ Hybridization*

166 The honey bee brains were prepared according to Olivier et al. (2008), with the
167 modification that each brain was fixed in 4 % paraformaldehyde (PFA, Sigma) at 4 °C
168 for 30 min, and dehydrated in ascending concentrations of ethanol, embedded in
169 paraffin, then sectioned 10 µm from the frontal side. In situ hybridization was
170 performed according to the kit instructions of BOSTER (# MK10197). The main steps
171 were as follows: the endogenous enzymes in the brain sections were firstly inactivated
172 with 3% H₂O₂; then the sections were treated with pepsin diluted with 3% citric acid
173 for 20 min at room temperature, and washed using PBS; each section was incubated
174 with 20 µl hybrid liquid of *ame-miR-279a* probe (5'ttaatgagtgtggatctagtca3')
175 overnight in 40°C; the reactions were blocked and sample incubated with biotinylated
176 anti-mouse digoxin. Colour development was carried out according to the instructions
177 of DAB kit. Finally, sections were dehydrated through a graded series of methanol,
178 soaked with xylene, mounted with neutral gum and examined with a TissueFAXS plus
179 microscope (TissueGnostics, Austria).

180

181 *2.6 Behavioural experiments*

182 Foragers (N=60-70) were captured from three independent typical colonies, with
183 over 20 foragers per colony. The bees were restrained as mentioned above. The
184 foragers were divided into two groups, one group was fed with *ame-miR-279a* mimic
185 (279a-M), and another one was fed with the mimic control nonsense sequences
186 (279aM-NS). Similarly, another group of foragers (N=60-70) was collected from the
187 same colonies. One half of these foragers were fed with *miR-279a* inhibitor (279aI),
188 another half were fed with the inhibitor control nonsense sequence (279aI-NS). Each

189 forager was fed with 4.5 μ l 50 % sucrose solution containing 1 μ g of each synthetic
190 reagent. The foragers were fed to satiety with 50 % sucrose solution after being fed
191 the reagents, then put back into the incubator. The bees were tested for sucrose
192 responsiveness using the proboscis extension reflex (PER) assay 24 h and 48 h after
193 treatment. Both antenna of foragers was touched with a droplet of ascending
194 concentrations of sucrose: 0.1, 0.3, 1, 3, 10 and 30 % (w: w) to test their sucrose
195 responsiveness according to previous studies (Pankiw et al., 2001; Page et al., 1998).
196 Analysis of variance (ANOVA) was used to analyze the data with PER response as a
197 dependent variable. PER response (%) was analyzed after arcsine-square root
198 transformation. Sugar concentration was treated as a repeated measures variable.

199 Bee brains in the 279aM and 279aM-NS groups were dissected immediately after
200 PER for total RNA extraction according to Section 2.3. The expression of
201 *ame-miR-279a* and *Mblk-1* were quantified using qRT-PCR with β -*actin* as a control
202 gene (Table 1).

203

204 2.7 Western blot

205 Proteins (90 μ g per samples) were extracted from 15 honey bee heads using the
206 Tissue or Cell Total Protein Extraction Kit (Sangon Biotech, Shanghai, China). The
207 protein samples were separated through a 5% denaturing polyacrylamide gel, and
208 transferred to nitrocellulose membranes (Pall Life Sciences, Shanghai, China).
209 Non-specific binding-sites on the membranes were blocked with 5 % nonfat milk in
210 TBST for 2 h at room temperature. The membrane was incubated with TBST
211 containing 5 % nonfat milk and diluted rabbit anti-*Mblk-1* polyclonal antibody (1: 200)
212 (SBS, Beijing, China) overnight at 4 °C. It was then washed, incubated with
213 horseradish peroxidase-labeled anti-rabbit IgG (1: 500) (BeyotimeBiotech, Shanghai,
214 China) for an hour at room temperature, and washed again. The immunological
215 detection was carried out according to instructions of the Enhanced HRP-DAB
216 Chromogenic Substrate Kit (Tiangen Biotech, Beijing, China).

217

218 2.8 S2 cell culture and luciferase reporter assay

219 A 421-bp fragment from *Mblk-1* 3'UTR and its mutant sequence and a 249-bp
220 coding region of *ame-miR-279a* were synthesized and amplified using 2×PCR Mix
221 (TaKaRa) (Fig. S3). The *Mblk-1* 3' UTR and its mutant were cloned into a
222 pAc5.1-firefly luciferase-V5-His vector respectively (Fig. S4A), and the
223 *ame-miR-279a* coding region was cloned into a pAc5.1-V5-His vector (Fig. S4B),
224 *XhoI* and *NotI* restriction sites were added to the 5' end of the forward and reverse
225 primers, respectively (Table. 2). *Drosophila* S₂ cells were cultured with 10% fetal
226 bovine serum (HyClone) in Schneider's Insect Medium (Invitrogen, Carlsbad, USA).
227 Cells were seeded at 1×10⁶ cells per well in a 12-well plate. One day later,
228 *ame-miR-279a* expression vector (pAc-*ame-miR-279a*) was co-transfected with either
229 pAc-fluc-*Mblk-13'*UTR, pAc-fluc-*Mblk-13'*UTRm, or an empty vector (pAc) in the
230 cells using the calcium phosphate transfection method as described by Tiscornia et al
231 (2006). In all cases, 12 μl CaCl₂ (2 M) and 6 μg transfer vector were mixed, and 1.5
232 μg of pCopia-Renilla luciferase was added as internal control. Forty eight hours after
233 transfection, luciferase assays were performed using a dual-specific luciferase assay
234 kit (#RG027, Biyuntian, Shanghai, China). Renilla luciferase activity provided
235 normalization for firefly luciferase activity.

236

237 2.9 Statistical analysis

238 Statistical analysis was conducted as indicated in the text and in figure legends.
239 All t-tests used were two tailed. All tests were done by SPSS 16.0.

240

241 3. Results

242 3.1 The expression of *ame-miR-279* paralogs in the brains of nurse and forager bees

243 We had previously detected a significantly higher expression level of
244 *ame-miR-279a* in the heads of nurses compared to foragers in normal colonies (Liu et
245 al., 2012), and *ame-miR-279b*, *ame-miR-279c*, *ame-miR-279d* were also detected in
246 honey bees (Chen et al., 2010; Qin et al., 2014). What might be the differences in
247 expression among these *miR-279* paralogs between nurses and foragers? As shown in
248 Fig. 1, there was a significantly higher level of *ame-miR-279a* in the brain of nurses

249 and foragers than *ame-miR-279b*, *ame-miR-279c*, *ame-miR-279d*. It reveals the
250 important role of *ame-miR-279a* in the brain function of the honey bee.

251

252 3.2 The expression pattern of *ame-miR-279a* in the brains of nurses and foragers

253 There was a significantly higher expression of *ame-miR-279a* in the heads of
254 nurses than in those of foragers in typical colonies (Liu et al., 2012), and it showed a
255 high degree of temporal specificity during the development of adult workers, with the
256 highest expression in the 12-day-old nurses and remaining stable in over 30-day-old
257 foragers (Shi et al., 2014). These suggest a possible important function of
258 *ame-miR-279a* in honey bee behavior plasticity. To confirm this hypothesis, the
259 expression and localization of *ame-miR-279a* in the brains of nurses and foragers
260 were investigated. We first measured the *ame-miR-279a* expression in the brains of
261 nurses and foragers exhibiting normal behavior in typical colonies. A t-test showed
262 that *ame-miR-279a* was significantly highly expression in 12-day-old nurses
263 compared to the 30-day-old foragers ($t=3.79$, $P<0.05$) (Fig. 2A). However, the
264 differential expression of *ame-miR-279a* between nurses and foragers may be
265 associated with their ages but not their different behavior. To resolve this question, we
266 created the single-cohort colonies, and tested *ame-miR-279a* expression in foragers
267 and nurses of the same age. As expected, this pattern stayed the same regardless
268 whether nurses and foragers were both young (12 days old) or both old (30 days old)
269 in single cohort colonies (Fig. 2B). The *ame-miR-279a* expression between nurses of
270 different ages (12 vs. 30 days old) was not significantly different, nor was it between
271 foragers of different ages (Fig. S5). Northern blot further confirmed that
272 *ame-miR-279a* had a higher expression in nurses than in foragers, regardless of
273 whether both groups were 12 days old or 30 days old (Fig. 2C).

274 To determine the localization of *ame-miR-279a* in adult honey bee brains (nurses
275 and foragers), in situ hybridization was performed using LNA (locked nucleic acid)
276 miRNA. The results showed that *ame-miR-279a* (brown staining) was predominantly
277 expressed in the Kenyon cells of the mushroom bodies (Fig. 3A, B) and in the lamina
278 of the optic lobes in nurse and forager (Fig. 3A, C). The blank control produced no

279 brown staining (Fig. 3D). Moreover, *ame-miR-279a* expression in the brain showed
280 no obvious spatial difference between the nurse bees and forager bees even when they
281 were of the same age (Fig. S6). Taken all together, these results confirmed the
282 important role of *ame-miR-279a* in the bee behavioral maturation.

283

284 *3.3 inhibition and overexpression of ame-miR-279a in the honey bee*

285 Considering the importance of *ame-miR-279a* in behavioral maturation, we
286 decided to overexpress and inhibit the miRNA in honey bees to examine possible
287 effects on behavior. The synthetic inhibitor (anti-miRNA) and mimic of
288 *ame-miR-279a* were fed to foragers together with 50% sucrose solution. The
289 qRT-PCR confirmed the overexpression and inhibition of *ame-miR-279a* in the brains
290 of honey bee in the presence of the mimic and inhibitor respectively. As shown in Fig.
291 4, the *ame-miR-279a* expression in foragers from the M group was significantly
292 higher than in the NS group, while *ame-miR-279a* expression in foragers from the I
293 group was significantly lower than that of the INS group.

294

295 *3.4 ame-miR-279a affects the sucrose responsiveness of foragers*

296 To further investigate the possible function of *ame-miR-279a* in the honey bees'
297 behavioral maturation, we tested the effect of *ame-miR-279a* on PER first by using a
298 mimic. As was no significant difference in PER between 24 and 48 h ($F=3.22$, $df = 1$,
299 48 ; $P = 0.08$), we analyzed the two sets of data together. PER response varied
300 significantly with sugar concentrations ($F=15.78$, $df = 5$, 48 ; $P<0.001$). PER response
301 was significantly lower in bees fed with a mimic (279aM) compared to a control
302 group fed with nonsense control (279aM-NS) ($F=13.12$, $df=1$, 5 ; $P < 0.001$, Fig. 5A).

303 We then tested the effect of *ame-miR-279a* on PER by using its inhibitor. There
304 was no significant difference in PER between 24 and 48 h ($F=1.07$, $df = 1$, 48 ; $P >$
305 0.1), and we analyzed the two sets of data together. PER response varied significantly
306 with sugar concentrations ($F=14.71$, $df = 5$, 48 ; $P<0.001$). PER response was
307 significantly higher in bees fed with an inhibitor (279aI) compared to a control group
308 fed with nonsense control (279aI-NS) ($F=4.96$, $df=1$, 5 ; $P < 0.04$, Fig. 5B).

309

310 *3.5 Quantification of the expression of ame-miR-279a and Mblk-1*

311 *Mblk-1* was predicted as the target of *ame-miR-279a* (Liu et al., 2012). In order
312 to confirm their interaction, we detected the expression of *ame-miR-279a* and *Mblk-1*
313 in the brains of honey bees from the experimental foragers above. As expected,
314 *ame-miR-279a* had much higher expression in the brains of foragers in group 279aM
315 than in group 279aM-NS ($t=14.924$, $P<0.05$) (Fig. 6), while *Mblk-1* had significantly
316 lower expression in the brains of foragers from the 279aM group than from the
317 279aM-NS group ($t=3.884$, $P<0.05$) 24 h after treatment (Fig. 6). The *Mblk-1* protein
318 level in forager heads from the corresponding honey bees was further examined by
319 western blot, as shown in Fig. 6. Honey bees in 279aM group showed a lower *Mblk-1*
320 protein level than the 279aM-NS group 24h after treatment (Fig. 6). Similar results
321 were obtained 48 h after treatment (Fig. S7).

322

323 *3.6 Confirmation of the interaction of ame-miR-279a with Mblk-1 using a luciferase*
324 *reporter assay*

325 To test whether *ame-miR-279a* actually targets the *Mblk-1* 3' UTR, we subcloned
326 a 421-bp fragment of the 3'UTR region of *Mblk-1* mRNA that included the predicted
327 *ame-miR-279a* recognition site (Fig. 7) into a luciferase reporter plasmid designated
328 as pAc-fluc-*Mblk-1*3'UTR (Fig. 8A). A sequence with mutations (m) was also
329 constructed as the negative control for the same reporter assay, named as
330 pAc-fluc-*Mblk-1*3'UTR-m. The coding region of *ame-miR-279a* was cloned into a
331 pAc5.1-V5-His vector designated as pAc-*ame-miR-279a*. When pAc-*ame-miR-279a*
332 was co-transfected with pAc-fluc-*Mblk-1*3'UTR in S_2 cells, the luciferase activity
333 significantly decreased compared to the assay involving co-transfection with
334 pAc-fluc-*Mblk-1*3'UTR m and pAc ($t=10.07$, $P<0.0001$, Fig. 8B). Moreover,
335 *ame-miR-279a* expression directly reduced the *Mblk-1* mRNA and protein levels (Fig.
336 4). All these results support the conclusion that *Mblk-1* is a direct target of
337 *ame-miR-279a*.

338

339 4. Discussion

340 The role of miRNA in insect behavior has been well established in recent years
341 (Lucas and Raikhel, 2013). The miR-iab4/iab8 locus controls self-righting behavior in
342 larvae of *Drosophila* by repressing the Hox gene *Ultrabithorax* (Picao-Osorio et al.,
343 2015). Ecdysone controls *let-7-Complex* to repress the circadian gene clockwork
344 orange to regulate the circadian rhythms of *Drosophila* (Chen et al., 2014).
345 *MicroRNA-133* inhibits the behavioral aggregation of locusts by controlling dopamine
346 (Yang et al., 2014). *MicroRNA-932* regulates the memory of honey bee by targeting
347 *actin* (Alexandre et al., 2014). *Dme-miR-279* regulates the JAK/STAT pathway to
348 drive the rest: activity rhythms in *Drosophila* (Luo and Sehgal, 2012). In this study,
349 we concentrated on *ame-miR-279a* since its expression was significantly higher in
350 nurses than that of foragers, and showed a high degree of temporal specificity in
351 typical colonies (Liu et al., 2012; Shi et al., 2014). However, it was not clear whether
352 the expression of *ame-miR-279a* was associated with task performance (nursing) or
353 age (young bees). We decoupled the task performance and age in honey bees by using
354 single cohort colonies, a method regularly used to accomplish this (e.g. Robinson et al.
355 1989, Ben-Shahar et al., 2002). We determined that the *ame-miR-279a* expression was
356 always higher in nurses than in foragers regardless of whether they were young
357 (typical nurses vs. precocious foragers), or were both old (overaged nurses vs. typical
358 foragers). These results are consistent with another study in honey bees, in which the
359 *foraging* gene was shown to regulate the behavioral transition between nurses and
360 foragers (Ben-Shahar et al., 2002). Thus, we deduced that there is a good correlation
361 between *ame-miR-279a* and honey bee behavioral changes.

362 Mushroom bodies (MBs) are higher-order brain centres thought to be important
363 for sensory integration, learning and memory formation in the honey bee (Giurfa,
364 2007; Menzel, 1999; Menzel, 2012). MBs have a high degree of structural plasticity
365 depending on caste and task performance, suggesting that they are associated with
366 honey bee social behaviours (Robinson et al., 1997; Withers et al., 1993). The MBs
367 are famous as important brain regions of olfactory learning in the vinegar fly,
368 *Drosophila melanogaster* (Hayashi et al., 2009). It has been reported that

369 *dme-miR-279* was detected with strongest expression in the head epidermis in regions
370 adjacent to where the sensory organ progenitors form in *Drosophila* (Stark et al.,
371 2005). A putative orphan receptor (HR38) homologue that mediates
372 ecdysteroid-signaling, showed higher expression in the MBs of forager brains
373 compared to nurse bees, suggesting its involvement in regulation of the division of
374 labour of the workers (Yamazaki et al., 2006). In this study, we demonstrated that
375 *ame-miR-279a* is expressed more in the Kenyon cells of the mushroom bodies,
376 suggesting that *ame-miR-279a* may play a role in social behaviour. However, there
377 were no obvious spatial differences between nurses and foragers when we used *in situ*
378 hybridization. This suggests that the differences in *ame-miR-279a* levels between
379 nurses and foragers detected with RT-qPCR may represent increased expression in the
380 same cells. This is consistent with the expression pattern of the *foraging* gene in nurse
381 and forager bees, which was proved to regulate the division of labour of honey bees
382 (Ben-Shahar et al., 2002).

383 It was reported that *dme-miR-279* can regulate the formation of carbon dioxide
384 (CO₂) neurons by targeting the transcription factor Nerfin-1 in *Drosophila*
385 (Cayirlioglu et al., 2008), and that *Prospero* restricts CO₂ neuron formation indirectly
386 via *miR-279* and directly by repressing the common targets, Nerfin-1 and Esg,
387 suggesting the importance of *dme-miR-279* in the neuron and olfactory system
388 development in *Drosophila* (Hartl et al., 2011). In this study, we found that
389 overexpression of *ame-miR-279a* attenuated the sucrose responsiveness of foragers
390 (Fig. 5A), while its reduction enhanced their sucrose responsiveness (Fig. 5B).
391 Responsiveness to sucrose is associated with foraging choices, as bees with high
392 sucrose responsiveness preferentially collect pollen or water while bees with low
393 sucrose responsiveness mainly collect nectar (Pankiw and Page, 1999; Scheiner et al.,
394 2001a), suggesting the importance of *ame-miR-279a* in regulating honey bee olfactory
395 behaviour. Moreover, we found that nurses always had higher expression of
396 *ame-miR-279a* than foragers regardless of their age (Fig. 2). It has been demonstrated
397 that nurse bees are less responsive than foragers to gustatory stimuli (Scheiner et al.,
398 2001a,b), and water foragers have higher responsiveness to sucrose than both of

399 pollen and nectar foragers (Pankiw, 2005). In our study, overexpression of
400 *ame-miR-279a* in foragers may make them physiologically similar to nurses, resulting
401 in lower sucrose responsiveness (Fig. 5A, B), and suggesting that *ame-miR-279a* may
402 modulate the honey bee behavioural transition from nurses to foragers, or stimulate
403 foragers to change their behaviour from nectar collection to water or pollen foraging
404 when colony conditions demand so.

405 We have previously predicted *Mblk-1* to be a possible target for *ame-miR-279a*
406 (Liu et al., 2012). The expression of *ame-miR-279a* is largely confined to the
407 mushroom body of the honey bee brain (Fig. 3), and overexpression of *ame-miR-279a*
408 significantly inhibited the mRNA and protein expression of *Mblk-1* in forager brains
409 (Fig. 6). Moreover, our luciferase assay confirmed that *ame-miR-279a* targets the
410 3'UTR of *Mblk-1* because transfection of pAc-fluc-*Mblk-1*3'UTR reduced the
411 luciferase activity and pAc-fluc-*Mblk-1*3'UTRm rescued this suppression to the same
412 level as that of the blank control (Fig. 8). These results strongly indicate that
413 *ame-miR-279a* directly targets *Mblk-1*. The *Mblk-1* gene, encoding a putative
414 transcription factor is also expressed preferentially in the large-type Kenyon cells of
415 honey bee MBs. It contains several motifs characteristic of transcription factors,
416 including RHF1 and RHF2, a nuclear localization signal and glutamine-run motifs
417 (Takeuchi et al., 2001). Thus, *Mblk-1* is thought to be involved in brain function by
418 regulating transcription of its target genes. It has been reported that *Mblk-1* may
419 function in MB neural circuits directly modulated by the Ras/MAPK pathway (Park et
420 al., 2003). E93, a homologue of *Mblk-1* in *Drosophila*, expressed highly in the brain
421 of the fly, has been shown to affect olfactory sensory neurons (Jafari et al., 2012).
422 MBR-1, another homologue of *Mblk-1* in the nematode *Caenorhabditis elegans*, was
423 also reported to have neuronal functions, in which it is required for the pruning of
424 specific neurites that occur during larval development (Kage et al., 2005). Moreover,
425 it was also shown that MBR-1 is required for olfactory plasticity in adult animals
426 (Hayashi et al., 2009; Takayanagi-Kiya et al., 2017). Taken together, we deduce that
427 *Mblk-1* may be involved in the regulation of behavioral plasticity of honey bee
428 through its target gene *ame-miR-279a* in the MBs.

429 In summary, we found that *ame-miR-279a* showed significantly higher expression
430 in nurses than in foragers regardless of their ages, and *ame-miR-279a* was primarily
431 localized in the Kenyon cells of the mushroom body of foragers and nurses;
432 overexpression of *ame-miR-279a* attenuated the sucrose responsiveness of foragers,
433 while its inhibition enhanced their sucrose responsiveness. Moreover, we determined
434 that *ame-miR-279a* directly targets the mRNA of *Mblk-1*. These findings suggest that
435 *ame-miR-279a* plays important roles in regulating honey bee division of labour.
436

437 **Acknowledgements**

438 This work was supported by grants of National Natural Science Foundation of
439 China (31302039), Education Department Research Project of Anhui Province
440 (2013SQRL018ZD). We thank Wenfeng Chen for his kindly provide with luciferase
441 reporter plasmid, and thank Tiande Liang for technical assistance in collecting honey
442 bees and preparing samples, Aung Si, Zhiguo Li and Shoujun Huang for critically
443 reading the manuscript.

444

445 **Author's contributions**

446 F.L. planned the experiments, performed *In Situ Hybridization*, the reporter assay,
447 data analysis and wrote the manuscript. T.F.S. performed RNA extraction, RT-PCR
448 and qRT-PCR *analysis*, western blot. W.Y., X.S. and L.Q. performed experimental
449 experiments. Z.Y.H. was involved in experimental design, data analysis and
450 manuscript revision. S.W.Z. and L.S.Y. performed manuscript revision. All authors
451 have read the final draft of the manuscript

452

453 **References**

- 454 Ambros, V., 2004. The functions of animal microRNAs. *Nature* 431, 350–355.
- 455 Bartel, D. P., 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116,
456 281–297.
- 457 Behura, S.K. and Whitfield, C.W., 2010. Correlated expression patterns of microRNA genes with
458 age-dependent behavioural changes in honeybee. *Insect Mol. Biol.* 19, 431–439.
- 459 Ben-Shahar, Y., 2005. The foraging gene, behavioral plasticity, and honeybee division of labor. *J.*
460 *Comp. Physiol. A* 191: 987–994
- 461 Ben-Shahar, Y., Dudek, N.L. and Robinson, G.E., 2004. Phenotypic deconstruction reveals
462 involvement of manganese transporter malvolio in honey bee division of labor. *J. Exp. Biol.*
463 207, 3281–3288.
- 464 Ben-Shahar, Y., Robichon, A., Sokolowski, M.B. and Robinson, G.E., 2002. Influence of gene
465 action across different time scales on behavior. *Science* 296, 741–744.
- 466 Cayirlioglu, P., Kadow, I. G., Zhan, X., Okamura, K., Suh, G. S. B., Gunning, D., Lai, E. C., and
467 Zipursky, S. L., 2008. Hybrid Neurons in a MicroRNA Mutant Are Putative Evolutionary
468 Intermediates in Insect CO₂ Sensory Systems. *Science* 319, 1256–1260.
- 469 Chen, X., Yu, X., Cai, Y., Zheng, H., Yu, D., Liu, G., Zhou, Q., Hu, S., and Hu, F. 2010.
470 Next-generation small RNA sequencing for microRNAs profiling in the honey bee *Apis*
471 *mellifera*. *Insect Mol. Biol.* 19, 799-805.
- 472 Cristino, A. S., Barchuk, A. R., Freitas, F.C.P., Narayanan, R. K., Biergans, S. D., Zhao, Z. et al.,
473 2014. Neuroligin-associated microRNA-932 targets actin and regulates memory in the
474 honeybee. *Nat. Commun.* 5, 5529.
- 475 Giurfa, M., 2007. Behavioral and neural analysis of associative learning in the honeybee: a taste
476 from the magic well. *J Comp Physiol A Neuroethol. Sens Neural. Behav. Physiol.* 193,
477 801–824.
- 478 Greenberg, J.K., Xia, J., Zhou, X., Thatcher, S.R., Gu, X., Ament, S.A. et al., 2012. Behavioral
479 plasticity in honey bees is associated with differences in brain microRNA transcriptome.
480 *Genes Brain Behav*, 11, 660–670.
- 481 Hartl, M., Loschek, L. F., Stephan, D., Siju, K. P., Knappmeyer, C., and Kadow, I. C. G., 2011. A
482 New Prospero and microRNA-279 Pathway Restricts CO₂ Receptor Neuron Formation. *J.*

- 483 Neurosci. 31, 15660–15673.
- 484 Hayashi, Y., Hirotsu, T., Iwata, R., Kage-Nakadai, E., Kunitomo, H., Ishihara, T., Iino, Y., and
485 Kubo, T. 2009. A trophic role for Wnt-Ror kinase signaling during developmental pruning in
486 *Caenorhabditis elegans*. Nat. Neurosci. 12, 981-987.
- 487 Heisenberg, M. Mushroom body memoir: From maps to models. Nat. Rev. Neurosci. 2003, 4,
488 266–275.
- 489 Hori, S., Kaneko, K., Saito, T. H., Takeuchi, H., and Kubo, T., 2010. Expression of two
490 microRNAs, ame-mir-276 and -1000, in the adult honeybee (*Apis mellifera*) brain.
491 Apidologie.
- 492 Huang, Z.Y. and Robinson, G.E., 1996. Regulation of honey bee division of labor by colony age
493 demography. Behav. Ecol. Sociobiol. 39, 147–158. Kage, E., Hayashi, Y., Takeuchi, H.,
494 Hirotsu, T., Kunitomo, H., Inoue, T., Arai, H., Iino, Y., and Kubo, T. (2005). MBR-1, a novel
495 helix-turn-helix transcription factor, is required for pruning excessive neurites in
496 *Caenorhabditis elegans*. *Current Biology* 15, 1554-1559.
- 497 Legeai, F., Rizk, G., Walsh, T., Edwards, O., Gordon, K., Lavenier, D., et al., 2010. Bioinformatic
498 prediction, deep sequencing of microRNAs and expression analysis during phenotypic
499 plasticity in the pea aphid, *Acyrtosiphon Pisum*. BMC Genomics 11, 281–290.
- 500 Liu, F., Peng, W., Li, Z.G., Li, W.F., Li, L., Pan, J. et al., 2012. Next-generation small RNA
501 sequencing for microRNAs profiling in *Apis mellifera*: comparison between nurses and
502 foragers. Insect Mol. Biol. 21, 297–303.
- 503 Livak, K.J. and Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
504 quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. Methods 25, 402–408.
- 505 Lucas, K., and Raikhel, A. S. (2013). Insect MicroRNAs: Biogenesis, expression profiling and
506 biological functions. Insect Mol. Biol. 43, 24–38.
- 507 Luo, W. and Sehgal, A., 2012. Regulation of Circadian Behavioral Output via a
508 MicroRNA-JAK/STAT Circuit. Cell 148, 765–779.
- 509 Menzel, R., 1999. Memory dynamics in the honeybee. J. Comp. Physiol. A 185, 323–340.
- 510 Menzel, R., 2012. The honeybee as a model for understanding the basis of cognition. Nat. Rev.
511 Neurosci. 13, 758–768.

- 512 Nunes, F.M.F., Ihle, K.E., Mutti, N.S., Simões, Z.L.P. and Amdam, G.V., 2013. The gene
513 vitellogenin affects microRNA regulation in honey bee (*Apis mellifera*) fat body and brain. J.
514 Exo. Biol. 216, 3724–3732.
- 515 Olivier, V., Massou, I., Celle, O., Blanchard, P., Schurr, F., Ribière, M., Gauthier, M., 2008. In
516 situ hybridization assays for localization of the chronic bee paralysis virus in the honey bee
517 (*Apis mellifera*) brain. J. Virol. Methods 153, 232–237.
- 518 Page, R. J., Erber, J. and Fondrk, M., 1998. The effect of genotype on response thresholds to
519 sucrose and foraging behavior of honey bees (*Apis mellifera* L.). J. Comp. Physiol. A 182,
520 489–500.
- 521 Pankiw, T., 2005. The honey bee foraging behavior syndrome: quantifying the response threshold
522 model of division of labor. IEEE Xplore Conference: Swarm Intelligence Symposium, DOI:
523 10.1109/SIS.2005.1501595
- 524 Pankiw, T. and Page Jr, R.E., 1999. The effect of genotype, age, sex, and caste on response
525 thresholds to sucrose and foraging behavior of honey bees (*Apis mellifera* L.). J. Comp.
526 Physiol. A 185, 207–213.
- 527 Pankiw, T., Waddington, K.D. and Page, R.E., 2001. Modulation of sucrose response thresholds in
528 honey bees (*Apis mellifera* L.): influence of genotype, feeding, and foraging
529 experience. J. Comp. Physiol. A 187, 293–301.
- 530 Park, J. M., Kunieda, T., and Kubo, T., 2003. The activity of Mblk-1, a mushroom body-selective
531 transcription factor from the honeybee, is modulated by the Ras/MAPK pathway. J. Biol.
532 CHEM. 278, 18689–18694.
- 533 Picao-Osorio, J., Johnston, J., Landgraf, M., Berni, J., and Alonso, C. R. (2015).
534 MicroRNA-encoded behavior in *Drosophila*. *Science* 350, 815–820.
- 535 Pillai, R. S., 2005. MicroRNA function: Multiple mechanisms for a tiny RNA? *Rna* 11,
536 1753–1761.
- 537 Qin, Q.-H., Wang, Z.-L., Tian, L.-Q., Gan, H.-Y., Zhang, S.-W., and Zeng, Z.-J. (2014). The
538 integrative analysis of microRNA and mRNA expression in *Apis mellifera* following
539 maze-based visual pattern learning. *Insect Sci.* 21, 619-636.
- 540 Ramkissoon, S.H., Mainwaring, L.A., Sloand, E.M., Young, N.S. and Kajigaya, S., 2006.
541 Nonisotopic detection of microRNA using digoxigenin labeled RNA probes. *Mol. Cell.*

- 542 Probe 20, 1–4.
- 543 Robinson, G.E., Fahrbach, S.E. and Winston, M.L., 1997. Insect societies and the molecular
544 biology of social behavior. *Bioessays* 19, 1099–1108.
- 545 Robinson, G.E., Grozinger, C.M. and Whitfield, C.W., 2005. Sociogenomics: social life in
546 molecular terms. *Nat. Rev. Genet.* 6, 257–270.
- 547 Robinson, G.E., 1992. Regulation of division of labor in insect societies. *Annu. Rev. Entomol.* 37,
548 637–665.
- 549 Robinson, G. E., and Page, R. E., 1989. Genetic determination of nectar foraging, pollen foraging,
550 and nest-site scouting in honey bee colonies. *Behav. Ecol. Sociobiol.* 24, 317–323.
- 551 Scheiner, R., Page, R.E. and Erber, J., 2001a. The effects of genotype, foraging role and sucrose
552 perception on the tactile learning performance of honey bees (*Apis mellifera* L.). *Neurobiol.*
553 *Learn. Mem.* 76, 138–150.
- 554 Scheiner, R., Page, R.E. and Erber, J., 2001b. Responsiveness to sucrose affects tactile and
555 olfactory learning in preforaging honey bees of two genetic strains. *Behav. Brain Res.* 120,
556 67–73.
- 557 Schmitz, G. G., Walter, T., Seibl, R., and Kessler, C., 1991. Nonradioactive labeling of
558 oligonucleotides in vitro with the hapten digoxigenin by tailing with terminal transferase.
559 *Analytical Biochemistry* 192, 222-231.
- 560 Shapira, M., Thompson, C.K. Soreq, H. and Robinson, G.E., 2001. Changes in neuronal
561 acetylcholinesterase gene expression and division of labor in honey bee colonies. *J. Mol.*
562 *Neurosci.* 17, 1–12.
- 563 Shi, T., Liu, F., Yu, L., Wang, T. and Qi, L., 2014. Expression levels of three miRNAs in the brain
564 of different day-old workers of *Apis mellifera ligustica* (Hymenoptera: Apidae). *Acta*
565 *Entomologica Sinica* 57, 1368–1374.
- 566 Stark, A., Brennecke, J., Bushati, N., Russell, R. B., and Cohen, S. M., 2005. Animal MicroRNAs
567 Confer Robustness to Gene Expression and Have a Significant Impact on 3'UTR Evolution.
568 *Cell* 123, 1133–1146.
- 569 Takayanagi-Kiya, S., Kiya, T., Kunieda, T., and Kubo, T. 2017. Mblk-1 Transcription Factor
570 Family: Its Roles in Various Animals and Regulation by NOL4 Splice Variants in Mammals.
571 *Int. J. Mol. Sci.* 18, 246.

- 572 Takeuchi, H., Kage, E., Sawata, M., Kamikouchi, A., Ohashi, K., Ohara, M. et al., 2001.
573 Identification of a novel gene, *Mblk-1*, that encodes a putative transcription factor expressed
574 preferentially in the large-type Kenyon cells of the honeybee brain, *Insect Mol. Biol.* 10,
575 487–494.
- 576 Tiscornia, G., Singer, O., and Verma, I. M., 2006. Production and purification of lentiviral vectors.
577 *Nat. Protoc.* 1, 241–245.
- 578 Toma, D.P., Moore, D., Bloch, G. and Robinson, G.E., 2000. Changes in period expression in the
579 brain and division of labor in honey bee colonies. *PNAS* 97, 6914–6919.
- 580 Tsuchimoto, M., Aoki, M., Takada, M., Kanou, Y., Sasagawa, H., Kitagawa, Y. & Kadowaki, T.,
581 2004. The changes of gene expression in honeybee (*Apis mellifera*) brains associated with
582 ages. *Zoolog. Sci.* 21, 23–28.
- 583 Vasudevan, S., Tong, Y., and Steitz, J. A., 2007. Switching from Repression to Activation:
584 MicroRNAs Can Up-Regulate Translation. *Science* 318, 1931–1934.
- 585 Whitfield, C.W., Cziko, A.M. and Robinson, G.E., 2003. Gene expression profiles in the brain
586 predict behavior in individual honey bees. *Science* 302, 296–299.
- 587 Winston, M.L., 1987. *The biology of the honeybee*. Harvard University Press, Cambridge, MA.
- 588 Withers, G.S., Fahrbach, S.E. and Robinson, G.E., 1993. Selective neuroanatomical plasticity and
589 division of labour in the honeybee. *Nature* 364, 238–240.
- 590 Wing, M.R., Bourdon, D.M. & Harden, T.K., 2003. PLC-epsilon: a shared effector protein in Ras-,
591 Rho-, and G alpha beta gamma-mediated signaling. *Mol. Interv.* 3, 273–280.
- 592 Yamazaki, Y., Shirai, K., Paul, R.K., Fujiyuki, T., Wakamoto, A., Takeuchi, H. and Kubo, T., 2006.
593 Differential expression of HR38 in the mushroom bodies of the honeybee brain depends on
594 the caste and division of labor. *Febs. Lett.* 580, 2667–2670.
- 595 Yang, M., Wei, Y., Jiang, F., Wang, Y., Guo, X., He, J., and Kang, L. (2014). MicroRNA-133
596 Inhibits Behavioral Aggregation by Controlling Dopamine Synthesis in Locusts. *PLOS Genet.*
597 10, e1004206.
- 598
- 599 **Table 1** Primer sequences used for qRT-PCR validation of *ame-miR-279a* and *Mblk-1*.
- 600
- 601 **Table 2** Primer sequences used for RT-PCR amplification of 3'UTR and

602 *pri-miR-279a*.603 **Table 1**

Primer	5' to 3'
<i>Mblk-1</i> -F	AACACCAAATACGACCCAAAAC
<i>Mblk-1</i> -R	CAACAGAGCCTTCTCCACTTCT
<i>ame-miR-279a</i> -F	CTTTCTAAGTATCAATAATG
<i>ame-miR-279a</i> --R	TCTTAAAATTCATATTCATA
β -actin-F	TGCCAACACTGTCCTTTCTG
β -actin-R	AGAATTGACCCACCAATCCA

604

605 **Table 2**

Primer	5' to 3'
<i>Mblk-1</i> 3'UTR-F	CGCCCGAAACCGCGAAAGAA
<i>Mblk-1</i> 3'UTR-R	GACGTCGAATCACGCCTTGT
<i>pri-miR-279a</i> -F	CTTTCTAAGTATCAATAATG
<i>pri-miR-279a</i> --R	TCTTAAAATTCATATTCATA

606

607

608 **Figure captions:**

609 **Fig. 1.** Expression levels of four *miR-279* paralogs (miR-279a, miR-279b, miR-279c,
610 miR-279d) in the brains of nurses and foragers.

611 **Fig. 2.** Expression levels (\pm SE) of *ame-miR-279a* in the brain of 12 and 30 days old
612 age-matched nurses and foragers from regular colonies (A) and single-cohort colonies
613 (B). Student t-test results were shown, with * denoting $P < 0.05$ and ** denoting
614 $P < 0.01$. Data based on three replicates (colonies). (C) Northern blot analysis of
615 *ame-miR-279a* in brains of age-matched 12-day-old young nurses (12N) and young
616 (“precocious”) foragers (12PF), and age-matched 30 days old foragers (30F) and old
617 (“overage”) nurses (30ON) from single-cohort colonies. 5s rRNA was used as a
618 reference.

619

620 **Fig. 3.** Expression of *ame-miR-279a* in the honey bee brain. OL, optic lobe; KC,
621 Kenyon cells. *ame-miR-279a* is highly expressed in the Kenyon cells of the
622 mushroom bodies and in the lamina of the optic lobes (brown colour) with the
623 positive probe (A). No brown labeling was seen in sections probed with a blank
624 control (D). Squares delineate regions in shown magnified in BC and EF. There were
625 no obvious spatial differences between nurses and foragers; these images are from a
626 nurse brain.

627

628 **Fig. 4.** *Ame-miR-279a* expression in the brains of foragers after oral feeding with
629 *mimic-mir-279a* (M) or nonsense sequence (NS), or *inhibitot-mir-279a* (I) or inhibitor
630 nonsense sequence (INS). An independent t-test result is shown, data represent the
631 mean from three independent experiments \pm s.e.m * means $P < 0.05$, ** means $P < 0.01$.

632

633 **Fig. 5.** Mean score (\pm SE) of bees responding to proboscis extension response to
634 various sugar concentrations after bees treated with a mimic (A) or inhibitor (B) of
635 *ame-miR-279a*. The effect of *ame-miR-279a* on foragers’ responsiveness to sucrose.
636 Responsiveness to sucrose was significantly lower ($P < 0.01$) in foragers fed on a
637 *miR-279a* mimic (279aM) compared to those fed with a nonsense sequence

638 (279aM-NS). Conversely, response to sucrose was significantly ($P < 0.01$) enhanced
 639 in foragers fed on a *miR-279a inhibitor* (279aI) compared to those fed with a
 640 nonsense sequence (279aI-NS). Data from three colonies were analyzed after
 641 arsine-square root transformation but presented here without transformation.

642

643 **Fig. 6.** Relative expression levels (\pm SE) of *ame-miR-279a* and *Mblk-1* from group
 644 279aM and 279aM-NS at 24h after treatment. Student t-test results are shown with *
 645 denoting $P < 0.05$, ** denoting $P < 0.01$. Data are from three replicates (colonies).

646 Western blot analysis of *Mblk-1* protein in foragers' heads from 279aM and
 647 279aM-NS at 24h after treatment, *β -actin* was used as the reference protein.

648

649 **Fig. 7.** Sequences of the interaction sites between *ame-miR-279a* and *Mblk-1-3'UTR*.
 650 Asterisks indicate mutated site, mutated nucleotide bases are shown in bold. Grey
 651 shaded areas indicate canonical 7mer "seed" region that aligns with the target site, the
 652 vertical lines indicate contiguous Watson-Crick pairing.

653

654 **Fig. 8. (A)** A schematic representation of the principle behind the luciferase assay.

655 **(B)** co-transfection of *pAc-fluc-Mblk-13'UTR* resulted in dramatic suppression of the
 656 luciferase activity. A normalized firefly/renilla luciferase value was plotted with
 657 \pm s.e.m.

658

659 **Fig. S1.** The 0.5 ml Eppendorf tube used in our experiment, A: A normal tube; B: The
 660 tube without lid; C: The tube with the part of top margin removed. C was used in the
 661 present study. D: Prepared tubes inside a rack with a pair of forceps and insulation. E:
 662 A restrained forager.

663

664 **Fig. S2.** During the hour of recovery, the honey bees were fed with 10 μ l sucrose
 665 solution (50%, in dH₂O) and kept in an incubator (28°C, 70% relative humidity) in
 666 total darkness. After the test, the honey bees were fed 50 μ l sucrose solution (50%, in
 667 dH₂O) and put back into the incubator to wait for the next test.

668

669 **Fig. S3.** A 421-bp fragment from *Mblk-1* 3'UTR (A) and a 249-bp coding region of
670 *ame-miR-279a* (B).

671

672 **Fig. S4.** Schematic overview of the pAc5.1-firefly luciferase-V5-His vector used to
673 construct pAc-fluc-*Mblk-1*3'UTR plasmids (A), and the pAc5.1-V5-His vector used
674 to construct pAc-*ame-miR-279a* plasmid (B).

675

676 **Fig. S5.** Mean (\pm SE) levels of *ame-miR-279a* expression in nurses and foragers. There
677 were no statistical differences between different ages. Data based on three replicates.
678 (Data are mean \pm s.e.m, n=3).

679


680 **Fig. S6.** Expression of *ame-miR-279a* in normal forager brain (A), 12 days old nurse
681 brain (D), 12 days old forager brain (H). OL, optic lobe; KC, Kenyon cells. Squares
682 delineate regions in shown magnified in BC, EF and IJ. *Ame-miR-279a* is highly
683 expressed in the Kenyon cells of the mushroom bodies and in the lamina of the optic
684 lobes (brown colour) with the positive probe.

685


686 **Fig. S7.** Relative expression levels (\pm SE) of *ame-miR-279a* and *Mblk-1* from group
687 279aM and 279aM-NS at 48h after treatment. Student t-test results are shown with *
688 denoting $P < 0.05$, ** denoting $P < 0.01$. Data are from three replicates (colonies).
689 Western blot analysis of *Mblk-1* protein in foragers' heads from 279aM and
690 279aM-NS at 48h after treatment, β -actin was used as the reference protein.




691

692




B







A




B




C




D





E





F













A



B



1. *Ame-miR-279a* shows significantly higher expression in the brains of nurse bees relative to forager bees regardless of their ages.
2. *Ame-miR-279a* is primarily localized in the Kenyon cells of the mushroom body in both foragers and nurses.
3. Overexpression of *ame-miR-279a* down-regulates the sucrose responsiveness of foragers, while its absence up-regulates their sucrose responsiveness.
4. *Ame-miR-279a* directly targets the mRNA of *Mblk-1*.