



Namulanda, G., Maisonet, M., Taylor, E., Flanders, W. D., Olson, D., Sjodin, A., ... Naeher, L. (2016). *In utero* exposure to organochlorine pesticides and early menarche in the Avon Longitudinal Study of Parents and Children. *Environment International*, 94, 467-472.
<https://doi.org/10.1016/j.envint.2016.06.001>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.envint.2016.06.001](https://doi.org/10.1016/j.envint.2016.06.001)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://www.sciencedirect.com/science/article/pii/S0160412016302197> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

Exposure to phytoestrogens *in utero* and age at menarche in a contemporary British cohort

Kristin J. Marks^{1,2}, Terryl J. Hartman^{1,2}, Ethel V. Taylor², Michael E. Rybak², Kate Northstone^{3,4}, Michele Marcus¹

¹ Epidemiology Department, Rollins School of Public Health, Emory University, 1518 Clifton Rd NE, Atlanta, Georgia 30322, United States

² National Center for Environmental Health, Centers for Disease Control and Prevention, 4770 Buford Hwy, Atlanta, Georgia 30341, United States

³ The National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care West (NIHR CLAHRC West) at University Hospitals Bristol NHS Foundation Trust, 9th Floor, White Friars, Lewins Mead, Bristol, BS1 2NT, United Kingdom

⁴ School of Social and Community Medicine, University of Bristol, 39 Whatley Road, Bristol, BS8 2PS, United Kingdom

Abstract

Phytoestrogens are estrogenic compounds that occur naturally in plants. Phytoestrogens can cross the placenta, and animal studies have found associations between *in utero* exposure to phytoestrogens and markers of early puberty. We investigated the association between *in utero* exposure to phytoestrogens and early menarche (defined as <11.5 years of age at onset) using data from a nested case-control study within the Avon Longitudinal Study of Parents and Children, a longitudinal study involving families living in the South West of England. Concentrations of six phytoestrogens were measured in maternal urine samples collected during pregnancy. Logistic regression was used to explore associations between tertiles of phytoestrogen concentrations and menarche status, with adjustment for maternal age at menarche, maternal education, pre-pregnancy body mass index (BMI), child birth order, and duration of breastfeeding. Among 367 mother-daughter dyads, maternal median (interquartile range) creatinine-corrected concentrations (in µg/g creatinine) were: daidzein 184.8 (88.8–383.7), enterodiol 76.1 (39.1–135.8), enterolactone 911.7 (448.1–1558.0), equol 4.3 (2.8–9.0), genistein 62.1 (27.1–160.9), and *O*-desmethylangolensin (*O*-DMA) 13.0 (4.4–34.5). In analyses comparing those in the highest tertile relative to those in the lowest tertile of *in utero* phytoestrogen exposure, higher enterodiol levels were inversely associated with early menarche (OR=0.45; 95% CI: 0.25–0.81), while higher *O*-DMA levels were associated with early menarche (odds ratio (OR) = 1.94; 95% confidence interval (CI): 1.07–3.51). These findings suggest that *in utero* exposure to phytoestrogens may be associated with earlier age at menarche, though the direction of association differs across phytoestrogens.

Keywords

ALSPAC, menarche, phytoestrogens, puberty, endocrine disruptors

Target journal: *Environment International*

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.

1. Introduction

Puberty is a crucial period of growth and development. The timing and patterning of pubertal events, such as age at menarche, can provide information on overall health and some previous exposures, while potentially forecasting future health outcomes (Christensen et al, 2011; Biro et al, 2001; Golub et al, 2008).

Age at menarche, on average, has decreased since the late 19th century (Wyshak & Frisch, 1982; Zacharias & Wurtman, 1969), and a secular trend towards earlier development of secondary sexual characteristics has been reported among girls in the Avon Longitudinal Study of Parents and Children (ALSPAC) based in the United Kingdom (Rubin et al, 2009). In the United States, recent estimates for average age at menarche (12.4 years) are almost a year younger than the average age at menarche of women born in the 1920s (13.3 years), and decreases in average age at menarche have been observed across all racial/ethnic groups (McDowell et al, 2007). While improvements in nutritional status since the 19th century and the increasing prevalence of childhood obesity may be in part responsible for this trend, exposure to endocrine disrupting chemicals (EDCs) may also lead to altered timing and patterning of pubertal development (Herman-Giddens et al, 1997; Buck Louis et al, 2008; Blanck et al, 2000; Biro et al, 2012; Christensen et al, 2011).

EDCs are chemicals that may affect the body's endocrine system and cause adverse developmental, reproductive, neurological, and immune effects in humans and animals. EDCs can be natural or man-made, and research suggests that EDCs may have the greatest impact during prenatal and early postnatal development when organ and neural systems are forming (National Institute of Environmental Health Sciences, 2015). Most EDCs have estrogenic and/or anti-androgenic actions (Daxenberger et al, 2001), which are thought to have puberty-inducing

effects in females (Mouritsen et al, 2010). Previous studies have examined the associations of *in utero* exposure to various EDCs with pubertal development, particularly age at menarche, with some conflicting results (Vasiliu et al, 2004; Blanck et al, 2000; Hatch et al, 2011; Christensen et al, 2011). Most studies were limited by the use of retrospectively-collected age at menarche data.

One potential class of naturally-occurring EDCs of interest is phytoestrogens.

Phytoestrogens are estrogenic compounds that occur naturally in plants, with the most common dietary source of phytoestrogens being soybean products (Kim & Park, 2012). Although exposure to phytoestrogens is mostly dietary, phytoestrogens can cross the placental barrier in humans (Foster et al, 2002). Phytoestrogen exposure may affect sexual development, including altered pubertal timing (Kim & Park, 2012).

Animal studies have reported the effects of phytoestrogens to be quite different according to time, dosage and route. Studies in rodents found that exposure to high doses of phytoestrogens (isoflavones) *in utero* and through diet in early life accelerated pubertal onset in female animals (Casanova et al, 1999; Takashima-Sasaki et al, 2006). In humans, the effect of soy-based infant formula on pubertal development has been studied to some extent, though this has yielded mixed results regarding an association with age at menarche (Adgent et al, 2011; Strom et al, 2001). However, there have been no human studies published to date that have investigated the association between *in utero* phytoestrogen exposure and age at menarche. Our aim was to do so, using maternal gestational levels of phytoestrogen exposure and prospectively-collected age at menarche data in a population-based nested case-control study.

2. Study design and methods

2.1 Study population

The Avon Longitudinal Study of Parents and Children (ALSPAC) is an ongoing prospective birth cohort of 14,541 pregnancies. ALSPAC enrolled pregnant women with an expected delivery date between April 1st, 1991 and December 31st, 1992 from three health districts in the former county of Avon, Great Britain. Information has been collected on these parents and children through interviews, mailed questionnaires, and clinic visits. Details on ALSPAC recruitment and study methods have been described elsewhere (Boyd et al, 2013).

A nested case-control study was conducted within the ALSPAC cohort to explore associations of prenatal maternal concentrations of various EDCs and age at menarche among the daughters. A 'Growing and Changing' questionnaire was developed to collect information on the offspring's pubertal development and distributed to participants annually between the ages of 8–17 years (1999–2008), with the exception of age 12 (2003). Menarche was determined through parental- or self-report of menarche status, and, if it had occurred, month and year of occurrence so that age could be computed. From the original base population of 14,062 live births, case and control series were selected from singleton (n=11,820) female subjects (n=5,756) who had completed at least two puberty staging questionnaires between the ages of 8 and 13 (5 possible questionnaires returned; n=3,682). Girls meeting eligibility criteria were ordered according to reported age at menarche when the 13-year old data became available. A cut-off of 11.5 years was established as defining 'early' menarche to satisfy sample size and power needed for the case-control study. Eligible cases could complete any two questionnaires in the series, provided that one was completed after menarche, while controls had to complete the 13-year old questionnaire in order to ascertain that menarche had not occurred by the cut-off of 11.5 years. Of the girls who reported menarche before the age of 11.5 (n=338), 59.8% (n=202) had a prenatal maternal urine sample available, and were considered potential cases. Among girls who

reported menarche at or after the age of 11.5, a random sample of 394 was chosen as potential controls, and of these, 61.2% (n=241) had a maternal urine sample available. After evaluating the integrity of the maternal urine samples, 86.1% (n=174) of potential case and 81.3% (n=196) of potential control samples were analyzed. Two cases and one control were excluded due to missing creatinine concentrations, leaving a total sample size of 367 mother-daughter dyads.

2.2 Laboratory analysis

Maternal urine samples were stored at -20 degrees Celsius before being transferred under controlled conditions to the National Center for Environmental Health, Centers for Disease Control and Prevention (Atlanta, GA) for analysis using high-performance liquid chromatography–tandem mass spectrometry. The analytical methods are described elsewhere (Rybak et al, 2008). Phytoestrogens (enterolactone, daidzein, genistein, enterodiol, *O*-desmethylangolensin, and equol) were measured in maternal first morning void urine samples collected at a median gestational age of 12 weeks (interquartile range 8–17 weeks). Phytoestrogen concentrations were creatinine-corrected (CDC, 2012). Maternal urine concentrations were used as a proxy for fetal exposure (Green & Marsit, 2015; Ahmed et al, 2011).

2.3 Statistical analysis

Potential confounders to be considered in the analyses were identified *a priori* based on previously published literature and biological plausibility. Covariates were collected at various time points. We considered the following as covariates: child race (white/non-white); maternal education (ordinally classified as <O-level (ordinary level: required, completed at age 16), O-level, or >O-level); maternal age at menarche (8–11 years / 12–15 years); maternal pre-pregnancy self-reported body mass index (BMI) (kg/m²); prenatal vegetarian diet (yes/no);

prenatal smoking (any/none); maternal age at delivery (years); child birth order (first born, second born, or third born or later); child birth weight (<2500 g / \geq 2500 g); breastfeeding duration (ordinally classified as not breastfed, <3 months, 3–5 months, \geq 6 months); use of infant soy formula (any/none); vegetarian diet during childhood (yes/no); and objectively measured childhood BMI Z-score at age 8 (if missing for age 8, used age 7, 9, or 10).

All data analysis was performed using SAS 9.3 (Cary, NC). Descriptive statistics were calculated for the sample comprised of 367 mother-daughter dyads; chi square and Fisher's exact tests were used to compare groups by menarche status. Medians and interquartile ranges were calculated for each phytoestrogen for the total sample and by menarche status, and the Wilcoxon rank sum test was used to compare groups by menarche status.

Prior to modeling, phytoestrogen concentrations were log transformed. Phytoestrogen concentrations were also divided into tertiles by using cut points based on the distribution among the controls. To investigate the association between maternal phytoestrogen concentration and earlier age at menarche, unconditional logistic regression models were used. Using the set of potential confounding variables selected *a priori* for consideration in multivariable regression models, the final model was achieved through hierarchical backwards elimination of insignificant variables (Kleinbaum et al, 1982). Maternal education, maternal age at menarche, maternal vegetarian diet, and childhood BMI were considered as potential effect modifiers. Given that 15% (n=56) of observations were missing data for covariates included in the model, multiple imputation using the fully conditional specification method was performed to address missing covariate data (Liu & De, 2015).

Please note that the ALSPAC study website contains details of all the data that are available through a fully searchable data dictionary: <http://www.bristol.ac.uk/>

alspac/researchers/access/ (University of Bristol, 2015). Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. The U.S. Centers for Disease Control and Prevention (CDC) Institutional Review Board assessed and approved human subjects protection. Mothers provided informed consent at time of enrollment.

3. Results

In the ALSPAC cohort, girls were predominantly born to white mothers, most of whom had ordinary levels of education or higher (Table 1). Cases were more likely to have mothers who had an earlier age at menarche; one-third of case mothers reported menarche between 8 and 11 years of age, compared to less than 14% of control mothers. Mothers of cases were more than twice as likely to have an overweight or obese pre-pregnancy BMI, and cases were more than twice as likely to have a childhood BMI that was more than one standard deviation above the mean. Cases were more likely to be the first born child (61.6% versus 51.6%) and more likely to have never been breastfed or breastfed for less than 3 months (48.5% versus 39.9%). The median age of menarche among cases was 11.0 years, while the median age of menarche among controls was almost two years later at 12.8 years.

Table 1. Characteristics of the Avon Longitudinal Study of Parents and Children (ALSPAC) nested case-control study population (N=367 mother-daughter dyads).

Characteristic	Menarche <11.5 years (N=172)		Menarche ≥11.5 years (N=195)		P-value for difference ^a
	N	%	N	%	
Child race					0.61
White	159	95.8	182	96.8	
Non-white	7	4.2	6	3.2	
Maternal education ^c					0.39
< O-level	36	21.7	32	16.8	
O-level	56	33.7	62	32.5	
>O-level	74	44.6	97	50.8	
Maternal age at menarche, years					<0.0001
8–11	50	33.3	23	13.9	

≥12	100	66.7	142	86.1	
Maternal pre-pregnancy BMI, kg/m ²					0.0004
<25 (under/normal weight)	113	72.4	159	87.8	
≥25 (overweight/obese)	43	27.6	22	12.2	
Prenatal vegetarian diet					0.92
Yes	10	6.1	11	5.9	
No	154	93.9	177	94.1	
Prenatal smoking					0.47
Any	29	17.3	27	14.4	
None	139	82.7	160	85.6	
Maternal age at delivery, years					0.63
<25	32	18.7	35	17.9	
25–29	71	41.5	73	37.4	
≥30	68	39.8	87	44.6	
Child birth order					0.03
First born	101	61.6	97	51.6	
Second born	36	22.0	66	35.1	
Third born or later	27	16.5	25	13.3	
Child birth weight, g ^d					1.00 ^b
<2500	<5	--	<5	--	
≥2500	166	--	188	--	
Breastfeeding duration, months					0.04
Not breastfed	26	16.0	38	20.8	
<3	53	32.5	35	19.1	
3–5	26	16.0	34	18.6	
≥6	58	35.6	76	41.5	
Use of infant soy formula ^d					0.43 ^b
Any	<5	--	<5	--	
None	163	--	187	--	
Vegetarian diet during childhood					0.65
Yes	8	4.8	7	3.8	
No	158	95.2	176	96.2	
Childhood BMI Z-score					<0.0001
<0	32	20.9	60	34.5	
0–1	50	32.7	77	44.3	
≥1	71	46.4	37	21.3	
Age at menarche, years	Median	IQR	Median	IQR	
	11.0	10.7-11.3	12.8	12.3-13.4	

Abbreviations: g, grams; kg/m², kilograms per meter-squared; IQR, interquartile range

^a Compared using chi-square tests unless otherwise noted

^b Compared using Fisher's exact test

^c <O-level=none, Certificate of Secondary Education, and vocational education, which are equivalent to no diploma or a GED in the United States. O-levels (ordinary levels) are required and completed at the age of 16. >O-level=A-levels (advanced levels) completed at 18, which are optional, but required to get into university; and a university degree.

^d Counts and percents suppressed due to small cell sizes

Median enterodiol concentrations were almost 15 µg/g creatinine lower among cases than among controls, while median *O*-desmethylangolensin (*O*-DMA) concentrations were more than

2 µg/g creatinine higher among cases than among controls (Table 2). There were no other differences evident.

Table 2. Gestational urinary phytoestrogen concentrations among mothers of girls with and without earlier age at menarche in the Avon Longitudinal Study of Parents and Children (ALSPAC) nested case-control study population (N=367 mother-daughter dyads).

Analyte ^a	Total		Menarche <11.5 years (N=172)		Menarche ≥11.5 years (N=195)		p-value ^b
	Median	IQR	Median	IQR	Median	IQR	
Enterolactone	911.7	448.1–1558.0	901.1	494.8–1478.2	923.8	440.6–1650.9	0.89
Daidzein	184.8	88.8–383.7	187.8	91.9–369.7	183.0	88.8–423.8	0.79
Genistein	62.1	27.1–160.9	60.2	26.6–157.6	65.1	27.2–162.7	0.58
Enterodiol	76.1	39.1–135.8	69.6	33.6–123.6	84.3	49.4–144.4	0.02
O-DMA	13.0	4.4–34.5	14.0	5.7–43.8	11.7	3.4–31.5	0.06
Equol	4.3	2.8–9.0	4.4	2.8–7.4	4.2	2.6–9.6	0.74

Abbreviations: IQR, interquartile range; O-DMA, O-Desmethylangolensin

^a Creatinine-corrected concentrations in µg/g creatinine

^b p-value for difference between cases and controls using the Wilcoxon Rank Sum Test

^c There are missing concentrations for genistein (n=1 where menarche <11.5 years), O-DMA (n=2 where menarche <11.5 years), and equol (n=1 where menarche <11.5 years and n=1 where menarche ≥11.5 years)

In the unadjusted model, decreased odds of early menarche were observed for those in the second and third tertiles of *in utero* enterodiol exposure compared to those in the lowest tertile (odds ratio (OR)_{second}=0.60; 95% confidence interval (CI)_{second}: 0.36–0.98; p-value_{second}=0.04; OR_{third}=0.58; 95% CI_{third}: 0.35–0.96; p-value_{third}=0.03; p-trend: 0.03) (Table 3). Increased odds of early menarche were observed for those in the second and third tertiles of *in utero* O-DMA exposure compared to those in the lowest tertile (OR_{second}=1.91; 95% CI_{second}: 1.12–3.27; p-value_{second}: 0.02; OR_{third}=1.97; 95% CI_{third}: 1.16–3.35; p-value_{third}: 0.02; p-trend: 0.02).

The results of the analyses adjusting for maternal age at menarche, maternal education, pre-pregnancy BMI, child birth order, and duration of breastfeeding were similar to those from the unadjusted analyses (Table 3). In the adjusted model, the odds of early menarche for each

unit increase of logged mothers' enterodiol concentration were OR=0.75 (95% CI: 0.59–0.96; p-trend: 0.02). When enterodiol was treated categorically, decreased odds of early menarche were observed for those in the second and third tertiles of *in utero* enterodiol exposure compared to those in the lowest tertile (OR_{second}=0.44; 95% CI_{second}: 0.25–0.77; p-value_{second}=0.005; OR_{third}=0.45; 95% CI_{third}: 0.25–0.81; p-value_{third}=0.008; p-trend: 0.007). The odds of early menarche for each unit increase of logged mothers' *O*-DMA concentration were OR=1.14 (95% CI: 1.00–1.29; p-trend: 0.05). When comparing those in the third tertile of *O*-DMA concentration to those in the lowest tertile, a 94% increase in the odds of early menarche was observed (OR_{third}=1.94; 95% CI_{third}: 1.07–3.51; p-value_{third}=0.03; p-trend: 0.03). No other significant associations were observed between phytoestrogen concentration and early menarche. There was no evidence of effect modification by maternal education, maternal age at menarche, maternal vegetarian diet, or childhood BMI.

Table 3. Associations of maternal urinary phytoestrogen concentrations with earlier age at menarche in the Avon Longitudinal Study of Parents and Children (ALSPAC) nested case-control study population (N=367 mother-daughter dyads).

Analyte ^c	Unadjusted ^a			Adjusted ^{ab}		
	OR (95% CI)	p value ^f	p for trend ^f	OR (95% CI)	p value ^f	p for trend ^f
Enterolactone						
Continuous ^d	1.04 (0.87–1.26)		0.65	1.23 (0.98–1.55)		0.07
Tertile 2 ^e (551.49–1314.96)	1.39 (0.85–2.28)	0.19		1.78 (1.02–3.10)	0.04	
Tertile 3 ^e (1314.96–8718.04)	0.98 (0.58–1.65)	0.93	0.94	1.65 (0.90–3.02)	0.10	0.10
Daidzein						
Continuous	1.03 (0.87–1.22)		0.70	1.02 (0.84–1.24)		0.83
Tertile 2 (110.09–319.03)	1.37 (0.83–2.25)	0.21		1.14 (0.66–1.99)	0.64	
Tertile 3 (319.03–21880.45)	1.00 (0.59–1.68)	0.99	0.99	0.85 (0.47–1.51)	0.57	0.58
Genistein						
Continuous	0.96 (0.83–1.11)		0.55	0.92 (0.78–1.08)		0.30
Tertile 2 (38.86–118.38)	1.02 (0.62–1.67)	0.94		0.90 (0.52–1.56)	0.70	
Tertile 3 (118.38–17916.60)	0.87 (0.52–1.44)	0.58	0.59	0.76 (0.43–1.35)	0.35	0.31
Enterodiol						
Continuous	0.79 (0.64–0.98)		0.03	0.75 (0.59–0.96)		0.02
Tertile 2 (56.49–117.99)	0.60 (0.36–0.98)	0.04		0.44 (0.25–0.77)	0.005	

Tertile 3 (117.99–1188.20)	0.58 (0.35–0.96)	0.03	0.03	0.45 (0.25–0.81)	0.008	0.007
O-DMA						
Continuous	1.12 (1.00–1.26)		0.05	1.14 (1.00–1.29)		0.05
Tertile 2 (4.77–21.04)	1.91 (1.12–3.27)	0.02		1.54 (0.85–2.77)	0.15	
Tertile 3 (21.04–1631.58)	1.97 (1.16–3.35)	0.02	0.02	1.94 (1.07–3.51)	0.03	0.03
Equol						
Continuous	0.97 (0.83–1.13)		0.70	0.98 (0.82–1.15)		0.78
Tertile 2 (3.19–6.93)	1.22 (0.74–1.99)	0.44		1.10 (0.63–1.91)	0.74	
Tertile 3 (6.93–9005.85)	0.86 (0.51–1.44)	0.55	0.57	0.90 (0.51–1.60)	0.72	0.73

^a Unconditional logistic regression

^b Adjusted for maternal age at menarche, maternal education, pre-pregnancy BMI, child birth order, and duration of breastfeeding

^c Creatinine-corrected concentrations in $\mu\text{g/g}$ creatinine

^d Continuous represents natural log transformed values of phytoestrogen concentration

^e Tertiles represent the comparison of the higher tertiles, tertiles 2 or 3, to the lowest tertile of phytoestrogen concentration

^f The p-value is for the comparison of tertile 2 or 3 to the lowest tertile of phytoestrogen concentration; the p for trend is for the trend across all three tertiles

4. Discussion

In this study, we observed strong associations between enterodiol and decreased odds of earlier age at menarche, and some evidence of an association between *O*-DMA and increased odds of earlier age at menarche.

Studies have suggested that lignans such as enterodiol and enterolactone exhibit biphasic effects (estrogenic and antiestrogenic effects), which are dependent on exposure level (Tang et al, 2015; Mousavi and Adlercreutz, 1992; Mueller et al, 2004; Pettersson and Gustafsson, 2001; Waters and Knowler, 1982; Welshons et al, 1987; Adlercreutz, 2007; Wang, 2002). At relatively low doses, some lignan exposures demonstrate estrogenic activity, stimulating cell growth, while at higher doses appear to behave as antiestrogenic agents, suppressing cell growth (Wang, 2002). The biphasic effects of lignans could potentially provide an explanation for the negative association observed between enterodiol and early menarche status.

O-DMA is an intestinal bacterial metabolite of daidzein, and about 90% of individuals harbor bacteria capable of metabolizing daidzein to *O*-DMA (Frankenfield, 2011). *O*-DMA is

less structurally similar to 17β -estradiol than its parent compound and therefore may exhibit different biological actions than daidzein. The underlying bacteria that metabolize daidzein to *O*-DMA may have a distinct physiological role; urinary excretion of *O*-DMA is a marker of harboring intestinal bacteria capable of C-ring cleavage, and therefore it is suspected that the role of the phenotype may extend beyond daidzein metabolism (Frankenfield, 2011).

To our knowledge, this is the first published study of associations of *in utero* phytoestrogen exposure with age at menarche. Although there were few significant associations found between phytoestrogen levels and age at menarche, there is biological plausibility for such an association. Exposures during pregnancy are extremely relevant to pubertal development, since this represents the period of initial organ development, including the brain, endocrine system, and reproductive tract. Furthermore, the fetus is more susceptible to such exposures due to smaller size, lack of a complete blood-brain barrier, and absence of metabolizing enzymes (Todaka et al, 2005). Studies have found that phytoestrogens can cross the placental barrier in humans, and one study (n=53) of Californian women undergoing amniocentesis found that 96% of second trimester amniotic fluid samples contained quantifiable amounts of dietary phytoestrogens (Foster et al, 2002). Based on evidence from animal studies, the main mechanism of action of phytoestrogens—the binding of phytoestrogens to estrogen receptors—may be particularly relevant for *in utero* exposure to phytoestrogens because of the timing of differentiation and development (Takagi et al, 2004; Takashima-Sasaki et al, 2006; Casanova et al, 1999). Studies in rodents have found that isoflavones administered through diet or subcutaneous injection during gestation or early life can lead to early vaginal opening (akin to early menarche in humans), irregular estrous cyclicality, and decreased GnRH activation (GnRH coordinates reproductive maturation and function) (Takagi et al, 2004; Takashima-Sasaki et al,

2006; Casanova et al, 1999; Kouki et al, 2003; Lewis et al, 2003; Bateman & Patisaul, 2008; Lee et al, 2009; Nagao et al, 2001).

To our knowledge, no previous studies have investigated *in utero* phytoestrogen exposure; therefore, we looked to previous studies on the effect of *in utero* exposure to other potential EDCs, which produced mixed results, as have previous studies on early life phytoestrogen exposure to soy infant formula. A cohort study (n=151) assessing *in utero* exposure to polychlorinated biphenyls (PCBs) and dichlorodiphenyldichloroethylene (DDE) with age at menarche found that increased exposure to DDE was associated with an earlier age at menarche, while exposure to PCBs was not associated (Vasiliu et al, 2004). A nested case-control study (n=448) found no association with *in utero* exposure to polyfluoroalkyl chemicals (PFCs) and age at menarche (Christensen et al, 2011). Studies on the association between soy-based infant formula and age at menarche are also inconclusive (the phytoestrogens found in soy are daidzein and genistein; *O*-DMA and equol are metabolites of daidzein). A retrospective cohort study (n=811) found no association between soy-based formula and self-reported recalled age at menarche (Strom et al, 2001). However, a prospective cohort study (n=2,028) in British girls of the ALSPAC study found a 53% increased risk of early menarche among those fed soy-based formula, when compared to cows' milk-based formula (Adgent et al, 2011); it should be noted that this paper examined the ALSPAC cohort as a whole, as opposed to the nested case-control study used in this paper. While it is difficult to compare across classes of EDCs and at different times of exposure, previous studies have yet to suggest a clear association between *in utero* and early life exposure to EDCs and age at menarche.

Strengths of this study are the inclusion of multiple phytoestrogen biomarkers, substantial covariate data available on mothers and children from multiple time points over gestation and

childhood, and outcome data generally collected in the year that the outcome occurred. Limitations of this study include a single spot urine measurement of phytoestrogen exposure which was not collected at a uniform time of gestation, the absence of a ‘Growing and Changing’ questionnaire at age 12, missing information on age at menarche among some controls, incomplete information on some covariates, and no assessment of phytoestrogen exposure or early childhood soy consumption (excluding soy formula) in the daughters, which might influence timing of menarche. Unlike some other EDCs that are estimated to have half-lives on the order of several years, peak rates of urinary excretion of phytoestrogens occur between 6 and 12 hours after ingestion (King & Bursill, 1998). Since phytoestrogens are excreted rather quickly, phytoestrogen exposure assessed through urinary excretion at a single time point may under- or overestimate intermittent phytoestrogen exposure. Age at menarche was obtained through self-report on ‘Growing and Changing’ puberty questionnaires completed every year by parents and/or children, depending on age. There is some potential for misclassification of the outcome, such as the completion of the questionnaire by a parent unaware of the child’s menarche status, or issues in the parent or child’s recall of the month and year menstruation began.

Urinary phytoestrogen concentrations during 1991–1992 among mothers of girls participating in the ALSPAC cohort were roughly two to three times higher for all phytoestrogens except equol (which was half as high) when compared to 2003–2006 National Health and Nutrition Examination Survey (NHANES) data for white women between 20 and 39 years old (CDC, 2012) (data not shown). It should be noted though that these samples were taken more than a decade apart.

It is also possible that the girls selected for this nested case-control study were not representative of the base cohort. When comparing ALSPAC girls who returned at least two ‘Growing and Changing’ questionnaires to those who did not return any questionnaires, non-respondents’ parents were more likely to have lower educational attainment. Compared to non-respondents, mothers of respondents were generally older at time of index birth. Further, non-respondents were more likely to be of non-white race. This could have affected our findings since socioeconomic status is related to age at menarche (Braithwaite et al, 2009); however, whether socioeconomic status is related to phytoestrogen concentrations is unclear. Although race is related to age at menarche (Biro et al, 2006; Biro et al, 2001; Wu et al, 2002; Freedman et al, 2002; Britton et al, 2004; McDowell et al, 2007) and was associated with maternal lignan concentrations in this study, we were not able to examine the effect of race due to the small number of non-white girls enrolled in ALSPAC. Furthermore, it has been suggested that several genes in Caucasians code for early menarche (Dvornyk and Waqar-ul-Haq, 2012), and since we did not include genetic data in this study, we do not know if our results could be affected. Last, due to a modest sample size, this study may have been underpowered to detect additional associations between *in utero* phytoestrogen exposure and age at menarche.

5. Conclusions

In summary, we compared exposure to phytoestrogens during pregnancy among mothers of girls who did and did not have earlier age at menarche in the ALSPAC cohort. We found an association between *O*-DMA and increased odds of earlier age at menarche, while decreased odds of earlier age at menarche were observed for enterodiols. As demonstrated in our study by the conflicting effects of phytoestrogens, plus the general lack of human studies on the associations between phytoestrogens and pubertal outcomes, there is a need for additional studies

to explore these associations in a variety of populations and to describe potential mechanisms of action for *O*-DMA and enterodiol.

Acknowledgments

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. The UK Medical Research Council and the Wellcome Trust (Grant ref: 092731) and the University of Bristol provide core support for ALSPAC. This work was specifically funded by the Centers for Disease Control and Prevention. The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.

References

- Adgent MA, Daniels JL, Rogan WJ, et al. Early-life soy exposure and age at menarche. *Paediatr Perinat Epidemiol*. 2011;26(2):163–175. doi: 10.1111/j.1365–3016.2011.01244.x
- Adlercreutz H. Lignans and human health. *Crit Rev Clin Lab Sci*. 2007;44(5–6):483–525. doi: 10.1080/10408360701612942
- Ahmed S, Mahabbat-e Khoda S, Rekha RS, et al. Arsenic-associated oxidative stress, inflammation, and immune disruption in human placenta and cord blood. *Environ Health Perspect*. 2011;119(2):258–264. doi: 10.1289.ehp.1002086
- Bateman HL, Patisaul HB. Disrupted female reproductive physiology following neonatal exposure to phytoestrogens or estrogen specific ligands is associated with decreased GnRH activation and kisspeptin fiber density in the hypothalamus. *Neurotoxicology*. 2008;29(6):988–997. doi: 10.1016/j.neuro.2008.06.008
- Biro FM, Greenspan LC, Galvez MP. Puberty in girls of the 21st century. *J Pediatr Adol Gynec*. 2012;25(5):289–294. doi: 10.1016/j.jpag.2012.05.009
- Biro FM, Huang B, Crawford PB, et al. Pubertal correlates in black and white girls. *J Pediatr*. 2006;148(2):234–240.
- Biro FM, McMahon RP, Striegel-Moore R, et al. Impact of timing of pubertal maturation on growth in black and white female adolescents: The National Heart, Lung, and Blood Institute Growth and Health Study. *J Pediatr*. 2001;138(5):636–643.
- Blanck HM, Marcus M, Tolbert PE, et al. Age at menarche and Tanner stage in girls exposed *in utero* and postnatally to polybrominated biphenyl. *Epidemiology*. 2000;11(6):641–647.
- Boyd A, Golding J, Macleod J, et al. Cohort Profile: the 'Children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol*. 2013;42(1):111–127. doi: 10.1093/ije/dys064
- Braithwaite D, Moore DH, Lustig RH, et al. Socioeconomic status in relation to early menarche among black and white girls. *Cancer Cause Control*. 2009;20(5):713–720. doi: 10.1007/s10552-008-9284-9
- Britton J, Wolff MS, Lapinski R, et al. Characteristics of pubertal development in a multi-ethnic population of nine-year-old girls. *Ann Epidemiol*. 2004;14(3):179–187.
- Buck Louis GM, Gray LE, Marcus M, et al. Environmental factors and puberty timing: Expert panel research needs. *Pediatrics*. 2008;121(Suppl 3):S192–S207. doi: 10.1542/peds.1813E
- Casanova M, You L, Gaido KW, et al. Developmental effects of dietary phytoestrogens in Sprague-Dawley rats and interactions of genistein and daidzein with rat estrogen receptors alpha and beta *in vitro*. *Toxicol Sci*. 1999;51(2):236–244.
- Centers for Disease Control and Prevention, National Center for Environmental Health. Second National Report on Biochemical Indicators of Diet and Nutrition in the U.S. Population. Atlanta: Centers for Disease Control and Prevention. 2012.
- Christensen KY, Maisonet M, Rubin C, et al. Exposure to polyfluoroalkyl chemicals during pregnancy is not associated with offspring age at menarche in a contemporary British cohort. *Environ Int*. 2011;37(1):129–135. doi: 10.1016/j.envint.2010.08.007
- Daxenberger A, Ibarreta D, Meyer HH. Possible health impact of animal oestrogens in food. *Hum Reprod Update*. 2001;7(3):340–355.
- Dvornyk V, Waqar-ul-Haq. Genetics of age at menarche: a systematic review. *Hum Reprod Update*. 2012;18(2):198–210. doi: 10.1093/humupd/dmr050

- Foster WG, Chan S, Platt L, et al. Detection of phytoestrogens in samples of second trimester human amniotic fluid. *Toxicol Lett.* 2002;129(3):199–205.
- Frankenfeld CL. *O*-Desmethylangolensin: The importance of equol's lesser known cousin to human health. *Adv Nutr.* 2011;2:317–324. doi: 10.3945/an.111.000539
- Freedman DS, Khan LK, Serdula MK, et al. Relation of age at menarche to race, time period, and anthropometric dimensions: The Bogalusa Heart Study. *Pediatrics.* 2002;110(4):e43.
- Golub MS, Collman GW, Foster PM, et al. Public health implications of altered puberty timing. *Pediatrics.* 2008;121(Suppl 3):S218-S230. doi: 10.1542/peds.2007-1813G
- Green BB, Marsit CJ. Select prenatal environmental exposures and subsequent alterations of gene-specific and repetitive element DNA methylation in fetal tissues. *Curr Environ Health Rep.* 2015; 2(2):126-136. doi: 10.1007/s40572-015-0045-0
- Hatch EE, Troisi R, Wise LA, et al. Preterm birth, fetal growth, and age at menarche among women exposed prenatally to diethylstilbestrol (DES). *Reprod Toxicol.* 2011;31(2):151–157. doi: 10.1016/j.reprotox.2010.11.006
- Herman-Giddens ME, Slora EJ, Wasserman RC, et al. Secondary sexual characteristics and menses in young girls seen in office practice: a study from the Pediatric Research in Office Settings Network. *Pediatrics.* 1997;99(4):505-512.
- Kim SH, Park MJ. Effects of phytoestrogen on sexual development. *Korean J Pediatr.* 2012;55(8):265-271. doi: 10.3345/kjp.2012.55.8.265
- King RA, Bursill DB. Plasma and urinary kinetics of the isoflavones daidzein and genistein after a single soy meal in humans. *Am J Clin Nutr.* 1998;67:867–872.
- Kleinbaum DG, Kupper LL, Morgenstern H. 1982. Epidemiologic research: Principles and quantitative methods: John Wiley & Sons.
- Kouki T, Kishitake M, Okamoto M, et al. Effects of neonatal treatment with phytoestrogens, genistein and daidzein, on sex difference in female rat brain function: estrous cycle and lordosis. *Horm Behav.* 2003;44(2):140–145.
- Lee W, Lee S-H, Ahn R-S, et al. Effect of genistein on the sexual maturation in immature female rats. *Korean J Pediatr.* 2009;52(1):111-118. doi: 10.3345/kjp.2009.52.1.111
- Lewis RW, Brooks N, Milburn GM, et al. The effects of the phytoestrogen genistein on the postnatal development of the rat. *Toxicol Sci.* 2003;71(1):74–83.
- Liu Y, De A. Multiple imputation by fully conditional specification for dealing with missing data in a large epidemiologic study. *Int J Stat Med Res.* 2015;4(3):287-295. doi: 10.6000/1929-6029.2015.04.03.7
- McDowell MA, Brody DJ, Hughes JP. Has age at menarche changed? Results from the National Health and Nutrition Examination Survey (NHANES) 1999–2004. *J Adolescent Health.* 2007;40(3):227–231.
- Mouritsen A, Aksglaede L, Sørensen K, et al. Hypothesis: exposure to endocrine-disrupting chemicals may interfere with timing of puberty. *Int J Androl.* 2010;33(2):346–359. doi: 10.1111/j.1365-2605.2010.01051.x.
- Mousavi Y, Adlercreutz H. Enterolactone and estradiol inhibit each other's proliferative effect on MCF-7 breast cancer cells in culture. *J Steroid Biochem Mol Biol.* 1992;41(3–8):615–619.
- Mueller SO, Simon S, Chae K, Metzler M, Korach KS. Phytoestrogens and their human metabolites show distinct agonistic and antagonistic properties on estrogen receptor alpha (ERalpha) and ERbeta in human cells. *Toxicol Sci.* 2004;80(1):14–25. doi: 10.1093/toxsci/kfh147

- Nagao T, Yoshimura S, Saito Y, et al. Reproductive effects in male and female rats of neonatal exposure to genistein. *Reprod Toxicol*. 2001;15(4):399–411.
- National Institutes of Health, National Institute of Environmental Health Sciences. Endocrine Disruptors. Research Triangle Park: National Institute of Environmental Health Sciences. 2015. Retrieved from: <http://www.niehs.nih.gov/health/topics/agents/endocrine/>
- Pettersson K, Gustafsson JA. Role of estrogen receptor beta in estrogen action. *Annu Rev Physiol*. 2001;63:165–192. doi: 10.1146/annurev.physiol.63.1.165
- Rubin C, Maisonet M, Kieszak S, et al. Timing of maturation and predictors of menarche in girls enrolled in a contemporary British cohort. *Paediatr Perinat Epidemiol*. 2009;23(5):492–504. doi: 10.1111/j.1365-3016.2009.01055.x
- Rybak ME, Parker DL, Pfeiffer CM. Determination of urinary phytoestrogens by HPLC-MS/MS: A comparison of atmospheric pressure chemical ionization (APCI) and electrospray ionization (ESI). *J Chromatogr B*. 2008;861:145–150.
- Setchell KDR. Phytoestrogens: the biochemistry, physiology, and implications for human health of soy isoflavones. *Am J Clin Nutr*. 1998;68(6 Suppl):1333S–1346S.
- Strom BL, Schinnar R, Ziegler EE, et al. Exposure to soy-based formula in infancy and endocrinological and reproductive outcomes in young adulthood. *J Am Med Assoc*. 2001;286(7):807–814.
- Takagi H, Shibutani M, Lee K-Y, et al. Lack of modifying effects of genistein on disruption of the reproductive system by perinatal dietary exposure to ethinylestradiol in rats. *Reprod Toxicol*. 2004;18(5):687–700.
- Takashima-Sasaki K, Komiyama M, Adachi T, et al. Effect of exposure to high isoflavone-containing diets on prenatal and postnatal offspring mice. *Biosci Biotech Biochem*. 2006;70(12):2874–2882.
- Tang R, Chen M, Zhou K, et al. Prenatal lignan exposures, pregnancy urine estrogen profiles and birth outcomes. *Environ Pollut*. 2015;205:261–268. doi: 10.1016/j.envpol.2015.06.006
- Todaka E, Sakurai K, Fukata H, et al. Fetal exposure to phytoestrogens—The difference in phytoestrogen status between mother and fetus. *Environ Res*. 2005;99(2):195–203.
- University of Bristol, Avon Longitudinal Study of Parents and Children. Accessing the resource. Bristol: University of Bristol. 2015. Retrieved from: <http://www.bristol.ac.uk/alspac/researchers/access/>
- Vasiliu O, Muttineni J, Karmaus W. *In utero* exposure to organochlorines and age at menarche. *Hum Reprod*. 2004;19(7):1506–1512.
- Wang LQ. Mammalian phytoestrogens: enterodiol and enterolactone. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2002;777(1–2):289–309.
- Waters AP, Knowler JT. Effect of a lignan (HPMF) on RNA synthesis in the rat uterus. *J Reprod Fertil*. 1982;66(1):379–381.
- Welshons WV, Murphy CS, Koch R, Calaf G, Jordan VC. Stimulation of breast cancer cells in vitro by the environmental estrogen enterolactone and the phytoestrogen equol. *Breast Cancer Res Treat*. 1987;10(2):169–175.
- Wu T, Mendola P, Buck GM. Ethnic differences in the presence of secondary sex characteristics and menarche among US girls: The Third National Health and Nutrition Examination Survey, 1988–1994. *Pediatrics*. 2002;110(4):752–757.
- Wyshak G, Frisch RE. Evidence for a secular trend in age of menarche. *N Engl J Med*. 1982;306(17):1033–1035.
- Zacharias L, Wurtman RJ. Age at menarche. *N Engl J Med*. 1969;280(16):868–875.