

## Article

# Caffeine, CYP1A2 genotype, and sports performance: is timing important?

Pickering, Craig

Available at <http://clock.uclan.ac.uk/22297/>

*Pickering, Craig (2018) Caffeine, CYP1A2 genotype, and sports performance: is timing important? Irish Journal of Medical Science . ISSN 0021-1265*

It is advisable to refer to the publisher's version if you intend to cite from the work.  
<http://dx.doi.org/10.1007/s11845-018-1811-4>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

**Title:** Letter to the Editor – Caffeine, CYP1A2 genotype, and sports performance; is timing important?

**Author:** Craig Pickering<sup>1,2</sup>

1. Institute of Coaching and Performance, University of Central Lancashire, Preston, UK
2. Exercise and Nutritional Genomics Research Centre, DNAFit Ltd, London, UK

Corresponding Author:

Craig Pickering

Institute of Coaching and Performance, University of Central Lancashire, Fylde Road, Preston, PR1 2HE, UK.

Email: [craig@dnafit.com](mailto:craig@dnafit.com).

**Disclosure Statement:**

Craig Pickering is an employee of DNAFit Ltd, a genetic testing company. He received no financial incentives for the preparation of this manuscript, and the views contained within are his, and not representative of DNAFit Ltd.

**Words:** 572

To The Editor

I read with great interest the recent papers authored by Rahimi [1], published in this journal, and by Guest and colleagues [2], published in *Medicine & Science in Sport & Exercise*, exploring the ergogenic effects of caffeine on different *CYP1A2* genotypes. Briefly, Rahimi reported 6mg/kg of caffeine did not enhance resistance exercise performance in C allele carriers in a cohort of males, but did for AA genotypes. Similarly, Guest et al. [2] found that caffeine was ergolytic for endurance performance at a dose of 4mg/kg for CC genotypes, was ergogenic at doses of both 2- and 4mg/kg for AA genotypes, and neutral for AC genotypes at both doses, as well as for CC genotypes at 2mg/kg. Since 2012, a number of papers have been published exploring the impact of this particular polymorphism on caffeine's performance benefits, with equivocal results [3,4]. These previous studies tended to be hindered by complex study designs [4], or low subject numbers [3]. The latter point is particularly important, given that the frequency of CC genotypes tends to be very low (~8%) [2], illustrated by the complete absence of CC genotypes available for group analysis in Rahimi's cohort [1]. As such, the additional insights garnered from both Rahimi [1] and Guest and colleagues [2] shed further insight into the inter-individual variation of ergogenic effects following caffeine ingestion.

The results, however, promote significant cognitive dissonance. As an AC genotype who used caffeine extensively as an ergogenic aid during my professional athletic career, do I really believe that caffeine had no performance benefits for me? It's hard for me to reach such a conclusion, even though the results of both Rahimi and Guest and colleagues' research [1,2] suggest as much. I'm sure many other practitioners feel the same, and indeed perhaps worry that the standard caffeine recommendations are harmful, or at best neutral, for a significant proportion of the athletes they work with [5].

The mechanism proposed by Guest et al. [2] is that, because C allele carriers metabolise caffeine at a slower rate than AA genotypes [6], they experience prolonged vasoconstriction, which is likely to be performance limiting in endurance events where the transfer of oxygen and nutrients to the working muscle is crucial. Additionally, Womack and colleagues [3] speculated that the metabolites of caffeine – paraxanthine, theobromine, and theophylline – have additional ergogenic effects; in this case, the presence of these ergogenic substances would be lower in C allele carriers at a given time point due to the slower metabolization of caffeine, reducing caffeine's performance benefits. This proposed mechanism was echoed by Rahimi [1].

If the above mechanisms are indeed correct, then there remains the possibility that caffeine can still be ergogenic for C allele carriers, but that such individuals need to consume it a greater amount of time prior to exercise. In the majority of studies – including Guest et al. [2] and Rahimi [1] – subjects consume caffeine ~60 minutes pre-exercise. However, for C allele carriers, could the ergogenic effects of caffeine be restored by utilizing a caffeine dose 90 or 120 minutes pre-exercise? Such an hypothesis is, of course, speculative, and requires testing – but it does represent a potential way by which caffeine can indeed be ergogenic for all. The resolution of whether caffeine is truly ergolytic or neutral for *CYP1A2* C allele carriers, or if it merely necessitates a different caffeine strategy, represents an important step on the journey towards more personalized sports nutrition guidelines.

**Compliance with Ethical Standards:**

Conflict of Interest: Craig Pickering is an employee of DNAFit Ltd, a genetic testing company. He received no financial incentives for the preparation of this manuscript, and the views contained within are his, and not representative of DNAFit Ltd.

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

## References:

1. Rahimi R. The effect of CYP1A2 genotype on the ergogenic properties of caffeine during resistance exercise: a randomized, double-blind, placebo-controlled, crossover study. *Ir J Med Sci*. 2018 doi: 10.1007/s11845-018-1780-7
2. Guest N, Corey P, Vescovi J, El-Sohemy A. Caffeine, CYP1A2 Genotype, and Endurance Performance in Athletes. *Med Sci Sports Exerc*. 2018. doi:10.1249/MSS.0000000000001596.
3. Womack CJ, Saunders MJ, Bechtel MK, Bolton DJ, Martin M, Luden ND, Dunham W, Hancock M. The influence of a CYP1A2 polymorphism on the ergogenic effects of caffeine. *J Int Soc Sports Nutr*. 2012;9(1):7.
4. Pataky MW, Womack CJ, Saunders MJ, Goffe JL, D'lugos AC, El-Sohemy A, Luden ND. *Scand J Med Sci Sports*. 2016;26(6):613-9.
5. Pickering C, Kiely J. Are the current guidelines on caffeine use in sport optimal for everyone? Inter-individual variation in caffeine ergogenicity, and a move towards personalised sports nutrition. *Sports Med*. 2018;48(1):7-16.
6. Sachse C, Brockmöller J, Bauer S, Roots I. Functional significance of a C→A polymorphism in intron 1 of the cytochrome P450 CYP1A2 gene tested with caffeine. *Br J Clin Pharmacol*. 1999;47(4):445-9.