

Accepted Manuscript

Intracellular acidification reduces L-arginine transport via system y⁺L but not via system y⁺/CATs and nitric oxide synthase activity in human umbilical vein endothelial cells

Marco A. Ramírez, Jorge Morales, Marcelo Cornejo, Elias H. Blanco, Edgardo Mancilla-Sierpe, Fernando Toledo, Ana R. Beltrán, Luis Sobrevia

PII: S0925-4439(18)30043-7

DOI: <https://doi.org/10.1016/j.bbadis.2018.01.032>

Reference: BBADIS 65048

To appear in:


Received date: 7 November 2017

Revised date: 15 January 2018

Accepted date: 31 January 2018

Please cite this article as: Marco A. Ramírez, Jorge Morales, Marcelo Cornejo, Elias H. Blanco, Edgardo Mancilla-Sierpe, Fernando Toledo, Ana R. Beltrán, Luis Sobrevia , Intracellular acidification reduces L-arginine transport via system y⁺L but not via system y⁺/CATs and nitric oxide synthase activity in human umbilical vein endothelial cells. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Bbadis(2018), <https://doi.org/10.1016/j.bbadis.2018.01.032>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Intracellular acidification reduces L-arginine transport via system y⁺L but not via system y⁺/CATs and nitric oxide synthase activity in human umbilical vein endothelial cells

Marco A Ramírez^{1,2,*}, Jorge Morales^{1,2}, Marcelo Cornejo², Elias H Blanco², Edgardo Mancilla-Sierpe^{2,3}, Fernando Toledo^{1,4}, Ana R Beltrán^{2,5}, Luis Sobrevia^{1,6,7,*}

¹ Cellular and Molecular Physiology Laboratory (CMPL), Division of Obstetrics and Gynaecology, School of Medicine, Faculty of Medicine, Pontificia Universidad Católica de Chile, Santiago 8330024, Chile. ² Cellular Physiology Laboratory, Biomedical Department, Faculty of Health Sciences, Universidad de Antofagasta, Antofagasta 1270300, Chile. ³ Unit of Pathological Anatomy, Hospital Regional de Antofagasta, Antofagasta 1270001, Chile. ⁴ Department of Basic Sciences, Faculty of Sciences, Universidad del Bío-Bío, Chillán 3780000, Chile. ⁵ Department of Education, Faculty of Education, Universidad de Antofagasta, Antofagasta 1270300, Chile. ⁶ Department of Physiology, Faculty of Pharmacy, Universidad de Sevilla, Seville E-41012, Spain. ⁷ University of Queensland Centre for Clinical Research (UQCCR), Faculty of Medicine and Biomedical Sciences, University of Queensland, Herston, QLD 4029, Queensland, Australia.

Running title: Modulation of L-arginine transport by intracellular pH

Correspondence: Dr Marco A Ramírez
Biomedical Department, Faculty of Health Sciences
Universidad de Antofagasta
Antofagasta 1270300, Chile
Tel. +5655-2637434, Fax +5655-2637259, E-mail:
marco.ramirez@uantof.cl

Professor Luis Sobrevia
Cellular and Molecular Physiology Laboratory (CMPL)
Division of Obstetrics and Gynaecology, School of Medicine
Faculty of Medicine, Pontificia Universidad Católica de Chile
Santiago 8330024, Chile.
Tel. +562-23548116, Fax +562-26321924, E-mail: lsobrevia@uc.cl

Abstract

L-Arginine is taken up via the cationic amino acid transporters (system y^+ /CATs) and system y^+L in human umbilical vein endothelial cells (HUVECs). L-Arginine is the substrate for endothelial NO synthase (eNOS) which is activated by intracellular alkalization, but nothing is known regarding modulation of system y^+ /CATs and system y^+L activity, and eNOS activity by the pHi in HUVECs. We studied whether an acidic pHi modulates L-arginine transport and eNOS activity in HUVECs. Cells loaded with a pH-sensitive probe were subjected to 0.1-20 mmol/L NH_4Cl pulse assay to generate pHi 7.13-6.55. Before pHi started to recover, L-arginine transport (0-20 or 0-1000 $\mu\text{mol/L}$, 10 s, 37°C) in the absence or presence of 200 $\mu\text{mol/L}$ *N*-ethylmaleimide (NEM) (system y^+ /CATs inhibitor) or 2 mmol/L L-leucine (system y^+L substrate) was measured. Protein abundance for eNOS and serine¹¹⁷⁷ or threonine⁴⁹⁵ phosphorylated eNOS was determined. The results show that intracellular acidification reduced system y^+L but not system y^+ /CATs mediated L-arginine maximal transport capacity due to reduced maximal velocity. Acidic pHi reduced NO synthesis and eNOS serine¹¹⁷⁷ phosphorylation. Thus, system y^+L activity is downregulated by an acidic pHi, a phenomenon that may result in reduced NO synthesis in HUVECs.

Keywords: L-arginine transport; intracellular pH; system y^+L ; system y^+ ; endothelium

Abbreviations

pH _o	Extracellular pH
pH _i	Intracellular pH
4F2hc	Heavy chain of the cell surface antigen 4F2
HUVECs	Human umbilical vein endothelial cells
HPAEC	Human pulmonary artery endothelial cells
RAEC	Rat aorta endothelial cells
hENT1/2	human equilibrative nucleoside transporters 1 and 2
GDM	Gestational diabetes mellitus
hCATs	Human cationic amino transporters
NO	Nitric oxide
NOS	Nitric oxide synthase
eNOS	Endothelial nitric oxide synthase
NH ₄ Cl	Ammonium chloride
DAF-FM	4-amino-5-methylamino-2',7'-difluorofluorescein
L-NAME	<i>N</i> ^G -nitro-L-arginine methyl ester
BCECF-AM	Bicarboxyethyl-5,6-carboxyfluorescein acetoxymethyl ester
NEM	<i>N</i> -ethylmaleimide

1. Introduction

A variety of membrane transport systems removing metabolic substrates from the extracellular medium are expressed in the foetoplacental endothelium [1-4]. The activity of some of these transport systems is modulated by changes in the extracellular (pHo) and intracellular (pHi) pH [5,6]. The cationic amino acid L-arginine, the substrate for the synthesis of nitric oxide (NO) via the endothelial NO synthase (eNOS) [7,8], is taken up mainly by the cationic amino acid transporters (CATs, also referred as system y^+ or system y^+ /CATs) family [4] and system y^+L in human umbilical vein endothelial cells (HUVECs) [2,9,10]. System y^+ /CATs corresponds to a family of five proteins of which mainly the high affinity ($K_m \sim 100$ - $250 \mu\text{mol/L}$) hCAT-1 and hCAT-2B isoforms are expressed in HUVECs [4,11]. System y^+L activity results from heterodimers formed by the interaction of the heavy chain of the cell surface antigen 4F2 (4F2hc) with the light chains 4F2-1c2 (or y^+LAT -1) or 4F2-1c3 (or y^+LAT -2) [1,11-14]. System y^+L activity accounts for L-arginine transport with a very high affinity ($K_m \sim 1$ - $20 \mu\text{mol/L}$) and small and large neutral amino acids, such as L-leucine, requiring extracellular sodium in HUVECs [4,9]. System y^+ /CATs and system y^+L activity are reported as independent of a change in pHo in mammalian cells [11,14]. However, there are no reports addressing whether the activity of these membrane transport systems is modulated by the pHi.

Increased L-arginine transport mediated by system y^+ /CAT-1 [15] and system y^+L [10] results in higher eNOS activity in HUVECs and other cell types [16,17]. Interestingly, intracellular alkalization activates eNOS in HUVECs [18], human pulmonary arterial endothelial cells (HPAECs) [19], and rat aorta endothelial cells (RAECs) [20]. However, it is unknown whether eNOS activation in response to a change in the pHi leading to an alkaline or acidic intracellular environment associated with system y^+ /CATs and system y^+L transport activity in human endothelial cells. Intracellular alkalization due to lower NHE1 activity

reduced the transport of the endogenous nucleoside adenosine in HUVECs [6]. Since adenosine is a vasodilator in most vascular beds including the foetoplacental circulation [21] via increasing the L-arginine transport and NO synthesis in HUVECs [22], and dysfunction of the foetoplacental vasculature is addressed as the cause of altered umbilical vein blood flow in growth restricted foetus [23,24], it is likely that changes in the pHi in HUVECs alters the dynamics of NO-dependent dilation mechanisms of the umbilical vein therefore limiting the delivery of nutrients to the foetus [25]. This study aimed to characterise the role of a change in pHi on L-arginine transport mediated via system y^+ /CATs and system y^+L and on NO synthesis in HUVECs.

2. Material and methods

2.1 Antibodies and materials

Primary monoclonal mouse *anti*-eNOS phosphorylated at serine¹¹⁷⁷, *anti*-eNOS phosphorylated at threonine⁴⁹⁵, and *anti*- β -actin were from Sigma Aldrich (St Louis, MO, USA). Primary monoclonal mouse *anti*-total eNOS antibody and secondary horseradish peroxidase-conjugated goat *anti*-mouse antibodies were from Santa Cruz Biotechnology (Santa Cruz, CA, USA). For isolation of HUVECs from umbilical cords, Collagenase Type II from *Clostridium histolyticum* (Boehringer, Mannheim, FRG) was used. Medium M199, newborn (NBCS) and fetal calf (FCS) sera, L-glutamine, and penicillin-streptomycin were from Gibco Life Technologies (Carlsbad, CA, USA). L-[³H]Arginine and D-[³H]mannitol were from NEN (Dreieich, FRG). N^G -Nitro-L-arginine methyl ester (L-NAME) was from Sigma Aldrich, Immobilon-P polyvinylidene difluoride membranes from BioRad Laboratories (Hertfordshire, UK), and the fluorescent dye 4-amino-5-methylamino-2',7'-difluorofluorescein (DAF-FM) from Molecular Probes (Leiden, The Netherlands).

2.2 *Study group*

This study included samples collected from 23 full-term normal pregnancies from the Hospital Clínico UC-CHRISTUS (HCUC-C) in Santiago de Chile and Clínica de la Mujer (CLM) in Antofagasta (Chile). Pregnant women included in this study did not smoke or consume drugs or alcohol and had no intrauterine infection or any other medical or obstetrical complications. The ethnicity of patients involved in this study was Hispanic. The investigation conforms to the principles outlined in the Declaration of Helsinki. Ethics Committee approvals from the Faculty of Medicine of the Pontificia Universidad Católica de Chile and CLM and informed written consent of patients were obtained.

2.3 *Human placenta and umbilical cords*

Placentas were collected at delivery on ice and transferred to the laboratory until use 15-30 min later. Middle sections of umbilical cords (100-120 mm length) were dissected into 200 mL phosphate-buffered saline (PBS) solution (mmol/L: 130 NaCl, 2.7 KCl, 0.8 Na₂HPO₄, 1.4 KH₂PO₄, pH 7.4, 4°C) until use 6-12 h later for isolation of endothelial cells [6,26].

2.4 *Cell culture*

This study was done in primary cultures of HUVECs from normal pregnancies. The reason why selecting this type of cells is because (i) they are from the umbilical vein which carries foetal blood after crossing the placenta circulatory bed towards the foetus body with the umbilical vein blood being rich in oxygen and nutrients and unloaded of toxins and waste from the foetus circulation, (ii) umbilical vein blood carries signalling molecules that are transferred from the mother through the placenta into the foetal circulation, (iii) molecules

synthesised and released within the placenta tissue are available at the umbilical vein blood thus transferring regulatory signals from the placenta to this vessel by changing, for example, the offering of nutrients to the growing foetus, and (iv) HUVECs release extracellular vesicles, including exosomes, that could potentially alter the downstream vasculature (i.e., the foetal circulation) altering or changing the function or phenotype of the endothelium in the foetal vascular bed [27,28]. HUVECs were isolated by collagenase digestion (0.25 mg/mL collagenase) from umbilical cords obtained at delivery from normal pregnancies and cultured (37°C, 5% CO₂) in 1% gelatin-coated Petri dishes (100 mm diameter) up to passage 3 in primary culture medium (PCM; M199 containing 5 mmol/L D-glucose, 10% NBCS, 10% FCS, 3.2 mmol/L L-glutamine and 100 U/mL penicillin-streptomycin) as reported [6,26]. Sixteen hours prior experiments the incubation medium was changed to M199 medium containing 0.25% NBCS and 0.25% FCS. Experiments were in the absence or presence of *N*^G-nitro-L-arginine methyl ester (L-NAME, 100 µmol/L, NOS inhibitor) and cell viability was assayed using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetra-zolium bromide assay (Sigma-Aldrich) as reported [6].

2.5 *pHi measurement and recovery*

Cells were loaded (10 min, 37°C) with the fluorescent pH-sensitive probe 2,7-bicarboxyethyl-5,6-carboxyfluorescein acetoxymethyl ester (BCECF-AM, 12 µmol/L) as described [6]. Probe excess was removed rinsing (x3) with control solution (CS) (mmol/L: NaCl 145, KCl 5, NaH₂PO₄ 1 Na₂SO₄ 1, CaCl₂ 1.8, MgCl₂ 1, HEPES 30, D-glucose 5, pH 7.4, 37°C). Fluorescence ratios were registered every 0.5-seconds interval. The pHi was estimated using standard calibration curves with 10 µmol/L nigericin and high-K⁺ in a calibrating

solution (pH 6.2, 7.2, 8.2) as described [6]. The pH_i recovery was examined by the NH_4Cl pulse technique [6]. After the basal pH_i was stabilized (~ 3 min) cells were exposed (2 min) to CS with 0.1, 1, or 20 mmol/L NH_4Cl ($\text{NH}_4\text{Cl}/\text{CS}$ solution). Cells were then rinsed with NH_4Cl -free CS, and cell viability assayed as above.

2.6 Uptake of L-arginine

Since pH_i recovery started after 25 s of removal of $\text{NH}_4\text{Cl}/\text{CS}$, transport assays in CS were performed at 20 s (37°C). To identify the involvement of system y^+/CATs and system y^+L on L-arginine transport the cells were incubated with CS with or without 200 $\mu\text{mol/L}$ *N*-ethylmaleimide (NEM) (a general inhibitor of system y^+/CATs) [1,11], 2 mmol/L L-leucine (a neutral amino acid that competes with L-arginine for system y^+L) [1,11], or NEM + L-leucine as previously described [10]. Overall uptake at 2 and 100 $\mu\text{mol/L}$ L-arginine (6 $\mu\text{Ci/mL}$ L- ^3H arginine, 20 s, 37°C) for system y^+L and system y^+/CATs , respectively, was measured in confluent cells in CS as described [10,26]. The fraction of uptake inhibited by NEM was considered as system y^+/CATs mediated, and the portion of uptake inhibited by L-leucine in cells coincubated with NEM to block system y^+/CATs contribution was regarded as system y^+L mediated [1,10,11].

2.7 Kinetics of L-arginine transport

Overall 0-20 $\mu\text{mol/L}$ or 0-1000 $\mu\text{mol/L}$ L-arginine transport (for system y^+L and system y^+/CATs , respectively) was measured in CS as above. Overall transport of L-arginine was defined as the sum of a saturable component plus a non-saturable, linear component of transport in the ranges of L-arginine concentrations used in this study (hereafter referred as a

K_D value defined by $m \cdot [Arg]$, where m corresponds to slopes of lineal phases of transport at a given L-arginine concentration $[Arg]$ [26]. Cell monolayers were rinsed with ice-cold CS to terminate tracer uptake.

The initial rate of transport (i.e., linear uptake up to 10 s) was derived from the slope of the linear phases of L-arginine transport. Values for transport were adjusted to the one phase exponential association equation considering the least squares fit:

$$v_i = V_m \cdot (1 - e^{-(k \cdot t)})$$

where v_i is initial velocity, V_m is mayor velocity at a given time (t) and L-arginine concentration, and e and k are constants. Overall L-arginine transport at initial rates was adjusted to the Michaelis-Menten hyperbola plus a nonsaturable, linear component (K_D) as described [26]. The saturable transport of L-arginine was derived by subtracting the $m \cdot [Arg]$ components from overall transport, and the transport kinetic parameters maximal velocity (V_{max}) and apparent Michaelis-Menten constant (K_m) of transport were calculated [26].

The relative contribution of system y^+L and system $y^+/CATs$ (y^+L/y^+F) to total transport (i.e., y^+L plus $y^+/CATs$ mediated transport) in cells non-treated ($-NH_4$) or treated ($+NH_4$) with NH_4Cl was estimated from V_{max}/K_m values by:

$$y^+L/y^+F_{-NH_4} = \frac{y^+L V_{max} \cdot y^+K_m}{y^+L K_m \cdot y^+V_{max}}$$

or

$$y^{+L}/y^{+}F_{+NH_4} = \frac{y^{+L}V_{max} \cdot y^{+}K_m}{y^{+L}K_m \cdot y^{+}V_{max}}$$

where $y^{+L}V_{max}$ and $y^{+L}K_m$ are kinetic parameters for system y^{+L} -saturable transport, and $y^{+}V_{max}$ and $y^{+}K_m$ for system y^{+} /CATs saturable transport.

The relative effect of NH_4Cl on transport activity via system y^{+L} ($1/^{-NH_4/+NH_4}F_{y^{+L}}$) or system y^{+} /CATs ($1/^{-NH_4/+NH_4}F_{y^{+}}$) was estimated by:

$$\frac{1}{^{-NH_4/+NH_4}F_{y^{+L}}} = \frac{^{+NH_4}V_{max} \cdot ^{-NH_4}K_m}{^{-NH_4}V_{max} \cdot ^{+NH_4}K_m}$$

or

$$\frac{1}{^{-NH_4/+NH_4}F_{y^{+}}} = \frac{^{+NH_4}V_{max} \cdot ^{-NH_4}K_m}{^{-NH_4}V_{max} \cdot ^{+NH_4}K_m}$$

where $^{-NH_4}V_{max}$ and $^{-NH_4}K_m$, or $^{+NH_4}V_{max}$ and $^{+NH_4}K_m$ are kinetic parameters for transport in cells non-treated or treated with NH_4Cl , respectively [6].

The efficiency of the effect of a change in the pHi (E_{pHi}) on the uptake at a fixed concentration ($^U E_{pHi}$) or the maximal transport capacity ($^{Vmax/Km} E_{pHi}$) for a range of concentrations of L-arginine via system y^{+} /CATs and system y^{+L} was estimated by:

$$^U E_{pHi} = \frac{^{-NH_4}U - ^{+NH_4}U}{^{-NH_4}pHi - ^{+NH_4}pHi}$$

or

$$v_{max/Km} E_{pHi} = \frac{-^{NH_4} V_{max}/K_m - ^{+NH_4} V_{max}/K_m}{-^{NH_4} pHi - ^{+NH_4} pHi}$$

where uptake (U) at a given concentration of L-arginine (2 or 100 $\mu\text{mol/L}$ in this study) was measured in the absence ($-^{NH_4}$) or presence ($+^{NH_4}$) of NH_4Cl at basal pHi ($-^{NH_4} pHi$) or pHi in the presence of NH_4Cl ($+^{NH_4} pHi$). For a range of concentrations of L-arginine (0-20 or 0-1000 $\mu\text{mol/L}$ in this study), the values for V_{max}/K_m were used. Each transport assay was run in duplicate with transport activity expressed as $\text{pmol}/\mu\text{g}$ protein/minute. Values for $^U E_{pHi}$ and $v_{max/Km} E_{pHi}$ are expressed as a change in the $\text{pmol}/\mu\text{g}$ protein/minute relative to 1. Radioactivity in 0.5 N KCl cell digests was determined by liquid scintillation counting, and uptake was corrected for D- ^3H]mannitol disintegrations per minute (d.p.m.) in the extracellular space [26].

2.8 *NOS activity*

NOS activity was assayed by quantification of the intracellular content of L-citrulline by high-performance liquid chromatography in confluent HUVECs in the absence or presence of 100 $\mu\text{mol/L}$ L-NAME, as reported [15,26].

2.9 *Western blot for eNOS*

Total protein was obtained from confluent cells washed twice with ice-cold PBS and harvested in 100 μL of lysis buffer composed by 63.7 mmol/L Tris/HCl (pH 6.8), 10% glycerol, 2% sodium dodecylsulphate, 1 mmol/L sodium orthovanadate, 50 mg/mL leupeptin, and 5% 2-mercaptoethanol, as described [26]. Cells were sonicated (6 cycles, 5 s, 100 Watts,

4°C), and total protein was separated by centrifugation (14000 g, 15 min, 4°C). Proteins (60 µg) were separated by polyacrylamide gel (10%) electrophoresis and transferred onto Immobilon-P polyvinylidene difluoride membranes. The proteins were then probed against total eNOS (1:500 dilution, 12 h, 4°C), eNOS phosphorylated at serine¹¹⁷⁷ (*P*-Ser¹¹⁷⁷-eNOS, 1:1000 dilution, 12 h, 4°C), eNOS phosphorylated at threonine⁴⁹⁵ (*P*-Thr⁴⁹⁵-eNOS, 1:1000 dilution, 12 h, 4°C), and β-actin (1:3000, 1 h, room temperature). Membranes were rinsed in Tris buffer saline-Tween (TBS-T) and incubated (1 h) in TBS-T/0.2% BSA containing secondary horseradish peroxidase-conjugated antibodies. Proteins were detected by enhanced chemiluminescence (film exposure time was 1 min) in a ChemiDoc-It 510 Imagen System (UVP, LCC Upland, CA, USA) and quantified by densitometry [26].

2.10 Statistical analysis

The sample size was estimated considering a power of 80% to detect a difference between groups (by a two-sided alpha level of 0.05). Values for clinical parameters are given as mean ± S.D. For *in vitro* assays the values were mean ± S.E.M., where *n* indicates the number of different biological samples and corresponding cell cultures with 3-4 replicates per experiment. Comparisons between two groups were performed using Student's unpaired *t*-test and between more than two groups by analysis of variance (ANOVA, two-ways). If the ANOVA demonstrated a significant interaction between variables, *post hoc* analyses were performed by the multiple-comparison Bonferroni test. The statistical software GraphPad InStat 3.1 and GraphPad Prism 7.0d (GraphPad Software Inc., San Diego, CA, USA) was used for data analysis. *P*<0.05 was considered statistically significant.

3. Results

3.1 Study group

Pregnant women included in this study were with normal pregnancy, normotensive, normal fasting glycaemia at delivery, singleton, and of similar age and height (Table 1). Weight and body mass index (BMI) at delivery were higher compared with the first determination early in pregnancy (9-16 weeks of gestation) where women were normoweight (BMI <25 kg/m²). The total gestational weight gain between early in pregnancy and delivery was 11.9 ± 0.7 kg with a BMI variation of 0.6 kg/m² reaching BMI values that were >25-29.9 kg/m².

3.1. Basal pHi

Exposure of cells to 20 mmol/L NH₄Cl increased the pHi value, and NH₄Cl removal caused rapid acidification (~1 second, pHi = 6.51 ± 0.04) lasting for ~25 s before a significant pHi recovery started reaching initial pHi value in ~6 min (Figure 1A). The basal pHi value (7.19 ± 0.03) was reduced by NH₄Cl in a concentration-dependent manner (half-maximal effective concentration (EC_{50}) = 1.29 ± 0.03 mmol/L NH₄Cl, equivalent to pHi = 6.75 ± 0.02) (Figure 1B), without altering the cell survival (97-99% alive cells between 0.1-20 mmol/L NH₄Cl) (not shown), confirming previous observations in this cell type [6].

3.2. System y⁺/CATs and system y⁺L mediated uptake of L-arginine

In the absence of NH₄Cl (i.e., at basal pHi), the overall uptake of 100 μmol/L L-arginine was inhibited mainly by NEM ($69 \pm 6\%$), with a minor inhibition caused by L-leucine ($25 \pm 3\%$) but blocked in cells coincubated with NEM + L-leucine (Figure 1C). In cells exposed to 20 mmol/L NH₄Cl and washed with CS solution (i.e., pHi ~6.5), the overall uptake

was reduced compared with cells in the absence of NH_4Cl in a proportion (0.14 ± 0.02 pmol/ μg protein/minute) that was similar in cells incubated with L-leucine in the absence of NH_4Cl (0.15 ± 0.06 pmol/ μg protein/minute). In the presence of NH_4Cl , the uptake of L-arginine was inhibited by NEM or NEM + L-leucine in a similar proportion (0.42 ± 0.02 and 0.44 ± 0.02 pmol/ μg protein/minute, respectively), but NH_4Cl did not alter the uptake of L-arginine inhibited by L-leucine in the absence of this salt. Uptake of L-arginine mediated by system y^+/CATs predominates over a minor contribution of system y^+L at $100 \mu\text{mol/L}$ L-arginine (Figure 1D).

The overall uptake at $2 \mu\text{mol/L}$ L-arginine (0.17 ± 0.05 pmol/ μg protein/minute) in the absence of NH_4Cl was lower ($67 \pm 3\%$) compared with $100 \mu\text{mol/L}$ L-arginine (Figure 1E). The uptake detected at this concentration of L-arginine was similar to the fraction of uptake inhibited by L-leucine in $100 \mu\text{mol/L}$ L-arginine (0.15 ± 0.06 pmol/ μg protein/minute). Overall uptake was unaltered by NEM but blocked by L-leucine or NEM + L-leucine. In the presence of NH_4Cl the $2 \mu\text{mol/L}$ L-arginine uptake was abolished in all experimental conditions. Uptake of L-arginine mediated by system y^+L accounted for $2 \mu\text{mol/L}$ L-arginine in HUVECs (Figure 1F). NH_4Cl did not alter 100 or $2 \mu\text{mol/L}$ L-arginine uptake via system y^+/CATs . However, the system y^+L activity in the presence of NH_4Cl at these two concentrations of L-arginine was abolished.

3.3. *pHi-dependent uptake of L-arginine via system y^+L/CATs and system y^+L*

Overall uptake of $100 \mu\text{mol/L}$ L-arginine was unaltered by 0.1 or 1 mmol/L NH_4Cl in the absence of NEM and by 0.1 mmol/L NH_4Cl in the presence of NEM, but reduced by 20 mmol/L NH_4Cl in the absence or presence of NEM (Figure 2A). Incubation of cells with NEM resulted in a NH_4Cl concentration-dependent inhibition of uptake ($EC_{50} = 0.32 \pm 0.04 \text{ mmol/L}$

NH_4Cl). Overall uptake of 2 $\mu\text{mol/L}$ L-arginine in the presence of NEM was inhibited by NH_4Cl ($EC_{50} = 0.31 \pm 0.03 \text{ mmol/L NH}_4\text{Cl}$), but uptake was unaltered in cells incubated with NEM + L-leucine (Figure 2B). Uptake mediated by system y^+/CATs was not significantly altered by NH_4Cl (Figure 2C) and independent of the resulting pHi (Figure 2D). However, uptake mediated by system y^+L was reduced in a concentration-dependent manner by NH_4Cl ($EC_{50} = 0.29 \pm 0.03 \text{ mmol/L NH}_4\text{Cl}$) and the resulting acidic pHi ($EC_{50} = 6.89 \pm 0.11 \text{ pHi}$).

The efficiency of inhibition of a change in pHi (ΔpHi) on 2 $\mu\text{mol/L}$ L-arginine uptake ($^U E_{\text{pHi}}$) mediated via system y^+L was higher at the smaller variation of pHi (ΔpHi 0.06 in this study) and less pronounced but reaching comparable values at higher variations of pHi (ΔpHi 0.38 and 0.69 in this study) from the basal pHi value in HUVECs (Figure 2E). However, the $^U E_{\text{pHi}}$ for uptake via system y^+/CATs was unaltered by the pHi. The $^U E_{\text{pHi}}$ for system y^+L compared with system y^+/CATs mediated uptake was higher at pHi 7.19 compared with uptake at pHi 6.81 or 6.5 (Figure 2F).

3.4. *pHi-dependent system y^+/CATs and system y^+L transport kinetics*

In the absence of NH_4Cl , the overall transport of L-arginine in the range of 0-1000 $\mu\text{mol/L}$ was semi-saturable, inhibited mainly by NEM but marginally inhibited by L-leucine, and blocked by NEM + L-leucine (not shown) as previously reported [10]. The derived transport obtained after subtracting the linear, non-saturable component from overall transport (i.e., K_D), was saturable in all experimental conditions and adjusted to a first order linear regression in Eadie-Hofstee plots as reported [10,15]. Incubation of cells with increasing concentrations of NH_4Cl did not alter the K_D for overall transport and the V_{max} , K_m , or V_{max}/K_m for saturable transport in this range of L-arginine concentrations (Table 2).

Overall transport of L-arginine in the range of 0-20 $\mu\text{mol/L}$ L-arginine was semi-saturable, unaltered by NEM but abolished by L-leucine or NEM + L-leucine (not shown) as previously reported [10]. The derived saturable L-arginine transport was unaffected by NEM but blocked by L-leucine and NEM + L-leucine (Figure 3A). A first-degree regression line well fitted saturable transport at different pHi values in Eadie-Hofstee plots (Figure 3B). Increasing concentrations of NH_4Cl did not alter the K_D for overall transport but reduced the V_{max} (Table 2) and V_{max}/K_m (Figure 3C) without changing the K_m for saturable transport.

The efficiency of inhibition of a given ΔpHi in 0-20 $\mu\text{mol/L}$ L-arginine transport kinetic parameters ($V_{\text{max}}/K_m E_{\text{pHi}}$) for system y^+L was also higher at the smaller variation of pHi (ΔpHi 0.06) and less pronounced but reaching comparable values at higher variations of pHi (ΔpHi 0.38 and 0.69) from the basal pHi (Figure 3D). The pHi unaltered the $V_{\text{max}}/K_m E_{\text{pHi}}$ for transport via system y^+/CATs . The $V_{\text{max}}/K_m E_{\text{pHi}}$ for system y^+L compared with system y^+/CATs mediated transport was higher at all pHi used in this study (Figure 3E).

3.6 NOS activity

In the absence of NH_4Cl , total synthesis of L-citrulline in cells incubated with 100 $\mu\text{mol/L}$ L-arginine was partially reduced by NEM and L-leucine but abolished by NEM + L-leucine (Figure 4A). NH_4Cl similarly inhibited L-citrulline synthesis in the absence or presence of L-leucine but blocked by NEM and NEM + L-leucine. Incubation of cells with L-NAME blocked L-citrulline synthesis in all experimental conditions. The NOS-dependent fraction of synthesis of L-citrulline was reduced by NEM or L-leucine but abolished by NEM + L-leucine (Figure 4C). In the presence of NH_4Cl , NOS-dependent L-citrulline synthesis was decreased partially reaching similar values to those in the presence of L-leucine. However, it was abolished in the presence of NEM or NEM + L-leucine.

In the absence of NH_4Cl , total synthesis of L-citrulline in cells incubated with 2 $\mu\text{mol/L}$ L-arginine was unaltered by NEM but abolished by L-leucine and NEM + L-leucine (Figure 4B). NH_4Cl and L-NAME also abolished L-citrulline synthesis. The NOS-dependent synthesis of L-citrulline in the absence of NH_4Cl was unaltered by NEM but blocked by L-leucine and NEM + L-leucine (Figure 4D). NH_4Cl reduced, but NEM, L-leucine, and NEM + L-leucine blocked the NOS-dependent synthesis of L-citrulline.

3.7 *eNOS* expression and activation

Total eNOS protein abundance was unaltered by an acidic pHi (Figure 5A,B). Phosphorylation of eNOS at Ser¹¹⁷⁷ was reduced in a pHi-dependent manner ($EC_{50} = 0.77 \pm 0.03$ pHi) (Figure 5C); however, eNOS phosphorylation at Thr⁴⁹⁵ was unaltered by acidic pHi (Figure 5D).

4. Discussion

This study shows that pHi is a factor that modulates the L-arginine transport in primary cultured HUVECs from normal pregnancies. Intracellular acidification causes a reduction in the L-arginine transport via system y^+L but not via system $y^+/CATs$, and in the activity of eNOS due to lower activator phosphorylation in Ser¹¹⁷⁷ at this enzyme. Since NO is involved in a broader range of biological effects other than regulation of vascular tone, intracellular acidification may have significant implications in diseases associated with endothelial dysfunction, such as gestational diabetes mellitus and cancer, where the pHo and pHi are altered [6,27–29].

HUVECs show a pHi ~7.19 as previously reported (pHi ~7.21) [6,30,31], and is close to the pH reported in the human umbilical vein blood (pH ~7.35) [25,32-34]. Intracellular acidification downregulates the transport activity of the human equilibrative nucleoside transporters 1 and 2 in HUVECs [6], and Na⁺/H⁺ exchanger 1 in human lymphoblasts [34] and the human colonic carcinoma T₈₄ cell line [35]. Since overall L-arginine transport was reduced as the pHi changed to acidic in HUVECs, L-arginine transport mechanisms are responsive to a change in the pHi in this cell type. Inhibition by the acidic pHi was partial and similar to that induced by L-leucine at basal or acidic pHi in the presence of 100 μmol/L L-arginine. Considering the apparent K_m for L-arginine uptake via hCAT-1 (K_m ~120 μmol/L) and hCAT-2B (K_m ~250 μmol/L) in this cell type [1,2,36] it is likely that these isoforms were involved in this phenomenon. L-Arginine transport via system y⁺/CATs is independent of pHo in mammalian cells [2,11,37]. However, there are no studies addressing modulation of L-arginine transport by pHi in endothelium or other cell types [2,11,34,37]. Our results suggest that system y⁺/CATs activity (likely hCAT-1 and hCAT-2B) is independent of intracellular acidification up to ΔpHi ~0.69 from the basal pHi since the effect of the ΔpHi on 100 μmol/L L-arginine uptake (${}^U E_{pHi}$ 0.022 ± 0.012 , range 0.01-0.06) or the V_{max}/K_m for transport (${}^{V_{max}/K_m} E_{pHi}$ 0.0037 ± 0.0011 , range 0.002-0.005) of this amino acid was unaltered.

Uptake of L-arginine is also mediated by system y⁺L in HUVECs [9,10]. Uptake of 2 μmol/L L-arginine was almost exclusively mediated via system y⁺L meanwhile at higher levels (100 μmol/L) transport was via system y⁺/CATs and system y⁺L. The results show that system y⁺L transport activity was sensitive to acidic pHi in HUVECs [34,38], an effect that was higher at smaller changes from the basal pHi. Thus, system y⁺L seems more efficiently modulated by a discrete change in pHi (ΔpHi 0.06 in this study) from the physiological pHi in

HUVECs. Similar changes were seen with 2 $\mu\text{mol/L}$ or a broader concentration of L-arginine ($U_{E_{\text{pHi}}} / V_{\text{max}}/K_m E_{\text{pHi}} \sim 0.8$). Thus, pHi modulation of system y^+L activity results from changing the V_{max}/K_m for system y^+L , an effect that results from reduced V_{max} . Several possibilities may explain this finding, i.e., (i) reduced number of transporters available at the plasma membrane due to lower expression or recycling with no change in their transport capacity, (ii) decreased transport capacity of a fixed number of membrane transporters, or (iii) both phenomena. Since pHi effect on transport was assayed for 15 s and system y^+L half-life is most likely unaltered at this incubation time, the reduced L-arginine transport may result from a lower activity rather than expression of system y^+L in HUVECs.

An increase in the activity of system y^+L associated with a higher synthesis of NO in HUVECs [1,10], human platelets [16,17], and rat cortical astrocytes [39]. Also, since system y^+L may be located close to eNOS in the plasma membrane [2], the pHi-decreased system y^+L transport activity may result in lower eNOS activity in HUVECs. Our results show that L-NAME-inhibited L-citrulline formation from L-arginine (index of NOS activity) [8] was lower at acidic pHi. This phenomenon was associated with a reduced activity of system y^+L , but not system $y^+/CATs$ and abolished at 2 $\mu\text{mol/L}$ but partially reduced at 100 $\mu\text{mol/L}$ L-arginine. Interestingly, eNOS activity seems linked to system $y^+/CATs$ (particularly hCAT-1 and hCAT-2) in HUVECs [2,40–42]. However, an intracellular L-arginine pool not fed from this amino acid extracellular content is also a supplying source for eNOS in this cell type [40,42]. Thus, CATs transport activity could be unaltered, up or downregulated and these changes will not necessarily lead to parallel changes in NOS activity. Indeed, HUVECs from late-onset preeclampsia show increased hCAT-1-mediated L-arginine transport but reduced eNOS activity [43,44]. Potential explanations for this phenomenon include the possibility that

system γ^+ /CATs activity may deliver L-arginine for NOS activity and NO generation, arginase activity for the synthesis of polyamines, or for protein synthesis [43,44].

The reduced NOS activity seen in HUVECs in an acidic pH_i was likely due to lower eNOS activation since its lower activator phosphorylation at serine¹¹⁷⁷ [8,18,26] instead of an increased inhibitory phosphorylation of threonine⁴⁹⁷ [8,18,26]. Since intracellular alkalization activates eNOS in HUVECs [18], and other endothelium including human pulmonary aortic [19] and rat aortic [20] endothelial cells, intracellular acidification may result in reduced NO synthesis in HUVECs. Interestingly, a change in pH_i from 7.5 to ~6.5 resulted in a more significant reduction of NOS activity compared with a pH_i shift from 6.5 to 5.5 [18], thus complementing similar findings for system γ^+ L activity in HUVECs. Thus, a change in pH_i causing intracellular acidification is a phenomenon involved in downregulation of the system γ^+ L/eNOS activity in HUVECs. The possibility that intracellular acidification inhibited NOS was not related to L-arginine uptake is unlikely since incubation of cells with NEM + L-leucine abolished L-arginine uptake and NO synthesis. This proposal is supported by studies in rat astrocytes knockdown for system γ^+ /LAT-2 expression where system γ^+ L activity and NO generation was reduced [39].

In summary, intracellular acidification results in reduced membrane transport of L-arginine mediated via system γ^+ L but not via system γ^+ /CATs in HUVECs (see Figure 6). Diminished transport resulted from lower maximal transport capacity due to reduced V_{\max} without significant alterations in the apparent K_m for transport. Therefore, an acidic pH_i seems not to alter the intrinsic properties of system γ^+ L but the activity of membrane transporters in HUVECs. Interestingly, the pH_i sensitivity of L-arginine transport was higher as smaller the change in the pH_i from the basal pH_i in this cell type. Additionally, intracellular acidification also reduced the synthesis of NO and activator phosphorylation of eNOS, which seems to

result from reduced system γ^+L activity and lower activation of eNOS. Interestingly, preliminary results show that basal pHi is alkaline in HUVECs exposed to an A_{2A} adenosine receptors antagonist (L Sobrevia, *unpublished*), suggesting that basal pHi is potentially maintained by activation of this type of adenosine receptors in HUVECs. We hypothesise that changing the pHi into an acidic intracellular environment is a phenomenon likely involved in the lower adenosine-mediated relaxation of foetoplacental vasculature via reducing the endothelial system γ^+L /eNOS activity as seen in diseases of pregnancy such as preeclampsia [44,45], obesity [46,47], or gestational diabetes mellitus [21,27,28].

Conflict of interest

There is no conflict of interest.

Acknowledgements

Authors thank Mrs Amparo Pacheco from the Cellular and Molecular Physiology Laboratory (CMPL) at Division of Obstetrics and Gynaecology, Faculty of Medicine, Pontificia Universidad Católica de Chile, for excellent technical and secretarial assistance, and the personnel of the Hospital Clínico UC-CHRISTUS (Santiago de Chile) and Clínica de la Mujer (Antofagasta, Chile) labour ward for the supply of placentas. This work was supported by the Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) [grant number 1150377] and Semillero Dirección de Investigación, Universidad de Antofagasta [grant number 5309, 5313], Chile.

References

- [1] R. Devés, C.A. Boyd, Transporters for cationic amino acids in animal cells: discovery, structure, and function, *Physiol. Rev.* 78 (1998) 487–545.
- [2] G.E. Mann, D.L. Yudilevich, L. Sobrevia, Regulation of amino acid and glucose transporters in endothelial and smooth muscle cells, *Physiol. Rev.* 83 (2003) 183–252, <http://dx.doi.org/10.1152/physrev.00022.2002>.
- [3] F. Gaccioli, I.L.M.H. Aye, S. Roos, S. Lager, V.I. Ramirez, Y. Kanai, T.L. Powell, T. Jansson, Expression and functional characterisation of System L amino acid transporters in the human term placenta, *Reprod. Biol. Endocrinol.* 13 (2015) 57, <http://dx.doi.org/10.1186/s12958-015-0054-8>.
- [4] E. Guzmán-Gutiérrez, A. Armella, F. Toledo, F. Pardo, A Leiva, L. Sobrevia, Insulin requires A₁ adenosine receptors expression to reverse gestational diabetes-increased L-arginine transport in human umbilical vein endothelium. *Purinergic Signal.* 12 (2016) 175–190, <http://dx.doi.org/10.1007/s11302-015-9491-2>.
- [5] K. Barnes, H. Dobrzynski, S. Foppolo, P.R. Beal, F. Ismat, E.R. Scullion, S.A. Baldwin, Distribution and functional characterization of equilibrative nucleoside transporter-4, a novel cardiac adenosine transporter activated at acidic pH. *Circ. Res.* 99 (2006) 510–519, <http://dx.doi.org/10.1161/01.RES.0000238359.18495.42>.
- [6] N. Celis, J. Araos, C. Sanhueza, F. Toledo, A.R. Beltran, F. Pardo, M. Ramírez, L. Sobrevia, Intracellular acidification increases adenosine transport in human umbilical vein endothelial cells, *Placenta* 51 (2017) 10–17, <http://dx.doi.org/10.1016/j.placenta.2017.01.120>.
- [7] G. Wu, S.M. Morris. Arginine metabolism: nitric oxide and beyond, *Biochem. J.* 336 (1998) 1–17.

- [8] I. Fleming. Molecular mechanisms underlying the activation of eNOS, *Pflügers Arch.* 459 (2010) 793–806, <http://dx.doi.org/10.1007/s00424-009-0767-7>.
- [9] R. Sala, B.M. Rotoli, E. Colla, R. Visigalli, A. Parolari, O. Bussolati, G.C. Gazzola, V. Dall'Asta, Two-way arginine transport in human endothelial cells: TNF- α stimulation is restricted to system y^+ . *Am. J. Physiol. Cell. Physiol.* 282 (2002) C134–C143.
- [10] Y. Arancibia-Garavilla, F. Toledo, P. Casanello, L. Sobrevia, Nitric oxide synthesis requires activity of the cationic and neutral amino acid transport system y^+L in human umbilical vein endothelium, *Exp. Physiol.* 88 (2003) 699–710, <http://dx.doi.org/10.1113/eph8802647>.
- [11] S. Bröer, A. Bröer, Amino acid homeostasis and signalling in mammalian cells and organisms, *Biochem. J.* 474 (2017) 1935–1963, <http://dx.doi.org/10.1042/BCJ20160822>.
- [12] B.F. Haynes, M.E. Hemler, D.L. Mann, G.S. Eisenbarth, J. Shelhamer, H.S. Mostowski, C.A. Thomas, J.L. Strominger, A.S. Fauci, Characterization of a monoclonal antibody (4F2) that binds to human monocytes and to a subset of activated lymphocytes, *J. Immunol.* 126 (1981) 1409–1414.
- [13] D. Torrents, R. Estévez, M. Pineda, E. Fernández, J. Lloberas, Y.B. Shi, Identification and characterization of a membrane protein (y^+L amino acid transporter-1) that associates with 4F2hc to encode the amino acid transport activity y^+L . A candidate gene for lysinuric protein intolerance, *J. Biol. Chem.* 273 (1998) 32437–32445, <http://dx.doi.org/10.1074/jbc.273.49.32437>.
- [14] F. Verrey, E.I. Closs, C.A. Wagner, M. Palacín, H. Endou, Y. Kanai, CATs and HATs: The SLC7 family of amino acid transporters, *Pflügers Arch.* 447 (2004) 532–542,

- <http://dx.doi.org/10.1007/s00424-003-1086-z>.
- [15] M. González, S. Rojas, P. Avila, L. Cabrera, R. Villalobos, C. Palma, C. Aguayo, E. Peña, V. Gallardo, E. Guzmán-Gutiérrez, T. Sáez, R. Salsoso, C. Sanhueza, F. Pardo, A. Leiva, L. Sobrevia, Insulin reverses d-glucose–increased nitric oxide and reactive oxygen species generation in human umbilical vein endothelial cells. *PLoS One* 10 (2015) e0122398, <http://dx.doi.org/10.1371/journal.pone.0122398>.
- [16] M.G. Signorello, R. Pascale, G. Leoncini, Transport of L-arginine and nitric oxide formation in human platelets, *Eur. J. Biochem.* 270 (2003) 2005–2012, <http://dx.doi.org/10.1046/j.1432-1033.2003.03572.x>.
- [17] M. Kakoki, H.S. Kim, C.J. Edgell, N. Maeda, O. Smithies, D.L. Mattson, Amino acids as modulators of endothelium-derived nitric oxide, *Am. J. Physiol. Renal Physiol.* 291 (2006) F297–F304, <http://dx.doi.org/10.1152/ajprenal.00417.2005>.
- [18] I. Fleming, M. Hecker, R. Busse, Intracellular alkalinization induced by bradykinin sustains activation of the constitutive nitric oxide synthase in endothelial cells, *Circ. Res.* 74 (1994) 1220–1226, <http://dx.doi.org/10.1161/01.RES.74.6.1220>.
- [19] S. Mizuno, Y. Demura, S. Ameshima, S. Okamura, I. Miyamori, T. Ishizaki. Alkalosis stimulates endothelial nitric oxide synthase in cultured human pulmonary arterial endothelial cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 283 (2002) L113–L119, <http://dx.doi.org/10.1152/ajplung.00436.2001>.
- [20] V.K. Capellini, C.B.A. Restini, L.M. Bendhack, P.R.B. Evora, A.C. Celotto, The effect of extracellular pH changes on intracellular pH and nitric oxide concentration in endothelial and smooth muscle cells from rat aorta. *PLoS One* 8 (2013) e62887, <http://dx.doi.org/10.1371/journal.pone.0062887>.

- [21] L. Silva, M. Subiabre, J. Araos, T. Sáez, R. Salsoso, F. Pardo, A. Leiva, R. San Martín, F. Toledo, L. Sobrevia, Insulin/adenosine axis linked signalling, *Mol. Aspects Med.* 55 (2017) 45–61, <http://dx.doi.org/10.1016/j.mam.2016.11.002>.
- [22] R. San Martín, L. Sobrevia, Gestational diabetes and the adenosine/L-arginine/nitric oxide (ALANO) pathway in human umbilical vein endothelium, *Placenta* 27 (2006) 1–10, <http://dx.doi.org/10.1016/j.placenta.2005.01.011>.
- [23] A.A. Baschat, Venous Doppler evaluation of the growth-restricted fetus, *Clin. Perinatol.* 38 (2011) 103–112, vi. <http://dx.doi.org/10.1016/j.clp.2010.12.001>.
- [24] A. Najafzadeh, J.E. Dickinson, Umbilical venous blood flow and its measurement in the human fetus, *J. Clin. Ultrasound.* 40 (2012) 502–511, <http://dx.doi.org/10.1002/jcu.21970>.
- [25] J. Araos, L. Silva, R. Salsoso, T. Sáez, E. Barros, F. Toledo, J. Gutiérrez, F. Pardo, A. Leiva, C. Sanhueza, L. Sobrevia, Intracellular and extracellular pH dynamics in the human placenta from diabetes mellitus, *Placenta* 43 (2016) 47–53, <http://dx.doi.org/10.1016/j.placenta.2016.05.003>.
- [26] M. Subiabre, L. Silva, R. Villalobos-Labra, F. Toledo, M. Paublo, M.A. López, R. Salsoso, F. Pardo, A. Leiva, L. Sobrevia, Maternal insulin therapy does not restore foetoplacental endothelial dysfunction in gestational diabetes mellitus, *Biochim. Biophys. Acta – Mol. Basis Dis.* 1863 (2017) 2987–2998, <http://dx.doi.org/10.1016/j.bbadis.2017.07.022>.
- [27] T. Sáez, P. De Vos, L. Sobrevia, M. M. Faas, Is there a role for exosomes in foetoplacental endothelial dysfunction in gestational diabetes mellitus? *Placenta* 61 (2018) 48–54, <http://dx.doi.org/10.1016/j.placenta.2017.11.007>.

- [28] T. Sáez, R. Salsoso, A. Leiva, F. Toledo, P. De Vos, M.M. Faas, L. Sobrevia, Human umbilical vein endothelium-derived exosomes play a role in foetoplacental endothelial dysfunction in gestational diabetes mellitus, *Biochim. Biophys. Acta* 1864 (2018) 499–508, <http://dx.doi.org/10.1016/j.bbadis.2017.11.010>.
- [29] C. Sanhueza, J. Araos, L. Naranjo, F. Toledo, A.R. Beltrán, M.A. Ramírez, J. Gutiérrez, F. Pardo, A. Leiva, L. Sobrevia, Sodium/proton exchanger isoform 1 regulates intracellular pH and cell proliferation in human ovarian cancer, *Biochim. Biophys. Acta – Mol. Basis Dis.* 1863 (2017) 81–91, <http://dx.doi.org/10.1016/j.bbadis.2016.10.013>.
- [30] T. Tamagaki, S. Sawada, H. Imamura, Y. Tada, S. Yamasaki, A. Toratani, T. Sato, S. Komatsu, N. Akamatsu, M. Yamagami, K. Kobayashi, K. Kato, K. Yamamoto, K. Shirai, K. Yamada, T. Higaki, K. Nakagawa, H. Tsuji, M. Nakagawa, Effects of high-density lipoproteins on intracellular pH and proliferation of human vascular endothelial cells, *Atherosclerosis* 123 (1996) 73–82, [http://dx.doi.org/10.1016/0021-9150\(95\)05774-9](http://dx.doi.org/10.1016/0021-9150(95)05774-9).
- [31] V. Huck, A. Niemeyer, T. Goerge, E.M. Schnaeker, R. Ossig, P. Rogge, M.F. Schneider, H. Oberleithner, S.W. Schneide, Delay of acute intracellular pH recovery after acidosis decreases endothelial cell activation, *J. Cell. Physiol.* 211 (2007) 399–409, <http://dx.doi.org/10.1002/jcp.20947>.
- [32] E.R. Yeomans, J.C. Hauth, L.C. Gilstrap, D.M. Strickland, Umbilical cord pH, pCO₂, and bicarbonate following uncomplicated term vaginal deliveries. *Am. J. Obstet. Gynecol.* 151 (1985) 798–800, [http://dx.doi.org/10.1016/0002-9378\(85\)90523-X](http://dx.doi.org/10.1016/0002-9378(85)90523-X).
- [33] M. Pietryga, J. Brązert, E. Wender-Oegowska, R. Biczysko, M. Dubiel, S. Gudmundsson, Abnormal uterine Doppler is related to vasculopathy in pregestational

- diabetes mellitus, *Circulation* 112 (2005) 2496–2500, <http://dx.doi.org/10.1161/CIRCULATIONAHA.104.492843>.
- [34] F.J. Bernardez-Zapata, C. Moreno-Rey, Normal values of gases in the vein of the umbilical cord during the postpartum period and postcesarea immediately in normal fetuses to term, *Ginecol. Obstet. Mex.* 82 (2014) 170–176.
- [35] A.R. Beltrán, L.R. Carraro-Lacroix, C.N.A. Bezerra, M. Cornejo, K. Norambuena, F. Toledo, J. Araos, F. Pardo, A. Leiva, C. Sanhueza, G. Malnic, L. Sobrevia, M. Ramirez. *Escherichia coli* heat-stable enterotoxin mediates Na^+/H^+ exchanger 4 inhibition involving cAMP in T₈₄ human intestinal epithelial cells. *PLoS One* 10 (2015) e0146042, <http://dx.doi.org/10.1371/journal.pone.0146042>.
- [36] R. San Martín, L. Sobrevia, Gestational diabetes and the adenosine/L-arginine/nitric oxide (ALANO) pathway in human umbilical vein endothelium, *Placenta* 27 (2006) 1–10, <http://dx.doi.org/10.1016/j.placenta.2005.01.011>.
- [37] E.I. Closs, A. Simon, N. Vékony, A. Rotmann, Plasma membrane transporters for arginine, *J. Nutr.* 134 (2004) 2752S–2759S.
- [38] D. Fotiadis, Y. Kanai, M. Palacín, The SLC3 and SLC7 families of amino acid transporters, *Mol. Aspects Med.* 34 (2013) 139–158, <http://dx.doi.org/10.1016/j.mam.2012.10.007>.
- [39] Zielinska 2015 M. Zielińska, K. Milewski, M. Skowrońska, A. Gajos, E. Ziemińska, A. Beręsewicz, J. Albrecht, Induction of inducible nitric oxide synthase expression in ammonia-exposed cultured astrocytes is coupled to increased arginine transport by upregulated $\gamma^+/\text{LAT2}$ transporter, *J. Neurochem.* 135 (2015) 1272–1281, <http://dx.doi.org/10.1111/jnc.13387>.

- [40] E.I. Closs, J.S. Scheld, M. Sharafi, U. Förstermann, Substrate supply for nitric-oxide synthase in macrophages and endothelial cells: role of cationic amino acid transporters, *Mol. Pharmacol.* 57 (2000) 68–74.
- [41] P. Casanello, L. Sobrevia, Intrauterine growth retardation is associated with reduced activity and expression of the cationic amino acid transport systems γ^+ /hCAT-1 and γ^+ /hCAT-2B and lower activity of nitric oxide synthase in human umbilical vein endothelial cells, *Circ. Res.* 91 (2002) 127–134, <http://dx.doi.org/10.1161/01.RES.0000027813.55750.E7>.
- [42] A. Simon, L. Plies, A. Habermeier, U. Martiné, M. Reining, E.I. Closs, Role of neutral amino acid transport and protein breakdown for substrate supply of nitric oxide synthase in human endothelial cells, *Circ. Res.* 93 (2003) 813–820, <http://dx.doi.org/10.1161/01.RES.0000097761.19223.0D>.
- [43] R. Salsoso, E. Guzmán-Gutiérrez, T. Sáez, K. Bugueño, M.A. Ramírez, M. Farías, F. Pardo, A. Leiva, C. Sanhueza, A. Mate, C. Vázquez, L. Sobrevia, Insulin restores L-arginine transport requiring adenosine receptors activation in umbilical vein endothelium from late-onset preeclampsia, *Placenta* 36 (2015) 287–296, <http://dx.doi.org/10.1016/j.placenta.2014.12.007>.
- [44] R. Salsoso, M. Farías, J. Gutiérrez, F. Pardo, D.I. Chiarello, F. Toledo, A. Leiva, A. Mate, C.M. Vázquez, L. Sobrevia, Adenosine and preeclampsia, *Mol. Aspects Med.* 55 (2017) 126–139, <http://dx.doi.org/10.1016/j.mam.2016.12.003>.
- [45] D.I. Chiarello, R. Salsoso, F. Toledo, A. Mate, C.M. Vázquez, L. Sobrevia, Foetoplacental communication via extracellular vesicles in normal pregnancy and preeclampsia, *Mol. Aspects Med.* 2018 (In Press), <http://dx.doi.org/10.1016/j.mam.2017.12.002>.

- [46] F. Pardo, R. Villalobos-Labra, D.I. Chiarello, R. Salsoso, F. Toledo, J. Gutiérrez, A. Leiva, L. Sobrevia, Molecular implications of adenosine in obesity, *Mol. Aspects Med.* 55 (2017) 90–101, <http://dx.doi.org/10.1016/j.mam.2017.01.003>.
- [47] F. Pardo, R. Villalobos-Labra, B. Sobrevia, F. Toledo, L. Sobrevia, Extracellular vesicles in obesity and diabetes mellitus, *Mol. Aspects Med.* 2018 (In Press), <http://dx.doi.org/10.1016/j.mam.2017.11.010>.

Figure 1. Effect of cell pHi on L-arginine uptake in HUVECs. **A.** Cells were preloaded with BCECF-AM and transferred into a spectrofluorometer. After basal pHi stabilisation, the cells were exposed (2 min) to a control solution containing 20 mmol/L NH₄Cl (+NH₄Cl). Cells were then rinsed with a NH₄Cl-free solution (–NH₄Cl) and left in this medium for pHi recovery (see Materials and methods). A typical record is shown. The *insert* indicates the data for the first 55 s after removal of NH₄Cl. The grey area indicates the time (10 s) used for overall uptake of 2 or 100 μmol/L L-arginine (6 μCi/mL L-[³H]arginine, 37°C). **B.** pHi values for cells exposed to a Na⁺-free solution without (0) or with NH₄Cl. **C.** Overall 100 μmol/L L-arginine uptake in +NH₄Cl or –NH₄Cl solution in the absence (–) or presence (+) of *N*-ethylmaleimide (NEM) or L-leucine. **D.** L-Arginine uptake mediated via system y⁺/CATs derived from data in C. **E.** Overall 2 μmol/L L-arginine uptake as in C. **F.** L-Arginine uptake mediated via system y⁺L derived from data in E. In B, **P*<0.05 versus without or with 0.1 mmol/L NH₄Cl. In C, **P*<0.05 versus all other values, †*P*<0.05 versus corresponding values in the presence of NEM or NEM + L-leucine. ‡*P*<0.05 versus values in –NH₄Cl in the presence of NEM. In D, **P*<0.03 versus corresponding values in 2 μmol/L L-arginine. In E, **P*<0.03 versus all other values except for –NH₄Cl in the presence of NEM. In F, **P*<0.03 versus corresponding values in +NH₄Cl. Values are mean ± S.E.M. (*n* = 18).

Figure 2. Effect of pHi on L-arginine uptake in HUVECs. **A.** L-Arginine (100 $\mu\text{mol/L}$) uptake (6 $\mu\text{Ci/mL}$ L-[^3H]arginine, 10 s, 37°C) in primary cultures of HUVECs non-treated (0) or treated with increasing concentrations of NH_4Cl as described in Materials and methods. Cells were in the absence (Control) or presence of 200 $\mu\text{mol/L}$ *N*-ethylmaleimide (NEM). **B.** L-Arginine (2 $\mu\text{mol/L}$) uptake as in **A** in the presence of 200 $\mu\text{mol/L}$ NEM (i.e., Control for this concentration of L-arginine) or NEM plus 2 mmol/L L-leucine (NEM + L-leucine). **C.** L-Arginine uptake via system y^+/CATs and system $y^+\text{L}$ derived from data in **A** and **B**, respectively, in the absence or presence of NH_4Cl . **D.** L-Arginine uptake against pHi values in cells as in **C**. **E.** Efficiency of a change in the pHi (ΔpHi) on the uptake of L-arginine ($^U E_{\text{pHi}}$) via system y^+/CAT and system $y^+\text{L}$ from data in **D**. Values for $^U E_{\text{pHi}}$ are expressed as a change in the $\text{pmol}/\mu\text{g}$ protein/minute relative to 1 (see Materials and methods). **F.** Relative $^U E_{\text{pHi}}$ for system y^+/CAT and system $y^+\text{L}$ from data in **E**. In **A**, **B**, and **C**, $*P < 0.05$ versus corresponding values without NH_4Cl . In **D**, $*P < 0.05$ versus corresponding values at pHi 7.13 and pHi 7.19. In **E**, $*P < 0.03$ versus all other corresponding values, $\dagger P < 0.05$ versus corresponding value at $\Delta\text{pHi} = 0.69$ pHi units. In **F**, $*P < 0.03$ versus all other values, $\dagger P < 0.05$ versus values at pHi = 6.5. Values are mean \pm S.E.M. ($n = 19$).

Figure 3. Effect of NH₄Cl on L-arginine saturable transport in HUVECs. **A.** Saturable L-arginine transport (6 μ Ci/mL L-[³H]arginine, 10 s, 37°C) was measured in primary cultures of HUVECs non-treated (Control, i.e., pHi 7.19) or treated with 0.1, 1, or 20 mmol/L NH₄Cl reaching pHi 7.13, 6.81, or 6.5, respectively (see Materials and methods). Cells at pHi 7.19 were in the absence or presence of 200 μ mol/L N-ethylmaleimide (NEM) or NEM plus 2 mmol/L L-leucine (NEM + L-leucine). Cells in pHi 7.13, 6.81, or 6.5 were exposed to NEM. **B.** Eadie-Hofstee plots for transport data in cells in the presence of NEM as in A. **C.** Maximal transport capacity (V_{max}/K_m) for systems $y^+/CATs$ and system y^+L from data in A (see also Table 2). **D.** The efficiency of a change in the pHi (ΔpHi) on maximal transport capacity for L-arginine ($^{Vmax/Km}E_{pHi}$) via system $y^+/CATs$ and system y^+L from data in A. Values for $^{Vmax/Km}E_{pHi}$ are expressed as a change in the pmol/ μ g protein/minute relative to 1 (see Materials and methods). **E.** Relative $^{Vmax/Km}E_{pHi}$ for system y^+/CAT and system y^+L from data in D. In C, * $P < 0.05$ versus corresponding values at pHi 7.13 and 7.19. In D, * $P < 0.05$ versus all other values. Values are mean \pm S.E.M. ($n = 19$).

Figure 4. pHi dependency of NOS activity in HUVECs. **A.** Intracellular L-citrulline level was determined by H.P.L.C. in primary cultures of HUVECs not treated ($-\text{NH}_4\text{Cl}$) or treated ($+\text{NH}_4\text{Cl}$) with 20 mmol/L NH_4Cl in the absence or presence of 100 $\mu\text{mol/L}$ N^G -nitro-L-arginine methyl ester (L-NAME) as described in Materials and Methods. Assays were in cells in 100 $\mu\text{mol/L}$ L-arginine in the absence or presence of 200 $\mu\text{mol/L}$ *N*-ethylmaleimide (NEM) or NEM plus 2 mmol/L L-leucine (NEM + L-leucine). **B.** The intracellular L-citrulline level in HUVECs in 2 $\mu\text{mol/L}$ L-arginine as in A. **C.** Nitric oxide synthase (NOS) activity-dependent L-citrulline synthesis derived from data in A. **D.** Nitric oxide synthase (NOS) activity-dependent L-citrulline synthesis derived from data in B. In A, * $P < 0.05$ versus all other values, † $P < 0.05$ versus corresponding values except in the presence of L-leucine. ‡ $P < 0.05$ versus all other corresponding values. All values in the presence of L-NAME are significantly different ($P < 0.03$) from values in the absence of this inhibitor. In B, * $P < 0.03$ versus all other values. In C, * $P < 0.05$ versus all other values, † $P < 0.05$ versus all other corresponding values except in the presence of L-leucine, ‡ $P < 0.05$ versus all other corresponding values except in the presence of NEM + L-leucine, § $P < 0.03$ versus corresponding values in the presence of L-leucine. In D, * $P < 0.05$ versus all other values except for $-\text{NH}_4\text{Cl}$ in the presence of NEM, † $P < 0.05$ versus all other corresponding values, ‡ $P < 0.05$ versus corresponding values in $-\text{NH}_4\text{Cl}$. Values are mean \pm S.E.M. ($n = 19$).

Figure 5. pHi dependency of eNOS expression and phosphorylation in HUVECs. A.

Western blot for total (Total eNOS) or phosphorylated at Serine¹¹⁷⁷ (P-Ser¹¹⁷⁷eNOS) or Threonine⁴⁹⁵ (P-Thr⁴⁹⁵eNOS) eNOS protein in primary cultures of HUVECs non-treated (Control, i.e., pHi = 7.19) or treated with 0.1, 1, or 20 mmol/L NH₄Cl reaching pHi 7.13, 6.81, or 6.5, respectively (see Materials and methods). β -Actin is the loading control. Total eNOS/ β -actin (**B**), P-Ser¹¹⁷⁷eNOS/Total eNOS (**C**) or P-Thr⁴⁹⁵eNOS/Total eNOS (**C**) protein ratios from cells as in A. * $P < 0.05$ versus values at pHi 7.19 and 7.13. Values are mean \pm S.E.M. ($n = 19$).

Figure 6. Modulation of L-arginine/NO signalling pathway by intracellular pH in the human foetoplacental endothelium. The physiological intracellular pH (pHi) value in human umbilical vein endothelial cells (HUVECs) from normal pregnancies (pHi 7.2) maintain the L-arginine transport via the very high affinity transport system y^+L with a maximal transport capacity (V_{max}/K_m) of ~ 0.2 pmol/ μ g protein/minute/ $(\mu$ mol/L). L-Arginine is metabolised by the endothelial nitric oxide synthase (eNOS) into L-citrulline and nitric oxide (NO). When pHi value is acidic (pHi 6.5), the V_{max}/K_m is reduced (\Downarrow) to ~ 0.02 pmol/ μ g protein/minute/ $(\mu$ mol/L) resulting in lower uptake of L-arginine (dotted lines). The reduced uptake in L-arginine transport and its subsequent lower bioavailability to eNOS leads to minor NO generation likely due to lower activator phosphorylation at serine 1177 (Ser¹¹⁷⁷) residue at eNOS.

Table 1. Clinical variables in pregnant women and newborns.

<i>Mother</i>	
Age (years)	31.4 ± 3.9 (26.2 – 37.5)
Height (cm)	161.7 ± 3.9 (154.1 – 163.2)
Weight (kg)	
9-16 weeks of gestation	56.2 ± 3.4 (54.6 – 59.5)
Delivery	68.1 ± 3.9 (64.2 – 69.1) *
BMI (kg/m ²)	
9-16 weeks of gestation	21.5 ± 1.7 (20.1 – 22.3)
Delivery	26.1 ± 0.9 (25.1 – 29.3) *
Mean arterial pressure (mm Hg)	
9-16 weeks of gestation	77.1 ± 3.5 (75.5 – 79.2)
Delivery	81.7 ± 6.1 (79.8 – 89.2)
Glycemia fasting (mg/dL)	83.1 ± 6.9 (74.9 – 89.5)
OGTT (mg/dL)	
Glycemia basal	82.7 ± 3.2 (78.2 – 85.3)
Glycemia 2 hours after glucose	83.3 ± 4.1 (82.1 – 89.0)
<i>Newborn</i>	
Sex (female/male)	13/10
Gestational age (weeks)	38.3 ± 0.7 (38.0 – 38.9)
Birth weight (grams)	3182 ± 210 (3051 – 3346)
Height (cm)	49.7 ± 1.2 (48.1 – 52.2)
Ponderal index (grams/cm ³ x 100)	2.59 ± 0.12 (2.12 – 2.63)

Legend for Table 1 in the next page

Legend for Table 1

Women that coursed with normal pregnancies ($n = 23$) were included in this study. Weight, body mass index (BMI), and blood pressure were determined at the first interview with the obstetrician (9-16 weeks of pregnancy) and at delivery. BMI was calculated by weight in kilograms divided by the square of the height in meters. Ponderal index was calculated by weight in grams divided by the cube of height in centimeters multiplied by 100. Oral glucose tolerance test (OGTT) was measured at the 1st trimester of pregnancy in all women with a normal glycaemia (see Materials and methods). * $P < 0.05$ versus corresponding values at 9-16 weeks of gestation. Values are mean \pm S.D. plus range in brackets.

Table 2. Effect of intracellular pH on the kinetic parameters for L-arginine transport in HUVECs

	<i>Saturable transport</i>			<i>Overall transport</i>	
	V_{max} (pmol/ μ g protein/ minute)	K_m (μ mol/L)	V_{max}/K_m (pmol/ μ g protein/ minute/(μ mol/L))	K_D (pmol/ μ g protein/ minute/(μ mol/L))	v_i (pmol/ μ g protein/ 0.5 seconds)
System $y^+/CATs$					
Without NH_4Cl	0.60 \pm	79 \pm 49	0.008 \pm 0.004	0.0025 \pm 0.0002	0.00279 \pm
pHi 7.19	0.18				0.00022
With NH_4Cl (mmol/L)					
0.1 pHi	0.61 \pm	97 \pm 29		0.0022 \pm 0.0002	0.00258 \pm
7.13	0.11		0.006 \pm 0.002		0.00022
1 pHi	0.58 \pm	73 \pm 31		0.0022 \pm 0.0002	0.00279 \pm
6.89	0.12		0.008 \pm 0.003		0.00018
20 pHi	0.53 \pm	84 \pm 32		0.0023 \pm 0.0003	0.00240 \pm
6.50	0.08		0.006 \pm 0.002		0.00021
System y^+L					
Without NH_4Cl	0.38 \pm	1.98 \pm	0.192 \pm 0.079	0.0492 \pm 0.0048	0.00159 \pm
pHi 7.19	0.11	1.06			0.00019
With NH_4Cl (mmol/L)					
0.1 pHi	0.33 \pm	2.01 \pm		0.0511 \pm 0.0041	0.00137 \pm
7.13	0.08	0.91	0.164 \pm 0.057		0.00019
1 pHi	0.21 \pm	2.12 \pm		0.0488 \pm 0.0052	0.00084 \pm
6.89	0.09 *	0.99	0.099 \pm 0.044 *		0.00011 *
20 pHi	0.04 \pm	1.99 \pm		0.0479 \pm 0.0048	0.00017 \pm
6.50	0.02 *†	0.12	0.020 \pm 0.003 *†		0.00008 *†

Legend for Table 2 in the next page.

Legend for Table 2

Transport of L-arginine (20 s, 37°C) was measured in HUVECs from normal pregnancies. Transport assays were done in cells not treated (*Without NH₄Cl*) or treated (*With NH₄Cl*) in a NH₄Cl-acid pulse as described in Materials and methods. The resulting intracellular pH (pHi) values are indicated. Maximal velocity (V_{\max}) and apparent Michaelis-Menten constant (K_m) of saturable transport in the range of 0-20 $\mu\text{mol/L}$ (for system y^+L) or 0-1000 (for system $y^+/CATs$) L-arginine were calculated assuming a single Michaelis-Menten hyperbola. V_{\max}/K_m represents maximal L-arginine transport capacity. The lineal phase of overall transport of L-arginine (K_D) was obtained from transport data fitted to a Michaelis-Menten equation increased in a lineal component. Initial velocity (v_i) was calculated for 0.5 s with 100 or 2 $\mu\text{mol/L}$ L-arginine transport. All values for V_{\max} , K_m , and v_i for system y^+L are lower ($P<0.05$) and values for V_{\max}/K_m and K_D were higher ($P<0.05$) than corresponding values for system $y^+/CATs$. * $P<0.05$ versus corresponding values in cells without or with 0.1 mmol/L NH₄Cl. † $P<0.05$ versus corresponding values in cells without or with 0.1 mmol/L NH₄Cl. *Without insulin*. ‡ $P<0.05$ versus corresponding values in cells with 1 mmol/L NH₄Cl. Values are mean \pm S.E.M. ($n = 19$).

Highlights

- HUVECs show a basal intracellular alkaline pH (pHi ~7.2).
- Intracellular acidification inhibits L-arginine transport via system y⁺L, but not the system y⁺/CATs activity.
- Nitric oxide synthesis (NO) is reduced by intracellular acidification.
- Reduced system y⁺L activity associated with lower endothelial NO synthase (eNOS) activation.
- Intracellular pH regulates system y⁺L/eNOS signalling in HUVECs.









Figure 1









Figure 2






Figure 3


A



B



C



D




Figure 4







Figure 5




Figure 6