

Evaluation of labile iron formation in the atmosphere

*A. Ito¹, S. Myriokefalitakis², M. Kanakidou³, N. Mahowald⁴, R. A. Scanza⁴, A. Baker⁵, T. Jickells⁵, M. Sarin⁶, S. Bikkina⁷, Y. Gao⁸, R. Shelley⁹, C. Buck¹⁰, W. Landing⁹, A. Bowie¹¹, M. Perron¹¹, N. Meskhidze¹², M. Johnson¹³, Y. Feng¹⁴, R. Duce¹⁵ (¹JAMSTEC, ²Utrecht Univ., ³Univ. of Crete, ⁴Cornell Univ., ⁵Univ. of East Anglia, ⁶Physical Research Laboratory, ⁷Stockholm Univ., ⁸Rutgers Univ., ⁹Florida State Univ., ¹⁰Univ. of Georgia, ¹¹Univ. of Tasmania, ¹²North Carolina State Univ., ¹³NASA Ames Research Center, ¹⁴Argonne National Laboratory, ¹⁵Texas AM Univ.)

Atmospheric deposition of labile iron (Fe) to the ocean has been suggested to modulate primary ocean productivity and thus indirectly affect the climate. A key process contributing to atmospheric sources of labile Fe is associated with atmospheric acidity, which leads to Fe transformation from insoluble to soluble forms. Significant progress has been made in our understanding of atmospheric inputs of labile Fe from natural and anthropogenic sources to the oceans. However, there are still large uncertainties regarding the relative importance of different sources of Fe and effects of atmospheric processing on the bioavailability of the delivered Fe.

Here, we investigate the effects of atmospheric processing on Fe solubility and contribution of different sources of Fe to labile Fe in the atmosphere. We compiled Fe loading and solubility in aerosols from four atmospheric chemistry transport models and a number of field measurements. Fe-containing aerosols from combustion sources are characterized by low loading and high solubility, compared to mineral dust. Therefore, labile Fe loading may be separately attributed to combustion and dust aerosols, assuming their distinct emission sources and atmospheric processes. The results suggest that combustion aerosols substantially contribute to labile Fe loading measured in the atmosphere. Thus, assessments of dust-borne Fe fertilization of the oceans should include Fe-containing mineral aerosols affected by combustion sources.