

## **Laser Raman and infrared spectra of some uracil derivatives\***

V K Rastogi, H P Mital<sup>†</sup> and S N Sharma<sup>††</sup>

Department of Physics, Lajpat Rai College, Sahibabad-201 005, India

Received 6 November 1989, accepted 25 April 1990

The N-heterocyclic molecules e.g. pyrimidine, cytosine, uracil and their derivatives are of considerable biological and pharmaceutical importance (Handschumacher and Welch 1960, Martin 1961). Recent spectroscopic studies of these compounds have been motivated because the vibrational spectra of free base molecules is very useful for the understanding of specific biological processes and in the analysis of relatively complex systems. Along the macromolecular double-helix chain, the vibrational modes of each base interact with those of other bases through hydrogen bonds or stacking effects and these interactions affect the ring vibrations. The vibrational spectra of various nucleic acid constituents including uracil have been studied in detail (Angel 1961, Lord and Thomas 1967, Susi and Ard 1971, Susi *et al* 1973, Bandekar and Zundel 1983, Singh *et al* 1987), but very little work on the vibrational spectra of derivatives of uracil has been done (Shimanouchi *et al* 1964, Sanyal *et al* 1977, Ferro *et al* 1980, Goel *et al* 1981, Aruna and Shanmugam 1985, Yadav *et al* 1988). In order to extend this study further, we have recorded the IR and laser Raman spectra of 5-chloro and 5-amino uracils for the first time and assigned the observed fundamental bands.

Pure solid samples of 5-chlorouracil (mp > 300 °C) and 5-aminouracil (mp > 300 °C) were obtained from M/S Sigma Chemical Co, USA and used as such without further purification. The IR absorption spectra have been recorded on Beckman IR-12 spectrophotometer in KBr pellet in the region 400-4000 cm<sup>-1</sup>. The laser Raman spectra in the region 50-2000 cm<sup>-1</sup> (Raman spectra beyond 2000 cm<sup>-1</sup> are not worth-mentioning) have been recorded on a Spex spectrophotometer using Argon laser source with exciting radiations 5145 Å.

<sup>†</sup>Department of Physics, Meerut College, Meerut-250 001, India.

<sup>††</sup>Indian Institute of Petroleum, Dehradun-248 005.

\*Presented at International Conference on Raman Spectroscopy, IACS, Calcutta, Nov. 2-7, 1988.

Figures 1-3 show the structures of the uracil and 5-chloro and 5-amino uracils molecules with classical numbering of the atoms respectively. The vibrational spectra (IR and Raman) have been interpreted assuming planar geometry and  $C_2$  point group symmetry for both the molecules and the observed fundamentals have been assigned to different normal modes of vibration. The vibrational modes are classified as transforming like the basis functions of the  $a'$  and  $a''$  irreducible representations. Thus the 30 fundamental modes of 5-chlorouracil may be classified into  $21a'$  symmetry species (in-plane) and  $9a''$  species (out-of-plane) and in




Figure 1. Uracil.




Figure 2. 5-Chlorouracil.




Figure 3. 5-Aminouracil.

the case of 5-aminouracil, the 36 fundamental modes may be classified into  $25a'$  and  $11a''$  symmetry species. The analysis of the bands and the assignment of the fundamental frequencies are made on the basis of magnitude and relative intensities of the observed bands and help has been taken from the assignments made by earlier workers in the similar molecules. The wavenumbers of the observed fundamental bands in IR and Raman spectra alongwith the proposed assignments are given in Table 1.

The frequencies due to CH/NH stretching modes are well separated from the frequencies due to other modes. The molecules under study contain only one hydrogen atom attached to the carbon atom of the ring (Figures 2 and 3) and thus involve only one C–H valence oscillation which have been identified at  $3032$  and  $3018\text{ cm}^{-1}$  in the IR spectra of the 5-chloro and 5-amino uracils respectively. These assignments are in agreement with those proposed by Ferro *et al* (1980) and Aruna and Shanmugam (1985) in the case of related molecules. The bands observed at  $1422\text{ cm}^{-1}$  (IR) and  $1410\text{ cm}^{-1}$  (Raman) in the case of 5-chlorouracil and at  $1425\text{ cm}^{-1}$  in 5-aminouracil have been assigned to C–H in-plane bending mode.

The N–H stretching was assigned by Susi and Ard (1971) at  $3145\text{ cm}^{-1}$  in 1-methyl uracil and at  $3146\text{ cm}^{-1}$  in 1-methyl thymine and by Sanyal *et al* (1977) at  $3178$ ,  $3098\text{ cm}^{-1}$  in 5-bromouracil, at  $3188$ ,  $3128\text{ cm}^{-1}$  in 5-methyl uracil and at  $3178$ ,  $3107\text{ cm}^{-1}$  in 4(6)-aminouracil. In the present case only one of the two N–H stretching modes could be observed at  $3059$  and  $3080\text{ cm}^{-1}$  in the IR spectra of 5-chloro and 5-amino uracils respectively.

The  $\text{NH}_2$  group has two (N-H) stretching frequencies, one being symmetric and other asymmetric. The frequency of asymmetric vibration is higher than that of

Table I. Assignment of fundamental frequencies (in  $\text{cm}^{-1}$ ) of 5-chloro and 5-amino uracils.

5-Chloro-uracil		5-Amino-uracil		Symmetry species	Assignment
IR	Raman	IR	Raman		
—	—	3370	—	$a'$	$\text{NH}_2$ asymmetric stretch
—	—	3295	—	$a'$	$\text{NH}_2$ symmetric stretch
3059	—	3080	—	$a'$	N-H stretch
3032	—	3018	—	$a'$	C-H stretch
1705	—	1740	—	$a'$	C=O stretch
1677	1676	1672	—	$a'$	C-O stretch
1632	1625	1615	1645	$a'$	C-C stretch
—	—	1588	—	$a'$	$\text{NH}_2$ scissor
1560	—	1556	—	$a'$	N-H in-plane bending
1542	—	1541	—	$a'$	N-H in-plane bending
1509	—	1506	1493	$a'$	C-N stretch
1456	1473	1462	—	$a'$	C-N stretch
1440	—	—	—	$a'$	ring stretch
1422	1410	1425	—	$a'$	C-H in-plane bending
—	—	1387	1400	$a'$	$\text{NH}_2$ rock
1340	1339	—	1334	$a'$	ring stretch
—	—	1276	—	$a'$	C-NH <sub>2</sub> stretch
—	—	1245	1251	$a'$	$\text{NH}_2$ rock
1230	—	1212	1215	$a'$	ring stretch
—	—	1108	—	$a'$	$\text{NH}_2$ twisting
1090	—	1077	—	$a'$	ring stretch
1008	—	1016	—	$a'$	ring in-plane bending
—	—	880	—	$a'$	C-N in-plane bending
781	784	—	—	$a'$	C-Cl stretch
742	734	—	—	$a'$	C=O in-plane bending
680	668	—	—	$a'$	ring in-plane bending
—	639	—	—	$a'$	C-O in-plane bending
610	618	—	—	$a'$	C-Cl in-plane bending
542	548	552	—	$a'$	ring in-plane bending
946	—	—	—	$a''$	N-H out-of-plane bending
865	—	863	—	$a''$	N-H out-of-plane bending
815	—	805	—	$a''$	C-H out-of-plane bending
756	—	767	—	$a''$	ring out-of-plane bending
—	—	749	—	$a''$	$\text{NH}_2$ wagging
—	—	488	—	$a''$	$\text{NH}_2$ torsion
445	—	438	—	$a''$	C=O out-of-plane bending
428	—	—	—	$a''$	ring out-of-plane bending

Table I. (Contd.)

5-Chloro-uracil		5-Amino-uracil		Symmetry species	Assignment
IR	Raman	IR	Raman		
412	420	418	—	$a''$	C=O out-of-plane bending
234	—	—	—	$a''$	C-N out-of-plane bending
—	141	—	134		lattice
—	125	—	—		lattice
—	101	—	100		lattice
—	90	—	—		lattice
—	78	—	—		lattice

symmetric one. The  $\text{NH}_2$  asymmetric stretch modes have been identified at 3370 and 3295  $\text{cm}^{-1}$  respectively, in the IR spectrum of 5-aminouracil in conformation with the assignments proposed by Aruna and Shanmugam (1985) for 6-amino-2-thiouracil.

The neutral form of uracil and its substituted derivatives have three double bonds and their stretching vibration are expected to give rise to three strong infrared bands in the region 1500-1700  $\text{cm}^{-1}$ . Out of the three, two strong bands at 1705 and 1677  $\text{cm}^{-1}$  in the IR spectrum of 5-chlorouracil and at 1740 and 1672  $\text{cm}^{-1}$  in the IR spectrum of 5-aminouracil are assigned to the two C=O stretching fundamentals. No such vibration was observed in the Raman spectrum of 5-aminouracil, whereas one of these modes could be observed at 1676  $\text{cm}^{-1}$  in the Raman spectrum of 5-chlorouracil. The third vibration at 1632 and 1625  $\text{cm}^{-1}$  in the IR and Raman spectra, respectively, of 5-chlorouracil and at 1615 and 1645  $\text{cm}^{-1}$  in the IR and Raman spectra respectively of 5-aminouracil have been associated with C=C stretching vibration. These assignments are in agreement with those available for uracil, (Susi and Ard 1971), and mono and disubstituted uracil (Sanyal et al 1977). In the IR spectra of 5-chlorouracil, the 1677  $\text{cm}^{-1}$  band appears to be a closely spaced doublet. This splitting may be due to crystal field effect as suggested by Sanyal et al (1977). The space group of the 5-chlorouracil crystal is  $P2_1/c$ , with  $a=8.450$ ,  $b=6.886$ ,  $c=11.419$  Å and  $\beta=123.93^\circ$ .

The  $\text{NH}_2$  group has scissoring, rocking, twisting and wagging modes of bending vibrations. The bands observed at 1588, 1387 and 1108 and 749  $\text{cm}^{-1}$  in IR spectrum of 5-aminouracil have been assigned as scissoring, rocking, twisting and wagging modes respectively. The band observed at 1400  $\text{cm}^{-1}$  in Raman spectrum of 5-aminouracil has been assigned to  $\text{NH}_2$  rocking mode. These assignments find support from the works of Aruna and Shanmugam (1985).

The N-H in-plane bending vibrations have been identified at 1508 and 1417  $\text{cm}^{-1}$  in uracil (Shimanouchi et al 1964), at 1590, 1525  $\text{cm}^{-1}$  in 6-amino-

uracil and at 1550, 1510  $\text{cm}^{-1}$  in 6-amino-2-thiouracil (Aruna and Shanmugam 1985). In consonance with these assignments the bands observed at 1560, 1542  $\text{cm}^{-1}$  in the IR spectrum of 5-chlorouracil and at 1556, 1541  $\text{cm}^{-1}$  in the IR spectrum of 5-aminouracil have been identified as N-H in-plane bending vibration. The N-H out-of-plane bending vibrations have been identified at 865 and 946  $\text{cm}^{-1}$  in the IR spectrum of 5-chlorouracil and at 863  $\text{cm}^{-1}$  in the IR spectrum of 5-aminouracil.

The C=O in-plane bending vibration could only be observed in 5-chlorouracil at 742  $\text{cm}^{-1}$  in the IR spectrum and at 734 and 639  $\text{cm}^{-1}$  in the Raman spectrum. The C=O out-of-plane bending modes could be identified at 445 and 412  $\text{cm}^{-1}$  in the IR spectrum and at 420  $\text{cm}^{-1}$  in the Raman spectrum of 5-chlorouracil and at 438 and 418  $\text{cm}^{-1}$  in the IR spectrum of 5-aminouracil.

The ring out-of-plane bending modes have been observed at 756 and 428  $\text{cm}^{-1}$  in 5-chlorouracil and at 767  $\text{cm}^{-1}$  in 5-aminouracil.

Below 200  $\text{cm}^{-1}$  five lines in the Raman spectrum of 5-chlorouracil and two lines in the Raman spectrum of 5-aminouracil have been observed, which are in a region where lattice modes associated with C=O...H-N hydrogen bonds are expected.

### Acknowledgments

The authors are thankful to Prof P S Naraynan, IISc, Bangalore for providing necessary facilities for recording laser Raman spectra and to UGC, New Delhi for providing financial assistance.

### References

- Angel C J L 1961 *J. Chem. Soc.* 504  
Aruna S and Shanmugam G 1985 *Spectrochim. Acta* **38A** 815  
Bandekar J and Zundel G 1982 *Spectrochim. Acta* **38A** 815  
Bandekar J and Zundel G 1983 *Spectrochim. Acta* **39A** 343  
Ferro D, Bencivenni L, Teghil R and Pelino M 1980 *J. Indian Chem. Soc.* **57** 626  
Goel R K, Gupta S K, Agarwal M L and Sharma S N 1981 *Indian J. Pure Appl. Phys.* **19** 501  
Handschumacher R E and Welch A D 1960 *The Nucleic Acids* Vol 4 (New York: Academic Press) 453  
Lord R C and Thomas G J (Jr) 1967 *Spectrochim. Acta* **23A** 2551  
Martin G T 1961 *Biological Antagonism* (New York: Blakiston)  
Rastogi V K, Mital H P and Sharma S N 1988 *Proc. International Conference on Raman Spectroscopy* and Sons London (New York: John Wiley) p 87  
Sanyal N K, Srivastava S L and Goel R K 1977 *Indian J. Phys.* **52B** 108  
Shimanouchi T, Tsuboi M and Kyogobu 1964 *Adv. Chem. Phys.* (New York: Interscience) **7** p 435  
Singh B, Khan Sawar Akram and Uday Bhanu 1987 *Indian J. Chem.* **26A** 1066  
Susi H and Ard J S 1971 *Spectrochim. Acta* **27A** 1549  
Susi H, Ard J S and Purcell J M 1973 *Spectrochim. Acta* **29A** 725  
Yadav R A, Yadav P N S and Yadav J S 1988 *Proc. Indian Acad. Sci.* **100** 69