

# BMJ Open

## Effectiveness of service models and organisational structures supporting tuberculosis identification and management in hard-to-reach populations in countries of low and medium tuberculosis incidence: A systematic review

|                                 |  |
|---------------------------------|--|
| Journal:                        | <i>BMJ Open</i>  |
| Manuscript ID                   | bmjopen-2017-019642.R1   |
| Article Type:                   | Research   |
| Date Submitted by the Author:   | n/a  |
| Complete List of Authors:       | <p>Heuvelings, Charlotte C.; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>Greve, Patrick F.; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>de Vries, Sophia G.; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>Jelle Visser, Benjamin ; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>Bélar, Sabine; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>Janssen, Saskia; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>Cremers, Anne L.; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> <p>Spijker, René; Academic Medical Center, University of Amsterdam, Medical Library</p> <p>Shaw, Elizabeth; National Institute for Health and Clinical Excellence</p> <p>Hill, Ruaraidh A.; University of Liverpool, Liverpool Reviews and Implementation Group, Health Services Research</p> <p>Zumla, Alimuddin; University College London, London, United Kingdom., Infection and Immunity</p> <p>Sandgren, Andreas; European Centre for Disease Prevention and Control</p> <p>van der Werf, MJ; European Centre for Disease Prevention and Control,</p> <p>Grobusch, Martin Peter; Academic Medical Center, University of Amsterdam, Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine</p> |
| <b>Primary Subject Heading</b>: | Public health  |
| Secondary Subject Heading:      | Evidence based practice, Global health, Health services research, Infectious   |

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

|           |  |
|-----------|--|
|           | diseases   |
| Keywords: | Tuberculosis < INFECTIOUS DISEASES, HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Public health < INFECTIOUS DISEASES |
|           |  |

SCHOLARONE™  
Manuscripts

For peer review only

1  
2  
3 **Effectiveness of service models and organisational structures supporting tuberculosis**  
4 **identification and management in hard-to-reach populations in countries of low and medium**  
5 **tuberculosis incidence: A systematic review**  
6  
7

8 Charlotte C Heuvelings<sup>1</sup>, Patrick F Greve<sup>1</sup>, Sophia G de Vries<sup>1</sup>, Benjamin J Visser<sup>1</sup>, Sabine Bélard<sup>1</sup>,  
9 Saskia Janssen<sup>1</sup>, Anne L. Cremers<sup>1</sup>, René Spijker<sup>2</sup>, Beth Shaw<sup>3</sup>, Ruairaidh A Hill<sup>4</sup>, Alimuddin Zumla<sup>5</sup>,  
10 Andreas Sandgren<sup>6</sup>, Marieke J van der Werf<sup>6</sup>, Martin P. Grobusch<sup>1</sup>  
11  
12

- 13  
14 1. Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases,  
15 Division of Internal Medicine, Academic Medical Center, University of Amsterdam,  
16 Meibergdreef 9, 1105 AZ, Amsterdam, the Netherlands  
17  
18 2. Medical Library, Academic Medical Center, University of Amsterdam, Meibergdreef 9, 1105  
19 AZ, Amsterdam, the Netherlands  
20  
21 3. National Institute for Health and Care Excellence, Level 1A, City Tower, Piccadilly Plaza,  
22 3Manchester, M1 4BT, United Kingdom  
23  
24 4. Liverpool Reviews and Implementation Group, Health Services Research, University of  
25 Liverpool, Liverpool, L69 3GB, United Kingdom  
26  
27 5. Division of Infection and Immunity, University College London, and NIHR Biomedical  
28 Research Centre at UCL Hospitals, London, United Kingdom  
29  
30 6. European Centre for Disease Prevention and Control, Tomtebodavägen 11a, 171 65 Solna,  
31 Sweden  
32  
33  
34  
35  
36  
37  
38

39 Word count abstract: 297

40 Word count main text: 3,536

41 Additional files: 2 Tables; 1 Figure; 1 Box; 7 Supplements  
42  
43

44 Correspondence to: M.J. van der Werf [marieke.vanderwerf@ecdc.europa.eu](mailto:marieke.vanderwerf@ecdc.europa.eu)  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

**Objective** To determine which service models and organisational structures are effective and cost-effective for delivering tuberculosis services to hard-to-reach populations.

**Design** Embase and MEDLINE (1990-2017) were searched in order to update and extend the 2011 systematic review commissioned by NICE, discussing interventions targeting service models and organisational structures for the identification and management of tuberculosis in hard-to-reach populations. The NICE and Cochrane Collaboration standards were followed.

**Setting** European Union, European Economic Area, European Union candidate countries and Organisation for Economic Co-operation and Development countries.

**Participants** Hard-to-reach populations, including migrants, homeless people, drug users, prisoners, sex workers, people living with HIV and children within vulnerable and hard-to-reach populations.

**Primary and secondary outcome measures** Effectiveness and cost-effectiveness of the interventions.

**Results** From the 19,720 citations found, five new studies were identified, in addition to the six discussed in the NICE review. Community health workers from the same migrant community, street teams and peers improved tuberculosis screening uptake by providing health education, promoting tuberculosis screening and organising contact tracing. Mobile tuberculosis clinics, specialised tuberculosis clinics and improved cooperation between health care services can be effective at identifying and treating active tuberculosis cases, and are likely to be cost-effective. No difference in treatment outcome was detected when directly observed therapy was delivered at a health clinic or at a convenient location in the community.

**Conclusions** Although evidence is limited due to the lack of high quality studies, interventions using peers and community health workers; mobile tuberculosis services, specialised tuberculosis clinics, and improved co-operation between health services can be effective to control tuberculosis in hard-to-reach populations. Future studies should evaluate the (cost-)effectiveness of interventions on TB identification and management in hard-to-reach populations and countries should be urged to publish the outcomes of their TB control systems.

**Systematic review registration** PROSPERO CRD42015017865.

### Strengths and limitations of this study

- PRISMA and Cochrane Collaboration reporting guidelines for systematic reviews were followed.
- The search was highly sensitive but we might have missed important information as many European countries do not publish their tuberculosis identification and management data in journals, our search focused on Embase and MEDLINE.
- We identified five studies and discuss the results together with the six studies identified by the NICE review to give the complete body of evidence.
- None of the included studies was of high quality and there was high heterogeneity across the studies prohibiting a meta-analysis.

### Introduction

Prevention and control of tuberculosis (TB) is based on early detection and diagnosis of TB followed by effective treatment. In 2015 there were an estimated 10.4 million incident TB cases worldwide, an estimated 4.3 million cases were either not diagnosed, or diagnosed but not reported to national TB programmes.<sup>1</sup> Trends for TB treatment are encouraging, with most notified TB cases completing their treatment successfully; although treatment success rates in some regions, such as the European region, were considerably below the WHO World Health Assembly target of 85%.<sup>1</sup>

In many countries with a low TB incidence (less than ten TB cases per 100,000 population),<sup>2</sup> TB prevails in the big cities where vulnerable and hard-to-reach (under-served) populations are concentrated.<sup>3</sup> These populations, such as people who are homeless (or have insecure accommodation), misuse drugs or are migrants, are at higher risk of contracting TB and are more likely unable or unwilling to seek medical care and comply with the long term TB treatment. Managing TB in those populations is therefore challenging, due to barriers caused by stigma, cultural barriers, poor access to health care services and low levels of accurate TB knowledge.<sup>4,5-7</sup> This therefore requires special efforts. Health care services need to be organised effectively to identify and diagnose TB cases, and to provide adequate treatment and support. This can be organised in different ways, e.g. mainly as hospital-based<sup>8</sup> or health centre-based;<sup>9</sup> including the public sector, private sector,<sup>10</sup> or civil society and other partners.<sup>11</sup> Sometimes, organisation of the services has proven ineffective in managing TB.<sup>12</sup>

The review question of this systematic review with a scoping component was: 'Which service models and organisational structures, including different types of healthcare workers and settings, are effective and cost-effective for delivering TB services to hard-to-reach populations in low- and medium-incidence countries?'

Findings of this review and the previously published review series<sup>4,13</sup> formed the base for the guidance document by the European Centre for Disease Prevention and Control (ECDC) on controlling TB in hard-to-reach and vulnerable populations.<sup>14</sup>

### Methods

1  
2  
3 In 2011, the Matrix Knowledge Group published a review, commissioned by the National Institute for  
4 Health and Clinical Excellence (NICE), on effectiveness and cost-effectiveness of service models or  
5 structures, focussing on the type of health care worker and setting, to identify and manage TB in hard-  
6 to-reach populations. We updated and extended the NICE review,<sup>15</sup> using the same methodology but  
7 adjusting the focus by excluding latent TB infection and including additional hard-to-reach  
8 populations. The review was conducted following standards described by the Cochrane Collaboration<sup>16</sup>  
9 and NICE methods guidelines.<sup>17</sup> Results are reported according to PRISMA guidelines.<sup>18</sup> The review  
10 protocol was registered in advance in the database of prospectively registered systematic reviews in  
11 health and social care, PROSPERO (CRD42015017865).  
12  
13  
14  
15

### 16 *Selection of studies and data management*

17 The same search strategy as for the previous NICE review<sup>15</sup> and the previous published review by  
18 Heuvelings et al.<sup>13</sup> was used, searching Embase and MEDLINE through the Ovid platform. The search  
19 was expanded by including all European Union (EU)/European Economic Area (EEA) and EU  
20 candidate countries to the Organisation for Economic Co-operation and Development (OECD)  
21 countries (see **Box 1**).<sup>15</sup> Two hard-to-reach populations (people living with HIV and children within  
22 vulnerable and hard-to-reach populations), were added in addition to the hard-to-reach populations  
23 included by the NICE review (migrants including refugees, asylum seekers and the Roma population,  
24 homeless people including rough sleepers and shelter users, drug users, prisoners and sex workers).<sup>15</sup>  
25 The update of the search conducted for the NICE review<sup>15</sup> covered the period 1 January 2010  
26 (overlapping the end of the search period of the NICE review<sup>15</sup> with a few months) to 24 February  
27 2017. The search for the expanded geographical area and newly included hard-to-reach populations  
28 covered a time period from 1 January 1990 (beginning of the search period used in the NICE review<sup>15</sup>)  
29 to 24 February 2017.  
30  
31  
32  
33  
34

35 Reference lists of relevant systematic reviews were scanned. No language restrictions were applied.  
36 Studies focussing on the effectiveness and/or cost-effectiveness of interventions for service models and  
37 organisational structures supporting TB identification and management of hard-to-reach populations  
38 (see **Box 1**) were included.  
39

40 Predefined interventions were using more convenient locations (like specialised TB centres, shelters  
41 for homeless people or drug users, needle exchange/methadone programme locations, port of arrival,  
42 schools or mobile clinics) and peers or health care workers with the same ethnic or cultural  
43 background; however, other interventions could also be included if they supported TB identification or  
44 management in hard-to-reach populations. TB identification tools, TB diagnostics, incentives, social  
45 support, directly observed therapy and treatment of co-morbidities are discussed in another review.<sup>13</sup>  
46 In this review we aim to identify the effectiveness of the type of health worker and setting, to identify and  
47 manage TB in hard to reach and vulnerable populations.  
48  
49  
50  
51

52 The comparator was defined during the review process; interventions were compared to a relevant  
53 comparator, for example usual care or no intervention, another intervention, or historical comparison.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Outcomes were defined as any measure of TB identification and management (for example, number of  
4 people screened, screening coverage, proportion receiving treatment and treatment completion rate).  
5 Effectiveness was defined as an improvement in any measure of TB identification and/or management.  
6 Randomised and non-randomised studies were eligible for inclusion.  
7

8 See **Supplementary Material I** for the **PROSPERO study protocol**, **Supplementary Material II**  
9 **for** PICOS (Population-Intervention-Comparator-Outcome-Study design) questions and  
10 **Supplementary Material III** for the complete search strategy and search results.  
11

#### 12 *Data extraction, data items, and synthesis*

13  
14 Identified citations were entered into an EndNote database, and duplicates removed (EndNote X7.1,  
15 Thomson Reuters 2014). The inclusion criteria were piloted and refined using the first 25 citations.  
16 Double screening was conducted by one reviewer screening 100% of the citations (CCH) while another  
17 two reviewers screened 50% of the citations each (PFG, SGdV) for inclusion on title and abstract.  
18 Disagreement was resolved by discussion. Full text files of included citations were retrieved;  
19 irretrievable articles (not available after attempts online, from the university library or through  
20 contacting authors) were excluded. Two reviewers assessed full text records for inclusion (CCH, PFG).  
21 Disagreement was resolved by discussion. Agreement after screening on title and abstract was 99.6%  
22 with an inter-rater reliability (Cohen's kappa) of  $\kappa = 0.985$ .  
23  
24  
25  
26  
27

28 Data extraction forms from the NICE review<sup>15</sup> were used to extract information on participant  
29 characteristics, settings, types of services/organisational structures, types of healthcare workers  
30 delivering the service, outcome measures, methods of analysis and results. For one study data  
31 extraction was conducted by two reviewers (CCH, PFG) independently. For the remaining studies, data  
32 extraction was conducted by one reviewer (CCH) and checked by a second (PFG); disagreement was  
33 resolved by discussion. In one case, the study author was contacted to verify data and obtain additional  
34 data.<sup>19</sup>  
35  
36  
37

38 To facilitate comparability, data synthesis was structured in a similar way to that of the NICE review.<sup>15</sup>  
39 Studies were divided into those examining service models and organisational structures for TB  
40 identification (screening) and those examining service models and organisational structures for TB  
41 management (treatment and support) in hard-to-reach populations. Data were analysed narratively, and  
42 appropriateness of meta-analysis considered. Findings were reported as stated by the study authors.  
43  
44  
45

#### 46 *Risk of bias in individual studies and overall strength of evidence*

47 The modified NICE Quality Assessment Tools<sup>17</sup> (based on the Graphical Appraisal Tool for  
48 Epidemiological studies) were used to assess quality and risk of bias of included studies. This included  
49 an assessment of selection of study sample, minimisation of selection bias and contamination,  
50 controlling confounding, outcome measurements, analytical methods and risk of bias. Two reviewers  
51 (CCH, PFG) assessed one study independently; the remaining studies were assessed by one reviewer  
52 (CCH), and checked by a second reviewer (PFG). Any disagreement was resolved by discussion.  
53 Studies were given a quality rating based on the quality assessment: high quality [++], medium quality  
54  
55  
56  
57  
58  
59  
60

[+], or low quality [-]. The strength of the evidence was assessed and reported as described in the previous NICE review<sup>15</sup> (**Supplementary Material IV**).

## Results

Of the 19,720 citations identified by the literature search five studies were included in this review (**Figure 1**).<sup>11,19-22</sup> These five studies are in addition to the six studies<sup>23-28</sup> included in the NICE review.<sup>15</sup> The results section in this paper focuses on the evidence of the five studies identified in our updated review. The evidence statements (presented in **Supplementary Material IV**) summarise evidence identified in terms of consistency, quality and applicability, combining evidence from the NICE review<sup>15</sup> and this update.

All five studies were conducted in the EU; two in the United Kingdom (UK),<sup>19,22</sup> one in Germany,<sup>20</sup> one in Portugal<sup>11</sup> and one in Spain.<sup>21</sup> Two studies focussed on homeless people,<sup>19,20</sup> one on homeless people and drug users,<sup>22</sup> one on drug users alone<sup>11</sup> and one on migrants.<sup>21</sup> Four studies<sup>5,19-21</sup> addressed the influence of the type of healthcare worker on TB identification and TB management and one study focussed on the influence of different settings on TB identification.<sup>22</sup> A variety of study designs were included; one study was a prospective cluster randomised controlled trial (RCT),<sup>19</sup> one was an economic evaluation using a compartmental model of treated and untreated active TB cases<sup>22</sup>, and three studies were retrospective comparison studies.<sup>11,20,21</sup> Study characteristics of included studies are described in **Table 1**. The data extraction forms by study are presented in **Supplementary Material V**.

None of the included studies in this review had a low risk of bias, three studies<sup>19,21,22</sup> had a medium risk of bias, the other two studies<sup>5,20</sup> were assessed as having a high risk of bias (**Supplementary Material VI**).

We did not perform a meta-analysis due to study heterogeneity. Results were synthesised narratively.<sup>29</sup>

Main outcomes for services structures and organisational models for TB identification among hard-to-reach populations, combined with the findings of the NICE review,<sup>15</sup> are summarised in **Table 2**. For full evidence statements, see **Supplementary Material IV**.

Three studies<sup>19-21</sup> compared the effect of the type of healthcare worker on TB identification.

In the UK, a cluster-randomised trial found that peer educators working together with shelter staff to encourage homeless people to participate in a TB screening programme using mobile X-ray units (MXU), did not improve screening uptake compared to encouragement by shelter staff only (respectively 40%, interquartile range [IQR] 25-61 versus 45%, IQR 33-55; adjusted risk ratio [aRR] 0.98, 95% confidence interval [95% CI] 0.80 to 1.20).<sup>19</sup> Control sites were not 'naïve' for peer intervention which could have caused contamination of the control sites and contributed to the negative finding.



1  
2  
3 In Germany, introduction of TB education and promotion of voluntary chest X-ray (CXR) screening at  
4 least once every two years by community health workers (CHWs) improved screening uptake in  
5 homeless people and drug users. Annual screening coverage increased from 10.0% at the beginning of  
6 the study period (2002-2004) to 15.0% during the middle part of the study period (2004-2006), the last  
7 part of the study period had a 13.4% annual screening coverage (2005-2007). Screening once every  
8 two years increased screening coverage from 18.0% (2002-2004) to 26.4% (2004-2006). Coverage was  
9 23.4% at the third and final study period (spanning 2005-2007).<sup>20</sup> The authors did not test for statistical  
10 significance, and denominator data (the number of homeless people and drug users in the study area)  
11 were estimated.  
12  
13  
14  
15

16 In Barcelona, Spain, contact tracing organised by CHWs coming from the same migrant community as  
17 the person diagnosed with TB, improved contact tracing among migrants to 66.2% (2003-2005)  
18 compared to 55.4% (2000-2002) in the period before the implementation of the intervention using  
19 CHWs (adjusted odds ratio [aOR] of an index case having their contacts screened before and after the  
20 intervention was 1.8, 95% CI 1.3 to 2.5,  $p < 0.001$ ).<sup>21</sup> Identification and tracing of at least one contact  
21 was taken as appropriate contact tracing, where all contacts at risk should be traced to detect and treat  
22 TB transmission early. The population characteristics varied, the age and country of origin was  
23 different between both periods. The importance of contact tracing is to identify cases early to reduce  
24 transmission, the authors did not report if any of the contacts traced had active TB.  
25  
26  
27  
28

29 Two studies<sup>11,22</sup> evaluated the effect of the type of healthcare worker and the setting on TB  
30 identification and TB management.  
31

32 In Portugal, improved co-operation of 'key partners' (street teams, TB clinics, drug user support  
33 centres, local public health department and local hospital) for TB identification and management in  
34 drug users was evaluated in a before-and-after study. Representatives of all 'key partners' (authors'  
35 term) worked on improving policies, clinic screening procedures and co-operation. Key partners were  
36 trained in identifying drug users in their population; and offering health promotion, notification cards,  
37 free transport to the TB clinic, free medical and substance abuse care, directly observed therapy (DOT)  
38 for active TB cases, identification of non-compliant patients and the cause of non-compliance, and  
39 tailor-made strategies to improve compliance. This resulted in an increase of TB screening uptake;  
40 from 52 drug users being screened before the intervention (2001-2003 when there was no active  
41 screening policy) to 465 drug users screened thereafter (2005-2007). Of all people misusing drugs  
42 taking-up screening, the proportion without TB symptoms increased from 41.6% to 93.5% (OR =  
43 21.76; 95% CI 13.03 – 36.33) indicating improved TB awareness and access to screening facilities for  
44 drug users. Of all drug users with active TB, the proportion identified by screening increased from  
45 13.4% to 61.0% (OR 10.1; 95% CI: 4.44 – 23.0). Treatment default rates decreased from 35.4% to  
46 10.2% (OR 0.21, 95% CI 0.08-0.54), compared to the period before the intervention (2001-2003)  
47 when TB treatment was not compulsory and compliance was stimulated by TB education and  
48 providing information on the importance of treatment completion.<sup>11</sup> Although the absolute number of  
49 drug users screened increased, information on the screening coverage was not available as denominator  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 data were not provided. Another limitation is that the results were not adjusted for confounding factors,  
4 baseline characteristics might have been different as the two cohorts were recruited over different time  
5 periods and participation was voluntary which may have led to selection bias.  
6  
7

8 In the United Kingdom, the effectiveness and cost-effectiveness of the 'Find and Treat' service (raising  
9 awareness of TB screening and providing a mobile TB screening and treatment service) for homeless  
10 people and drug users was evaluated and compared to people (with a history of homelessness,  
11 imprisonment, drug abuse or mental health problems) self-presenting to a London TB clinic receiving  
12 standard TB care at the clinic.<sup>22</sup> The authors estimated that 22.9% of the patients detected by the 'Find  
13 and Treat' service with the longest first symptom-to-detection time would not have self-presented plus  
14 35.4% were asymptomatic at time of detection and would not have self-presented, only part of the  
15 asymptomatic patients would self-present to a TB clinic at a later stage when symptoms would have  
16 developed. The 'Find and Treat' service had a higher treatment completion rate (67.1% versus 56.8%),  
17 and a lower lost to follow-up rate (2.1% versus 17.2%) compared to the control group receiving  
18 standard TB care at a TB clinic. The authors concluded that the 'Find and Treat' service was cost-  
19 effective, when using the threshold used by NICE of £20,000 to £30,000/QALY gained, with an  
20 incremental cost ratio of £18,000 per QALY gained for the TB screening service and £4,100 per  
21 QALY gained for the TB management service. This study has a few limitations: firstly, it is a non-  
22 randomised study, secondly the 'Find and Treat' service identifies extremely hard-to-reach populations  
23 of which some would never self-present therefore the findings could be even better in less hard-to-  
24 reach populations, and thirdly the economical evaluation is based on a compartmental model that does  
25 not take secondary transmission and drug-resistance into account.  
26  
27  
28  
29  
30  
31  
32  
33

### 34 Discussion

35 To tackle TB and disrupt transmission in high-income, low TB incidence settings, improvement of TB  
36 care in hard-to-reach populations is of vital importance. In this updated review, five studies,<sup>11,19-22</sup>  
37 published between 1 January 2010 and 24 February 2017, evaluating effectiveness of services models  
38 and organisational structures supporting TB identification and management of hard-to-reach  
39 populations, were identified in addition to the six studies considering active TB<sup>23-28</sup> identified by the  
40 NICE review.<sup>15</sup> Only one study<sup>22</sup> evaluated cost-effectiveness. Although the evidence from two  
41 reviews is limited, it highlights those interventions that are likely to be effective and those that have no  
42 clear evidence of being effective (**Table 2**). For development of the ECDC guidance document,<sup>14</sup> a  
43 scientific panel compiled by ECDC carefully considered these findings. Their main suggestions for  
44 action were to involve CHWs or peers to improve TB screening uptake and TB treatment completion  
45 among homeless people<sup>20</sup> and drug users;<sup>5,20,23</sup> to use outreach teams to improve TB screening uptake  
46 and TB treatment completion among vulnerable populations;<sup>22</sup> and to strengthen relationships and good  
47 collaboration between health care workers, peers, communities, and patients to improve treatment  
48 outcome among vulnerable populations.<sup>5,20,22,23</sup> The updated systematic review provided evidence for  
49 all suggestions except for using peers to improve screening uptake. This is in contrast to an American  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 study<sup>23</sup> included in the original NICE review,<sup>15</sup> which showed that peers improved contact tracing and  
4 treatment adherence among drug users.  
5

### 6 7 *Strengths and limitations*

8 PRISMA and Cochrane Collaboration reporting guidelines for systematic reviews were followed.  
9 Established screening protocols were used, including double screening, and the search was highly  
10 sensitive. The methodology from the previous NICE review<sup>15</sup> was followed, in order to connect this  
11 update and, so, describe the full body of relevant evidence. High quality evidence is lacking. Only  
12 one<sup>23</sup> study from the NICE review<sup>15</sup> was considered to be of high quality, all other studies had some  
13 risk of bias (five medium risk<sup>19,21,22,24,26</sup> and five high risk<sup>11,20,25,27,28</sup>). Therefore, only limited  
14 conclusions can be drawn. Most studies lacked identification and adjustment for confounding factors  
15 and the use of appropriate analytical methods. In addition, many studies were biased, particularly with  
16 regard to potential selection bias. A meta-analysis could not be performed, because of heterogeneity  
17 across the studies. Gaps in evidence exist; no studies focussing on children within vulnerable and hard-  
18 to-reach populations, or on people living with HIV or sex workers were identified. Only three studies  
19 provided economic data; one study identified by this review,<sup>22</sup> and two<sup>25,27</sup> by the NICE review.<sup>15</sup>

20 Our search focussed on publications in databases Embase and MEDLINE. Many European countries  
21 have strong organisational structures for TB identification and management, but these countries did not  
22 publish their data on these organisational structures in journals, which may have caused a publication  
23 bias. Comparing findings of the NICE review<sup>15</sup> with this review comes with some limitations. For the  
24 NICE review only 10% of the citations were double screened,<sup>15</sup> compared to 100% for this updated  
25 review, therefore studies conducted between 1990-2010 might have been missed. The NICE review  
26 focussed their recommendations on the population in the United Kingdom,<sup>15</sup> and this review focussed  
27 on populations in high-income, low TB incidence countries. Further methodology was identical.  
28  
29  
30  
31  
32  
33  
34  
35

36 The evidence identified by this review and the previous NICE review<sup>15</sup> along with evidence presented  
37 in a review series covering the barriers and facilitators of seeking TB care,<sup>6</sup> and the effectiveness of  
38 interventions for TB identification and management in hard-to-reach populations,<sup>13</sup> was used to  
39 develop the ECDC guidance on improving TB identification and management among hard-to-reach  
40 and vulnerable populations in Europe.<sup>14</sup> ECDC recommended that implementation of the interventions  
41 is context-specific; it depends on the setting, target population, resources available and health-care  
42 systems in place. Interventions focussing on one specific hard-to-reach population might not work in  
43 another hard-to-reach population, therefore, the interventions have to be adapted and re-assessed per  
44 target population.<sup>14</sup> Given the scope of this review, considering settings across Europe, findings  
45 presented here are potentially relevant to any low-incidence region, and are relevant to other  
46 institutions/governmental organisations seeking to improve service structures for TB identification and  
47 management among hard-to-reach populations.  
48  
49  
50  
51  
52  
53

54 Characteristics of different hard-to-reach populations and their TB epidemiology vary per country and  
55 setting. Challenges in identification and management of TB should be identified and targeted, tailored  
56  
57  
58  
59  
60

1  
2  
3 to the specific setting and hard-to-reach population. These TB interventions could be integrated within  
4 broader programmes targeting specific populations. A follow-up systematic review should include  
5 information from national public health services about their organisational structures for TB  
6 identification and management. National public health services are urged to regularly analyse their  
7 organisational structures for TB identification and management and publish this data.

8  
9 Efforts to improve quality of research on service models and organisational structures should be made,  
10 even though it is often challenging to perform 'clean', unbiased, and un-confounded trials in hard-to-  
11 reach populations, as attrition rates are often high, and confounding factors are plentiful. This includes  
12 conducting (cluster) randomised controlled trials and before-and-after studies where appropriate,  
13 recruiting an adequate number of participants, using relevant control groups, and minimising selection  
14 bias. Standardised case definitions for hard-to-reach populations should be created. Feasibility,  
15 effectiveness, cost-effectiveness and impact of interventions should be evaluated. Mathematical  
16 economic models can be used to evaluate costs.<sup>14</sup>

### 21 **Conclusions**

22 Identification and management of TB in hard-to-reach populations is suboptimal.<sup>2</sup> Therefore, service  
23 models and organisational structures to identify and manage TB in hard-to-reach populations should be  
24 improved and evaluated regularly.

25  
26 Our systematic review, in conjunction with the original NICE review<sup>15</sup> provides limited evidence, due  
27 to the lack of high quality studies, that interventions such as using peers and CHWs; mobile TB  
28 services, specialised TB clinics, screening, or active case finding in non-healthcare settings, as well as  
29 improved co-operation between key services can help to improve TB identification and management.

30  
31 Further research should be undertaken to evaluate other effective and cost-effective ways to identify  
32 and manage TB in hard-to-reach populations and countries with good TB control systems are urged to  
33 evaluate their system and publish the data.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Competing interests

All authors have completed the ICMJE uniform disclosure form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) and all except for MPG and BS declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work. MPG reports grants from ECDC, for the conduct of part of the study. BS reports that NICE - her employing organisation - has published guidance in this area.

### Authors' contribution

RS conducted the literature search. CCH, SGdV and PFG performed the study selection. CCH, and PFG collected the data, performed quality/risk of bias assessment and synthesised the data. CCH prepared the manuscript and supplementary files. MPG and MvdW supervised the whole process and contributed to the work. All authors were involved in interpretation of the data, and the writing of the manuscript. All authors contributed to, and endorsed the final version of the manuscript. MPG is guarantor.

### Ethics

Ethics approval was not required for this systematic review.

### Funding

This study was funded by the European Centre for Disease Prevention and Control (contract reference OJ/02/05/2014-PROC/2014/014). None of the authors have received payment from a pharmaceutical company or other agency to write this article.

### Statement of independence of researchers from funders

The funder of the study was involved in study design, data interpretation, and reporting. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

### Data sharing


Data extraction forms and quality assessment forms are available from **Supplementary files V and VI**.

## References

1. World Health Organization (WHO). Global Tuberculosis report 2016. Geneva: WHO 2016 (WHO/HTM/TB/2016.13).
2. World Health Organization (WHO) and European Respiratory Society. Towards TB elimination: an action framework for low-incidence countries. Geneva; WHO 2014 (WHO/HTM/TB/2014.13).
3. de Vries G, Aldridge RW, Cayla JA, et al. Epidemiology of tuberculosis in big cities of the European Union and European Economic Area countries. *Euro Surveill* 2014; **19**(9).
4. de Vries SG, Cremers AL, Heuvelings CC, et al. (in press). Barriers and facilitators for the uptake of tuberculosis diagnostic and treatment services by hard-to-reach populations in low- and medium-incidence countries: A systematic review of qualitative literature. *Lancet Infect Dis* 2017; published online March 17. [http://dx.doi.org/10.1016/S1473-3099\(16\)30531-X](http://dx.doi.org/10.1016/S1473-3099(16)30531-X).
5. Kamholz SL. Resurgence of tuberculosis: the perspective a dozen years later. *J Assoc Academic Minor Phys* 1996; **7**(3): 83-6.
6. de Vries G, van Hest RA, Richardus JH. Impact of mobile radiographic screening on tuberculosis among drug users and homeless persons. *Am J Respir Crit Care Med* 2007; **176**(2): 201-7.
7. van Hest NA, Aldridge RW, de Vries G, et al. Tuberculosis control in big cities and urban risk groups in the European Union: a consensus statement. *Euro surveill* 2014; **19**(9).
8. Belling R, McLaren S, Boudioni M, Woods L. Pan-London tuberculosis services: a service evaluation. *BMC Health Serv Res* 2012; **12**: 203.
9. Ohkado A, Williams G, Ishikawa N, Shimouchi A, Simon C. The management for tuberculosis control in Greater London in comparison with that in Osaka City: lessons for improvement of TB control management in Osaka City urban setting. *Health Policy* 2005; **73**(1): 104-23.
10. Lei X, Liu Q, Escobar E, et al. Public-private mix for tuberculosis care and control: a systematic review. *Int J Tuberc Lung Dis* 2015; **34**: 20-32.
11. Duarte R, Santos A, Mota M, Carvalho A, Marques A, Barros H. Involving community partners in the management of tuberculosis among drug users. *Public Health* 2011; **125**(1): 60-2.
12. Frieden TR, Fujiwara PI, Washko RM, Hamburg MA. Tuberculosis in New York City--turning the tide. *N Engl J Med* 1995; **333**(4): 229-33.
13. Heuvelings C, de Vries S, Greve PF, et al. (in press). Effectiveness of interventions aiming at identifying and managing tuberculosis among hard-to-reach populations: A systematic review. *Lancet Infect Dis* 2017; published online March 17. [http://dx.doi.org/10.1016/S1473-3099\(16\)30532-1](http://dx.doi.org/10.1016/S1473-3099(16)30532-1).
14. European Centre for Disease Prevention and Control (ECDC). Guidance on tuberculosis control in vulnerable and hard-to-reach populations. Stockholm: ECDC 2016.
15. Rizzo M, Martin A, Jamal F, et al. Evidence review on the effectiveness and cost-effectiveness of service models or structures to manage tuberculosis in hard-to-reach groups. Matrix evidence: NICE – National Institute for Health and Clinical Excellence, 2011.
16. Higgins J, Green S, (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]: The Cochrane Collaboration; 2011.
17. National Institute for Health and Clinical Excellence, London. Methods for the development of NICE public health guidance (second edition). 2009.
18. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009; **339**: b2535.
19. Aldridge RW, Hayward AC, Hemming S, et al. Effectiveness of peer educators on the uptake of mobile X-ray tuberculosis screening at homeless hostels: a cluster randomised controlled trial. *BMJ Open* 2015; **5**(9): e008050.
20. Goetsch U, Bellinger OK, Buettel KL, Gottschalk R. Tuberculosis among drug users and homeless persons: impact of voluntary X-ray investigation on active case finding. *Infection* 2012; **40**(4): 389-95.

- 1  
2  
3 21. Ospina JE, Orcau A, Millet JP, Sanchez F, Casals M, Cayla JA. Community health workers  
4 improve contact tracing among immigrants with tuberculosis in Barcelona. *BMC Public Health* 2012;  
5 **12**: 158.
- 6 22. Jit M, Stagg H, Aldridge R, White P, Abubakar I. Dedicated outreach service for hard to reach  
7 patients with tuberculosis in London: observational study and economic evaluation. *BMJ* 2011; **343**:  
8 d5376.
- 9 23. Ricks PM. Tuberculosis control among substance users: The indigenous leadership outreach  
10 model vs. standard care. Chicago, IL: University of Illinois, 2008.
- 11 24. El-Hamad I, Casalini C, Matteelli A, et al. Screening for tuberculosis and latent tuberculosis  
12 infection among undocumented immigrants at an unspecialised health service unit. *Int J Tuberc Lung*  
13 *Dis* 2001; **5**(8): 712-6.
- 14 25. Bothamley GH, Rowan JP, Griffiths CJ, et al. Screening for tuberculosis: the port of arrival  
15 scheme compared with screening in general practice and the homeless *Thorax* 2002; **57**(1): 45-9.
- 16 26. Miller TL, Hilsenrath P, Lykens K, McNabb SJ, Moonan PK, Weis SE. Using cost and health  
17 impacts to prioritize the targeted testing of tuberculosis in the United States. *Ann Epidemiol* 2006;  
18 **16**(4): 305-12.
- 19 27. Mor Z, Lerman Y, Leventhal A. Pre-immigration screening process and pulmonary  
20 tuberculosis among Ethiopian migrants in Israel. *Eur Respir J* 2008; **32**(2): 413 - 8.
- 21 28. Dèruaz J, Zellweger J, P. Directly observed therapy for tuberculosis in a low prevalence  
22 region: first experience at the Tuberculosis Dispensary in Lausanne. *Swiss Med Wkly* 2004; **134**: 552-8.
- 23 29. Popay J, Roberts H, Sowden A, et al. Guidance on the conduct of narrative synthesis in  
24 systematic reviews. A product from the ESRC methods programme *Lancaster* 2006.
- 25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1. Study selection process



For the Organisation for Economic Co-operation and Development (OECD) countries and hard-to-reach populations discussed in the previous NICE review,<sup>13</sup> the study period covers 1 January 2010 to 24 February 2017. For the newly included European Union (EU)/European Economic Area/EU candidate countries, and the newly included hard-to-reach populations (people living with Human Immunodeficiency Virus (HIV) and children within vulnerable and hard-to-reach populations, the study period covers 1 January 1990 to 24 February 2017.



**Table 1. Characteristics of studies applying different service models and organisational structures to improve TB identification and TB management**

| First Author (year) country                                  | Population                     | Aims  | Intervention   | Comparator   | Study design                               | Outcome measure  | Quality score |
|--|--------------------------------|---|--|--|--|--|---------------|
| <b>TB identification (studies identified by this review)</b> |                                |   |  |  |  |  |               |
| Jit et al. (2011) <sup>22</sup><br>United Kingdom            | Homeless people and drug users | To assess the effectiveness and cost-effectiveness of the Find and Treat service for diagnosing and managing hard-to-reach individuals with active TB in London.  | Period 2007-2010: Find and Treat service:<br>- screening by MXU<br>- peers raising awareness<br>- treatment support  | Passive case detection and standard treatment at a London TB clinic  | Observational and cost-effectiveness study | Identified TB cases, treatment completion, lost to follow-up, incremental costs from healthcare taxpayer perspective | +             |
| Duarte et al. (2011) <sup>11</sup><br>Portugal               | Drug users                     | To evaluate the effect of an intervention with key partners (TB clinic, drug users support centres, shelters, street teams, public health department and hospital) delivering promotion of health-seeking behaviour, eliminating potential barriers for TB screening at a chest clinic and DOT, on identifying TB cases and treatment compliance. | Improved co-operation of key partners (2005-2007):<br>- health education and screening promotion<br>- improved screening procedures<br>- implementation of DOT<br>- free TB care and transport<br>- providing medical and drug abuse treatment<br>- active follow-up of non-compliant patients, the key partners worked together to reach the patient, identify the cause and organise suitable treatment strategies | Period before the intervention (2001-2003):<br><br>- no active screening policy<br>- referral to chest clinic after discharge from hospital<br>- treatment not compulsory<br>- information about disease and treatment given to improve compliance<br>- psychosocial support<br>- free TB treatment, transport and breakfast | Before-after study                         | Identified TB cases and treatment compliance   | -             |
| Goetsch et al. (2012) <sup>20</sup><br>Germany               | Homeless people and drug users | To estimate the coverage of a low-threshold CXR screening programme for pulmonary TB among illicit drug users and homeless persons.   | CHWs providing TB education and promoting voluntary CXR screening 1-2x/year  | Comparing the beginning of the 5 year intervention period with the end (2002-2007)   | Retrospective effectiveness study          | Screening coverage   | -             |
| Ospina et al. (2012) <sup>21</sup><br>Spain                  | Migrants                       | To evaluate the effectiveness of an intervention with community health workers to improve contact tracing among migrants  | CHWs active follow up of cases and contacts, including visits of the cases at home, accompanying at outpatient appointments, providing counselling and information on treatments (2003-2005)   | Pre-intervention period (2000-2002)  | Before-after study                         | Number of migrants that were included in contact tracing   | +             |

|  |                 |   |   |   |                                     |                  |   |
|--|-----------------|---|---|---|-------------------------------------|------------------|---|
| Aldridge et al. (2015) <sup>19</sup><br>United Kingdom | Homeless people | To compare TB screening uptake between current practice of encouraging homeless people by shelter staff and encouragement by shelter staff plus volunteer peer educators. | Encouragement of TB screening by peers in addition to shelter staff | Encouragement of TB screening by shelter staff only | Cluster randomised controlled trial | Screening uptake | + |
|--|-----------------|---|---|---|-------------------------------------|------------------|---|

**TB identification (studies identified by the previous NICE review<sup>15</sup>)**

|   |  |  |  |   |   |   |    |
|---|--|--|--|---|---|---|----|
| El-Hamad et al. (2001) <sup>24</sup><br>Italy           | Migrants   | To compare the completion rates of screening procedures for TB infection among undocumented migrants at specialised TB units and non-specialised health clinics                          | TB screening at specialised TB clinic                              | TB screening at a general health service for migrants   | Prospective cohort                                      | Screening completion                              | +  |
| Bothamley et al. (2002) <sup>25</sup><br>United Kingdom | Migrants and homeless people                                   | To compare the yield and costs of TB screening in three settings: a new entrants' clinic within the port of arrival (POA) scheme; a large general practice; and centres for the homeless | TB screening at a general practice (GP)                            | TB screening at POA and at homeless centres   | Cost analysis   | Cost per person screened per case of TB prevented | -  |
| Deruaz & Zellweger (2004) <sup>28</sup><br>Switzerland  | Migrants, alcohol or drug users, homeless people and prisoners | Evaluation of first experience of the DOT programme for TB introduced in the Canton of Vaud in 1997  | 1. Full DOT<br>2. DOT delivered at TB clinic                       | 1. Partial DOT (DOT only first 2 months of treatment)<br>2. DOT delivered at social outreach site | Before-after study                                      | Adherence to treatment and outcome                | -  |
| Miller et al. (2006) <sup>26</sup><br>United States     | Homeless people and prisoners                                  | To evaluate and compare the efficiency of a non-state-law-mandated TB screening programme for homeless persons with a state-law-mandated TB screening programme for prisoners            | Non-state-law-mandated TB screening programme for homeless persons | State-law-mandated TB screening programme for prisoners   | Retrospective comparison of the cost and health impacts | TB cases averted and cost                         | +  |
| Ricks (2008) <sup>23</sup><br>United States             | Drug users   | To compare the effectiveness of using peers versus 'standard' public health workers to coordinate TB treatment   | Enhanced case management by peers                                  | Limited case management by health care professionals  | Randomised controlled trial                             | Adherence to treatment                            | ++ |

|   |          |  |                         |                          |                               |   |   |
|---|----------|--|-------------------------|--------------------------|-------------------------------|---|---|
| Mor et al. (2008) <sup>27</sup><br>Israel | Migrants | To examine the effectiveness and cost-effectiveness of pre-migration screening and post-migration screening at POA | Pre-migration screening | Post-migration screening | Retrospective cohort analysis | Active TB cases, time between migration and diagnosis, cost-savings | - |
|---|----------|--|-------------------------|--------------------------|-------------------------------|---|---|

### List of Abbreviations

CHW = community health worker; CXR = chest X-ray; C = comparator group; DOT = direct observed treatment; GP = general practice; I = intervention group; IRIS = immune reconstitution inflammatory syndrome; IVDU = intravenous drug users; MXU = mobile X-ray unit; *n* = number of participants; QALYs = quality adjusted life years; POA = port-of-arrival; RCT = randomised controlled trial; TDM = therapeutic drug monitoring; TB = tuberculosis

Study quality: high quality [++], medium quality [+], or low quality [-]

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

**Table 2. Effectiveness of service models and organisational structures interventions to improve TB identification and TB management**

| Population                        | Intervention (I)  | Comparator (C)   | Studies (first author, year, country)  | No. of participants<br>I C |      | Comparison   | Outcome  | Risk of Bias                               |
|-----------------------------------|---|--|--|----------------------------|------|--|--|--|
| Homeless people<br><br>Drug users | Health/TB education and promotion of screening by street teams, drug users support centres, shelters and CHWs | Beginning of the intervention when CHWs were just introduced<br><br>No active screening policy | Goetsch et al. 2012 (Germany) <sup>20</sup><br><br>Duarte et al. 2011 (Portugal) <sup>11</sup> | 465                        | 125  | Retrospective comparison over intervention period<br><br>Retrospective before-after comparison | Improved annual TB screening uptake among homeless people and drug users (from 10.0% to 15.0% at the peak). <sup>20</sup> The percentage of all drug users with active TB identified by screening increased from 13.4% to 61.0% (OR 10.1 [95%CI 4.44-23.0]). <sup>11</sup>   | High <sub>2</sub><br><br>High <sub>3</sub> |
| Homeless people                   | TB education and promotion of screening by peers and shelter staff  | TB education and promotion of screening by shelter staff only                                  | Aldridge et al. 2015 (United Kingdom) <sup>19</sup>  | 1150                       | 1192 | Comparing randomised intervention cluster with comparator cluster                              | No difference in screening uptake (I = 40% [IQR 25-61] versus C = 45% [IQR 33-55], aRR= 0.98 [95%CI 0.80-1.20]).   | Medium <sub>4</sub>                        |
| Migrants                          | Pre-migration screening   | Post-migration screening at POA  | Mor et al. 2008, cited in the NICE review (Israel) <sup>27</sup>                               | 162                        | 105  | Retrospective Intervention versus comparator comparison  | Reduced the risk of developing TB in the new country and was cost-effective (0.28% of the pre-migration versus 0.32% of the post-migration screening migrants developed TB; RR 0.82 p<0.01). The detection period was shorter as well (193 days versus 487 days between entry and diagnosis; OR=0.72 [95%CI 0.59-0.89] p=0.002). | High <sub>8</sub>                          |
| Prisoners and homeless people     | TB screening in a prison  | TB screening at a homeless centre  | Miller et al. 2006, cited in the NICE review (United States) <sup>26</sup>                     | 22920                      | 822  | Retrospective comparison of two cohorts  | No difference in screening uptake (94.7% in prison versus 95% in homeless centre p=0.179) but higher proportion of active TB cases were identified at the homeless centre (1.2% versus 0.03% at a prison setting, p<0.001)   | Medium <sub>9</sub>                        |
| Homeless people and migrants      | Active case finding by symptom-based questionnaire at homeless centres  | Active case finding by symptom-based questionnaire at POA                                      | Bothamley et al. 2002, cited in the NICE review (United Kingdom) <sup>25</sup>                 | 262                        | 199  | Cost analysis  | Active case finding at POA was most cost-effective (costs per person screened for every case prevented at POA £10.00, at homeless centre £23.00).  | High <sub>10</sub>                         |
| Migrants                          | Active case finding at a specialised TB clinic using 2 visits   | Active case finding at a general primary care clinic, with referral for CXR, using 3 visits    | El-Hamad et al. 2001, cited in the NICE review (Italy) <sup>24</sup>                           | 749                        | 483  | Prospective intervention versus comparator comparison  | Improved screening completion among migrants (85.6% in TB clinic versus 71.4% at primary care clinic, p = not reported; OR=2.57 [95%CI 1.92-3.42]).  | Medium <sub>5</sub>                        |

|   |   |  |   |     |     |  |  |                     |
|---|---|--|---|-----|-----|--|--|---------------------|
| Drug users  | Contact tracing by peers or CHWs from the same migrant community                  | Peers versus other health care workers   | Ricks 2008, cited in the NICE review (United States) <sup>23</sup>            | 48  | 46  | RCT  | Improved contact tracing among drug users (75% by peers versus 47% by healthcare workers, $p = 0.03$ ) <sup>23</sup> and migrants (from 55.4% without CHWs to 66.2% with CHWs; aOR 1.8 [95%CI 1.3-2.5] $p < 0.001$ ). <sup>21</sup>  | Low                 |
| Migrants  |   | Normal practice before introducing CHWs  | Ospina et al. 2012 (Spain) <sup>21</sup>                                      | 388 | 572 | Before-after comparison  |  | Medium <sub>1</sub> |
| Drug users and homeless people                      | Mobile TB screening and treatment service at convenient location in the community | Passive case detection and management at a TB clinic   | Jit et al. 2011 (United Kingdom) <sup>22</sup>                                | 48  | 252 | Prospective intervention versus comparator comparison plus economic evaluation | Improved TB identification among homeless people and drug users; particularly in asymptomatic patients (35.4% extra identified) and those that delay seeking health care (22.2% extra identified). Higher treatment completion rate (67.1% versus 56.8%) and lower lost to follow-up rate (2.1% versus 17.2%). Both parts of the service are cost-effective (screening = £18,000/QALY gained, treatment is £4,100/QALY gained) | Medium <sup>6</sup> |
| Drug users  | Enhanced case management by peers   | Limited case management by regular health care workers   | Ricks 2008, cited in the NICE review (United States) <sup>23</sup>            | 48  | 46  | RCT  | Improved treatment completion in drug users (85% by peers versus 61% by health care workers, RR=2.68 [95%CI 1.24-5.82] $p=0.01$ ).   | Low                 |
| Drug users  | DOT and active follow-up of non-compliant patients by 'key partners'              | Non-compulsory TB treatment and education about TB disease and treatment to improve compliance | Duarte et al. 2011 (Portugal) <sup>11</sup>                                   | 465 | 125 | Retrospective before-after comparison  | Reduced treatment default rates (from 35.4% to 10.2%; OR 0.21 [95%CI 0.08-0.54]).  | High <sub>2</sub>   |
| Migrants, drug users, homeless people and prisoners | DOT at a convenient location in the community                                     | DOT at a health clinic   | Deruaz & Zellweger 2004, cited in the NICE review (Switzerland) <sup>28</sup> | 36  | 18  | Retrospective before-after comparison  | No significant difference in successful treatment outcome, treatment completion and cure rate (85.2% at convenient location versus 92.6% at health clinic, $p=0.67$ )  | High <sub>7</sub>   |

### List of Abbreviations

aOR = adjusted odds ratio, aRR = adjusted risk ratio, C = comparator group, CHWs = community health workers, DOT = directly observed treatment, I = intervention group, IQR = interquartile range, OR = odds ratio, POA = port-of-arrival, RCT = randomised controlled trial, TB = tuberculosis, 95%CI = 95% confidence interval

### Footnotes Risk of Bias:

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47
- <sub>1</sub> = Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods). Contact tracing of only one contact was enough to be called contact tracing, and the ultimate aim of contact tracing (increase case detection and reduce transmission) was not analysed in this study.
- <sub>2</sub> = Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods). Denominator not given therefore unable to calculate screening coverage.
- <sub>3</sub> = Risk of selection bias as participation was voluntary. Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods). No statistical test used to show statistical significance of the findings, an estimated number was used for the denominator.
- <sub>4</sub> = Most comparator sites were not naïve for peer intervention, no individual information of the participants was collected, the characteristics between the two groups might have been significantly different.
- <sub>5</sub> = Not adjusted for difference in baseline characteristics.
- <sub>6</sub> = Study was designed to evaluate the cost-effectiveness, no statistical test used to evaluate statistical significant findings. The 'Find and Treat' service identifies extremely hard-to-reach populations that would never self-present, the findings would underestimate the benefit of the service. The economical evaluation is based on a compartmental model that does not take secondary transmission and drug-resistance into account,
- <sub>7</sub> = Risk of bias due to difference in collecting treatment adherence outcome at the health clinic a nurse recorded treatment adherence at time of visit, in the social outreach group a health care worker was interviewed up to 6 months after treatment completion and was asked about the treatment adherence, risk of recall bias. Not recorded how many people per setting received 6 months of DOT (full DOT) and how many received 2 months of DOT and 4 months of self-treatment (partial DOT), what was another intervention in this study. Allocation to setting was based on needs of participants what might have caused bias.
- <sub>8</sub> = Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods), pre-migration group had a shorter follow-up period than post-migration group what may have influenced the detection of number of TB cases in the pre-migration group.
- <sub>9</sub> = Unclear if the differences in outcome was caused by the setting or by the different methods or to differences in TB prevalence in the different populations.
- <sub>10</sub> = TB prevalence might be different in the different populations as the costs are calculated per active case detected this is a major issue, there were only 3 active TB cases detected, all in the POA group. The economic perspective used was not reported and the costs of identification were not discounted.

**Box 1: Inclusion/exclusion criteria for this review**

## Inclusion criteria:

- Discussing service models and organisational structures, different types of healthcare workers and settings for delivering TB services to hard-to-reach populations;
- Having been conducted in any of the EU/EEA countries (only updated review), the candidate countries\* (only updated review) and the other OECD countries\*\*
- Having been published in 2010 or later for the OECD countries\*\*
- Having been published in 1990 or later for the EU/EEA countries and the EU candidate countries\* not being one of the OECD countries (only updated review)
- Including data from any hard-to-reach population:
  - homeless people
  - people who abuse drugs or alcohol
  - sex workers
  - prisoners or people with a history of imprisonment
  - migrants, including vulnerable migrant populations such as asylum seekers, refugees and the Roma population
  - children within vulnerable and hard-to-reach populations (only updated review)
  - people living with HIV (only updated review)
- Present quantitative empirical data
- Being a (cost)-effectiveness study, or any other type of quantitative primary research, discussing (cost)-effectiveness

## Exclusion criteria:

- Latent TB infection (only updated review)
- Systematic review (only used for reference searching)

*EU/EEA = European Union, European Economic Area; OECD = Organisation for Economic Co-operation and Development; TB = Tuberculosis*

*\*EU candidate countries = Albania, Montenegro, Serbia, the former Yugoslav Republic of Macedonia, and Turkey*

*\*\* OECD countries = Australia, Austria, Belgium, Canada, Chile, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Israel, Italy, Japan, Korea, Luxembourg, Mexico, the Netherlands, New Zealand, Norway, Poland, Portugal, Slovak Republic, Slovenia, Spain, Sweden, Switzerland, Turkey, United Kingdom, United States*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Supplementary Material**

**Supplementary Material I:** Original study protocol

**Supplementary Material II.** PICOS

**Supplementary Material III.** Search strategy

**Supplementary Material IV.** Evidence statements

**Supplementary Material V.** Data extraction forms

**Supplementary Material VI.** Quality Assessment

**Supplementary Material VII.** PRISMA checklist

For peer review only



1  
2  
3 **Effectiveness of service models and organisational structures supporting tuberculosis**  
4 **identification and management in hard-to-reach populations in countries of low and medium**  
5 **tuberculosis incidence: A systematic review**  
6  
7

8 Charlotte C Heuvelings<sup>1</sup>, Patrick F Greve<sup>1</sup>, Sophia G de Vries<sup>1</sup>, Benjamin J Visser<sup>1</sup>, Sabine Bélard<sup>1</sup>,  
9 Saskia Janssen<sup>1</sup>, Anne L. Cremers<sup>1</sup>, René Spijker<sup>2</sup>, Beth Shaw<sup>3</sup>, Ruairaidh A Hill<sup>4</sup>, Alimuddin Zumla<sup>5</sup>,  
10 Andreas Sandgren<sup>6</sup>, Marieke J van der Werf<sup>6</sup>, Martin P. Grobusch<sup>1</sup>  
11  
12

- 13  
14 1. Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases,  
15 Division of Internal Medicine, Academic Medical Center, University of Amsterdam,  
16 Meibergdreef 9, 1105 AZ, Amsterdam, the Netherlands  
17  
18 2. Medical Library, Academic Medical Center, University of Amsterdam, Meibergdreef 9, 1105  
19 AZ, Amsterdam, the Netherlands  
20  
21 3. National Institute for Health and Care Excellence, Level 1A, City Tower, Piccadilly Plaza,  
22 3Manchester, M1 4BT, United Kingdom  
23  
24 4. Liverpool Reviews and Implementation Group, Health Services Research, University of  
25 Liverpool, Liverpool, L69 3GB, United Kingdom  
26  
27 5. Division of Infection and Immunity, University College London, and NIHR Biomedical  
28 Research Centre at UCL Hospitals, London, United Kingdom  
29  
30 6. European Centre for Disease Prevention and Control, Tomtebodavägen 11a, 171 65 Solna,  
31 Sweden  
32  
33  
34  
35  
36  
37  
38

39 Word count abstract: 297

40 Word count main text: 3,536

41 Additional files: 2 Tables; 1 Figure; 1 Box; 7 Supplements  
42  
43

44 Correspondence to: M.J. van der Werf [marieke.vanderwerf@ecdc.europa.eu](mailto:marieke.vanderwerf@ecdc.europa.eu)  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

**Objective** To determine which service models and organisational structures are effective and cost-effective for delivering tuberculosis services to hard-to-reach populations.

**Design** Embase and MEDLINE (1990-2017) were searched in order to update and extend the 2011 systematic review commissioned by NICE, discussing interventions targeting service models and organisational structures for the identification and management of tuberculosis in hard-to-reach populations. The NICE and Cochrane Collaboration standards were followed.

**Setting** European Union, European Economic Area, European Union candidate countries and Organisation for Economic Co-operation and Development countries.

**Participants** Hard-to-reach populations, including migrants, homeless people, drug users, prisoners, sex workers, people living with HIV and children within vulnerable and hard-to-reach populations.

**Primary and secondary outcome measures** Effectiveness and cost-effectiveness of the interventions.

**Results** From the 19,720 citations found, five new studies were identified, in addition to the six discussed in the NICE review. Community health workers from the same migrant community, street teams and peers improved tuberculosis screening uptake by providing health education, promoting tuberculosis screening and organising contact tracing. Mobile tuberculosis clinics, specialised tuberculosis clinics and improved cooperation between health care services can be effective at identifying and treating active tuberculosis cases, and are likely to be cost-effective. No difference in treatment outcome was detected when directly observed therapy was delivered at a health clinic or at a convenient location in the community.

**Conclusions** Although evidence is limited due to the lack of high quality studies, interventions using peers and community health workers; mobile tuberculosis services, specialised tuberculosis clinics, and improved co-operation between health services can be effective to control tuberculosis in hard-to-reach populations. Future studies should evaluate the (cost-)effectiveness of interventions on TB identification and management in hard-to-reach populations and countries should be urged to publish the outcomes of their TB control systems.

**Systematic review registration** PROSPERO CRD42015017865.

### Strengths and limitations of this study

- PRISMA and Cochrane Collaboration reporting guidelines for systematic reviews were followed.
- The search was highly sensitive but we might have missed important information as many European countries do not publish their tuberculosis identification and management data in journals, our search focused on Embase and MEDLINE.
- We identified five studies and discuss the results together with the six studies identified by the NICE review to give the complete body of evidence.
- None of the included studies was of high quality and there was high heterogeneity across the studies prohibiting a meta-analysis.

### Introduction

Prevention and control of tuberculosis (TB) is based on early detection and diagnosis of TB followed by effective treatment. In 2015 there were an estimated 10.4 million incident TB cases worldwide, an estimated 4.3 million cases were either not diagnosed, or diagnosed but not reported to national TB programmes.<sup>1</sup> Trends for TB treatment are encouraging, with most notified TB cases completing their treatment successfully; although treatment success rates in some regions, such as the European region, were considerably below the WHO World Health Assembly target of 85%.<sup>1</sup>

In many countries with a low TB incidence (less than ten TB cases per 100,000 population),<sup>2</sup> TB prevails in the big cities where vulnerable and hard-to-reach (under-served) populations are concentrated.<sup>3</sup> These populations, such as people who are homeless (or have insecure accommodation), misuse drugs or are migrants, are at higher risk of contracting TB and are more likely unable or unwilling to seek medical care and comply with the long term TB treatment. Managing TB in those populations is therefore challenging, due to barriers caused by stigma, cultural barriers, poor access to health care services and low levels of accurate TB knowledge.<sup>4,5-7</sup> This therefore requires special efforts. Health care services need to be organised effectively to identify and diagnose TB cases, and to provide adequate treatment and support. This can be organised in different ways, e.g. mainly as hospital-based<sup>8</sup> or health centre-based,<sup>9</sup> including the public sector, private sector,<sup>10</sup> or civil society and other partners.<sup>11</sup> Sometimes, organisation of the services has proven ineffective in managing TB.<sup>12</sup>

The review question of this systematic review with a scoping component was: 'Which service models and organisational structures, including different types of healthcare workers and settings, are effective and cost-effective for delivering TB services to hard-to-reach populations in low- and medium-incidence countries?'

Findings of this review and the previously published review series<sup>4,13</sup> formed the base for the guidance document by the European Centre for Disease Prevention and Control (ECDC) on controlling TB in hard-to-reach and vulnerable populations.<sup>14</sup>

### Methods

1  
2  
3 In 2011, the Matrix Knowledge Group published a review, commissioned by the National Institute for  
4 Health and Clinical Excellence (NICE), on effectiveness and cost-effectiveness of service models or  
5 structures, focussing on the type of health care worker and setting, to identify and manage TB in hard-  
6 to-reach populations. We updated and extended the NICE review,<sup>15</sup> using the same methodology but  
7 adjusting the focus by excluding latent TB infection and including additional hard-to-reach  
8 populations. The review was conducted following standards described by the Cochrane Collaboration<sup>16</sup>  
9 and NICE methods guidelines.<sup>17</sup> Results are reported according to PRISMA guidelines.<sup>18</sup> The review  
10 protocol was registered in advance in the database of prospectively registered systematic reviews in  
11 health and social care, PROSPERO (CRD42015017865).  
12  
13  
14  
15

### 16 *Selection of studies and data management*

17 The same search strategy as for the previous NICE review<sup>15</sup> [and the previous published review by](#)  
18 [Heuvelings et al.](#)<sup>13</sup> was used, searching Embase and MEDLINE through the Ovid platform. The search  
19 was expanded by including all European Union (EU)/European Economic Area (EEA) and EU  
20 candidate countries to the Organisation for Economic Co-operation and Development (OECD)  
21 countries (see **Box 1**).<sup>15</sup> Two hard-to-reach populations (people living with HIV and children within  
22 vulnerable and hard-to-reach populations), were added in addition to the hard-to-reach populations  
23 included by the NICE review (migrants including refugees, asylum seekers and the Roma population,  
24 homeless people including rough sleepers and shelter users, drug users, prisoners and sex workers).<sup>15</sup>  
25 The update of the search conducted for the NICE review<sup>15</sup> covered the period 1 January 2010  
26 (overlapping the end of the search period of the NICE review<sup>15</sup> with a few months) to 24 February  
27 2017. The search for the expanded geographical area and newly included hard-to-reach populations  
28 covered a time period from 1 January 1990 (beginning of the search period used in the NICE review<sup>15</sup>)  
29 to 24 February 2017.  
30  
31  
32  
33  
34

35 Reference lists of relevant systematic reviews were scanned. No language restrictions were applied.  
36 Studies focussing on the effectiveness and/or cost-effectiveness of interventions for service models and  
37 organisational structures supporting TB identification and management of hard-to-reach populations  
38 (see **Box 1**) were included.  
39

40 Predefined interventions were using more convenient locations (like specialised TB centres, shelters  
41 for homeless people or drug users, needle exchange/methadone programme locations, port of arrival,  
42 schools or mobile clinics) and peers or health care workers with the same ethnic or cultural  
43 background; however, other interventions could also be included if they supported TB identification or  
44 management in hard-to-reach populations. [TB identification tools, TB diagnostics, incentives, social](#)  
45 [support, directly observed therapy and treatment of co-morbidities are discussed in another review.](#)<sup>13</sup> [In](#)  
46 [this review we aim to identify the effectiveness of the type of health worker and setting, to identify and](#)  
47 [manage TB in hard to reach and vulnerable populations.](#)  
48  
49  
50  
51

52 [The comparator was defined during the review process: interventions were compared to a relevant](#)  
53 [comparator, for example usual care or no intervention, another intervention, or historical comparison.](#)  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Outcomes were defined as any measure of TB identification and management (for example, number of  
4 people screened, screening coverage, proportion receiving treatment and treatment completion rate).  
5 Effectiveness was defined as an improvement in any measure of TB identification and/or management.  
6 Randomised and non-randomised studies were eligible for inclusion.  
7

8 See **Supplementary Material I** for the **PROSPERO study protocol**, **Supplementary Material II**  
9 **for** PICOS (Population-Intervention-Comparator-Outcome-Study design) questions and  
10 **Supplementary Material III** for the complete search strategy and search results.  
11

#### 12 *Data extraction, data items, and synthesis*

13  
14 Identified citations were entered into an EndNote database, and duplicates removed (EndNote X7.1,  
15 Thomson Reuters 2014). The inclusion criteria were piloted and refined using the first 25 citations.  
16 Double screening was conducted by one reviewer screening 100% of the citations (CCH) while another  
17 two reviewers screened 50% of the citations each (PFG, SGdV) for inclusion on title and abstract.  
18 Disagreement was resolved by discussion. Full text files of included citations were retrieved;  
19 irretrievable articles (not available after attempts online, from the university library or through  
20 contacting authors) were excluded. Two reviewers assessed full text records for inclusion (CCH, PFG).  
21 Disagreement was resolved by discussion. Agreement after screening on title and abstract was 99.6%  
22 with an inter-rater reliability (Cohen's kappa) of  $\kappa = 0.985$ .  
23  
24  
25  
26  
27

28 Data extraction forms from the NICE review<sup>15</sup> were used to extract information on participant  
29 characteristics, settings, types of services/organisational structures, types of healthcare workers  
30 delivering the service, outcome measures, methods of analysis and results. For one study data  
31 extraction was conducted by two reviewers (CCH, PFG) independently. For the remaining studies, data  
32 extraction was conducted by one reviewer (CCH) and checked by a second (PFG); disagreement was  
33 resolved by discussion. In one case, the study author was contacted to verify data and obtain additional  
34 data.<sup>19</sup>  
35  
36  
37

38 To facilitate comparability, data synthesis was structured in a similar way to that of the NICE review.<sup>15</sup>  
39 Studies were divided into those examining service models and organisational structures for TB  
40 identification (screening) and those examining service models and organisational structures for TB  
41 management (treatment and support) in hard-to-reach populations. Data were analysed narratively, and  
42 appropriateness of meta-analysis considered. Findings were reported as stated by the study authors.  
43  
44  
45

#### 46 *Risk of bias in individual studies and overall strength of evidence*

47 The modified NICE Quality Assessment Tools<sup>17</sup> (based on the Graphical Appraisal Tool for  
48 Epidemiological studies) were used to assess quality and risk of bias of included studies. This included  
49 an assessment of selection of study sample, minimisation of selection bias and contamination,  
50 controlling confounding, outcome measurements, analytical methods and risk of bias. Two reviewers  
51 (CCH, PFG) assessed one study independently; the remaining studies were assessed by one reviewer  
52 (CCH), and checked by a second reviewer (PFG). Any disagreement was resolved by discussion.  
53 Studies were given a quality rating based on the quality assessment: high quality [++], medium quality  
54  
55  
56  
57  
58  
59  
60

[+], or low quality [-]. The strength of the evidence was assessed and reported as described in the previous NICE review<sup>15</sup> (**Supplementary Material IV**).

## Results

Of the 19,720 citations identified by the literature search five studies were included in this review (**Figure 1**).<sup>11,19-22</sup> These five studies are in addition to the six studies<sup>23-28</sup> included in the NICE review.<sup>15</sup> The results section in this paper focuses on the evidence of the five studies identified in our updated review. The evidence statements (presented in **Supplementary Material IV**) summarise evidence identified in terms of consistency, quality and applicability, combining evidence from the NICE review<sup>15</sup> and this update.

All five studies were conducted in the EU; two in the United Kingdom (UK),<sup>19,22</sup> one in Germany,<sup>20</sup> one in Portugal<sup>11</sup> and one in Spain.<sup>21</sup> Two studies focussed on homeless people,<sup>19,20</sup> one on homeless people and drug users,<sup>22</sup> one on drug users alone<sup>11</sup> and one on migrants.<sup>21</sup> Four studies<sup>5,19-21</sup> addressed the influence of the type of healthcare worker on TB identification and TB management and one study focussed on the influence of different settings on TB identification.<sup>22</sup> A variety of study designs were included; one study was a prospective cluster randomised controlled trial (RCT),<sup>19</sup> one was an economic evaluation using a compartmental model of treated and untreated active TB cases<sup>22</sup>, and three studies were retrospective comparison studies.<sup>11,20,21</sup> Study characteristics of included studies are described in **Table 1**. The data extraction forms by study are presented in **Supplementary Material V**.

None of the included studies in this review had a low risk of bias, three studies<sup>19,21,22</sup> had a medium risk of bias, the other two studies<sup>5,20</sup> were assessed as having a high risk of bias (**Supplementary Material VI**).

We did not perform a meta-analysis due to study heterogeneity. Results were synthesised narratively.<sup>29</sup>

Main outcomes for services structures and organisational models for TB identification among hard-to-reach populations, combined with the findings of the NICE review,<sup>15</sup> are summarised in **Table 2**. For full evidence statements, see **Supplementary Material IV**.

Three studies<sup>19-21</sup> compared the effect of the type of healthcare worker on TB identification.

In the UK, a cluster-randomised trial found that peer educators working together with shelter staff to encourage homeless people to participate in a TB screening programme using mobile X-ray units (MXU), did not improve screening uptake compared to encouragement by shelter staff only (respectively 40%, interquartile range [IQR] 25-61 versus 45%, IQR 33-55; adjusted risk ratio [aRR] 0.98, 95% confidence interval [95% CI] 0.80 to 1.20).<sup>19</sup> Control sites were not 'naïve' for peer intervention which could have caused contamination of the control sites and contributed to the negative finding.

1  
2  
3 In Germany, introduction of TB education and promotion of voluntary chest X-ray (CXR) screening at  
4 least once every two years by community health workers (CHWs) improved screening uptake in  
5 homeless people and drug users. Annual screening coverage increased from 10.0% at the beginning of  
6 the study period (2002-2004) to 15.0% during the middle part of the study period (2004-2006), the last  
7 part of the study period had a 13.4% annual screening coverage (2005-2007). Screening once every  
8 two years increased screening coverage from 18.0% (2002-2004) to 26.4% (2004-2006). Coverage was  
9 23.4% at the third and final study period (spanning 2005-2007).<sup>20</sup> The authors did not test for statistical  
10 significance, and denominator data (the number of homeless people and drug users in the study area)  
11 were estimated.  
12  
13  
14  
15

16 In Barcelona, Spain, contact tracing organised by CHWs coming from the same migrant community as  
17 the person diagnosed with TB, improved contact tracing among migrants to 66.2% (2003-2005)  
18 compared to 55.4% (2000-2002) in the period before the implementation of the intervention using  
19 CHWs (adjusted odds ratio [aOR] of an index case having their contacts screened before and after the  
20 intervention was 1.8, 95% CI 1.3 to 2.5,  $p < 0.001$ ).<sup>21</sup> Identification and tracing of at least one contact  
21 was taken as appropriate contact tracing, where all contacts at risk should be traced to detect and treat  
22 TB transmission early. The population characteristics varied, the age and country of origin was  
23 different between both periods. The importance of contact tracing is to identify cases early to reduce  
24 transmission, the authors did not report if any of the contacts traced had active TB.  
25  
26  
27  
28

29 Two studies<sup>11,22</sup> evaluated the effect of the type of healthcare worker and the setting on TB  
30 identification and TB management.  
31

32 In Portugal, improved co-operation of 'key partners' (street teams, TB clinics, drug user support  
33 centres, local public health department and local hospital) for TB identification and management in  
34 drug users was evaluated in a before-and-after study. Representatives of all 'key partners' (authors'  
35 term) worked on improving policies, clinic screening procedures and co-operation. Key partners were  
36 trained in identifying drug users in their population; and offering health promotion, notification cards,  
37 free transport to the TB clinic, free medical and substance abuse care, directly observed therapy (DOT)  
38 for active TB cases, identification of non-compliant patients and the cause of non-compliance, and  
39 tailor-made strategies to improve compliance. This resulted in an increase of TB screening uptake;  
40 from 52 drug users being screened before the intervention (2001-2003 when there was no active  
41 screening policy) to 465 drug users screened thereafter (2005-2007). Of all people misusing drugs  
42 taking-up screening, the proportion without TB symptoms increased from 41.6% to 93.5% (OR =  
43 21.76; 95% CI 13.03 – 36.33) indicating improved TB awareness and access to screening facilities for  
44 drug users. Of all drug users with active TB, the proportion identified by screening increased from  
45 13.4% to 61.0% (OR 10.1; 95% CI: 4.44 – 23.0). Treatment default rates decreased from 35.4% to  
46 10.2% (OR 0.21, 95% CI 0.08-0.54), compared to the period before the intervention (2001-2003)  
47 when TB treatment was not compulsory and compliance was stimulated by TB education and  
48 providing information on the importance of treatment completion.<sup>11</sup> Although the absolute number of  
49 drug users screened increased, information on the screening coverage was not available as denominator  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 data were not provided. Another limitation is that the results were not adjusted for confounding factors,  
4 baseline characteristics might have been different as the two cohorts were recruited over different time  
5 periods and participation was voluntary which may have led to selection bias.  
6  
7

8 In the United Kingdom, the effectiveness and cost-effectiveness of the 'Find and Treat' service (raising  
9 awareness of TB screening and providing a mobile TB screening and treatment service) for homeless  
10 people and drug users was evaluated and compared to people (with a history of homelessness,  
11 imprisonment, drug abuse or mental health problems) self-presenting to a London TB clinic receiving  
12 standard TB care at the clinic.<sup>22</sup> The authors estimated that 22.9% of the patients detected by the 'Find  
13 and Treat' service with the longest first symptom-to-detection time would not have self-presented plus  
14 35.4% were asymptomatic at time of detection and would not have self-presented, only part of the  
15 asymptomatic patients would self-present to a TB clinic at a later stage when symptoms would have  
16 developed. The 'Find and Treat' service had a higher treatment completion rate (67.1% versus 56.8%),  
17 and a lower lost to follow-up rate (2.1% versus 17.2%) compared to the control group receiving  
18 standard TB care at a TB clinic. The authors concluded that the 'Find and Treat' service was cost-  
19 effective, when using the threshold used by NICE of £20,000 to £30,000/QALY gained, with an  
20 incremental cost ratio of £18,000 per QALY gained for the TB screening service and £4,100 per  
21 QALY gained for the TB management service. This study has a few limitations: firstly, it is a non-  
22 randomised study, secondly the 'Find and Treat' service identifies extremely hard-to-reach populations  
23 of which some would never self-present therefore the findings could be even better in less hard-to-  
24 reach populations, and thirdly the economical evaluation is based on a compartmental model that does  
25 not take secondary transmission and drug-resistance into account.  
26  
27  
28  
29  
30  
31  
32

### 33 **Discussion**

34 To tackle TB and disrupt transmission in high-income, low TB incidence settings, improvement of TB  
35 care in hard-to-reach populations is of vital importance. In this updated review, five studies,<sup>11,19-22</sup>  
36 published between 1 January 2010 and 24 February 2017, evaluating effectiveness of services models  
37 and organisational structures supporting TB identification and management of hard-to-reach  
38 populations, were identified in addition to the six studies considering active TB<sup>23-28</sup> identified by the  
39 NICE review.<sup>15</sup> Only one study<sup>22</sup> evaluated cost-effectiveness. Although the evidence from [two](#)  
40 [reviews](#) is limited, it highlights those interventions that are likely to be effective and those that have no  
41 clear evidence of being effective (**Table 2**). For development of the ECDC guidance document,<sup>14</sup> a  
42 scientific panel compiled by ECDC carefully considered these findings. Their main suggestions for  
43 action were to involve CHWs or peers to improve TB screening uptake and TB treatment completion  
44 among homeless people<sup>20</sup> and drug users;<sup>5,20,23</sup> to use outreach teams to improve TB screening uptake  
45 and TB treatment completion among vulnerable populations;<sup>22</sup> and to strengthen relationships and good  
46 collaboration between health care workers, peers, communities, and patients to improve treatment  
47 outcome among vulnerable populations.<sup>5,20,22,23</sup> The updated systematic review provided evidence for  
48 all suggestions except for using peers to improve screening uptake. [This is in contrast to](#) an American  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



study<sup>23</sup> included in the original NICE review,<sup>15</sup> [which](#) showed that peers improved contact tracing and treatment adherence among drug users.

### *Strengths and limitations*

PRISMA and Cochrane Collaboration reporting guidelines for systematic reviews were followed. Established screening protocols were used, including double screening, and the search was highly sensitive. The methodology from the previous NICE review<sup>15</sup> was followed, in order to connect this update and, so, [describe](#) the full body of relevant evidence. High quality evidence is lacking. Only one<sup>23</sup> study from the NICE review<sup>15</sup> was considered to be of high quality, all other studies had some risk of bias (five medium risk<sup>19,21,22,24,26</sup> and five high risk<sup>11,20,25,27,28</sup>). Therefore, only limited conclusions can be drawn. Most studies lacked identification and adjustment for confounding factors and the use of appropriate analytical methods. In addition, many studies were biased, particularly with regard to potential selection bias. A meta-analysis could not be performed, because of heterogeneity across the studies. Gaps in evidence exist; no studies focussing on children within vulnerable and hard-to-reach populations, or on people living with HIV or sex workers were identified. Only three studies provided economic data; one study identified by this review,<sup>22</sup> and two<sup>25,27</sup> by the NICE review.<sup>15</sup>

Our search focussed on publications in databases Embase and MEDLINE. Many European countries have strong organisational structures for TB identification and management, but these countries did not publish their data on these organisational structures in journals, which may have caused a publication bias. [Comparing](#) findings of the NICE review<sup>15</sup> with this review comes with some limitations. For the NICE review only 10% of the citations were double screened,<sup>15</sup> compared to 100% for this updated review, therefore studies conducted between 1990-2010 might have been missed. The NICE review focussed their recommendations on the population in the United Kingdom,<sup>15</sup> and this review focussed on populations in high-income, low TB incidence countries. Further methodology was identical.

The evidence identified by this review and the previous NICE review<sup>15</sup> along with evidence presented in a review series covering the barriers and facilitators of seeking TB care,<sup>6</sup> and the effectiveness of interventions for TB identification and management in hard-to-reach populations,<sup>13</sup> was used to develop the ECDC guidance on improving TB identification and management among hard-to-reach and vulnerable populations in Europe.<sup>14</sup> ECDC recommended that implementation of the interventions is context-specific; it depends on the setting, target population, resources available and health-care systems in place. Interventions focussing on one specific hard-to-reach population might not work in another hard-to-reach population, therefore, the interventions have to be adapted and re-assessed per target population.<sup>14</sup> Given the scope of this review, considering settings across Europe, findings presented here are potentially relevant to any low-incidence region, and are relevant to other institutions/governmental organisations seeking to improve service structures for TB identification and management among hard-to-reach populations.

Characteristics of different hard-to-reach populations and their TB epidemiology vary per country and setting. Challenges in identification and management of TB should be identified and targeted, tailored

1  
2  
3 to the specific setting and hard-to-reach population. These TB interventions could be integrated within  
4 broader programmes targeting specific populations. A follow-up systematic review should include  
5 information from national public health services about their organisational structures for TB  
6 identification and management. National public health services are urged to regularly analyse their  
7 organisational structures for TB identification and management and publish this data.

8  
9 Efforts to improve quality of research on service models and organisational structures should be made,  
10 even though it is often challenging to perform 'clean', unbiased, and un-confounded trials in hard-to-  
11 reach populations, as attrition rates are often high, and confounding factors are plentiful. This includes  
12 conducting (cluster) randomised controlled trials and before-and-after studies where appropriate,  
13 recruiting an adequate number of participants, using relevant control groups, and minimising selection  
14 bias. Standardised case definitions for hard-to-reach populations should be created. Feasibility,  
15 effectiveness, cost-effectiveness and impact of interventions should be evaluated. Mathematical  
16 economic models can be used to evaluate costs.<sup>14</sup>

### 21 **Conclusions**

22 Identification and management of TB in hard-to-reach populations is suboptimal.<sup>2</sup> Therefore, service  
23 models and organisational structures to identify and manage TB in hard-to-reach populations should be  
24 improved and evaluated regularly.

25  
26  
27 Our systematic review, in conjunction with the original NICE review<sup>15</sup> provides limited evidence, due  
28 to the lack of high quality studies, that interventions such as using peers and CHWs; mobile TB  
29 services, specialised TB clinics, screening, or active case finding in non-healthcare settings, as well as  
30 improved co-operation between key services can help to improve TB identification and management.

31 Further research should be undertaken to evaluate other effective and cost-effective ways to identify  
32 and manage TB in hard-to-reach populations and countries with good TB control systems are urged to  
33 evaluate their system and publish the data.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Competing interests

All authors have completed the ICMJE uniform disclosure form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) and all except for MPG and BS declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work. MPG reports grants from ECDC, for the conduct of part of the study. BS reports that NICE - her employing organisation - has published guidance in this area.

### Authors' contribution

RS conducted the literature search. CCH, SGdV and PFG performed the study selection. CCH, and PFG collected the data, performed quality/risk of bias assessment and synthesised the data. CCH prepared the manuscript and supplementary files. MPG and MvdW supervised the whole process and contributed to the work. All authors were involved in interpretation of the data, and the writing of the manuscript. All authors contributed to, and endorsed the final version of the manuscript. MPG is guarantor.

### Ethics

Ethics approval was not required for this systematic review.

### Funding

This study was funded by the European Centre for Disease Prevention and Control (contract reference OJ/02/05/2014-PROC/2014/014). None of the authors have received payment from a pharmaceutical company or other agency to write this article.

### Statement of independence of researchers from funders

The funder of the study was involved in study design, data interpretation, and reporting. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

### Data sharing


Data extraction forms and quality assessment forms are available from **Supplementary files V and VI**.

## References

1. World Health Organization (WHO). Global Tuberculosis report 2016. Geneva: WHO 2016 (WHO/HTM/TB/2016.13).
2. World Health Organization (WHO) and European Respiratory Society. Towards TB elimination: an action framework for low-incidence countries. Geneva; WHO 2014 (WHO/HTM/TB/2014.13).
3. de Vries G, Aldridge RW, Cayla JA, et al. Epidemiology of tuberculosis in big cities of the European Union and European Economic Area countries. *Euro Surveill* 2014; **19**(9).
4. de Vries SG, Cremers AL, Heuvelings CC, et al. (in press). Barriers and facilitators for the uptake of tuberculosis diagnostic and treatment services by hard-to-reach populations in low- and medium-incidence countries: A systematic review of qualitative literature. *Lancet Infect Dis* 2017; published online March 17. [http://dx.doi.org/10.1016/S1473-3099\(16\)30531-X](http://dx.doi.org/10.1016/S1473-3099(16)30531-X).
5. Kamholz SL. Resurgence of tuberculosis: the perspective a dozen years later. *J Assoc Academic Minor Phys* 1996; **7**(3): 83-6.
6. de Vries G, van Hest RA, Richardus JH. Impact of mobile radiographic screening on tuberculosis among drug users and homeless persons. *Am J Respir Crit Care Med* 2007; **176**(2): 201-7.
7. van Hest NA, Aldridge RW, de Vries G, et al. Tuberculosis control in big cities and urban risk groups in the European Union: a consensus statement. *Euro surveill* 2014; **19**(9).
8. Belling R, McLaren S, Boudioni M, Woods L. Pan-London tuberculosis services: a service evaluation. *BMC Health Serv Res* 2012; **12**: 203.
9. Ohkado A, Williams G, Ishikawa N, Shimouchi A, Simon C. The management for tuberculosis control in Greater London in comparison with that in Osaka City: lessons for improvement of TB control management in Osaka City urban setting. *Health Policy* 2005; **73**(1): 104-23.
10. Lei X, Liu Q, Escobar E, et al. Public-private mix for tuberculosis care and control: a systematic review. *Int J Tuberc Lung Dis* 2015; **34**: 20-32.
11. Duarte R, Santos A, Mota M, Carvalho A, Marques A, Barros H. Involving community partners in the management of tuberculosis among drug users. *Public Health* 2011; **125**(1): 60-2.
12. Frieden TR, Fujiwara PI, Washko RM, Hamburg MA. Tuberculosis in New York City--turning the tide. *N Engl J Med* 1995; **333**(4): 229-33.
13. Heuvelings C, de Vries S, Greve PF, et al. (in press). Effectiveness of interventions aiming at identifying and managing tuberculosis among hard-to-reach populations: A systematic review. *Lancet Infect Dis* 2017; published online March 17. [http://dx.doi.org/10.1016/S1473-3099\(16\)30532-1](http://dx.doi.org/10.1016/S1473-3099(16)30532-1).
14. European Centre for Disease Prevention and Control (ECDC). Guidance on tuberculosis control in vulnerable and hard-to-reach populations. Stockholm: ECDC 2016.
15. Rizzo M, Martin A, Jamal F, et al. Evidence review on the effectiveness and cost-effectiveness of service models or structures to manage tuberculosis in hard-to-reach groups. Matrix evidence: NICE – National Institute for Health and Clinical Excellence, 2011.
16. Higgins J, Green S, (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]: The Cochrane Collaboration; 2011.
17. National Institute for Health and Clinical Excellence, London. Methods for the development of NICE public health guidance (second edition). 2009.
18. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009; **339**: b2535.
19. Aldridge RW, Hayward AC, Hemming S, et al. Effectiveness of peer educators on the uptake of mobile X-ray tuberculosis screening at homeless hostels: a cluster randomised controlled trial. *BMJ Open* 2015; **5**(9): e008050.
20. Goetsch U, Bellinger OK, Buettel KL, Gottschalk R. Tuberculosis among drug users and homeless persons: impact of voluntary X-ray investigation on active case finding. *Infection* 2012; **40**(4): 389-95.

- 1  
2  
3 21. Ospina JE, Orcau A, Millet JP, Sanchez F, Casals M, Cayla JA. Community health workers  
4 improve contact tracing among immigrants with tuberculosis in Barcelona. *BMC Public Health* 2012;  
5 **12**: 158.
- 6 22. Jit M, Stagg H, Aldridge R, White P, Abubakar I. Dedicated outreach service for hard to reach  
7 patients with tuberculosis in London: observational study and economic evaluation. *BMJ* 2011; **343**:  
8 d5376.
- 9 23. Ricks PM. Tuberculosis control among substance users: The indigenous leadership outreach  
10 model vs. standard care. Chicago, IL: University of Illinois, 2008.
- 11 24. El-Hamad I, Casalini C, Matteelli A, et al. Screening for tuberculosis and latent tuberculosis  
12 infection among undocumented immigrants at an unspecialised health service unit. *Int J Tuberc Lung*  
13 *Dis* 2001; **5**(8): 712-6.
- 14 25. Bothamley GH, Rowan JP, Griffiths CJ, et al. Screening for tuberculosis: the port of arrival  
15 scheme compared with screening in general practice and the homeless *Thorax* 2002; **57**(1): 45-9.
- 16 26. Miller TL, Hilsenrath P, Lykens K, McNabb SJ, Moonan PK, Weis SE. Using cost and health  
17 impacts to prioritize the targeted testing of tuberculosis in the United States. *Ann Epidemiol* 2006;  
18 **16**(4): 305-12.
- 19 27. Mor Z, Lerman Y, Leventhal A. Pre-immigration screening process and pulmonary  
20 tuberculosis among Ethiopian migrants in Israel. *Eur Respir J* 2008; **32**(2): 413 - 8.
- 21 28. Dèruaz J, Zellweger J, P. Directly observed therapy for tuberculosis in a low prevalence  
22 region: first experience at the Tuberculosis Dispensary in Lausanne. *Swiss Med Wkly* 2004; **134**: 552-8.
- 23 29. Popay J, Roberts H, Sowden A, et al. Guidance on the conduct of narrative synthesis in  
24 systematic reviews. A product from the ESRC methods programme *Lancaster* 2006.
- 25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1. Study selection process



For the Organisation for Economic Co-operation and Development (OECD) countries and hard-to-reach populations discussed in the previous NICE review,<sup>13</sup> the study period covers 1 January 2010 to 24 February 2017. For the newly included European Union (EU)/European Economic Area/EU candidate countries, and the newly included hard-to-reach populations (people living with Human Immunodeficiency Virus (HIV) and children within vulnerable and hard-to-reach populations, the study period covers 1 January 1990 to 24 February 2017.

**Table 1. Characteristics of studies applying different service models and organisational structures to improve TB identification and TB management**

| First Author (year) country                                  | Population                     | Aims  | Intervention   | Comparator   | Study design                               | Outcome measure  | Quality score |
|--|--------------------------------|---|--|--|--|--|---------------|
| <b>TB identification (studies identified by this review)</b> |                                |   |  |  |  |  |               |
| Jit et al. (2011) <sup>22</sup><br>United Kingdom            | Homeless people and drug users | To assess the effectiveness and cost-effectiveness of the Find and Treat service for diagnosing and managing hard-to-reach individuals with active TB in London.  | Period 2007-2010: Find and Treat service:<br>- screening by MXU<br>- peers raising awareness<br>- treatment support  | Passive case detection and standard treatment at a London TB clinic  | Observational and cost-effectiveness study | Identified TB cases, treatment completion, lost to follow-up, incremental costs from healthcare taxpayer perspective | +             |
| Duarte et al. (2011) <sup>11</sup><br>Portugal               | Drug users                     | To evaluate the effect of an intervention with key partners (TB clinic, drug users support centres, shelters, street teams, public health department and hospital) delivering promotion of health-seeking behaviour, eliminating potential barriers for TB screening at a chest clinic and DOT, on identifying TB cases and treatment compliance. | Improved co-operation of key partners (2005-2007):<br>- health education and screening promotion<br>- improved screening procedures<br>- implementation of DOT<br>- free TB care and transport<br>- providing medical and drug abuse treatment<br>- active follow-up of non-compliant patients, the key partners worked together to reach the patient, identify the cause and organise suitable treatment strategies | Period before the intervention (2001-2003):<br><br>- no active screening policy<br>- referral to chest clinic after discharge from hospital<br>- treatment not compulsory<br>- information about disease and treatment given to improve compliance<br>- psychosocial support<br>- free TB treatment, transport and breakfast | Before-after study                         | Identified TB cases and treatment compliance   | -             |
| Goetsch et al. (2012) <sup>20</sup><br>Germany               | Homeless people and drug users | To estimate the coverage of a low-threshold CXR screening programme for pulmonary TB among illicit drug users and homeless persons.   | CHWs providing TB education and promoting voluntary CXR screening 1-2x/year  | Comparing the beginning of the 5 year intervention period with the end (2002-2007)   | Retrospective effectiveness study          | Screening coverage   | -             |
| Ospina et al. (2012) <sup>21</sup><br>Spain                  | Migrants                       | To evaluate the effectiveness of an intervention with community health workers to improve contact tracing among migrants  | CHWs active follow up of cases and contacts, including visits of the cases at home, accompanying at outpatient appointments, providing counselling and information on treatments (2003-2005)   | Pre-intervention period (2000-2002)  | Before-after study                         | Number of migrants that were included in contact tracing   | +             |

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |                 |   |   |   |                                     |                  |   |
|--|-----------------|---|---|---|-------------------------------------|------------------|---|
| Aldridge et al. (2015) <sup>19</sup><br>United Kingdom | Homeless people | To compare TB screening uptake between current practice of encouraging homeless people by shelter staff and encouragement by shelter staff plus volunteer peer educators. | Encouragement of TB screening by peers in addition to shelter staff | Encouragement of TB screening by shelter staff only | Cluster randomised controlled trial | Screening uptake | + |
|--|-----------------|---|---|---|-------------------------------------|------------------|---|

**TB identification (studies identified by the previous NICE review<sup>15</sup>)**

|   |  |  |  |   |   |   |    |
|---|--|--|--|---|---|---|----|
| El-Hamad et al. (2001) <sup>24</sup><br>Italy           | Migrants   | To compare the completion rates of screening procedures for TB infection among undocumented migrants at specialised TB units and non-specialised health clinics                          | TB screening at specialised TB clinic                              | TB screening at a general health service for migrants   | Prospective cohort                                      | Screening completion                              | +  |
| Bothamley et al. (2002) <sup>25</sup><br>United Kingdom | Migrants and homeless people                                   | To compare the yield and costs of TB screening in three settings: a new entrants' clinic within the port of arrival (POA) scheme; a large general practice; and centres for the homeless | TB screening at a general practice (GP)                            | TB screening at POA and at homeless centres   | Cost analysis   | Cost per person screened per case of TB prevented | -  |
| Deruaz & Zellweger (2004) <sup>28</sup><br>Switzerland  | Migrants, alcohol or drug users, homeless people and prisoners | Evaluation of first experience of the DOT programme for TB introduced in the Canton of Vaud in 1997  | 1. Full DOT<br>2. DOT delivered at TB clinic                       | 1. Partial DOT (DOT only first 2 months of treatment)<br>2. DOT delivered at social outreach site | Before-after study                                      | Adherence to treatment and outcome                | -  |
| Miller et al. (2006) <sup>26</sup><br>United States     | Homeless people and prisoners                                  | To evaluate and compare the efficiency of a non-state-law-mandated TB screening programme for homeless persons with a state-law-mandated TB screening programme for prisoners            | Non-state-law-mandated TB screening programme for homeless persons | State-law-mandated TB screening programme for prisoners   | Retrospective comparison of the cost and health impacts | TB cases averted and cost                         | +  |
| Ricks (2008) <sup>23</sup><br>United States             | Drug users   | To compare the effectiveness of using peers versus 'standard' public health workers to coordinate TB treatment   | Enhanced case management by peers                                  | Limited case management by health care professionals  | Randomised controlled trial                             | Adherence to treatment                            | ++ |



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|   |          |  |                         |                          |                               |   |   |
|---|----------|--|-------------------------|--------------------------|-------------------------------|---|---|
| Mor et al. (2008) <sup>27</sup><br>Israel | Migrants | To examine the effectiveness and cost-effectiveness of pre-migration screening and post-migration screening at POA | Pre-migration screening | Post-migration screening | Retrospective cohort analysis | Active TB cases, time between migration and diagnosis, cost-savings | - |
|---|----------|--|-------------------------|--------------------------|-------------------------------|---|---|

**List of Abbreviations**

CHW = community health worker; CXR = chest X-ray; C = comparator group; DOT = direct observed treatment; GP = general practice; I = intervention group; IRIS = immune reconstitution inflammatory syndrome; IVDU = intravenous drug users; MXU = mobile X-ray unit; *n* = number of participants; QALYs = quality adjusted life years; POA = port-of-arrival; RCT = randomised controlled trial; TDM = therapeutic drug monitoring; TB = tuberculosis

Study quality: high quality [++], medium quality [+], or low quality [-]

For peer review only

**Table 2. Effectiveness of service models and organisational structures interventions to improve TB identification and TB management**

| <u>Population</u>                               | <u>Intervention (I)</u>  | <u>Comparator (C)</u>  | <u>Studies (first author, year, country)</u>   | <u>No. of participants</u><br><u>I            C</u> |             | <u>Comparison</u>  | <u>Outcome</u>   | <u>Risk of Bias</u>                                    |
|---|--|--|--|---|-------------|--|--|--|
| <u>Homeless people</u><br><br><u>Drug users</u> | <u>Health/TB education and promotion of screening by street teams, drug users support centres, shelters and CHWs</u> | <u>Beginning of the intervention when CHWs were just introduced</u><br><br><u>No active screening policy</u> | <u>Goetsch et al. 2012 (Germany)<sup>20</sup></u><br><br><u>Duarte et al. 2011 (Portugal)<sup>11</sup></u> | <u>465</u>  | <u>125</u>  | <u>Retrospective comparison over intervention period</u><br><br><u>Retrospective before-after comparison</u> | <u>Improved annual TB screening uptake among homeless people and drug users (from 10.0% to 15.0% at the peak).<sup>20</sup> The percentage of all drug users with active TB identified by screening increased from 13.4% to 61.0% (OR 10.1 [95%CI 4-44-23-0]).<sup>11</sup></u>  | <u>High<sub>2</sub></u><br><br><u>High<sub>3</sub></u> |
| <u>Homeless people</u>                          | <u>TB education and promotion of screening by peers and shelter staff</u>  | <u>TB education and promotion of screening by shelter staff only</u>   | <u>Aldridge et al. 2015 (United Kingdom)<sup>19</sup></u>  | <u>1150</u>   | <u>1192</u> | <u>Comparing randomised intervention cluster with comparator cluster</u>                                     | <u>No difference in screening uptake (I = 40% [IQR 25-61] versus C = 45% [IQR 33-55], aRR= 0.98 [95%CI 0.80-1.20]).</u>  | <u>Medium<sub>4</sub></u>                              |
| <u>Migrants</u>                                 | <u>Pre-migration screening</u>   | <u>Post-migration screening at POA</u>   | <u>Mor et al. 2008, cited in the NICE review (Israel)<sup>27</sup></u>                                     | <u>162</u>  | <u>105</u>  | <u>Retrospective Intervention versus comparator comparison</u>   | <u>Reduced the risk of developing TB in the new country and was cost-effective (0.28% of the pre-migration versus 0.32% of the post-migration screening migrants developed TB; RR 0.82 p&lt;0.01). The detection period was shorter as well (193 days versus 487 days between entry and diagnosis; OR=0.72 [95%CI 0.59-0.89] p=0.002).</u> | <u>High<sub>8</sub></u>                                |
| <u>Prisoners and homeless people</u>            | <u>TB screening in a prison</u>  | <u>TB screening at a homeless centre</u>   | <u>Miller et al. 2006, cited in the NICE review (United States)<sup>26</sup></u>                           | <u>22920</u>  | <u>822</u>  | <u>Retrospective comparison of two cohorts</u>   | <u>No difference in screening uptake (94.7% in prison versus 95% in homeless centre p=0.179) but higher proportion of active TB cases were identified at the homeless centre (1.2% versus 0.03% at a prison setting, p&lt;0.001)</u>   | <u>Medium<sub>9</sub></u>                              |
| <u>Homeless people and migrants</u>             | <u>Active case finding by symptom-based questionnaire at homeless centres</u>  | <u>Active case finding by symptom-based questionnaire at POA</u>   | <u>Bothamley et al. 2002, cited in the NICE review (United Kingdom)<sup>25</sup></u>                       | <u>262</u>  | <u>199</u>  | <u>Cost analysis</u>   | <u>Active case finding at POA was most cost-effective (costs per person screened for every case prevented at POA £10.00, at homeless centre £23.00).</u>   | <u>High<sub>10</sub></u>                               |
| <u>Migrants</u>                                 | <u>Active case finding at a specialised TB clinic using 2 visits</u>   | <u>Active case finding at a general primary care clinic, with referral for CXR, using 3 visits</u>           | <u>El-Hamad et al. 2001, cited in the NICE review (Italy)<sup>24</sup></u>                                 | <u>749</u>  | <u>483</u>  | <u>Prospective intervention versus comparator comparison</u>   | <u>Improved screening completion among migrants (85.6% in TB clinic versus 71.4% at primary care clinic, p = not reported; OR=2.57 [95%CI 1.92-3.42]).</u>   | <u>Medium<sub>5</sub></u>                              |

|   |   |  |  |                     |                     |  |  |                                    |
|---|---|--|--|---------------------|---------------------|--|--|------------------------------------|
| <a href="#">Drug users</a>  | <a href="#">Contact tracing by peers or CHWs from the same migrant community</a>                  | <a href="#">Peers versus other health care workers</a>   | <a href="#">Ricks 2008, cited in the NICE review (United States)<sup>23</sup></a>                | <a href="#">48</a>  | <a href="#">46</a>  | <a href="#">RCT</a>  | <a href="#">Improved contact tracing among drug users (75% by peers versus 47% by healthcare workers, p = 0.03)<sup>23</sup> and migrants (from 55.4% without CHWs to 66.2% with CHWs; aOR 1.8 [95%CI 1.3-2.5] p&lt;0.001).<sup>21</sup></a>   | <a href="#">Low</a>                |
| <a href="#">Migrants</a>  |   | <a href="#">Normal practice before introducing CHWs</a>  | <a href="#">Ospina et al. 2012 (Spain)<sup>21</sup></a>  | <a href="#">388</a> | <a href="#">572</a> | <a href="#">Before-after comparison</a>  |  | <a href="#">Medium<sub>1</sub></a> |
| <a href="#">Drug users and homeless people</a>                      | <a href="#">Mobile TB screening and treatment service at convenient location in the community</a> | <a href="#">Passive case detection and management at a TB clinic</a>   | <a href="#">Jit et al. 2011 (United Kingdom)<sup>22</sup></a>                                    | <a href="#">48</a>  | <a href="#">252</a> | <a href="#">Prospective intervention versus comparator comparison plus economic evaluation</a> | <a href="#">Improved TB identification among homeless people and drug users; particularly in asymptomatic patients (35.4% extra identified) and those that delay seeking health care (22.2% extra identified). Higher treatment completion rate (67.1% versus 56.8%) and lower lost to follow-up rate (2.1% versus 17.2%). Both parts of the service are cost-effective (screening = £18,000/QALY gained, treatment is £4,100/QALY gained)</a> | <a href="#">Medium<sup>6</sup></a> |
| <a href="#">Drug users</a>  | <a href="#">Enhanced case management by peers</a>   | <a href="#">Limited case management by regular health care workers</a>   | <a href="#">Ricks 2008, cited in the NICE review (United States)<sup>23</sup></a>                | <a href="#">48</a>  | <a href="#">46</a>  | <a href="#">RCT</a>  | <a href="#">Improved treatment completion in drug users (85% by peers versus 61% by health care workers, RR=2.68 [95%CI 1.24-5.82] p=0.01).</a>  | <a href="#">Low</a>                |
| <a href="#">Drug users</a>  | <a href="#">DOT and active follow-up of non-compliant patients by 'key partners'</a>              | <a href="#">Non-compulsory TB treatment and education about TB disease and treatment to improve compliance</a> | <a href="#">Duarte et al. 2011 (Portugal)<sup>11</sup></a>                                       | <a href="#">465</a> | <a href="#">125</a> | <a href="#">Retrospective before-after comparison</a>  | <a href="#">Reduced treatment default rates (from 35.4% to 10.2%; OR 0.21 [95%CI 0.08-0.54]).</a>  | <a href="#">High<sub>2</sub></a>   |
| <a href="#">Migrants, drug users, homeless people and prisoners</a> | <a href="#">DOT at a convenient location in the community</a>                                     | <a href="#">DOT at a health clinic</a>   | <a href="#">Deruaz &amp; Zellweger 2004, cited in the NICE review (Switzerland)<sup>28</sup></a> | <a href="#">36</a>  | <a href="#">18</a>  | <a href="#">Retrospective before-after comparison</a>  | <a href="#">No significant difference in successful treatment outcome, treatment completion and cure rate (85.2% at convenient location versus 92.6% at health clinic, p=0.67)</a>   | <a href="#">High<sub>7</sub></a>   |

### List of Abbreviations

aOR = adjusted odds ratio, aRR = adjusted risk ratio, C = comparator group, CHWs = community health workers, DOT = directly observed treatment, I = intervention group, IQR = interquartile range, OR = odds ratio, POA = port-of-arrival, RCT = randomised controlled trial, TB = tuberculosis, 95%CI = 95% confidence interval

### Footnotes Risk of Bias:

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47
- <sub>1</sub> = Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods). Contact tracing of only one contact was enough to be called contact tracing, and the ultimate aim of contact tracing (increase case detection and reduce transmission) was not analysed in this study.
- <sub>2</sub> = Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods). Denominator not given therefore unable to calculate screening coverage.
- <sub>3</sub> = Risk of selection bias as participation was voluntary. Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods). No statistical test used to show statistical significance of the findings, an estimated number was used for the denominator.
- <sub>4</sub> = Most comparator sites were not naïve for peer intervention, no individual information of the participants was collected, the characteristics between the two groups might have been significantly different.
- <sub>5</sub> = Not adjusted for difference in baseline characteristics.
- <sub>6</sub> = Study was designed to evaluate the cost-effectiveness, no statistical test used to evaluate statistical significant findings. The 'Find and Treat' service identifies extremely hard-to-reach populations that would never self-present, the findings would underestimate the benefit of the service. The economical evaluation is based on a compartmental model that does not take secondary transmission and drug-resistance into account,
- <sub>7</sub> = Risk of bias due to difference in collecting treatment adherence outcome at the health clinic a nurse recorded treatment adherence at time of visit, in the social outreach group a health care worker was interviewed up to 6 months after treatment completion and was asked about the treatment adherence, risk of recall bias. Not recorded how many people per setting received 6 months of DOT (full DOT) and how many received 2 months of DOT and 4 months of self-treatment (partial DOT), what was another intervention in this study. Allocation to setting was based on needs of participants what might have caused bias.
- <sub>8</sub> = Not adjusted for important confounding factors (intervention and comparator group were recruited over different time periods), pre-migration group had a shorter follow-up period than post-migration group what may have influenced the detection of number of TB cases in the pre-migration group.
- <sub>9</sub> = Unclear if the differences in outcome was caused by the setting or by the different methods or to differences in TB prevalence in the different populations.
- <sub>10</sub> = TB prevalence might be different in the different populations as the costs are calculated per active case detected this is a major issue, there were only 3 active TB cases detected, all in the POA group. The economic perspective used was not reported and the costs of identification were not discounted.

**Box 1: Inclusion/exclusion criteria for this review**

## Inclusion criteria:

- Discussing service models and organisational structures, different types of healthcare workers and settings for delivering TB services to hard-to-reach populations;
- Having been conducted in any of the EU/EEA countries (only updated review), the candidate countries\* (only updated review) and the other OECD countries\*\*
- Having been published in 2010 or later for the OECD countries\*\*
- Having been published in 1990 or later for the EU/EEA countries and the EU candidate countries\* not being one of the OECD countries (only updated review)
- Including data from any hard-to-reach population:
  - homeless people
  - people who abuse drugs or alcohol
  - sex workers
  - prisoners or people with a history of imprisonment
  - migrants, including vulnerable migrant populations such as asylum seekers, refugees and the Roma population
  - children within vulnerable and hard-to-reach populations (only updated review)
  - people living with HIV (only updated review)
- Present quantitative empirical data
- Being a (cost)-effectiveness study, or any other type of quantitative primary research, discussing (cost)-effectiveness

## Exclusion criteria:

- Latent TB infection (only updated review)
- Systematic review (only used for reference searching)

*EU/EEA = European Union, European Economic Area; OECD = Organisation for Economic Co-operation and Development; TB = Tuberculosis*

*\*EU candidate countries = Albania, Montenegro, Serbia, the former Yugoslav Republic of Macedonia, and Turkey*

*\*\* OECD countries = Australia, Austria, Belgium, Canada, Chile, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Israel, Italy, Japan, Korea, Luxembourg, Mexico, the Netherlands, New Zealand, Norway, Poland, Portugal, Slovak Republic, Slovenia, Spain, Sweden, Switzerland, Turkey, United Kingdom, United States*

1  
2  
3 **Supplementary Material**

4 **Supplementary Material I:** Original study protocol

5  
6 **Supplementary Material II.** PICOS

7  
8 **Supplementary Material III.** Search strategy

9  
10 **Supplementary Material IV.** Evidence statements

11  
12 **Supplementary Material V.** Data extraction forms

13  
14 **Supplementary Material VI.** Quality Assessment

15  
16 **Supplementary Material VII.** PRISMA checklist

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## PROSPERO International prospective register of systematic reviews

### Evidence review of the effectiveness and cost effectiveness of service models or structures aiming to manage tuberculosis in hard-to-reach groups, including a qualitative description of these service models or structures

Charlotte Heuvelings, Sophia de Vries, Patrick Greve, Benjamin Jelle Visser, Saskia Janssen, Sabine Belard, Lianne Cremers, Rene Spijker, Martin Grobusch

#### Citation

Charlotte Heuvelings, Sophia de Vries, Patrick Greve, Benjamin Jelle Visser, Saskia Janssen, Sabine Belard, Lianne Cremers, Rene Spijker, Martin Grobusch. Evidence review of the effectiveness and cost effectiveness of service models or structures aiming to manage tuberculosis in hard-to-reach groups, including a qualitative description of these service models or structures. PROSPERO 2015:CRD42015017865 Available from [http://www.crd.york.ac.uk/PROSPERO\\_REBRANDING/display\\_record.asp?ID=CRD42015017865](http://www.crd.york.ac.uk/PROSPERO_REBRANDING/display_record.asp?ID=CRD42015017865)

#### Review question(s)

Which service models and service structures are most effective and cost-effective at supporting TB identification and management of hard-to-reach groups? How can these service models and organisational structures be qualitatively described?

The following service structures will be explored:

- Type of healthcare worker (any person who was used to deliver the intervention) used to identify and manage TB in hard-to-reach groups.
- Setting used to identify and manage TB in hard-to-reach groups.

Who is responsible for the commissioning and delivery of TB services?

What (if any) theories or conceptual models underpin the service models/organisational structures?

What specific individuals or populations are targeted by the interventions?

How does engagement in various service models/organisational structures differ by group/subgroup (in terms of hard-to-reach group, age, or gender)?

#### Searches

Data Bases Health:

- EMBASE
- MEDLINE
- MEDLINE In-Process

Search terms

1. 'tuberculosis / TBC/ TB' 2. 'hard to reach groups / population / people' 3. 'vulnerable groups / population / people / poor / poverty / overcrowded / excluded populations'
4. 'homeless / hostels / shelters / poor / poverty' 5. 'drug\* / substance\* use\* / abuse\* / addict\* / dependent / problem\* /

disorder\*' 6. 'Alcohol\* use\* / abuse\* / problem\* / addict\*' 7. 'sex-worker / prostitute\* / transactional sex' 8. 'prison\* / inmate / detainee\* / detention / jail / imprisonment' 9. '\*migrant\* / illegal / Roma\* / gips\* / gyps\* / international student / expat\*' 10. 'child\*' 11. 'HIV / AIDS co-infection' 12. 'RCT / \*Controlled Trial' 13. 'Cohort stud\*' 14. '\*review\*' 15. '\*report\*' 16. 'Case stud\* / report\*' 17. 'intervention\*' 18. 'case finding (active or passive)' 19. 'case management / strategy / polic\*' 20. 'prevention' 21. 'control' 22. 'management' 23. 'treat\*' 24. 'Service model\* / delivery / health care service\* / urban health service\*' 25. 'General practice / physician / GP / family practice / nurse lead' 26. 'Social work\* / outreach / volunteer\*' 27. 'Rehab\* centre / pharmacy / ambulatory care / mobile clinic' 28. 'DOTS / directly observed therapy / adherence / patient compliance' 29. 'Telemedicine / telephone medicine' 30. 'Incentives / support' 31. 'Screening / counselling' 32. 'Chest X-ray/ radiography / CXR / diagnostic test / Mantoux / skin test / TST / GeneXpert / sputum smear / culture' 33. 'Health education / promotion' 34. 'Service provider' 35. 'Provider initiated / voluntary / patient initiated'

We will conduct five separate search strategies; the first two search strategies (one qualitative and one quantitative search) will be exactly the same searches as performed for the evidence reviews for the NICE guidelines. For these OECD countries, the searches will span from August 2010 till present (December 2014, with an update performed in the last contract month), without language barriers.

### Types of study to be included

(cost-)effectiveness study, or any other type of quantitative/qualitative primary research, or a systematic

### Condition or domain being studied

Tuberculosis public health interventions

### Participants/ population

Hard to reach groups in European Union and OECD, like (illegal) migrants, sex workers, homeless, substance abusers, prisoners, HIV positive people and the children of these groups

### Intervention(s), exposure(s)

The reviews aim to collect evidence on all areas of interventions and service models related to identification and management of TB in vulnerable and hard-to-reach populations, as defined in the systematic reviews by NICE.

- Improve coverage and uptake of screening and active case finding by:

- using more convenient locations (like specialised TB clinic, centres for the homeless, large general practise, port of arrival, schools, syringe exchange programmes, mobile clinic)
- using peers or staff from the same hard to reach group (as case managers or health advisors)
- using small monetary incentives or food vouchers
- identifying more members of hard to reach groups

- IPT combined with methadone treatment for drug users



- Directly Observed Therapy (DOT(S)) by peers and in more convenient locations
- Family based DOT(S) programme
- Legal detention to manage active TB
- Social care support
- Enhanced case management
- Directly Observed Preventive Therapy (DOPT) plus incentives

Additionally, we aim at considering the following additional interventions:

- General interventions to improve the situation for the vulnerable and socially disadvantaged groups, e.g. provision of housing, nutritional programmes, addressing challenges related to immigration from high-TB burden countries, addressing inequalities and socioeconomic deprivation
- Case finding, contact tracing and screening programmes, by classic or innovative interventions for active case finding to reduce delays in case detection and limit further transmission
- Case holding and treatment interventions to maximise treatment initiation, adherence and completion
- Use of peer support, incentives, and other means to increase effectiveness of any interventions aimed at case finding, case holding or case management
- The existence of programs aimed collaborations with, or interventions aimed at, alternative, traditional, and / or spiritual medicine in TB treatment. This could be a problem that could be an obstacle in the control in TB in mainly immigrants and asylum seekers originating from non-western countries
- Programmes aimed at detection of patients from vulnerable or hard-to-reach populations who were lost to follow-up will be included as well.
- Treat comorbidities including HIV, diabetes, hepatitis, substance use disorders
- Continuity of care in the public sector for prisoners released from prison
- Improve living conditions and nutrition for prisoners
- General policies, health system organisation and structure of services that are aiming at enhancing access of care of specific targeted vulnerable groups, to make health care (including TB services) less hard to reach for these individuals

As this is a follow-up on the review done by NICE the following inclusion criteria will be used:

- Have a focus on TB services of any kind;
- Are conducted in any of the EU/EEA countries, the 5 candidate countries and the other OECD countries
- Are published in 2010 or later for the OECD countries;
- Are published in 1990 or later for the EU/EEA countries and the 5 candidate countries;
- Include data from any hard-to-reach group
- Present qualitative and/or quantitative empirical data;

1  
2  
3  
4 • Discuss an intervention relating to one of the following: identifying TB cases; managing TB cases; design of service  
5 models, and:

6 • is a (cost)-effectiveness study, or any other type of quantitative primary research, or a systematic review.

7  
8 Studies focussing on the detection and management of latent TB infection are not in the scope of this review.

### 10 **Comparator(s)/ control**

11 Not relevant

### 13 **Context**

14 The research needs to take place in an EU or OECD country and focus on any of the earlier mentioned hard to reach  
15 groups

### 17 **Outcome(s)**

#### 18 **Primary outcomes**

19 The most (cost)-effective service models and service structures at supporting TB identification and management of  
20 hard-to-reach groups

#### 22 **Secondary outcomes**

23 Who are responsible for putting these service structures in place and organise them.

### 25 **Data extraction, (selection and coding)**

26 Following PRISMA guidelines, study selection will be performed by two separate review authors. Three authors will  
27 screen titles and abstracts independently in parallel for matching our research objectives. Consequently, full versions  
28 of potentially relevant articles will be retrieved to assess eligibility. All full-text articles will be double screened. Data  
29 will be collected from each publication and captured using the data extraction forms, according to the type of study.  
30 Data will be extracted from text, tables and figures. The data extraction forms will be completed independently by  
31 two review authors for a randomly selected sample of 10% of records. For the other records, the tool will be  
32 completed by one reviewer and checked by another, with any disagreements resolved by discussion or consultation of  
33 a senior study investigator.

### 35 **Risk of bias (quality) assessment**

36 In concordance with PRISMA guidelines and the NICE guidance document, methodological quality and risk of bias  
37 will be assessed separately for each eligible study. Two review authors will pilot-test the tool independently on five  
38 randomly selected studies. Quality assessments will be performed independently by two review authors for 10% of  
39 the included studies, for the remaining 90% of the records the tool will be completed by one reviewer and checked by  
40 another, disagreements will be resolved by discussion or consulting the third review author.

### 42 **Strategy for data synthesis**

43 We will decide to perform a meta-analysis according to the quality assessment scores and risk of bias assessments. If  
44 studies do not support meta-analysis, data will be synthesized narratively. Overall strength of the evidence will be  
45 assessed and reported in an additional file. Strength analysis will be performed by the reviewer who performed the  
46 data extraction and quality assessment for the respective studies.

### 48 **Analysis of subgroups or subsets**

49 Every subgroup will be analysed separately (see previous answer)

### 51 **Contact details for further information**

52 Charlotte Heuvelings

53 Baljuwplein 29-II

54 3033 XB Rotterdam

55 The Netherlands

lotjeh@hotmail.com

### Organisational affiliation of the review

Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine, Academic Medical Center, University of Amsterdam, Meibergdreef 9, 1105 AZ, Amsterdam, the Netherlands

www.amc.nl

### Review team

Dr Charlotte Heuvelings, AMC  
Dr Sophia de Vries, AMC  
Dr Patrick Greve, AMC  
Dr Benjamin Jelle Visser, AMC  
Dr Saskia Janssen, AMC  
Dr Sabine Belard, AMC  
Miss Lianne Cremers, AMC  
Mr Rene Spijker, AMC  
Professor Martin Grobusch, AMC

### Collaborators

Mr Andreas Sandgren, ECDC

### Details of any existing review of the same topic by the same authors

Not by the same authors but this review is a follow up of:

Identifying and managing tuberculosis among hard-to-reach groups. Public health guidance, PH37 - Issued:

March 2012. NICE – National Institute for Health and Care Excellence. Available at:  
<http://guidance.nice.org.uk/PH37>

### Anticipated or actual start date

15 December 2014

### Anticipated completion date

01 June 2015

### Funding sources/sponsors

European Centre of Disease Prevention and Control

### Conflicts of interest

None known

### Language

English

### Country

Netherlands, Sweden

### Subject index terms status

Subject indexing assigned by CRD

### Subject index terms

Cost-Benefit Analysis; Disease Management; Humans; Tuberculosis

### Stage of review

Completed and published

**Date of registration in PROSPERO**

24 March 2015

**Date of publication of this revision**

11 April 2017

**Details of final report/publication(s)**

Effectiveness of interventions for diagnosis and treatment of tuberculosis in hard-to-reach populations in countries of low and medium tuberculosis incidence: a systematic review

Heuvelings, Charlotte C et al.

The Lancet Infectious Diseases , 2017, [http://dx.doi.org/10.1016/S1473-3099\(16\)30532-1](http://dx.doi.org/10.1016/S1473-3099(16)30532-1).  
[http://thelancet.com/journals/laninf/article/PIIS1473-3099\(16\)30532-1/fulltext](http://thelancet.com/journals/laninf/article/PIIS1473-3099(16)30532-1/fulltext)

**DOI**

10.15124/CRD42015017865

**Stage of review at time of this submission**

|   | <b>Started</b> | <b>Completed</b> |
|---|----------------|------------------|
| Preliminary searches  | Yes            | Yes              |
| Piloting of the study selection process                         | Yes            | Yes              |
| Formal screening of search results against eligibility criteria | Yes            | Yes              |
| Data extraction   | Yes            | Yes              |
| Risk of bias (quality) assessment                               | Yes            | Yes              |
| Data analysis   | Yes            | Yes              |

**PROSPERO**

**International prospective register of systematic reviews**

The information in this record has been provided by the named contact for this review. CRD has accepted this information in good faith and registered the review in PROSPERO. CRD bears no responsibility or liability for the content of this registration record, any associated files or external websites.

## Supplementary Material II: PICOS (Population – Intervention – Comparator – Outcome – Study design)

### 1. Review question

Which service models and organisational structures, including different types of healthcare workers and settings, are effective and cost-effective for delivering TB services to hard-to-reach populations?

### 2. PICOS

#### Population

Hard-to-reach groups in low incidence countries, like:

- homeless people including rough sleepers and shelter users
- people who abuse drugs or alcohol
- sex workers
- prisoners or people with a history of imprisonment
- migrants, including vulnerable migrant populations such as asylum seekers, refugees and the Roma population
- children within vulnerable and hard-to-reach populations
- people living with HIV

Studies focusing on hard-to-reach populations from Organisation for Economic Co-operation and Development (OECD) countries, European Union, European Economic Area (EU/EEA) countries and the EU candidate countries were included.

#### EU/EEA and candidate countries

1. Albania
2. Austria
3. Belgium
4. Bulgaria
5. Croatia
6. Cyprus
7. Czech Republic
8. Denmark
9. Estonia
10. Finland
11. France
12. Germany
13. Greece
14. Hungary
15. Iceland
16. Ireland
17. Italy

#### OECD countries

1. Australia
2. Austria
3. Belgium
4. Canada
5. Chile
6. Czech Republic
7. Denmark
8. Estonia
9. Finland
10. France
11. Germany
12. Greece
13. Hungary
14. Iceland
15. Ireland
16. Israel
17. Italy

|   |                     |
|---|---------------------|
| 18. Latvia                                    | 18. Japan           |
| 19. Liechtenstein                             | 19. Korea           |
| 20. Lithuania                                 | 20. Luxembourg      |
| 21. Luxembourg                                | 21. Mexico          |
| 22. Malta                                     | 22. Netherlands     |
| 23. Montenegro                                | 23. New Zealand     |
| 24. Netherlands                               | 24. Norway          |
| 25. Norway                                    | 25. Poland          |
| 26. Poland                                    | 26. Portugal        |
| 27. Portugal                                  | 27. Slovak Republic |
| 28. Romania                                   | 28. Slovenia        |
| 29. Serbia                                    | 29. Spain           |
| 30. Slovakia                                  | 30. Sweden          |
| 31. Slovenia                                  | 31. Switzerland     |
| 32. Spain                                     | 32. Turkey          |
| 33. Sweden                                    | 33. United Kingdom  |
| 34. The former Yugoslav Republic of Macedonia | 34. United States   |
| 35. Turkey                                    |                     |
| 36. United Kingdom                            |                     |

Studies that did not specifically look at any of these target populations or were conducted in a different geographical area were excluded.

### **Intervention**

This review aimed to collect evidence on all areas of interventions targeting service structures and service models for the identification and management of TB in vulnerable and hard-to-reach populations. These interventions can cover activities to improve TB services for screening, active case finding or TB management. Predefined interventions included in the protocol were:

- Using different locations like:

- Hospitals
- Specialised TB clinics
- Shelters for homeless people
- Needle/syringe exchange programmes
- Locations that combine care: providing treatment for drug addiction, medical care, HIV care
- Large general practises
- Port of arrival
- Schools
- Mobile clinics
- Prisons

- Using different type of health care workers like:

- Doctors
  - Nurses
  - Community health workers
  - Peers
  - People from the same background
- Others
- Family based DOTS programs
  - Social care support and general interventions
    - Provision of housing
    - Nutritional programmes
    - Addressing challenges related to immigration from high-TB burden countries
    - Addressing inequalities and socioeconomic deprivation
  - Collaboration with alternative, traditional, and / or spiritual medicine

### **Comparator**

Not relevant.

The comparator was re-defined during the review process into:

Standard care.

### **Outcome**

The outcome measures were quantitative outcomes focusing on the effectiveness and/or cost-effectiveness of service structures and service models aiming to identify and manage TB in hard-to-reach populations in low incidence countries, including a qualitative description of these interventions.

### **Study design**

Randomised controlled trials (RCTs) focusing on service models and organisational structures on the selected hard-to-reach populations were included. Since it was very likely that few RCTs would be identified, we also included non-randomised studies like, but not exclusively, case-control studies, cohort studies, cross-sectional studies and observational studies. Quantitative and qualitative studies were included. Systematic reviews were included for reference checking only.

## **3. Further notes on PICOS**

For this systematic review of interventions with a scoping component, a very broad and sensitive search was conducted to cover a wide range of interventions. Predefined interventions were included in our registered protocol but the list of interventions was not exclusive and interventions were added to the list during the review process. Supplementary Material I reflects the registered protocol. Changes made during the implementation of the systematic review protocol are stated at the end of each section.

### Supplementary Material III: Search strategies

The previous NICE review<sup>1</sup> on the same topic was used as a framework for the search strategy and extended to the non-Organisation for Economic Co-operation and Development (OECD), countries of the European Union and European Economic Area and to the two newly included hard-to-reach groups (people living with HIV co-infected with TB and children within vulnerable and hard-to-reach populations). The search for the NICE review<sup>1</sup> was subtracted from our search to prevent double screening of records. The search was conducted by René Spijker, clinical librarian at the Academic Medical Center in Amsterdam, the Netherlands. All studies identified by the search were imported to an Endnote database. The original search was done on the 10<sup>th</sup> of December 2014 and updated on the 10<sup>th</sup> of April 2015 and 24<sup>th</sup> of February 2017.

The following two databases were used for the search:

- MEDLINE(R) In-Process & Other Non-Indexed Citations (OvidSP)
- Embase Classic + Embase

| Database                     | Hits          |
|------------------------------|---------------|
| Medline + Medline In-Process | 12,519        |
| Embase                       | 16,079        |
| <b>Total</b>                 | <b>19,720</b> |

### References:

1. Rizzo M, Martin A, Jamal F, et al. Evidence review on the effectiveness and cost-effectiveness of service models or structures to manage tuberculosis in hard-to-reach groups. London: Matrix evidence/National Institute for Health and Clinical Excellence 2011. <https://www.nice.org.uk/guidance/PH37/documents/review-4-evidence-review-on-the-effectiveness-and-cost-effectiveness-of-service-models-or-structures-to-manage-tuberculosis-in-hardtoeach-groups-2> (last assessed March 2017).



**1. Search in Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations and Ovid MEDLINE(R) 1946 to Present**

Hits: 12,519

|    |   |
|----|---|
| 1  | exp *tuberculosis/ or (tuberculosis or tb).ti,ab.   |
| 2  | ((hard\$ adj2 reach) or (hard\$ adj2 locate) or (hard\$ adj2 find) or (hard\$ adj2 treat) or (difficult adj2 locate) or (difficult adj2 engage) or social\$ exclu\$ or social inequalit\$ or (difficult\$ adj2 reach) or (difficult\$ adj2 find) or (difficult\$ adj2 treat) or (christian* or church* or chapel* or priest* or vicar* or catholic* or catholicism or protestant* or methodist* or baptist* or Jehovah* or presbyterian* or anglican* or pentecostal*) or (muslim* or islam* or mosque* or imam*)).ti,ab. or exp *Jew/ or (jew* or judaism* or synagogue*).ti,ab. or exp *religion/ or (christian* or church* or chapel* or priest* or vicar* or catholic* or catholicism or protestant* or methodist* or baptist* or Jehovah* or presbyterian* or anglican* or pentecostal*).ti,ab. or (jew* or judaism* or synagogue*).ti,ab. or (sikh* or hindu* or buddhis* or temple*).ti,ab. or ((religion* or religious* or faith*) and (people* or person* or group* or population or neighbour* or neighbor* or patient* or communit*)).ti,ab. |
| 3  | ((geograph\$ or transport\$ or physical) and barrier\$).ti,ab.  |
| 4  | ((low\$ or poor\$ or negative) and (quality adj2 life)).ti,ab.  |
| 5  | ((vulnerable or disadvantaged or at risk or high risk or low socioeconomic status or neglect\$ or affected or marginal\$ or forgotten or non-associative or unengaged or hidden or excluded or transient or inaccessible or underserved or stigma\$ or inequitable) and (people or population\$ or communit\$ or neighbourhood\$1 or neighborhood\$1 or group\$ or area\$1 or demograph\$ or patient\$ or social\$)).ti,ab. or exp *vulnerable population/  |
| 6  | *poverty/   |
| 7  | (refuser\$1 or nonuser\$1 or non-user\$1 or non user\$1 or discriminat\$ or shame or prejud\$ or racism or racial discriminat\$).ti,ab.   |
| 8  | *social support/ or exp *social status/ or *social stigma/ or exp *social isolation/ or exp *"quality of life"/ or exp *prejudice/ or exp *socioeconomics/  |
| 9  | prisoner\$1.ti,ab.  |
| 10 | (recent\$ adj2 release\$ adj2 (inmate\$ or prison\$ or detainee\$ or felon\$ or offender\$ or convict\$ or custod\$ or detention or incarcerat\$ or correctional or jail\$ or penitentiary\$)).ti,ab.   |
| 11 | ((prison\$ or penal or penitentiary\$ or correctional facilit\$ or jail\$ or detention centre\$ or detention center\$) and (guard\$1 or population or inmate\$ or system\$ or remand or detainee\$ or felon\$ or offender\$1 or convict\$ or abscond\$)).ti,ab.   |
| 12 | (parole or probation).ti,ab.  |
| 13 | exp *prisoner/  |
| 14 | ((custodial adj (care or sentence)) or (incarceration or incarcerated or imprisonment)).ti,ab.  |
| 15 | (immobile or (disabled and (house bound or home bound)) or ((house or home) adj3 bound)).ti,ab. or exp *homebound patient/  |
| 16 | ((hous\$ and (quality or damp\$ or standard\$ or afford\$ or condition\$ or dilapidat\$)) or ((emergency or temporary or inadequate or poor\$ or overcrowd\$ or over-crowd\$ or over-subscribed) and (hous\$ or accommodation or shelter\$ or hostel\$ or dwelling\$))).ti,ab. or exp *housing/   |
| 17 | (rough sleep\$ or runaway\$1 or ((homeless\$ or street or destitut\$) and (population or person\$1 or people or group\$ or individual\$1 or shelter\$ or hostel\$ or accommodation\$1))).ti,ab. or exp *homelessness/   |
| 18 | ((drug\$ or substance) and (illegal or misus\$ or abuse or intravenous or IV or problem use\$ or illicit use\$ or addict\$ or dependen\$ or dependant or delinquency)).ti,ab. or exp *addiction/  |
| 19 | ((alcohol\$ and (misus\$ or abuse or problem\$ use\$ or problem drink\$ or illicit use\$ or addict\$ or dependen\$ or delinquency)) or alcoholic\$1).ti,ab.   |
| 20 | (prostitution or sex work\$ or transactional sex\$ or prostitute\$1).ti,ab. or Prostitution/  |
| 21 | (poverty or deprivation or financial hardship\$).ti,ab.   |
| 22 | ((low-income or low income or low pay or low paid or poor or deprived or debt\$ or arrear\$) and (people or person\$1 or population\$1 or communit\$ or group\$ or social group\$ or neighbourhood\$1 or neighborhood\$1 or famil\$)).ti,ab. or exp *lowest income group/   |
| 23 | *poverty/   |
| 24 | (low\$ and social class\$).ti,ab.   |

|    |   |
|----|---|
| 25 | (traveller\$1 or gypsies or gypsy or Romany or roma).ti,ab. or exp *"Romani (people)"/  |
| 26 | (mental\$ and (health or ill or illness)).ti,ab. or *mental patient/ or exp *mental health/   |
| 27 | (health care worker\$1 or (health care adj2 service provi\$) or (health-care adj2 provi\$) or (((community adj1 leader\$) or (community adj1 (Manag\$ or advocat\$ or champion\$))) and (engag\$ or involv\$)).ti,ab.   |
| 28 | (complex adj2 (patient\$ or Need\$)).ti,ab.   |
| 29 | (outreach adj2 worker\$1).ti,ab. or exp *health auxiliary/  |
| 30 | (support adj2 worker\$1).ti,ab.   |
| 31 | (case adj2 worker\$1).ti,ab.  |
| 32 | (social adj2 worker\$1).ti,ab.  |
| 33 | social care professional\$1.ti,ab.  |
| 34 | ((social care adj2 service provi\$) or (social-care adj2 provi\$)).ti,ab.   |
| 35 | ((language\$ or communicat\$) and (barrier\$ or understand\$ or strateg\$ or proficien\$)) or translat\$ or interpret\$ or (cultur\$ and competen\$).ti,ab. or *language ability/   |
| 36 | (immigrant\$ or migrant\$ or asylum or refugee\$ or undocumented or foreign born or (born adj overseas) or (displaced and (people or person\$1))).ti,ab. or exp *refugee/   |
| 37 | exp *migrant/   |
| 38 | *immigration/   |
| 39 | or/2-38   |
| 40 | Intervention\$.ti,ab. or exp *crisis intervention/  |
| 41 | ((early or primary) adj2 Intervention\$).ti,ab.   |
| 42 | ((person\$ or individual or local\$ or community or cultural or structural or supported or indicated or target\$ or multi?component or comprehensive or pilot or media) and Intervention\$).ti,ab.  |
| 43 | ((midstream or mid-stream) and intervention\$).ti,ab.   |
| 44 | (Identify\$ or find or finding or locat\$ or trac\$ or contact\$ or discover\$ or detect or recruit\$ or attract\$).ti,ab.  |
| 45 | (case finding or ((active or passive) adj3 case finding)).ti,ab.  |
| 46 | ((program\$ or scheme\$1 or service\$1 or campaign\$ or mobili?ation or strateg\$ or measure or policy or policies) and (tuberculosis or tb)).ti,ab.  |
| 47 | ((case adj3 management) or case-managed).ti,ab. or *case management/ or *patient care planning/ or *case management/ or exp *health care management/  |
| 48 | (case adj3 manag\$ adj3 strategy).ti,ab. or continuity of * patient care/   |
| 49 | ((treat\$ or diagnosis) and management).ti,ab.  |
| 50 | ((active or passive) and (Case adj3 Management)).ti,ab.   |
| 51 | (risk assess\$ or risk profile or risk Indicator or care plan\$).ti,ab.   |
| 52 | (service and (model\$ or deliver\$)).ti,ab. or delivery of * health care/ or *health service/   |
| 53 | ((primary adj3 healthcare) or ((primary adj3 health\$) or care)).ti,ab. or exp *primary health care/  |
| 54 | (nurse or ((general or family) adj3 (practice\$ or practitioner\$ or physicians\$ or doctor\$))).ti,ab. or exp *nurse/ or (exp *tuberculosis/ or (tuberculosis or tb).ti,ab.) or exp *general practice/   |
| 55 | ((health or extension or multi-disciplinary or multidisciplinary) and (professional\$ or personal\$ or practitioner or worker\$ or partner\$ or promot\$ or provider or care team or care provider or unit or casework\$ or (case adj2 work\$))).ti,ab. or *health care personnel/ or exp *nursing assistant/ |
| 56 | (social adj2 (work\$ or Support\$ or Outreach)).ti,ab. or *social work/ or *social support/   |
| 57 | (volunteer\$ or voluntary or charit\$ or third sector).ti,ab. or *voluntary worker/ or exp *health care organization/   |
| 58 | (health adj1 (center\$1 or centre\$1 or facilit\$ or service\$ or clinic\$1 or hospital\$1 or program\$1)).ti,ab. or *public health/ or *residential care/  |
| 59 | ((day adj2 (care or hospital\$ or patient\$)) or workshop\$).ti,ab. or *day care/   |
| 60 | rehab\$.ti,ab. or *rehabilitation center/   |
| 61 | ((dedicated or permanent or rapid access or fixed or TB or tuberculosis) and (clinic\$1 or centre\$1 or center\$1 or program\$)).ti,ab.   |
| 62 | ((drug adj2 dependency) or substance abuse or HIV) and (unit\$ or clinic\$1 or centre\$1 or center\$1 or program\$) and (tuberculosis or tb)).ti,ab.  |
| 63 | (pharmac\$ or dispensary).ti,ab. or *pharmacy/  |

|    |   |
|----|---|
| 64 | (communit\$ or (support\$ adj2 communit\$)).ti,ab. or *community care/ or *health auxiliary/ or *public relations/ or *community hospital/ or *community health nursing/  |
| 65 | (directly observed treatment or directly observed therapy or (supervised adj2 treatment) or (coerc\$ adj2 (treat\$ or therapy))).ti,ab. or Directly Observed Therapy/   |
| 66 | (ambulatory adj2 care).ti,ab. or exp *ambulatory care/  |
| 67 | ((mobile or travel\$ or transport\$ or workplace or work-place or tertiary) and (health adj3 (care or work\$ or practitioner\$ or professional\$ or service\$ or center\$1 or centre\$1 or unit\$1 or program\$))).ti,ab. or *preventive health service/  |
| 68 | ((mobile or travel\$ or transport\$ or workplace or work-place or tertiary) and (nurs\$ or doctor\$)).ti,ab.  |
| 69 | ((out adj3 hours) or (after adj3 hours) or telephone or telemedicine).ti,ab. or after-hours care/ or exp *telehealth/ or *emergency care/ or *health care delivery/   |
| 70 | ((walk-in or walkin or walk in) adj2 (center\$1 or centre\$1 or service or program\$ or Clinic\$1 or Session or Assessment\$1)).ti,ab.  |
| 71 | (drop\$ adj1 in adj2 (center\$1 or centre\$1 or service or program\$ or clinic\$1 or session or meeting or assesment\$1)).ti,ab.  |
| 72 | ((health or home\$ or house\$) and (call\$ or visit\$)) or (home-care or home-based or (support\$ adj1 hous\$)).ti,ab. or Home Health Aides/ or *health auxiliary/ or exp *home care/   |
| 73 | ((early adj2 discharge) or (recent\$ adj2 discharged) or (out adj2 patient)).ti,ab. or *patient care/ or *outpatient department/  |
| 74 | (counselling or counseling or counsellor or counselor or (integrated counselling adj1 testing centre\$1) or (integrated counselling adj1 testing center\$1) or ICTC).ti,ab. or *counseling/ or *directive counseling/   |
| 75 | ((help adj2 group\$) or (self adj2 help) or support\$ or (peer adj2 peer)).ti,ab. or *self help/  |
| 76 | (collaborat\$ or shared or (integrated adj1 care\$) or ICP or network\$ or co-locat\$ or (one adj1 stop)).ti,ab. or *integrated health care system/   |
| 77 | ((health adj2 education) or (skill adj2 mix) or (role adj2 develop\$) or leadership or ((interdisciplinary or inter-team or Professional or team) adj2 communicate\$)).ti,ab. or exp *health education/ or exp *interdisciplinary communication/ or *leadership/ or *doctor patient relation/ or *nurse patient relationship/ or patient relationship*.ti,ab.   |
| 78 | ((outreach or mobile\$ or satellite\$ or hub or spoke or rural or urban or street or pavement\$1 or sidewalk\$1 or corner or shelter or hostel or sanatorium or sanitorium or sanitarium) and (tuberculosis or tb)).ti,ab.  |
| 79 | ((outreach or mobile\$ or satellite\$ or hub or spoke or rural or urban or street or pavement\$1 or sidewalk\$1 or corner or shelter or hostel or sanatorium or sanitorium or sanitarium) and (tuberculosis or tb)).ti,ab.  |
| 80 | or/40-79  |
| 81 | test\$.ti,ab.   |
| 82 | (examination\$1 or assessment\$1 or identification or assay\$ or detection).ti,ab.  |
| 83 | diagnosi\$.ti,ab. or *diagnostic test/  |
| 84 | ((chest adj2 x?ray) or chest radiograph or MXU).ti,ab. or *thorax radiography/  |
| 85 | (screen\$ or (new\$ adj1 screen\$)).ti,ab.  |
| 86 | (monitor\$ or sampling).ti,ab.  |
| 87 | ((target\$ or focus\$ or community or population or individual\$ or person\$ or opportunistic or coerc\$ or voluntary or initiated) and (test\$ or diagnosis or screen\$ or assay\$ or detection)).ti,ab.   |
| 88 | PIT.ti,ab.  |
| 89 | provider initiated test\$.ti,ab.  |
| 90 | ((rapid or prompt or quick\$ or earl\$ or (point adj2 care)) and (test\$ or screen\$ or diagnosi\$ or assay\$ or detection)).ti,ab.   |
| 91 | ((provider or anonymous or accurate or support\$ or incentiv\$ or counsel\$) and (test\$ or diagnosis or screen\$ or assay\$)).ti,ab. or *anonymous testing/  |
| 92 | (test\$ adj2 (center\$1 or centre\$1 or unit\$1 or setting)).ti,ab.   |
| 93 | or/81-92  |
| 94 | (acceptability or acceptable or attend\$ or access\$ or availab\$ or non-attend\$ or increas\$ or promot\$ or opt\$ or particip\$ or adhere\$ or involvement or uptake or take-up or utiliz\$ or utilis\$ or refus\$ or referr\$ or self-referr\$ or self-report\$ or barrier\$ or decreas\$ or isolation or interven\$ or aware\$ or opportunit\$ or advice or information or incentiv\$ or recruit\$ or |

|     |   |
|-----|---|
|     | find or finding or compliance or comply or retain or retention or provision or encour\$ or usage).ti,ab.  |
| 95  | (socio sanitary support or reimburs\$ or (social adj2 support) or ((cash or financial or money or monetary or economic or voucher or credit or drug\$1 or methadone or telephone) adj2 (benefit\$ or support or incentive or assist\$ or credit))).ti,ab. |
| 96  | (((lifestyle or behavio?r) adj2 (therapy or modif\$ or chang\$ or adapt\$ or adopt\$)) and (tuberculosis or tb)).ti,ab. or *social marketing/   |
| 97  | *marketing/   |
| 98  | *attitude to health/  |
| 99  | *health care delivery/  |
| 100 | *access to information/   |
| 101 | *confidentiality/   |
| 102 | *Health education/  |
| 103 | *health promotion/  |
| 104 | *patient compliance/  |
| 105 | *motivation/  |
| 106 | Stigma.ti,ab.   |
| 107 | *prevalence/  |
| 108 | *patient participation/   |
| 109 | *patient attitude/ or *refusal to participate/ or *treatment refusal/   |
| 110 | or/94-109   |
| 111 | treat\$.ti,ab. or Treatment Outcome/  |
| 112 | (directly observed treatment or directly observed therapy or (supervised adj2 treatment) or (coerc\$ adj2 (treat\$ or therapy))).ti,ab. or *directly observed therapy/  |
| 113 | (disease management or (treat\$ and (management or control))).ti,ab.  |
| 114 | ((adherence or compli\$ or non-compli\$ or default\$ or finish\$ or Retention or attrition or (drop adj1 out) or disappear\$ or abscond\$) and treat\$.ti,ab. or exp *patient compliance/   |
| 115 | ((referr\$ or self-referr\$ or (self adj diagnos\$)) and treat\$.ti,ab.   |
| 116 | ((suitab\$ or eligib\$) and treat\$.ti,ab.  |
| 117 | ((follow adj1 up) or discharge).ti,ab. or *follow up/   |
| 118 | ((positive or negative) and test).ti,ab.  |
| 119 | ((interrupt\$ or relapse\$ or stop\$ or cessation or with?ld\$ or avoidance or (lost adj2 follow)) and treat\$.ti,ab. or *treatment withdrawal/   |
| 120 | ((medicine\$1 or drug or treat\$) and (regimen or adherence)).ti,ab. or exp *self care/   |
| 121 | (treat\$ and (appointment\$ or Schedule\$)).ti,ab. or *patient scheduling/  |
| 122 | ((care adj2 seeking) and pathway\$.ti,ab.   |
| 123 | ((case adj3 management) or case-managed).ti,ab. or Case Management/ or *patient care planning/ or *health insurance/  |
| 124 | (case adj3 manag\$ adj3 strategy).ti,ab. or continuity.mp. or *patient care/ [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword]                    |
| 125 | ((case or treat\$ or diagnosis) and management).ti,ab.  |
| 126 | ((risk assessment or care plan\$) and (case adj3 management)).ti,ab.  |
| 127 | ((active or passive) and (case adj3 management)).ti,ab.   |
| 128 | or/111-127  |
| 129 | 1 and 39 and (80 or (93 and (110 or 128)))  |
| 130 | limit 129 to yr="1990 -Current"   |
| 131 | limit 130 to "english language"   |
| 132 | (exp animal/ or animal.hw. or nonhuman/) not (exp human/ or human cell/ or (human or humans).ti.)   |
| 133 | 131 not 132   |
| 134 | limit 133 to yr="1990 - 2010"   |
| 135 | (albania or bulgaria or cyprus or croatia or latvia or lithuania or luxembourg or malta or montenegro or romania or serbia or yugoslav or turkey).ti,ab,hw,in.  |
| 136 | 1 and 135 and (80 or (93 and (110 or 128)))   |
| 137 | limit 136 to yr="1990 -Current"   |
| 138 | 137 not 132   |

|     |             |
|-----|-------------|
| 139 | 138 not 130 |
| 140 | 133 not 134 |
| 141 | 139 or 140  |

## 2. Search in Embase Classic + Embase 1947 to 2014 December 2008

Ovid

Hits: 16,079

|    |   |
|----|---|
| 1  | <b>exp *tuberculosis/ or (tuberculosis or tb).ti,ab.</b>  |
| 2  | ((hard\$ adj2 reach) or (hard\$ adj2 locate) or (hard\$ adj2 find) or (hard\$ adj2 treat) or (difficult adj2 locate) or (difficult adj2 engage) or social\$ exclu\$ or social inequalit\$ or (difficult\$ adj2 reach) or (difficult\$ adj2 find) or (difficult\$ adj2 treat) or (christian* or church* or chapel* or priest* or vicar* or catholic* or catholicism* or protestant* or methodist* or baptist* or Jehovah* or presbyterian* or anglican* or pentecostal*) or (muslim* or islam* or mosque* or imam*).ti,ab. or exp *Jew/ or (jew* or judaism* or synagogue*).ti,ab. or exp *religion/ or (christian* or church* or chapel* or priest* or vicar* or catholic* or catholicism* or protestant* or methodist* or baptist* or Jehovah* or presbyterian* or anglican* or pentecostal*).ti,ab. or (jew* or judaism* or synagogue*).ti,ab. or (sikh* or hindu* or buddhis* or temple*).ti,ab. or ((religion* or religious* or faith*) and (people* or person* or group* or population or neighbour* or neighbor* or patient* or communit*).ti,ab. |
| 3  | ((geograph\$ or transport\$ or physical) and barrier\$.ti,ab.   |
| 4  | ((low\$ or poor\$ or negative) and (quality adj2 life)).ti,ab.  |
| 5  | ((vulnerable or disadvantaged or at risk or high risk or low socioeconomic status or neglect\$ or affected or marginal\$ or forgotten or non-associative or unengaged or hidden or excluded or transient or inaccessible or underserved or stigma\$ or inequitable) and (people or population\$ or communit\$ or neighbourhood\$1 or neighborhood\$1 or group\$ or area\$1 or demograph\$ or patient\$ or social\$)).ti,ab. or exp *vulnerable population/  |
| 6  | *poverty/   |
| 7  | (refuser\$1 or nonuser\$1 or non-user\$1 or non user\$1 or discriminat\$ or shame or prejud\$ or racism or racial discriminat\$.ti,ab.  |
| 8  | *social support/ or exp *social status/ or *social stigma/ or exp *social isolation/ or exp *"quality of life"/ or exp *prejudice/ or exp *socioeconomics/  |
| 9  | prisoner\$1.ti,ab.  |
| 10 | (recent\$ adj2 release\$ adj2 (inmate\$ or prison\$ or detainee\$ or felon\$ or offender\$ or convict\$ or custod\$ or detention or incarcerat\$ or correctional or jail\$ or penitentiari\$)).ti,ab.   |
| 11 | ((prison\$ or penal or penitentiari\$ or correctional facilit\$ or jail\$ or detention centre\$ or detention center\$) and (guard\$1 or population or inmate\$ or system\$ or remand or detainee\$ or felon\$ or offender\$1 or convict\$ or abscond\$)).ti,ab.   |
| 12 | (parole or probation).ti,ab.  |
| 13 | exp *prisoner/  |
| 14 | ((custodial adj (care or sentence)) or (incarceration or incarcerated or imprisonment)).ti,ab.  |
| 15 | (immobile or (disabled and (house bound or home bound)) or ((house or home) adj3 bound)).ti,ab. or exp *homebound patient/  |
| 16 | ((hou\$ and (quality or damp\$ or standard\$ or afford\$ or condition\$ or dilapidat\$)) or ((emergency or temporary or inadequate or poor\$ or overcrowd\$ or over-crowd\$ or over-subscribed) and (hou\$ or accommodation or shelter\$ or hostel\$ or dwelling\$)).ti,ab. or exp *housing/  |
| 17 | (rough sleep\$ or runaway\$1 or ((homeless\$ or street or destitut\$) and (population or person\$1 or people or group\$ or individual\$1 or shelter\$ or hostel\$ or accommodation\$1))).ti,ab. or exp *homelessness/   |
| 18 | ((drug\$ or substance) and (illegal or misus\$ or abuse or intravenous or IV or problem use\$ or illicit use\$ or addict\$ or dependen\$ or dependant or delinquency)).ti,ab. or exp *addiction/  |

|    |  |
|----|--|
| 19 | ((alcohol\$ and (misuse\$ or abuse or problem\$ use\$ or problem drink\$ or illicit use\$ or addict\$ or dependen\$ or delinquency)) or alcoholic\$1).ti,ab.   |
| 20 | (prostitution or sex work\$ or transactional sex\$ or prostitute\$1).ti,ab. or Prostitution/   |
| 21 | (poverty or deprivation or financial hardship\$).ti,ab.  |
| 22 | ((low-income or low income or low pay or low paid or poor or deprived or debt\$ or arrear\$) and (people or person\$1 or population\$1 or community\$ or group\$ or social group\$ or neighbourhood\$1 or neighborhood\$1 or famil\$)).ti,ab. or exp *lowest income group/ |
| 23 | *poverty/  |
| 24 | (low\$ and social class\$).ti,ab.  |
| 25 | (traveller\$1 or gypsies or gypsy or Romany or roma).ti,ab. or exp *"Romani (people)"/   |
| 26 | (mental\$ and (health or ill or illness)).ti,ab. or *mental patient/ or exp *mental health/  |
| 27 | (health care worker\$1 or (health care adj2 service provi\$) or (health-care adj2 provi\$) or (((community adj1 leader\$) or (community adj1 (Manag\$ or advocat\$ or champion\$))) and (engag\$ or involv\$)).ti,ab.  |
| 28 | (complex adj2 (patient\$ or Need\$)).ti,ab.  |
| 29 | (outreach adj2 worker\$1).ti,ab. or exp *health auxiliary/   |
| 30 | (support adj2 worker\$1).ti,ab.  |
| 31 | (case adj2 worker\$1).ti,ab.   |
| 32 | (social adj2 worker\$1).ti,ab.   |
| 33 | social care professional\$1.ti,ab.   |
| 34 | ((social care adj2 service provi\$) or (social-care adj2 provi\$)).ti,ab.  |
| 35 | ((language\$ or communicat\$) and (barrier\$ or understand\$ or strateg\$ or proficien\$)) or translat\$ or interpret\$ or (cultur\$ and competen\$)).ti,ab. or *language ability/   |
| 36 | (immigrant\$ or migrant\$ or asylum or refugee\$ or undocumented or foreign born or (born adj overseas) or (displaced and (people or person\$1))).ti,ab. Or exp *refugee/  |
| 37 | exp *migrant/  |
| 38 | *immigration/  |
| 39 | or/2-38  |
| 40 | Intervention\$.ti,ab. or exp *crisis intervention/   |
| 41 | ((early or primary) adj2 Intervention\$).ti,ab.  |
| 42 | ((person\$ or individual or local\$ or community or cultural or structural or supported or indicated or target\$ or multi?component or comprehensive or pilot or media) and Intervention\$).ti,ab.   |
| 43 | ((midstream or mid-stream) and intervention\$).ti,ab.  |
| 44 | (Identify\$ or find or finding or locat\$ or trac\$ or contact\$ or discover\$ or detect or recruit\$ or attract\$).ti,ab.   |
| 45 | (case finding or ((active or passive) adj3 case finding)).ti,ab.   |
| 46 | ((program\$ or scheme\$1 or service\$1 or campaign\$ or mobili?ation or strateg\$ or measure or policy or policies) and (tuberculosis or tb)).ti,ab.   |
| 47 | ((case adj3 management) or case-managed).ti,ab. or *case management/ or *patient care planning/ or *case management/ or exp *health care management/   |
| 48 | (case adj3 manag\$ adj3 strategy).ti,ab. or continuity of *patient care/   |
| 49 | ((treat\$ or diagnosis) and management).ti,ab.   |
| 50 | ((active or passive) and (Case adj3 Management)).ti,ab.  |
| 51 | (risk assess\$ or risk profile or risk Indicator or care plan\$).ti,ab.  |
| 52 | (service and (model\$ or deliver\$)).ti,ab. or delivery of *health care/ or *health service/   |
| 53 | ((primary adj3 healthcare) or ((primary adj3 health\$) or care)).ti,ab. or exp *primary health care/   |
| 54 | (nurse or ((general or family) adj3 (practice\$ or practitioner\$ or physicians\$ or doctor\$))).ti,ab. Or exp *nurse/ or (exp *tuberculosis/ or (tuberculosis or tb).ti,ab.)) or exp *general practice/   |
| 55 | ((health or extension or multi-disciplinary or multidisciplinary) and (professional\$ or personal\$ or practitioner or worker\$ or partner\$ or promot\$ or provider or care team or care provider or unit or casework\$ or (case adj2 work\$))).ti,ab. or *health care    |

|    |   |
|----|---|
|    | personnel/ or exp *nursing assistant/   |
| 56 | (social adj2 (work\$ or Support\$ or Outreach)).ti,ab. or *social work/ or *social support/   |
| 57 | ((lay or allied or link) and (professional\$ or practitioner\$1 or worker\$1 or advocate\$1 or personnel)).ti,ab. or *paramedical personnel//   |
| 58 | (volunteer\$ or voluntary or charit\$ or third sector).ti,ab. or *voluntary worker/ or exp *health care organization/   |
| 59 | (health adj1 (center\$1 or centre\$1 or facilit\$ or service\$ or clinic\$1 or hospital\$1 or program\$1)).ti,ab. or *public health/ or *residential care/  |
| 60 | ((day adj2 (care or hospital\$ or patient\$)) or workshop\$).ti,ab. or *day care/   |
| 61 | rehab\$.ti,ab. or *rehabilitation center/   |
| 62 | ((dedicated or permanent or rapid access or fixed or TB or tuberculosis) and (clinic\$1 or centre\$1 or center\$1 or program\$)).ti,ab.   |
| 63 | ((drug adj2 dependency) or substance abuse or HIV) and (unit\$ or clinic\$1 or centre\$1 or center\$1 or program\$) and (tuberculosis or tb)).ti,ab.  |
| 64 | (pharmac\$ or dispensary).ti,ab. Or *pharmacy/  |
| 65 | (communit\$ or (support\$ adj2 communit\$)).ti,ab. or *community care/ or *health auxiliary/ or *public relations/ or *community hospital/ or *community health nursing/  |
| 66 | (directly observed treatment or directly observed therapy or (supervised adj2 treatment) or (coerc\$ adj2 (treat\$ or therapy))).ti,ab. or Directly Observed Therapy/   |
| 67 | (ambulatory adj2 care).ti,ab. or exp *ambulatory care/  |
| 68 | ((mobile or travel\$ or transport\$ or workplace or work-place or tertiary) and (health adj3 (care or work\$ or practitioner\$ or professional\$ or service\$ or center\$1 or centre\$1 or unit\$1 or program\$))).ti,ab. or *preventive health service/  |
| 69 | ((mobile or travel\$ or transport\$ or workplace or work-place or tertiary) and (nurs\$ or doctor\$)).ti,ab.  |
| 70 | ((out adj3 hours) or (after adj3 hours) or telephone or telemedicine).ti,ab. or after-hours care/ or exp *telehealth/ or *emergency care/ or *health care delivery/   |
| 71 | ((walk-in or walkin or walk in) adj2 (center\$1 or centre\$1 or service or program\$ or Clinic\$1 or Session or Assesment\$1)).ti,ab.   |
| 72 | (drop\$ adj1 in adj2 (center\$1 or centre\$1 or service or program\$ or clinic\$1 or session or meeting or assesment\$1)).ti,ab.  |
| 73 | ((health or home\$ or house\$) and (call\$ or visit\$)) or (home-care or home-based or (support\$ adj1 hous\$)).ti,ab. or Home Health Aides/ or *health auxiliary/ or exp *home care/   |
| 74 | ((early adj2 discharge) or (recent\$ adj2 discharged) or (out adj2 patient)).ti,ab. or *patient care/ or *outpatient department/  |
| 75 | (counselling or counseling or counsellor or counselor or (integrated counselling adj1 testing centre\$1) or (integrated counselling adj1 testing center\$1) or ICTC).ti,ab. or *counseling/ or *directive counseling/   |
| 76 | ((help adj2 group\$) or (self adj2 help) or support\$ or (peer adj2 peer)).ti,ab. Or *self help/  |
| 77 | (collaborat\$ or shared or (integrated adj1 care\$) or ICP or network\$ or co-locat\$ or (one adj1 stop)).ti,ab. or *integrated health care system/   |
| 78 | ((health adj2 education) or (skill adj2 mix) or (role adj2 develop\$) or leadership or ((interdisciplinary or inter-team or Professional or team) adj2 communicate\$)).ti,ab. or exp *health education/ or exp *interdisciplinary communication/ or *leadership/ or *doctor patient relation/ or *nurse patient relationship/ or patient relationship*.ti,ab. |
| 79 | ((outreach or mobile\$ or satellite\$ or hub or spoke or rural or urban or street or pavement\$1 or sidewalk\$1 or corner or shelter or hostel or sanatorium or sanitorium or sanitarium) and (tuberculosis or tb)).ti,ab.  |
| 80 | or/40-79  |
| 81 | test\$.ti,ab.   |
| 82 | (examination\$1 or assessment\$1 or identification or assay\$ or detection).ti,ab.  |
| 83 | diagnosi\$.ti,ab. or *diagnostic test/  |

|     |  |
|-----|--|
| 84  | ((chest adj2 x?ray) or chest radiograph or MXU).ti,ab. or *thorax radiography/   |
| 85  | (screen\$ or (new\$ adj1 screen\$)).ti,ab.   |
| 86  | (monitor\$ or sampling).ti,ab.   |
| 87  | ((target\$ or focus\$ or community or population or individual\$ or person\$ or opportunistic or coer\$ or voluntary or initiated) and (test\$ or diagnosis or screen\$ or assay\$ or detection)).ti,ab.   |
| 88  | PIT.ti,ab.   |
| 89  | provider initiated test\$.ti,ab.   |
| 90  | ((rapid or prompt or quick\$ or earl\$ or (point adj2 care)) and (test\$ or screen\$ or diagnosi\$ or assay\$ or detection)).ti,ab.  |
| 91  | ((provider or anonymous or accurate or support\$ or incentiv\$ or counsel\$) and (test\$ or diagnosis or screen\$ or assay\$)).ti,ab. or *anonymous testing/   |
| 92  | (test\$ adj2 (center\$1 or centre\$1 or unit\$1 or setting)).ti,ab.  |
| 93  | or/81-92   |
| 94  | (acceptability or acceptable or attend\$ or access\$ or availab\$ or non-attend\$ or increas\$ or promot\$ or opt\$ or particip\$ or adhere\$ or involvement or uptake or take-up or utiliz\$ or utilis\$ or refus\$ or referr\$ or self-referr\$ or self-report\$ or barrier\$ or decreas\$ or isolation or interven\$ or aware\$ or opportunit\$ or advice or information or incentiv\$ or recruit\$ or find or finding or compliance or comply or retain or retention or provision or encour\$ or usage).ti,ab. |
| 95  | (socio sanitary support or reimburs\$ or (social adj2 support) or ((cash or financial or money or monetary or economic or voucher or credit or drug\$1 or methadone or telephone) adj2 (benefit\$ or support or incentive or assist\$ or credit))).ti,ab.  |
| 96  | ((lifestyle or behavio?r) adj2 (therapy or modif\$ or chang\$ or adapt\$ or adopt\$)) and (tuberculosis or tb)).ti,ab. or *social marketing/   |
| 97  | *marketing/  |
| 98  | *attitude to health/   |
| 99  | *health care delivery/   |
| 100 | *access to information/  |
| 101 | *confidentiality/  |
| 102 | *Health education/   |
| 103 | *health promotion/   |
| 104 | *patient attitude/ or *refusal to participate/ or *treatment refusal//   |
| 105 | *patient compliance/   |
| 106 | *motivation/   |
| 107 | Stigma.ti,ab.  |
| 108 | *prevalence/   |
| 109 | *patient participation/  |
| 110 | or/94-109  |
| 111 | treat\$.ti,ab. or Treatment Outcome/   |
| 112 | (directly observed treatment or directly observed therapy or (supervised adj2 treatment) or (coerc\$ adj2 (treat\$ or therapy))).ti,ab. Or *directly observed therapy/   |
| 113 | (disease management or (treat\$ and (management or control))).ti,ab.   |
| 114 | ((adherence or compli\$ or non-compli\$ or default\$ or finish\$ or Retention or attrition or (drop adj1 out) or disappear\$ or abscond\$) and treat\$).ti,ab. or exp *patient compliance/   |
| 115 | ((referr\$ or self-referr\$ or (self adj diagnos\$)) and treat\$).ti,ab.   |
| 116 | ((suitab\$ or eligib\$) and treat\$).ti,ab.  |
| 117 | ((follow adj1 up) or discharge).ti,ab. or *follow up/  |
| 118 | ((positive or negative) and test).ti,ab.   |
| 119 | ((interrupt\$ or relapse\$ or stop\$ or cessation or with?ld\$ or avoidance or (lost adj2 follow)) and treat\$).ti,ab. or *treatment withdrawal/   |
| 120 | ((medicine\$1 or drug or treat\$) and (regimen or adherence)).ti,ab. or exp *self care/  |
| 121 | (treat\$ and (appointment\$ or Schedule\$)).ti,ab. or *patient scheduling/   |
| 122 | ((care adj2 seeking) and pathway\$).ti,ab.   |
| 123 | ((case adj3 management) or case-managed).ti,ab. or Case Management/ or *patient care planning/ or *health insurance/   |
| 124 | (case adj3 manag\$ adj3 strategy).ti,ab. or continuity or *patient care/   |



|     |  |
|-----|--|
| 125 | ((case or treat\$ or diagnosis) and management).ti,ab.   |
| 126 | ((active or passive) and (case adj3 management)).ti,ab.  |
| 127 | ((risk assessment or care plan\$) and (case adj3 management)).ti,ab.   |
| 128 | or/111-127   |
| 129 | 1 and 39 and (80 or (93 and (110 or 128)))   |
| 130 | limit 129 to yr="1990 -Current"  |
| 131 | limit 130 to "english language"  |
| 132 | (animal\$ or badger\$ or Cow\$ or Cattle or bovine).ti,ab. or (animals/ not humans/)   |
| 133 | 131 not 132  |
| 134 | limit 133 to yr="1990 - 2010"  |
| 135 | 130 not 132  |
| 136 | 135 not 134  |
| 137 | (albania or bulgaria or cyprus or croatia or latvia or lithuania or luxembourg or malta or montenegro or romania or serbia or yugoslav or turkey).ti,ab,hw,in. |
| 138 | 1 and 135 and (80 or (93 and (110 or 128)))  |
| 139 | limit 138 to yr="1990 -Current"  |
| 140 | 139 not 132  |
| 141 | 140 not 135  |
| 142 | 136 or 141   |

## Supplementary Material IV: Evidence Statements

### *Grading of evidence*

**No evidence** – no evidence or clear conclusions from any of the studies;

**Weak evidence** – no clear or strong evidence/conclusions from high quality studies and only tentative evidence/conclusions from moderate quality studies or clear evidence/conclusions from low quality studies;

**Moderate evidence** – tentative evidence/conclusions from multiple high quality studies, or clear evidence/conclusions from one high quality study or multiple medium quality studies, with minimal inconsistencies across all studies;

**Strong evidence** – clear conclusions from multiple high quality studies.

### *Tuberculosis identification*

#### **Evidence statement 1: Effectiveness of using different types of healthcare workers on improving identification of active TB**

1-1 **Conflicting evidence** arose concerning types of healthcare workers to improve TB screening. Aldridge et al., 2015 [+] showed that the uptake of TB screening did not improve when peers encourage and educate homeless people, compared to shelter personnel.<sup>1</sup> On the contrary, Duarte et al. 2011 [-]<sup>2</sup> and Goetsch et al. 2012 [-]<sup>3</sup> showed an increase in uptake of TB screening amongst homeless people and drug users after involving community health workers (CHW) and key partners in the education and promotion of TB screening. The results demonstrated by Goetsch et al. fluctuated over the study period and both studies retrospectively compared the findings over different time periods, which might be an important source of bias.

1-2 **Moderate evidence** from two studies focussing on the identification of TB contacts by using peers. One study identified by this review, Ospina et al. 2012 [+], showed that contact tracing among migrants improved significantly by using CHWs from the same migrant community to coordinate contact tracing.<sup>4</sup> The comparison group was recruited over a different time period, which reduced the quality of this study.

Ricks, 2008 [++],<sup>5</sup> identified by the NICE review,<sup>6</sup> suggested that contact tracing of drug users with active TB improved by using former drug users compared to healthcare workers. It was unclear if the improvement could be contributed to the use of peers, as the study also used enhanced case management for the peer-led intervention group, the control group received limited case management.

#### **Evidence statement 2: Effectiveness of using different settings on improving identification of active TB**

2-1 **Weak evidence** from Jit et al. 2011 [+] showed that a mobile TB service is an effective intervention to identify hard-to-reach individuals with TB as 35% of the cases identified by this mobile

TB service were asymptomatic.<sup>7</sup>

Four studies were identified by the NICE review,<sup>6</sup> providing the following evidence:

2-2 **Weak evidence** from Mor et al., 2008 [-] suggesting that pre-migration screening in Ethiopian migrants reduced the risk of developing active TB in Israel and reduced the time from entry into the host country (Israel) until TB diagnosis compared to post-migration screening.<sup>8</sup> The study compared two migrant groups recruited over two different time period what limited the quality of the findings.

2-3 **Weak evidence** from El-Hamad et al., 2001 [+] suggesting that active case finding conducted in a TB clinic improved TB screening completion among migrants compared to TB screening conducted in a non-specialist primary care facility.<sup>9</sup> However, the study did not adjust for the differences in baseline characteristics.

2-4 **Weak evidence** from Bothamley et al., 2002 [-] showing that TB screening by symptom-based questionnaire was less useful in port-of-arrival clinics compared to homeless centres.<sup>10</sup> In 90.9% of the migrants screened by questionnaire further TB screening was conducted compared to 100% of the questionnaire screened homeless people. The study findings were limited as the study compared two different hard-to-reach groups in two different settings.

2-5 **Weak evidence** from Miller et al., 2006 [+] suggesting that there was no significant statistical difference in screening uptake in the homeless centre where screening was optional and a prison where screening was mandatory, however, the homeless people received incentives for TB screening.<sup>11</sup> In the homeless centre more cases were detected but different methods were used. The evidence was weak as the study compared two groups with possibly different baseline characteristics, different TB prevalence and different tests were offered.

### **Evidence statement 3: Cost-effectiveness of using different settings on improving identification of active TB**

3-1 **Weak evidence** from Jit et al. 2011 [+] that a mobile TB service is cost-effective, the incremental cost ratio for the screening service was £18,000 per Quality of Life Year (QALY) gained.<sup>7</sup>

A further two studies were identified by the NICE review.<sup>6</sup>

3-2 **Weak evidence** from Mor et al., 2008 [-] suggesting that pre-migration screening in Ethiopian migrants would save Israel \$449,817 over a five year period.<sup>8</sup> Different sources were used for the cost analysis, with varying reliability, limiting the findings.

3-3 **Weak evidence** from Bothamley et al., 2002 [-] showed that TB screening among migrants as part

of the port-of-arrival programme had a cost-saving of £12.70 per person screened, it would cost an additional £0.50 per person screened at a homeless centre and £7.00 per migrant screened at a general practice.<sup>10</sup> The findings of this study are limited as the study compared two different populations with possible different characteristics and TB prevalence.

### *Tuberculosis management*

#### **Evidence statement 4: Effectiveness of using different types of healthcare workers on improving management of active TB**

**Moderate evidence** from one study identified by the NICE review<sup>6</sup>, Ricks, 2008 [++], suggested that drug users receiving enhanced case management by peers were more likely to complete TB treatment than drug users who received limited case management by healthcare workers (RR = 2.68, 95%CI 1.24-5.82;  $p = 0.01$ ).<sup>5</sup> The findings of this study are limited as the intervention group received enhanced case management and were managed by peers, therefore it is unclear which of the two contributed (more) to the improved treatment outcome.

#### **Evidence statement 5: Effectiveness of using different settings on improving management of active TB**

5-1 **Weak evidence** from Jit et al. 2011 [+] that treatment completion rate in patients treated by the mobile 'Find and Treat' service was higher (67.1%) than in the patients receiving standard care at a TB clinic (56.8%) and lost to follow-up rate in the patients treated by the mobile 'Find and Treat' service was lower (2.1%) than in the patients receiving standard care at a TB clinic (17.2%).<sup>7</sup>

5-2 **Weak evidence** from Deruaz & Zellweger, 2004 [-], showed that there was no statistical difference in treatment outcome when directly observed treatment was delivered at the healthcare facility or at a convenient site in the community in mixed hard-to-reach groups.<sup>12</sup> The findings were limited by a potential selection bias and the way the treatment outcome was collected differed in both groups.

#### **Evidence statement 6: Cost-effectiveness of using different settings on improving management of active TB**

6-1 **Weak evidence** from Jit et al. 2011 [+] that a mobile TB management service is cost-effective, the incremental cost ratio for the service was £4,100 per Quality of Life Year (QALY) gained.<sup>7</sup>

## References

1. Aldridge R, Yates S, Hemming S, et al. Impact of peer educators on uptake of mobile x-ray tuberculosis screening at homeless hostels: A cluster randomised controlled trial. *Thorax* 2014; **69**: A44-A5.
2. Duarte R, Santos A, Mota M, Carvalho A, Marques A, Barros H. Involving community partners in the management of tuberculosis among drug users. *Public Health* 2011; **125**(1): 60-2.
3. Goetsch U, Bellinger OK, Buettel KL, Gottschalk R. Tuberculosis among drug users and homeless persons: impact of voluntary X-ray investigation on active case finding. *Infection* 2012; **40**(4): 389-95.
4. Ospina JE, Orcau A, Millet JP, Sanchez F, Casals M, Cayla JA. Community health workers improve contact tracing among immigrants with tuberculosis in Barcelona. *BMC public health* 2012; **12**: 158.
5. Ricks PM. Tuberculosis control among substance users: The indigenous leadership outreach model vs. standard care: University of Illinois at Chicago; 2008.
6. Rizzo M, Martin A, Jamal F, et al. Evidence review on the effectiveness and cost-effectiveness of service models or structures to manage tuberculosis in hard-to-reach groups. Matrix evidence: NICE – National Institute for Health and Care Excellence, 2011.
7. Jit M, Stagg H, Aldridge R, White P, Abubakar I. Dedicated outreach service for hard to reach patients with tuberculosis in London: observational study and economic evaluation. *BMJ* 2011; **343**: d5376.
8. Mor Z, Lerman Y, Leventhal A. Pre-immigration screening process and pulmonary tuberculosis among Ethiopian migrants in Israel. *European Respiratory Journal* 2008; **32**(2): 413 - 8.
9. El-Hamad I, Casalini C, Matteelli A, et al. Screening for tuberculosis and latent tuberculosis infection among undocumented immigrants at an unspecialised health service unit. *International Journal of Tuberculosis & Lung Disease* 2001; **5**(8): 712-6.
10. Bothamley GH, Rowan JP, Griffiths CJ, et al. Screening for tuberculosis: the port of arrival scheme compared with screening in general practice and the homeless *Thorax* 2002; **57**(1): 45-9.
11. Miller TL, Hilsenrath P, Lykens K, McNabb SJ, Moonan PK, Weis SE. Using cost and health impacts to prioritize the targeted testing of tuberculosis in the United States. *Annals of epidemiology* 2006; **16**(4): 305-12.
12. Dèruaz J, Zellweger J, P. . Directly observed therapy for tuberculosis in a low prevalence region: first experience at the Tuberculosis Dispensary in Lausanne. *Swiss medical weekly* 2004; **134**: 552-8.

Supplementary Material V: Data extraction forms

| Study details   | Population and setting   | Method of allocation to intervention/control  | Outcomes and methods of analysis   | Results  | Note by review team  |
|---|--|---|--|--|--|
| <p><b>Year:</b><br/>2015</p> <p><b>Country:</b><br/>United Kingdom (UK)</p> <p><b>Authors:</b><br/>Aldridge R.W., Hayward A.C., Hemming S. et al.<sup>1</sup></p> <p><b>Citation:</b><br/>Aldridge R.W., Hayward A.C., Hemming S. et al. Effectiveness of peer educators on the uptake of mobile X-ray tuberculosis screening at homeless hostels: a cluster randomised controlled trial. <i>BMJ open</i> 2015; 5: e008050.</p> <p><b>Aim of study:</b></p> | <p><b>Source population(s):</b><br/>Homeless people</p> <p><b>Eligible population:</b><br/>Hostels for homeless people in London, UK</p> <p><b>Selected population:</b><br/>Homeless people at hostels in London, UK</p> <p>All hostels in London who participated in 2 previous screening sessions were eligible for inclusion. Eligible hostels had not been screened in the 6 months prior to the scheduled mobile X-ray Unit (MXU) screening session.</p> <p><b>Excluded population:</b></p> | <p><b>Method of allocation:</b><br/>Sites were randomized to intervention or control group by Sealed Envelope™ <a href="http://www.sealedenvelope.com/">http://www.sealedenvelope.com/</a> which ensured allocation concealment until interventions were assigned. To ensure comparability between intervention and control arms, hostels were stratified on the basis of their size (binary variable indicating whether hostels had more than 43 beds) and previous screening uptake level (binary variable indicating whether hostels had greater than 50% historical</p> | <p><b>Primary outcomes:</b><br/>The number of eligible clients at a hostel venue screened for active pulmonary tuberculosis by MXU</p> <p><b>Secondary outcomes:</b><br/>Secondary analysis was also conducted by subgroups, for large and small hostels separately, and low and high previous screening uptake level using the binary categorical variables described previously for these analyses.</p> <p><b>Method of analysis:</b><br/>The study statistician conducted analysis blinded to the allocation of</p> | <p><b>Primary results:</b><br/><u>Control sites:</u><br/>- 1,192 residents<br/>- Median uptake of 45% (inter-quartile range (IQR):33,55)</p> <p><u>Intervention sites:</u><br/>-1,150 eligible residents<br/>- Median uptake of 40% (IQR:25,61)</p> <p>Poisson regression to account for the clustered study design, hostel size and historical screening levels, there was no evidence that peer educators increased uptake (adjusted risk ratio 0.98; 95% confidence interval (95%CI): 0.80,1.20).</p> <p><b>Secondary results:</b><br/>No evidence was found for peer educators</p> | <p><b>Limitations identified by author:</b><br/>- The study design was not powered to detect a difference in tuberculosis cases identified by the two arms as this would require considerably larger sample sizes and would have meant repeated screening at hostels, potentially diluting the effect of the intervention during the study<br/>- Unable to collect individual data as part of the study as this would have required individual consent and would have been challenging logistically given that screening took place within an operational setting where any data collection would have</p> |

|  |   |  |  |  |  |
|--|---|--|--|--|--|
| <p>To compare current practice for encouraging homeless people to be screened for tuberculosis by a mobile digital X-ray unit in London, UK, with the additional use of volunteer peer educators who have direct experience of tuberculosis and homelessness.</p> <p><b>Study design:</b><br/>Cluster randomised controlled trial</p> <p><b>Quality score:</b><br/>+</p> <p><b>Applicability:</b><br/>++</p> | <p>- Hostels with uptake levels over 80% were excluded from the study – as low chance of intervention having impact</p> <p>- Not allowing peers on the venue</p> <p><b>Setting:</b><br/>Homeless hostels in London, UK</p> <p><b>Sample characteristics:</b><br/>Majority of hostels had 43 or less beds, reported a historical screening uptake of &gt;50% and provided no incentives for screening.</p> | <p>uptake).</p> <p>A cluster randomised design was chosen as the intervention was aimed at the hostel sites rather than individuals clients</p> <p><b>Intervention(s) description:</b><br/>Peers moved around the hostel in conjunction with hostel staff and spoke to residents in order to encourage them to attend screening.</p> <p><b>Comparator/control(s) description:</b><br/>Encouragement by hostel staff</p> <p><b>Baseline comparisons:</b><br/>Proportion of people screened for TB</p> <p><b>Study sufficiently powered?:</b><br/>Yes, power calculation done before hand, met the sample size</p> | <p>intervention or control arms.</p> <p>Poisson regression analysis was used to analyse outcome events at screening hostels. Bed occupancy level was included as the exposure variable, screening uptake as the outcome (or indicator) variable, and hostel venue as a random effect to account for clustering at each site. The analysis was adjusted by inclusion of the randomisation stratification factors of historical uptake rates and bed size.</p> <p><b>Modelling method and assumptions:</b><br/>Clustered analysis, assuming that characteristics of the populations at respective hostels were comparable.</p> <p><b>Time horizon:</b><br/>February 2012 and</p> | <p>increasing uptake of screening for any of the other secondary analyses. The study team noted no adverse events.</p> | <p>interrupted the flow of screening and caused unacceptable delays for service users</p> <p>- Peers may lack technical knowledge and the confidence with which to challenge some of the client misconceptions and concerns that reduce screening uptake</p> <p>- Most effective when complemented i.e. delivered in conjunction with professionals</p> <p><b>Limitations identified by review team:</b></p> <p>- Hostels with high uptake were excluded = bias risk</p> <p>- Most sites were not naïve to peer intervention</p> <p><b>Evidence gaps and/or recommendations for future research:</b></p> <p>- Any difference in uptake with previous screening sessions?</p> <p>- Effectiveness of using</p> |
|--|---|--|--|--|--|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |  |  |              |  |  |
|--|--|--|--------------|--|--|
|  |  | requirement of 1045 individuals, or approximately 21 hostels in each arm | October 2013 |  | <p>peers as a standalone intervention versus peers working alongside professionals and this remains an important research question.</p> <p><b>Source of funding:</b><br/>NIHR and Wellcome Trust.</p> <p><b>Conflict of interests:</b><br/>None for most authors<br/>- AS (last author) is clinical lead for the Find and Treat service including the mobile digital X-ray unit.</p> |
|--|--|--|--------------|--|--|

| Study details   | Population and setting   | Method of allocation to intervention/control   | Outcomes and methods of analysis  | Results  | Note by review team   |
|---|--|--|---|--|---|
| <p><b>Year:</b><br/>2012</p> <p><b>Country:</b><br/>Germany<br/>(Frankfurt/Main)</p> <p><b>Authors:</b><br/>Goetsch U., Bellinger O.K., Buettel K.L., Gottschalk R.<sup>2</sup></p> | <p><b>Source population(s):</b><br/>Homeless &amp; drug users</p> <p><b>Eligible population:</b><br/>Homeless &amp; drug users recruited from homeless and drug services in Frankfurt/Main</p> | <p><b>Method of allocation:</b><br/>Not applicable (NA)</p> <p><b>Intervention(s) description:</b><br/>Community health worker educated staff and users of services for homeless and drug users about TB</p> | <p><b>Primary outcomes:</b><br/>Feasibility and sustainability of a TB programme focussing on TB education and voluntary X-ray investigation in homeless and drug users</p> <p><b>Secondary</b></p> | <p><b>Primary results:</b><br/>It is feasible when included in already existing public health services</p> <p><b>Secondary results:</b><br/>- No. CXR: 10/month in homeless &amp; 9/month in drug users<br/>After intervention</p> | <p><b>Limitations identified by author:</b><br/>- Selection bias, illegal immigrants might avoid authorities<br/>- Small number of TB patients makes it difficult to say anything about age and gender differences<br/>- No data on length of</p> |



|   |   |  |   |  |   |
|---|---|--|---|--|---|
| <p><b>Citation:</b><br/>Goetsch U., Bellinger O.K., Buettel K.L., Gottschalk R. Tuberculosis among drug users and homeless persons: impact of voluntary X-ray investigation on active case finding. <i>Infection</i> 2012; <b>40</b>: 389-95.</p> <p><b>Aim of study:</b><br/>To evaluate the feasibility and sustainability of the program, its coverage and both the case-finding rates and characteristics of cases. Also to assess the treatment outcomes</p> <p><b>Study design:</b><br/>Before and after comparison</p> <p><b>Quality score:</b><br/>-</p> <p><b>Applicability:</b><br/>+</p> | <p><b>Selected population:</b><br/>All subjects seen at the Public Health Department for chest X-rays (CXR) and fulfilling the criteria for homeless (stayed at shelter for &gt;2 nights) /drug users (attend day-care facilities, night shelter for drug users or needle exchange programme)</p> <p><b>Excluded population:</b><br/>Patients with tuberculosis (TB) symptoms detected in clinics and notified through the Protection against Infection Act</p> <p><b>Setting:</b><br/>Community health worker (CHW) went to services to promote CXR – CXR performed at Public Health Department</p> <p><b>Sample</b></p> | <p>transmission and promoted voluntary CXR at Public Health Department 1x/year or at least 1x/2years</p> <p>Community Health Worker obtained the medical history through standardised questionnaire</p> <p>CXR read by TB physician – referral and follow up test in a clinic could be initiated immediately</p> <p>Suspicion for active TB – CHW took care of further diagnostics and follow up</p> <p>Active TB needed hospitalisation for treatment</p> <p>CHW kept contact with doctors/social workers 2x/month, later monthly</p> <p>Contact tracing in shelter</p> <p>HIV was only notified in active TB</p> | <p><b>outcomes:</b><br/>Estimate the coverage of the programme, assess other risk factors and determine TB rates &amp; treatment outcome in the 2 groups</p> <p><b>Method of analysis:</b><br/>- t-test or analysis of variance for continuous variables<br/>- chi-square test or Fisher's exact test for categorical data</p> <p><b>Modelling method and assumptions:</b><br/>- Multivariate logistic regression effect of risk groups, birth place, age &amp; gender</p> <p><b>Time horizon:</b><br/>1 May 2002 – 30 April 2007</p> | <p>46/month in homeless &amp; 25/month in drug users</p> <p><b>Coverage:</b><br/>- Screening 1x/2 years: 18% of drug users &amp; 26% of homeless<br/>- Screening every year: 10% of drug users &amp; 15% of homeless (based on the range of drug users &amp; homeless people between 6,416 and 9,000 individuals in Frankfurt/Main)<br/>- Chao's heterogeneity model: 18-26.3%<br/>1 CXR/2 years<br/>- 2002-2004: 18.0%<br/>- 2003-2005: 19.3%<br/>- 2004-2006: 26.4%<br/>- 2005-2007: 23.4%<br/>and 10-15% CXR/year<br/>- 2002-2004: 10.0%<br/>- 2003-2005: 10.7%<br/>- 2004-2006: 15.0%<br/>- 2005-2007: 23.4%</p> <p><b>Case finding:</b><br/>- 39 TB cases in 5 years: 14 drug users &amp; 25 homeless = 8.7% of total TB cases in Frankfurt<br/>- 19 cases smear +, 7</p> | <p>drug use and homelessness<br/>- The impact of HIV can't be estimated<br/>- Unknown fluctuations of the study population make the denominator unstable</p> <p><b>Limitations identified by review team:</b><br/>- Patients had to travel to the public health department<br/>- Comparison over time, important confounder<br/>- Not adjusted for distance from service to public health department</p> <p><b>Evidence gaps and/or recommendations for future research:</b><br/>Use a control group and use of mobile CXR unit to increase screening uptake</p> <p><b>Source of funding:</b><br/>Not recorded (NR)</p> <p><b>Conflict of interests:</b><br/>None</p> |
|---|---|--|---|--|---|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |  |   |  |  |  |
|--|--|---|--|--|--|
|  | <p><b>characteristics:</b><br/>4,529 CXRs in 3,477 people<br/>- 66% homeless<br/>- 34% drug users</p> <p><u>Homeless:</u><br/>- 40.9 years ± 12.5 years<br/>- 90.1% male<br/>- 29.6% foreign born</p> <p><u>Drug users:</u><br/>- 35.8 years ± 8.3 years<br/>- 76.2% male<br/>- 28.0% foreign born (increased over study period → 2002: 15%, 2007:37%)</p> | <p>patients</p> <p><b>Comparator/control(s) description:</b><br/>Before intervention – no CHW who gave TB education and promoted CXR</p> <p><b>Baseline comparisons:</b><br/>Coverage of CXR screening before and after intervention</p> <p><b>Study sufficiently powered?:</b><br/>Low number of active TB cases</p> |  | <p>smear –ve but culture +ve<br/>- 13 cases clinical/ radiological diagnosis<br/>- Case finding rate 861/100 000 CXRs<br/>- Drug users 10/14 HIV +ve, homeless 1/25 HIV +ve (p&lt;0.001)</p> <p>No significant difference in case-finding rates according to gender (906/100,000 for men vs. 601/100,000 for women; p = 0.43), place of origin (906/100,000 for natives vs. 752/100,000 for foreign born; p = 0.61) or between the risk groups (888/100,000 for the homeless vs. illegal drug users (816/100,000; p = 0.80).</p> <p>Logistic regression model with the risk factors age, gender, risk group and place of origin: age was the only variable to be associated with the risk of TB and the variable risk group was an effect modifier</p> |  |
|--|--|---|--|--|--|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |  |  |  |  |  |
|--|--|--|--|--|--|
|  |  |  |  | <p><u>Treatment completion</u><br/>                 - 76.3% (29/38) completed treatment (one patient left Frankfurt to unknown destination before completion of treatment, excluded from analysis of treatment completion) – of which 19 homeless people and 10 drug users<br/>                 - 5 people needed admission because of non-compliance (3 drug users, 2 homeless)<br/>                 - 5 people died of other causes than TB (3 homeless and 2 drug users)<br/>                 - 4 people stopped treatment (lack of compliance) – 10.5%</p> <p>No difference in the treatment outcome between the two risk groups (p = 0.96), and age, gender or HIV-status did not have any significant effect on treatment outcome.</p> |  |
|--|--|--|--|--|--|

| Study details  | Population and setting   | Method of allocation to intervention/control   | Outcomes and methods of analysis  | Results  | Note by review team  |
|--|--|--|---|--|--|
| <p><b>Year:</b><br/>2012</p> <p><b>Country:</b><br/>Spain, Barcelona</p> <p><b>Authors:</b><br/>Ospina J.E., Orcau A., Millet J-P. et al.<sup>3</sup></p> <p><b>Citation:</b><br/>Ospina J.E., Orcau A., Millet J-P. et al. Community health workers improve contact tracing among immigrants with tuberculosis in Barcelona. <i>BMC public health</i> 2012; <b>12</b>: 158.</p> <p><b>Aim of study:</b><br/>To evaluate the effectiveness of an intervention with CHW to improve contact tracing among immigrants</p> <p><b>Study design:</b></p> | <p><b>Source population(s):</b><br/>Migrants</p> <p><b>Eligible population:</b><br/>Migrants in Barcelona</p> <p><b>Selected population:</b><br/>All TB cases registered by the TBPCP (the Barcelona TB Control Program) between January 1st 2000 and December 31st 2005, and resident in the city of Barcelona were included</p> <p><b>Excluded population:</b><br/>Not registered TB patients</p> <p><b>Setting:</b><br/>Barcelona TB program</p> <p><b>Sample</b></p> | <p><b>Method of allocation:</b><br/>All TB cases registered at TBPCP</p> <p><b>Intervention(s) description:</b><br/>Community health workers (CHW) actively follow up the cases and contacts, visit the cases at home, accompany case to outpatient appointments, provide counselling and information on treatments. Educational sessions in health care centres, private homes and immigrant associations. Assistance with obtaining residence permits, housing, food banks, public dining halls and health card application.</p> | <p><b>Primary outcomes:</b><br/>The influence on contact tracing of CHW intervention</p> <p><b>Secondary outcomes:</b><br/>The influence on contact tracing of other variables: sex, age, hospital of diagnosis, district of residence, birthplace, HIV, homeless</p> <p><b>Method of analysis:</b><br/>- Descriptive analysis was performed by calculating proportions<br/>- The median and interquartile range were calculated for quantitative variables<br/>- Categorical variables were compared using the X<sup>2</sup> test. Odds ratio (OR) and 95%CI were calculated as a measure of</p> | <p><b>Primary results:</b><br/>- The increase in contact tracing coverage of contacts of smear positive PTB and all clinical forms of TB in the intervention period was statistically significant<br/>* 81% (post) vs 65.7% (pre)<br/>*OR = 1.6 (95%CI = 1.2-2.0)<br/>* Adjusted OR 1.8 (95%CI: 1.3-2.5)<br/>p &lt; 0.001</p> <p>- CHW intervention had an aOR of 2.4 (95%CI: 1.3-4.3)<br/>p=0.005 to fail contact tracing for smear-positive cases and an aOR of 1.8 (95%CI: 1.3-2.5); p = 0.001 for all TB cases</p> <p><b>Secondary results:</b><br/>- Factors associated with failure to conduct contact tracing for</p> | <p><b>Limitations identified by author:</b><br/>- Variation in characteristics between both periods; an increase of cases between 25-39 years of age, from Latin America and India, Pakistan and from inner-city in the CHW group. The increase in immigrants would most likely have worsened contact tracing and therefore our figures may have underestimated the benefit of the CHW intervention.</p> <p><b>Limitations identified by review team:</b><br/>- Wide 95% CI's smear +ve making the OR less precise for:<br/>* Hospital D<br/>* Area of origin<br/>* District of residence unknown<br/>*HIV +ve</p> |

|  |  |  |  |   |  |
|--|--|--|--|---|--|
| <p>Quasi-experimental study with historical pre-post comparison</p> <p><b>Quality score:</b><br/>+</p> <p><b>Applicability:</b><br/>++</p> | <p><b>characteristics:</b><br/>572 cases pre-intervention &amp; 388 post-intervention</p> <p><b>Pre-intervention</b><br/>- 202 (35.3%) were from Latin American countries<br/>- 136 (23.8%) from India or Pakistan<br/>- 92 (16.1%) from North Africa<br/>- 142 cases (24.8%) from other countries</p> <p>- Majority men<br/>- 25-39 years<br/>- 72.2% Pulmonary TB<br/>- 35.3% smear +ve<br/>- 34.4% lived in inner city</p> <p><b>Post intervention</b><br/>- 152 (39.2%) were from Latin American countries<br/>- 112 (28.9%) from India or Pakistan<br/>- 42 (10.8%) from North Africa<br/>- 16 (4.1%) from Sub-Saharan Africa<br/>- 66 cases (17%) from other countries</p> | <p><b>Comparator/control(s) description:</b><br/>Pre-intervention</p> <p><b>Baseline comparisons:</b><br/>Pre- /post-intervention contact tracing was compared</p> <p><b>Study sufficiently powered?</b><br/>Yes; 95% CI sufficiently narrow</p> | <p>association.</p> <p><b>Modelling method and assumptions:</b><br/>Stepwise backward logistic regression; variables lacking significant association measures in univariate analysis were assumed not to be confounders.</p> <p><b>Time horizon:</b><br/>1st 2000 and December 31st 2005</p> | <p>smear +ve TB include:<br/>* Diagnosed in hospitals B and D<br/>* Birthplace in India-Pakistan or North Africa<br/>* Unknown district of residence<br/>* HIV infection<br/>* Homeless</p> <p>- Factors associated with failure to conduct contact tracing for all forms of TB include:<br/>* Male<br/>* Hospitals B and D<br/>* Birthplace other than Latin American countries<br/>* Unknown district of residence<br/>* Incarceration history<br/>* Homeless<br/>* Index who had culture-negative or extra-pulmonary TB or had a normal chest X-ray</p> <p>- Of all TB cases identified among foreign born people, 79.4% were seen by CHWs, 12.4% were contacted directly by</p> | <p>* Homeless</p> <p>- Wide 95% CI's all TB: making the OR less precise for:<br/>* District of residence unknown<br/>* Incarceration<br/>* Homeless</p> <p>95% CI's crossing 1 in hospital D</p> <p><b>Evidence gaps and/or recommendations for future research:</b><br/>- Studies on cost-effectiveness of the CHW interventions in the TB programs</p> <p><b>Source of funding:</b><br/>NR</p> <p><b>Conflict of interests:</b><br/>None</p> |
|--|--|--|--|---|--|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |  |  |  |  |  |
|--|--|--|--|--|--|
|  | <ul style="list-style-type: none"> <li>- Majority men</li> <li>- 25-39 years</li> <li>- 73.2% Pulmonary TB</li> <li>- 39.2% smear +ve</li> <li>- Almost 50% lived in inner city</li> </ul> |  |  | <p>the public health nurse and the remaining 8.2% could not be contacted</p> <ul style="list-style-type: none"> <li>- Active follow up in 194 TB cases and contact census, a total of 293 counselling sessions, 147 linguistic mediation session, 264 individualised and 97 group educational sessions about TB, 280 home visits, 70 hospital visits and 5,935 telephone calls (a median of 15.3 calls per case) were performed</li> </ul> |  |
|--|--|--|--|--|--|

| Study details  | Population and setting   | Method of allocation to intervention/control  | Outcomes and methods of analysis   | Results   | Note by review team   |
|--|--|---|--|---|---|
| <p><b>Year:</b> 2011</p> <p><b>Country:</b> Portugal</p> <p><b>Authors:</b> Duarte R., Santos A., Mota M. et al.<sup>4</sup></p> | <p><b>Source population(s):</b> Drug users in Vila Nova de Gaia, Portugal</p> <p><b>Eligible population:</b> Drug users in Vila Nova de Gaia</p> | <p><b>Method of allocation:</b> Before and after 2004 – intervention was implemented in 2004</p> <p><b>Intervention(s) description:</b> After 2004:</p> | <p><b>Primary outcomes:</b> Diagnosis of active TB, treatment compliance before and after intervention</p> <p><b>Secondary outcomes:</b> OR and 95%CI to measure association</p> | <p><b>Primary results:</b> 2001-2003:</p> <ul style="list-style-type: none"> <li>- 125 drug users observed in CDP of which:</li> <li>* 52 drug users were screened for TB (100% male, mean age 32 years)</li> <li>* 73 drug users were</li> </ul> | <p><b>Limitations identified by author:</b></p> <ul style="list-style-type: none"> <li>- Not a controlled trial</li> <li>- Risk for bias</li> <li>- Unknown what part of the intervention contributed to the outcome</li> </ul> |

|   |  |   |   |  |   |
|---|--|---|---|--|---|
| <p><b>Citation:</b><br/>Duarte R., Santos A., Mota M. et al. Involving community partners in the management of tuberculosis among drug users. <i>Public Health</i> 2011; <b>125</b>: 60-2.</p> <p><b>Aim of study:</b><br/>To evaluate the effect of the intervention on diagnosis of TB and treatment compliance</p> <p><b>Study design:</b><br/>Retrospective review of records<br/>Comparison before and after intervention (2004)</p> <p><b>Quality score:</b><br/>-</p> <p><b>Applicability:</b><br/>+</p> | <p>(VNdG), Portugal</p> <p><u>Population VNdG:</u><br/>290,000</p> <p>Estimated number of drug users = 4.3-6.4 per 1,000 inhabitants</p> <p><b>Selected population:</b><br/>Screening and treatment records for all drug users visiting Chest Disease Centre (CDP) between 2001-2007</p> <p><b>Excluded population:</b><br/>NR</p> <p><b>Setting:</b><br/>All drug users screened and treated at the outpatient TB clinic (Chest Disease Centre) 2001-2007 were reviewed</p> <p><b>Sample characteristics:</b><br/><u>2001-2003:</u><br/>- 125 drug users @CDP<br/>- 52 screened (100%</p> | <p>Intervention to improve early identification and treatment of drug users with TB.</p> <p>The key partners (outpatient TB clinic, drug users support centres, shelters and street teams, local public health department and the local hospital) identified drug users in their population</p> <ul style="list-style-type: none"> <li>- Promotion of health-seeking behaviour</li> <li>- Notification card for screening in CDP</li> <li>- Elimination of potential barriers: <ul style="list-style-type: none"> <li>* Street teams offered free transport</li> <li>* All care at CDP free of charge</li> </ul> </li> <li>- Encouraged referral but tried to manage TB screening locally</li> <li>- Seriously ill: immediate referral to CDP/local hospital (with transport and attendance)</li> </ul> | <p><b>Method of analysis:</b><br/>OR and 95%CI were used to measure the strength of the association</p> <p><b>Modelling method and assumptions:</b><br/>Improve early identification and treatment of drug users with TB</p> <p><b>Time horizon:</b><br/>2001-2003<br/>Intervention<br/>2005-2007</p> | <p>referred to CDP because of symptoms or following discharge from hospital with diagnosis TB</p> <p>Of all drug users observed at CDP; 82 drug users (65.6%) were diagnosed with active TB of which 11 drug users (13.4%) were identified by screening</p> <p><u>Treatment compliance:</u><br/>- 47.6% poor compliance<br/>- 35.4% stopped TB treatment</p> <p><u>TB in VNdG 2001-2003:</u><br/>- 515 TB cases of which 82 were drug users<br/>- 32 TB cases died of which 15 were drug users (OR 4.66, 95%CI 2.24 - 9.70).<br/>- TB/HIV co-infection: 63 (71%)</p> <p><u>2005-2007:</u> (after implementation of the programme)<br/>- 465 drug users were screened for TB (86%</p> | <p><b>Limitations identified by review team:</b></p> <ul style="list-style-type: none"> <li>- Retrospective design = risk of bias</li> <li>- Methods not well described</li> <li>- Unknown what percentage did not come for screening (how many people recruited for screening)</li> <li>- Difference in time zone = risk for confounders, might have been on the political agenda, been on the news etc. = bias</li> <li>- Low precision of estimates of effects (indicated by wide 95%CI) probably due to small sample size</li> </ul> <p><b>Evidence gaps and/or recommendations for future research:</b></p> <ul style="list-style-type: none"> <li>- Case-control trial to compare 2 different cities (one with intervention other without intervention)</li> <li>- Check cost-effectiveness</li> </ul> <p><b>Source of funding:</b></p> |
|---|--|---|---|--|---|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |  |   |  |   |  |
|--|--|---|--|---|--|
|  | <p>male, mean age 32 years)<br/>- 73 for symptoms or following discharge with diagnosis TB</p> <p><u>2005-2007:</u><br/>- 465 screened (86% male, mean age 36 years)<br/>- 30 for symptoms or following discharge with diagnosis TB</p> <p><b>Study definitions:</b><br/>Active TB: culture <i>M. tuberculosis</i> or clinical &amp; radiology criteria<br/>Latent TB: asymptomatic individuals with normal chest radiography and positive TST (TST &gt; 5 mm in immune-compromised persons, TST &gt; 10 mm in immune-competent persons)</p> | <p><u>At CDP:</u><br/>- Screening: symptom questionnaire, tuberculin skin test (TST) &amp; CXR: annual screening/after contact/symptoms<br/>- Directly observed therapy short-course (DOTS) at CDP, combined with other medical treatment/<br/>drug abuse treatment<br/>- CDP offered HIV testing in case of active TB</p> <p><b>Comparator/control(s) description:</b><br/><u>Before 2004:</u><br/>- Drug users referred to CDP with a diagnosis of TB after diagnosis from hospital<br/>- Treatment was not compulsory<br/>- To improve compliance: info was provided, psychosocial support, full treatment to patient and family (if</p> |  | <p>male, mean age 36 years)<br/>- 30 drug users were referred to CDP because of symptoms or following discharge from hospital with diagnosis TB</p> <p>Of all drug users seen at the CDP, 59 drug users (11.9%) were diagnosed with active TB of which 36 drug users (61%) were identified by screening</p> <p><u>Treatment compliance:</u><br/>- 23.7% poor compliance<br/>- 10.2% stopped treatment</p> <p><u>TB in VNdG 2005-2007:</u><br/>- 386 TB cases of which 59 were drug users<br/>- 19 TB cases died of which 8 were drug users (OR 4.66, 95%CI 1.79 - 12.15)<br/>- TB/HIV co-infection: 37 (64%)</p> <p><u>Conclusion:</u><br/>The number of screened</p> | <p>None</p> <p><b>Conflict of interests:</b><br/>None</p> <p><b>Ethical approval:</b><br/>Yes, approved by the CDP de Vila Nova de Gaia body</p> |
|--|--|---|--|---|--|



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |  |  |  |  |  |
|--|--|--|--|--|--|
|  |  | <p>needed), transport &amp; free breakfast.<br/>- No active screening policy</p> <p><b>Baseline comparisons:</b><br/>Number of TB cases screened</p> <p><b>Study sufficiently powered:</b><br/>NR but wide 95%CI</p> |  | <p>drug users had increased, therapy was available to a higher proportion of TB cases and active TB treatment compliance had improved significantly</p> <p><b>Secondary results:</b><br/>- Drug users screened for TB without symptoms: OR 21.8; 95%CI 13.0-36.3<br/>- Drug users with active TB: OR 10.1; 95%CI 4.44-23.0<br/>- Poor compliance: OR 0.34; 95%CI 0.16-0.72<br/>- Treatment stopped OR 0.21; 95%CI 0.08-0.54<br/>- % of drug users under TB cases OR 0.95; 95%CI 0.66-1.37</p> <p>- Mortality due to TB decreased from 18.3% in 2001- 2003 to 13.6% in 2005-2007 among drug users (OR 0.7, 95%CI 0.28 - 1.78).</p> <p>- TB/HIV co-infection decreased from 63 (71%) in 2001-2003 to</p> |  |
|--|--|--|--|--|--|

|  |  |  |  |   |  |
|--|--|--|--|---|--|
|  |  |  |  | 37 (64%) in 2005-2007 (OR 1.37, 95%CI 0.68-2.78). |  |
|--|--|--|--|---|--|

| Study details  | Population and setting  | Intervention/comparator  | Outcomes and methods of analysis  | Results  | Note by review team  |
|--|---|--|---|--|--|
| <p><b>Year:</b><br/>2011</p> <p><b>Country:</b><br/>UK</p> <p><b>Authors:</b><br/>Jit M. Stagg H.R.,<br/>Aldridge R. et al.<sup>3</sup></p> <p><b>Citation:</b><br/>Jit M. Stagg H.R.,<br/>Aldridge R. et al.<br/>Dedicated outreach<br/>service for hard to<br/>reach patients with<br/>tuberculosis in<br/>London:<br/>observational study<br/>and economic<br/>evaluation. <i>BMJ</i><br/>2011; <b>343</b>: d5376.</p> <p><b>Aim of study:</b><br/>To assess the cost<br/>effectiveness of the<br/>Find and Treat</p> | <p><b>Source population(s):</b><br/>Hard-to-reach<br/>individuals</p> <p><b>Eligible population:</b><br/>Hard-to-reach<br/>individuals with<br/>active pulmonary<br/>tuberculosis</p> <p><b>Selected population:</b><br/>Hard to reach<br/>individuals with<br/>active pulmonary<br/>tuberculosis screened<br/>or managed by the<br/>Find and Treat<br/>service</p> <p><b>Excluded population:</b><br/>- Cases with extra-<br/>pulmonary<br/>tuberculosis<br/>- Latent tuberculosis<br/>- Suspected</p> | <p><b>Method of allocation:</b><br/>NA</p> <p><b>Intervention(s) description:</b><br/>All individuals are<br/>screened on<br/>voluntary basis.<br/>1. Mobile X-ray<br/>screening clinic<br/>visited locations<br/>where high-risk<br/>groups could be<br/>found (homeless<br/>shelters, drug<br/>treatment centres,<br/>criminal services,<br/>street outreach etc.)<br/>2. Raise awareness<br/>3. Under take case<br/>holding<br/>4. Provide support<br/>for treatment<br/>completion<br/>(supported by peer<br/>workers)</p> | <p><b>Primary outcomes:</b><br/>Incremental costs,<br/>quality adjusted life<br/>year (QALY), for the<br/>Find and Treat<br/>service.</p> <p><b>Secondary outcomes:</b><br/>Cost-effectiveness<br/>ratios for the Find<br/>and Treat service</p> <p><b>Method of analysis:</b><br/>NR</p> <p><b>Modelling method and assumptions:</b><br/>- Discrete, multiple<br/>age cohort,<br/>compartmental<br/>model to model a<br/>population of<br/>individuals with<br/>active tuberculosis</p> <p>4 groups:</p> | <p><b>Primary results:</b><br/>The model estimated<br/>that, on average, the<br/>Find and Treat service<br/>identifies 16 and<br/>manages 123 active<br/>cases of tuberculosis<br/>each year in hard to<br/>reach groups in London.<br/>The service has a net<br/>cost of £1.4 million/year<br/>and, under conservative<br/>assumptions, gains 220<br/>QALYs. The<br/>incremental cost<br/>effectiveness ratio was<br/>£6,400-£10,000/QALY<br/>gained (about €7300-<br/>€11,000 or \$10,000-<br/>\$16,000 in September<br/>2011).</p> <p>- 22.9% of patients<br/>detected by the mobile<br/>screening unit with the<br/>longest delays between<br/>symptom onset and</p> | <p><b>Limitations identified by author:</b><br/>- Absence of<br/>randomisation of<br/>tuberculosis cases to be<br/>either managed or not<br/>managed by the Find<br/>and Treat service<br/>- The service also<br/>manages extremely<br/>hard to reach<br/>individuals, who are<br/>often already lost to<br/>follow-up at the time of<br/>referral or who would<br/>never present for care<br/>without the mobile<br/>screening unit. Hence<br/>the comparison of<br/>cases with<br/>retrospective controls<br/>probably<br/>underestimates the<br/>incremental benefit of<br/>the service, although<br/>we cannot be certain<br/>without a randomised</p> |

|   |   |   |   |  |  |
|---|---|---|---|--|--|
| <p>service for diagnosing and managing hard to reach individuals with active tuberculosis in London</p> <p><b>Study design:</b><br/>Economic evaluation using a discrete, multiple age cohort, compartmental model of treated and untreated cases of active tuberculosis.</p> <p><b>Type of economic analysis:</b><br/>Cost-effectiveness</p> <p><b>Economic perspective:</b><br/>Healthcare taxpayer perspective</p> <p><b>Internal validity:</b><br/>Yes</p> <p><b>Quality score:</b><br/>++</p> <p><b>Applicability:</b><br/>+</p> | <p>tuberculosis</p> <ul style="list-style-type: none"> <li>- Cases merely receiving prophylaxis (and hence unlikely to have active tuberculosis)</li> <li>- Cases for which the diagnostic delay could not be calculated</li> <li>- Cases younger than 16 years</li> </ul> <p><b>Setting:</b><br/>London, United Kingdom.</p> <p><b>Sample characteristics:</b></p> <ul style="list-style-type: none"> <li>- 48 mobile screening unit cases</li> <li>- 188 cases referred for case management support</li> <li>- 180 cases referred for loss to follow-up</li> <li>- 252 control cases, TB cases that presented to a TB clinic themselves, without intervention of the Find and Treat service (current method of passive case finding)</li> </ul> | <p><b>Comparator/control(s) description:</b><br/>Controls: passively detected cases with active pulmonary tuberculosis (individuals who presented to London tuberculosis services on their own accord without screening and referral to the Find and Treat service - notified to the Health Protection Agency's enhanced tuberculosis surveillance system between 1 January 2009 (when the system began recording risk factor information) and 9 August 2010. Controls were age matched with actively detected cases (within five year age categories) and had to display one or more risk factors (a history of homelessness or imprisonment, drug</p> | <ul style="list-style-type: none"> <li>- Active untreated tuberculosis</li> <li>- Active treated tuberculosis with up to 125 days of continuous treatment</li> <li>- Active treated tuberculosis with more than 125 days of continuous treatment</li> <li>- Lost to follow-up</li> </ul> <p>4 final outcomes:</p> <ul style="list-style-type: none"> <li>- Completion of treatment</li> <li>- Death due to tuberculosis related causes</li> <li>- Death due to other causes</li> <li>- Other final outcomes that the Find and Treat service is not expected to change (such as patients being transferred out of London or stopping treatment for clinical reasons).</li> </ul> <p><u>Assumptions:</u></p> <ul style="list-style-type: none"> <li>- The cost of a new mobile unit £600,000 were added to the</li> </ul> | <p>treatment presentation were unlikely to present for treatment without the activities of the Find and Treat service</p> <ul style="list-style-type: none"> <li>- 35.4% of mobile screening unit patients were asymptomatic on detection, and hence would not have presented for treatment without the unit.</li> <li>- Once on treatment, mobile screening unit cases managed by the Find and Treat service had a much lower risk of loss to follow-up than passively presenting controls (loss to follow-up probability after one year: 2.1% for cases, 17.2% for controls)</li> <li>- Cases referred to Find and Treat because of complex case management issues had higher rates of completing treatment (61.2% after one year) and lower rates of loss to follow-up (3.3% after one year) than controls</li> </ul> <p><b>Secondary results:</b></p> <ul style="list-style-type: none"> <li>- Every year the service</li> </ul> | <p>study</p> <ul style="list-style-type: none"> <li>- Did not incorporate secondary transmission into the economic evaluation, even though the mobile screening unit in particular probably averts several secondary cases by finding highly infectious individuals.</li> <li>- Did not measure the effect of the Find and Treat service on reducing the likelihood of patients developing and transmitting acquired drug resistance (as a result of poor treatment adherence). Drug resistance increases the duration and costs of treatment, as well as the risk of severe disease, thus prevention could be an important benefit of the service.</li> </ul> <p><b>Limitations identified by review team:</b><br/>Small group of PTB in intervention group</p> |
|---|---|---|---|--|--|

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|  |   |   |   |   |  |
|--|---|---|---|---|--|
|  | <p><b>Economic analysis data source:</b><br/>Find and Treat database for information (including risk factors and clinical information) of individuals, diagnosed with PTB (between Sep 2007- Sep 2010)</p> <p>The control group (TB cases that presented on their own accord to a TB clinic) was obtained from the Health Protection Agency between Jan 2009 and Aug 2010. Risk factors and clinical information for the controls were obtained from the enhanced tuberculosis surveillance system.</p> | <p>or alcohol abuse, or mental health problems).</p> <p><b>Baseline comparisons:</b><br/>Compared:<br/>- Having no Find and Treat service,<br/>- Having only one part of the service (the mobile screening unit or the case management component)<br/>- Having both parts of the service</p> <p><b>Study sufficiently powered:</b><br/>NR but a small number of PTB cases in the Find and Treat group</p> | <p>costs of the first year of the service, with discounted costs and outcomes totalled over five years</p> <p>- Costs of £8,300 and £75,000 for treatment of DS-TB and MDR-TB</p> <p>- Only 50% of asymptomatic cases with a positive result from the mobile screening unit would progress to symptomatic disease</p> <p>- Find and Treat cases would be lost to follow-up at the same rate as enhanced tuberculosis surveillance controls (17.2% per year) in the absence of the service, rather than at the higher rate we estimated for this extremely hard to reach group (34.7% per year).</p> <p>- Even without Find and Treat involvement, these cases could still passively re-engage</p> | <p>has a net cost of £1.4 million and gained 220 QALYs</p> <p>- Incremental cost effectiveness of the Find and Treat service was £6,400/QALY gained</p> <p>- Both components of the service are cost-effective at the same threshold. The mobile screening unit had an incremental ratio of £18,000/QALY gained, whereas the case management component had an incremental ratio of £4,100/QALY gained (In the most unfavourable (and highly unlikely) scenario, which combined all the unfavourable assumptions, the mobile screening unit and case management components had incremental ratios of £26,000/QALY gained and £6,800/QALY gained, respectively)</p> <p>0.5% of mobile screening unit patients</p> | <p><b>Evidence gaps and/or recommendations for future research:</b><br/>Include a larger intervention group, longer follow up study</p> <p><b>Source of funding:</b><br/>Grant from the English Department of Health grant reference number 0150305</p> <p>PJW was partly funded by centre funding from the Medical Research Council. IA and HS are partly funded by the National Institute for Health Research.</p> <p><b>Conflict of interests:</b><br/>None</p> |
|--|---|---|---|---|--|

|  |  |  |  |   |  |
|--|--|--|--|---|--|
|  |  |  | with treatment at the same rate as enhanced tuberculosis surveillance controls (51% per year). | and 5.3% of other Find and Treat patients had multidrug or extensively drug resistant infection |  |
|  |  |  | <b>Time horizon:</b><br>Sep 2007 – July 2010   |   |  |

### List of Abbreviations

aOR = adjusted Odds Ratio; CDP = Chest Disease Centre; CHW = Community Health Worker; CXR = Chest X-Ray; HIV = Human Immunodeficiency Virus; IQR = Inter-Quartile Range; MXU = Mobile X-ray Unit; NA = Not Applicable; NIHR = National Institute for Health Research; NR = Not Recorded; OR = Odds Ratio; p = p-value; PTB = Pulmonary Tuberculosis; QALY = Quality Adjusted Life Year; TB = Tuberculosis; TBTCP = The Barcelona TB Control Program; TST = Tuberculin Skin Test; UK = United Kingdom; VNdg = Vila Nova de Gaia; 95%CI = 95% Confidence Interval

### References

1. Aldridge RW, Hayward AC, Hemming S, et al. Effectiveness of peer educators on the uptake of mobile X-ray tuberculosis screening at homeless hostels: a cluster randomised controlled trial. *BMJ open* 2015; **5**: e008050.
2. Goetsch U, Bellinger OK, Buettel KL, Gottschalk R. Tuberculosis among drug users and homeless persons: impact of voluntary X-ray investigation on active case finding. *Infection* 2012; **40**: 389-95.
3. Ospina JE, Orcau A, Millet JP, Sanchez F, Casals M, Cayla JA. Community health workers improve contact tracing among immigrants with tuberculosis in Barcelona. *BMC public health* 2012; **12**: 158.
4. Duarte R, Santos A, Mota M, Carvalho A, Marques A, Barros H. Involving community partners in the management of tuberculosis among drug users. *Public Health* 2011; **125**: 60-2.
5. Jit M, Stagg H, Aldridge R, White P, Abubakar I. Dedicated outreach service for hard to reach patients with tuberculosis in London: observational study and economic evaluation. *BMJ* 2011; **343**: d5376.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

**Supplementary Material VI: Quality Assessment**

**Quality assessment of included effectiveness studies**

| Questions about:   |                                  | Population |     |     | Method of selection |     |     |     |     | Outcomes |     |     |     |     | Analysis |     |     | Summary |     | Score |     |
|--|----------------------------------|------------|-----|-----|---------------------|-----|-----|-----|-----|----------|-----|-----|-----|-----|----------|-----|-----|---------|-----|-------|-----|
| Year   | First author (year)              | 1·1        | 1·2 | 1·3 | 2·1                 | 2·2 | 2·3 | 2·4 | 2·5 | 3·1      | 3·2 | 3·3 | 3·4 | 3·5 | 4·1      | 4·2 | 4·3 | 4·4     | 5·1 |       | 5·2 |
| <b>Studies identified for this review</b>                  |                                  |            |     |     |                     |     |     |     |     |          |     |     |     |     |          |     |     |         |     |       |     |
| 2011   | Duarte et al. <sup>5</sup>       | ++         | +   | -   | NA                  | +   | ++  | -   | ++  | +        | +   | -   | +   | ++  | NA       | +   | -   | +       | -   | -     | -   |
| 2012   | Goetsch et al. <sup>20</sup>     | ++         | +   | -   | NA                  | +   | ++  | -   | ++  | ++       | -   | -   | NA  | NA  | NR       | ++  | -   | -       | -   | -     | -   |
| 2012   | Ospina et al. <sup>21</sup>      | ++         | ++  | ++  | ++                  | ++  | ++  | -   | ++  | +        | ++  | -   | NA  | NA  | ++       | ++  | -   | ++      | -   | ++    | +   |
| 2015   | Aldridge et al. <sup>19</sup>    | ++         | ++  | ++  | ++                  | ++  | -   | -   | ++  | +        | ++  | +   | NA  | NA  | -        | +   | +   | +       | +   | +     | +   |
| <b>Studies identified for the NICE review<sup>12</sup></b> |                                  |            |     |     |                     |     |     |     |     |          |     |     |     |     |          |     |     |         |     |       |     |
| 2001   | El-Hamad et al. <sup>24</sup>    | ++         | +   | ++  | NA                  | ++  | ++  | +   | +   | ++       | ++  | +   | ++  | ++  | NR       | ++  | +   | ++      | +   | +     | +   |
| 2004   | Deruaz & Zellweger <sup>28</sup> | ++         | +   | +   | -                   | +   | -   | -   | +   | -        | ++  | ++  | ++  | ++  | NR       | -   | ++  | +       | -   | +     | -   |
| 2008   | Ricks <sup>23</sup>              | ++         | +   | ++  | ++                  | ++  | ++  | ++  | +   | +        | ++  | ++  | ++  | +   | NR       | ++  | ++  | ++      | ++  | +     | ++  |

|    |  |
|----|--|
| ++ | well designed study, minimal risk of bias                  |
| +  | study may not have addressed all potential sources of bias |
| -  | significant risk of bias                                   |
| NA | not applicable   |
| NR | not reported   |

Quality assessment questions for effectiveness studies:

- 1·1 Is the source population or source area well described?
- 1·2 Is the eligible population or area representative of the source population or area?
- 1·3 Do the selected participants or areas represent the eligible population or area?
- 2·1 Selection of exposure (and comparison) group. How was selection bias minimised?
- 2·2 Was the selection of explanatory variables based on a sound theoretical basis?
- 2·3 Was the contamination acceptably low?
- 2·4 How well were likely confounding factors identified and controlled?
- 2·5 Is the setting applicable to Europe?
- 3·1 Were the outcome measures and procedures reliable?
- 3·2 Were the outcome measurements complete?
- 3·3 Were all the important outcomes assessed?
- 3·4 Was there a similar follow-up time in exposure and comparison groups?
- 3·5 Was follow-up time meaningful?
- 4·1 Was the study sufficiently powered to detect an intervention effect (if one exists)?
- 4·2 Were multiple explanatory variables considered in the analyses?
- 4·3 Were the analytical methods appropriate?
- 4·4 Was the precision of association given or calculable? Is association meaningful?
- 5·1 Are the study results internally valid (i.e. unbiased)?
- 5·2 Are the findings generalisable to the source population (i.e. externally valid)?

Quality assessment of included cost-effectiveness studies

| Questions about: |              | Applicability |     |     |     |     |     |     |     |     | Study limitations |     |     |     |     |     |     |     |      |      |  | Overall |
|------------------|--------------|---------------|-----|-----|-----|-----|-----|-----|-----|-----|-------------------|-----|-----|-----|-----|-----|-----|-----|------|------|--|---------|
| Year             | First author | 1·1           | 1·2 | 1·3 | 1·4 | 1·5 | 1·6 | 1·7 | 1·8 | 2·1 | 2·2               | 2·3 | 2·4 | 2·5 | 2·6 | 2·7 | 2·8 | 2·9 | 2·10 | 2·11 |  |         |

Studies identified for this review

|      |                          |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |   |                                   |
|------|--------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|-----------------------------------|
| 2011 | Jit et al. <sup>21</sup> | Y | Y | Y | Y | N | N | Y | N | Y | Y | N | Y | Y | Y | Y | Y | Y | Y | Y | N | Potential serious limitations [+] |
|------|--------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|-----------------------------------|

Studies identified for the NICE review<sup>12</sup>

|      |                                |    |   |    |   |    |   |   |    |    |     |    |    |    |    |     |     |   |    |   |                                   |
|------|--------------------------------|----|---|----|---|----|---|---|----|----|-----|----|----|----|----|-----|-----|---|----|---|-----------------------------------|
| 2002 | Bothamley et al. <sup>25</sup> | PA | Y | Y  | N | PA | N | N | PA | PA | U/C | PA | N  | N  | Y  | N   | N   | Y | PA | N | Very serious limitations [-]      |
| 2006 | Miller et al. <sup>26</sup>    | Y  | Y | PA | N | PA | N | N | PA | PA | PA  | PA | PA | PA | PA | PA  | PA  | Y | Y  | N | Potential serious limitations [+] |
| 2008 | Mor et al. <sup>27</sup>       | PA | Y | PA | N | PA | N | N | PA | PA | PA  | PA | N  | PA | PA | U/C | U/C | Y | N  | N | Very serious limitations [-]      |

|     |                      |
|-----|----------------------|
| Y   | Yes to question      |
| N   | No to question       |
| PA  | Partially applicable |
| NA  | Not Applicable       |
| U/C | Unclear              |



Quality assessment questions for cost-effectiveness studies:

- 1·1 Is the study population appropriate for the topic being evaluated?
- 1·2 Are the interventions appropriate for the topic being evaluated?
- 1·3 Is the system in which the study was conducted sufficiently similar to the current European context?
- 1·4 Was/were the perspective(s) clearly stated and what were they?
- 1·5 Are all direct health effects on individuals included, and are all other effects included where they are material?
- 1·6 Are all future costs and outcomes discounted appropriately?
- 1·7 Is the value of health effects expressed in terms of quality-adjusted life years (QALYs)?
- 1·8 Are costs and outcomes from other sectors fully and appropriately measured and valued?
- 2·1 Does the model structure adequately reflect the nature of the topic under evaluation?
- 2·2 Is the time horizon sufficiently long to reflect all important differences in costs and outcomes?
- 2·3 Are all important and relevant outcomes included?
- 2·4 Are the estimates of baseline outcomes from the best available source?
- 2·5 Are the estimates of relative 'treatment' effects from the best available source?
- 2·6 Are all important and relevant costs included?
- 2·7 Are the estimates of resource use from the best available source?
- 2·8 Are the unit costs of resources from the best available source?
- 2·9 Is an appropriate incremental analysis presented or can it be calculated from the data?
- 2·10 Are all important parameters whose values are uncertain subjected to appropriate sensitivity analysis?
- 2·11 Is there any potential conflict of interest?



# PRISMA 2009 Checklist

| Section/topic                      | #  | Checklist item  | Reported on page # |
|------------------------------------|----|---|--------------------|
| <b>TITLE</b>                       |    |   |                    |
| Title                              | 1  | Identify the report as a systematic review, meta-analysis, or both.   | 1                  |
| <b>ABSTRACT</b>                    |    |   |                    |
| Structured summary                 | 2  | Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number. | 2                  |
| <b>INTRODUCTION</b>                |    |   |                    |
| Rationale                          | 3  | Describe the rationale for the review in the context of what is already known.  | 3                  |
| Objectives                         | 4  | Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).  | 3 and Supp Mat II  |
| <b>METHODS</b>                     |    |   |                    |
| Protocol and registration          | 5  | Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.   | 3                  |
| Eligibility criteria               | 6  | Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.  | 4                  |
| Information sources                | 7  | Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.  | 4                  |
| Search                             | 8  | Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.   | Sup mat III        |
| Study selection                    | 9  | State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).   | 4                  |
| Data collection process            | 10 | Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.  | 4                  |
| Data items                         | 11 | List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.   | 4                  |
| Risk of bias in individual studies | 12 | Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.  | 5                  |
| Summary measures                   | 13 | State the principal summary measures (e.g., risk ratio, difference in means).   | 5                  |



# PRISMA 2009 Checklist

|                      |    |   |   |
|----------------------|----|---|---|
| Synthesis of results | 14 | Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ ) for each meta-analysis. | 5 |
|----------------------|----|---|---|

Page 1 of 2

| Section/topic                 | #  | Checklist item   | Reported on page #       |
|-------------------------------|----|--|--------------------------|
| Risk of bias across studies   | 15 | Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).   | -                        |
| Additional analyses           | 16 | Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.   | -                        |
| <b>RESULTS</b>                |    |  |                          |
| Study selection               | 17 | Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.  | 5                        |
| Study characteristics         | 18 | For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.   | Table 1, page 17         |
| Risk of bias within studies   | 19 | Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).  | Sup mat VI and page 6, 7 |
| Results of individual studies | 20 | For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot. | 6, 7                     |
| Synthesis of results          | 21 | Present results of each meta-analysis done, including confidence intervals and measures of consistency.  | -                        |
| Risk of bias across studies   | 22 | Present results of any assessment of risk of bias across studies (see Item 15).  | Sup mat VI               |
| Additional analysis           | 23 | Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).  | -                        |
| <b>DISCUSSION</b>             |    |  |                          |
| Summary of evidence           | 24 | Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).                     | 8                        |
| Limitations                   | 25 | Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).  | 8                        |
| Conclusions                   | 26 | Provide a general interpretation of the results in the context of other evidence, and implications for future research.  | 9                        |
| <b>FUNDING</b>                |    |  |                          |



# PRISMA 2009 Checklist

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

|         |    |  |    |
|---------|----|--|----|
| Funding | 27 | Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. | 12 |
|---------|----|--|----|

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: [www.prisma-statement.org](http://www.prisma-statement.org).

Page 2 of 2

For peer review only