

Geophysical Research Abstracts
Vol. 18, EGU2016-14310, 2016
EGU General Assembly 2016
© Author(s) 2016. CC Attribution 3.0 License.



High resolution isoprene emissions over Europe in past and future climate

Trissevgeni Stavrakou (1), Jean-François Müller (1), Maite Bauwens (1), Julie Berckmans (2), Steven Caluwaerts (3), Rozemien De Troch (2), Lesley De Cruz (2), Olivier Giot (2), Rafiq Hamdi (2), Piet Termonia (2), and Bert Schaeysbroeck (2)

(1) Royal Belgian Institute for Space Aeronomy, Brussels, Belgium (jenny@oma.be), (2) Royal Meteorological Institute, Brussels, Belgium, (3) Department of Physics and Astronomy, Ghent University, Ghent, Belgium

Isoprene is a key tropospheric species, well known as the dominant biogenic hydrocarbon emitted in the atmosphere, with global annual emissions of about 400-600 Tg (Guenther et al. 2006). It is highly reactive, enhances tropospheric ozone formation in polluted environments and contributes to secondary aerosol formation. The emissions of isoprene (and other biogenic VOCs) depend on the type and abundance of plants, and are modulated by meteorological parameters. Climate changes can therefore affect the spatiotemporal and interannual variation of these emissions. In this study we propose to use the MEGAN-MOHYCAN model (Muller et al. 2008, Stavrakou et al. 2014) to calculate the isoprene fluxes emitted by vegetation in past and future climate over the European (EURO-CORDEX) domain at a resolution of 12.5 km. More specifically, isoprene emissions over 1979-2012 will be calculated based either on ECMWF reanalysis data or on meteorological output from the ALARO-0 climate model (Giot et al., 2015). The impact of solar radiation changes, observed at European stations (Sanchez-Lorenzo et al., 2015), on the modelled emissions will also be investigated. MEGAN-MOHYCAN simulations will be performed over the period 2040-2100 based on input from ALARO-0 model driven by the RCP8.5 scenario. The isoprene interannual variability and emission trends will be derived and thoroughly discussed.

Giot, O. et al.: Validation of the ALARO-0 model within the EURO-CORDEX framework, *Geosci. Model Dev. Discuss.*, 8, 8387-8409, 2015.

Guenther, A. et al.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, 2006.

Müller, J.-F. et al.: Global isoprene emissions estimated using MEGAN, ECMWF analyses and a detailed canopy environmental model, *Atmos. Chem. Phys.*, 8, 1329-1341, 2008.

Sanchez-Lorenzo, A. et al.: Reassessment and update of long-term trends in downward surface shortwave radiation over Europe (1939-2012), *J. Geophys. Res.*, 120, doi:10.1002/2015JD023321, 2015.

Stavrakou, T. et al.: Isoprene emissions over Asia 1979-2012 : impact of climate and land use changes, *Atmos. Chem. Phys.*, 14, 4587-4605, 2014.