



**UCC Library and UCC researchers have made this item openly available.  
Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	Investigating the perceived benefits, barriers and beliefs towards physical activity in pregnancy among women with Gestational Diabetes Mellitus
<b>Author(s)</b>	Flannery, Caragh; Byrne, M.; Kearney, Patricia M.; Dunne, Fidelma P.; O'Riordan, M.; Walsh, J. C.; Hurst, C.
<b>Publication date</b>	2017-09
<b>Original citation</b>	Hurst, C., Flannery, C., Byrne, M., Kearney, P. M., Dunne, F., O'Riordan, M. and Walsh, J. C. (2017) 'Investigating the perceived benefits, barriers and beliefs towards physical activity in pregnancy among women with Gestational Diabetes Mellitus', Irish Medical Journal, 110(8), 617 (10pp). doi:10147/622573
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://dx.doi.org/10147/622573">http://dx.doi.org/10147/622573</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2017, Irish Medical Journal. All rights reserved.
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/5890">http://hdl.handle.net/10468/5890</a>

Downloaded on 2019-12-02T14:30:09Z



[This Month](#)

[Archive](#)

[Subscribe](#)

[For Authors](#)

[About Us](#)

[My CPD](#)

[Hosted Content](#)

[Home](#)

# Investigating the Perceived Benefits, Barriers and Beliefs towards Physical Activity in Pregnancy among Women with Gestational Diabetes Mellitus

C Hurst<sup>1</sup>, C Flannery<sup>2\*</sup>, M Byrne<sup>2</sup>, P M. Kearney<sup>3</sup>, F Dunne<sup>2</sup>, M O'Riordan<sup>3</sup>, J C. Walsh<sup>2</sup>.

<sup>1</sup>University of Limerick

<sup>2</sup>National University of Ireland, Galway

<sup>3</sup>University College Cork

## **Abstract**

***Gestational Diabetes Mellitus (GDM) is a growing concern and poses serious health risks to both mother and child<sup>1</sup>. The current study explores the psychological determinants of exercise behaviour in a sample of pregnant women with GDM. A cross-sectional survey design was employed to examine exercise behaviour, illness perceptions, perceived barriers and benefits, exercise beliefs, and exercise self-efficacy using validated questionnaires. A sample of 46 pregnant women was recruited from University College Hospital Galway, Letterkenny General Hospital, Cork University Hospital and Mayo General Hospital in Castlebar. Participant's varied; age (22-44 years), body mass index (19-41). High mean scores for Personal Control (24.5) and Treatment Control (30.2) subscales indicated strongly held positive beliefs in relation to***

***controllability of the illness. Total MET-min/week score was not related to any psychological variables. Analysis of the IPQ-R data revealed 'diet' (n=37, 80.4%) as the most referred to cause of diabetes. Exercise belief data identified "managing weight gain" (n= 21, 45.7%), and "losing baby weight" (n= 31, 67.4%) as the most frequent beliefs for engaging in physical activity during pregnancy and post pregnancy. Further research on the psychological determinants of physical activity behaviour among this population group is needed in order to create successful intervention strategies.***

## **Introduction**

Gestational Diabetes Mellitus (GDM) is increasing in prevalence, with rising levels of maternal overweight and obesity a major contributor<sup>2</sup>. GDM increases the likelihood of caesarean delivery, macrosomia, neonatal abnormalities and birth trauma<sup>3</sup>. The future health of the mother and baby are also at an increased risk of being compromised following diabetes in pregnancy, with the risk of developing type-2 diabetes seven times greater than in normal glycaemic pregnancies<sup>2</sup>. Children born to women with gestational diabetes are also at an increased risk of developing respiratory distress syndrome, jaundice, childhood obesity and metabolic syndrome when compared with children of non-diabetic women<sup>4</sup>. Research has found that being physically active prior to becoming pregnant as well as during pregnancy and postpartum, has the potential to reduce the risk of progression from gestational diabetes to type 2-diabetes<sup>5</sup>.

In the past, it has been speculated that exercise during pregnancy could lead to adverse events such as overheating, impaired delivery of oxygen and nutrients to the foetus and premature labour<sup>6</sup>. However, evidence to support these assertions is lacking and research indicates that exercise in pregnancy is in fact safe<sup>7</sup>. Physical activity reduces the risk of pregnancy complications such as preeclampsia, preterm delivery and gestational diabetes along with reducing discomfort, increasing fitness for labour and increasing the likelihood of vaginal delivery<sup>8,9</sup>. The American College of Obstetricians and Gynaecologists (ACOG) recommend 30 minutes of moderate intensity physical activity for pregnant women without obstetrical complications<sup>10</sup>. However, despite this information encouraging an active lifestyle during pregnancy remains a challenge. Regarding physical activity levels, the United States reported that only 15.8% of women engaged in exercise during pregnancy at the recommended level, while 34% of pregnant women had not engaged in any moderate or vigorous activity<sup>11</sup>. Similarly, low levels of physical activity (21.5%) were found in an Irish cohort of pregnant women<sup>12</sup> and in a sample of Danish pregnant

women, moderate-to-heavy activities decreased over the three trimesters of pregnancy<sup>13</sup>. Furthermore the results from a study conducted in Brazil showed alarmingly low rates of physical activity, with only 4.7% of pregnant women being active<sup>14</sup>.

Pregnancy is a time when health behaviours such as physical activity can be challenged in order to improve not only the mother's health but the health of the baby<sup>15</sup>. The likelihood of someone making a substantial behaviour change is dependent on a host of factors such as; beliefs, values, perceived barriers, and exercise self-efficacy<sup>16</sup>. The development and future success of exercise interventions among pregnant women with GDM hinges on having an accurate insight into these psychological determinants. There is a need to understand the facilitators and barriers to exercise in pregnancy and to understand the relationships between psychological predictors in order to develop effective interventions and strategies<sup>1,17</sup>. It is evident that prior to promoting exercise to this particular group of women, the specific exercise determinants need to be examined. Therefore the aim of this study is to explore the psychological determinants of exercise behaviour in pregnancy in women with GDM.

## Method

A cross-sectional study was conducted with a sample of pregnant women with GDM (N=46) in June 2014. Ethical approval was obtained from the National University of Ireland, Galway Research Ethics Committee and the University College Hospital Galway ethics committee. Participants were pregnant women diagnosed with GDM who were attending the diabetes clinics in four participating hospitals. Exclusion criteria included not having sufficient level of English language to complete the questionnaire, and the presence of any co-morbidity or pre-existing condition other than diabetes. Participants were recruited from the diabetes clinics and maternity outpatients of four large hospitals. Women were approached and provided with information detailing the study aims. Those willing to participate were given the choice of completing the survey on site, or were given a stamped addressed envelope to post the completed survey back. Participation was voluntary, and women were free to withdraw at any point. Informed consent was obtained prior to the completion of the survey. Women attending the clinic on the days when the researcher was not present were sent the survey via post along with a stamped addressed envelope for its return. A total of 101 surveys were distributed, based on previous research in the area<sup>6,16</sup>. Basic demographic information was obtained including ethnicity, relationship status (single/married); education level (secondary/tertiary); employment status (employed/unemployed); parity (first pregnancy/two or more) height; weight; smoking (yes/no) and drinking behaviour (number of units). Body mass index was calculated using height and weight. A number of validated questionnaire were used including International Physical Activity Questionnaire (IPAQ) Long version, The Revised Illness Perception Questionnaire (IPQ-R) Diabetes Version, Health Value Scale (HVS), Exercise Benefits and

Barriers Scale (EBBS), Exercise Belief Questionnaire (EBQ) Symons, Self-Efficacy for Exercise (SEE) Scale<sup>18-23</sup>.

Analysis was conducted using IBM SPSS Statistics, Version 21. Associations between physical activity scores on the IPAQ and the psychological factors were explored using Chi-squared tests, Kruskal-Wallis, Mann-Whitney U and Pearson's correlation. Pearson's Product Moment Correlation Coefficient was conducted to examine the relationship between demographic variables and Total MET-min/week, or the Leisure-Time Physical Activity (LTPA). Kruskal-Wallis test between recruitment location and both Total MET-min/week and LTPA. Mann-Whitney U tests were conducted between physical activity and smoking behaviour, education level, and employment status.

## Results

### *Descriptive Statistics*

Forty-seven questionnaires were returned with one incomplete survey excluded, thus 46 (45.5%) surveys were analysed. Participants ages ranged from 23 to 44 years (M=36.13, SD=4.05, N=38), with participant's BMI's ranging from 18.93 to 40.96 (M=29.77, SD=5.28, N=34). Descriptive characteristics are outlined in Table 1.

Variable	N (%)
<b>Relationship Status</b>	
Single	4 (8.7)
Married/In committed relationship	42 (91.3)
<b>Education Level</b>	
Completed Secondary School	16 (34.8)
Tertiary Qualification	30 (65.2)
<b>Employment Status</b>	
Employed	37 (80.4)
Unemployed	9 (19.6)
<b>Parity</b>	
First Pregnancy	15 (32.6)
Second or more	31 (67.4)
<b>Smoking Behaviour</b>	
Yes	8 (17.4)
No	38 (82.6)
<b>Alcohol Consumption</b>	
None	35 (76.1)
<2 Units	9 (19.6)
3 - 8 Units	1 (2.2)
>8 Units	1 (2.2)

### *Characteristics of participants reporting physical activity*

The analyses revealed no significant relationships between age and Total MET-min/week ( $r=0.06$ ,  $p=0.75$ ), BMI and Total MET-min/week ( $r=0.04$ ,  $p=0.83$ ), age and LTPA ( $r =0.05$ ,  $p=0.77$ ) or BMI and LTPA ( $r=-0.25$ ,  $p=0.15$ ). Recruitment, location and Total MET-min/week ( $\chi^2=1.38$ ,  $df=3$ ,  $p=0.71$ ) and LTPA ( $\chi^2=5.7$ ,  $df=3$ ,  $p=0.13$ ) found no significant differences between locations. No significant differences were found between the other demographic variables (alcohol consumption, and relationship status) on the physical activity scores. Smoking behaviour, education level, and employment status found no significant difference with the physical activity measures.

#### *Correlation analysis between physical activity scores on the IPAQ and the psychological factors*

Correlational analyses were conducted between both MET and LTPA physical activity scores, and each of the psychological measures; Health Value Scale, Exercise Benefits and Barriers Scale Total (EBBS), Benefits subscale, Barriers subscale, and the Self-Efficacy in Exercise Scale (SEE). Results revealed no significant correlations between any of the variables and Total MET-min/week scores. Significant results were found between the LTPA score, presented in table 2.

**Table 2. Summary of inter-correlations between IPAQ Leisure-Time Physical Activity Domain Score (LTPA) and Relevant Psychological Variables**

Variable	1.	2.	3.	4.	5.	6.
1. LTPA	-					
2. HVS	.19	-				
3. EBBS Total	.42**	.40**	-			
4. SEE	.32*	.42**	.54*	-		
5. EBBS Barrier subscale	-.30*	-.51**	-.83**	.62**	-	
6. EBBS Benefit subscale	.43**	.32*	.96**	.42**	-.60**	-

Note: HVS= Health Value Scale, EBBS=Exercise Benefits and Barriers Scale, SEE= Self-Efficacy for Exercise Scale

\* $p<.05$ , \*\* $p<.01$

The IPQ-R data, representing the participants' views about their illness, was analysed by comparing the mean scores on each of the seven subscales. The higher mean scores on the Personal Control and Treatment Control subscales indicated strongly held positive beliefs by participants in relation to the controllability of the illness. Lower mean scores were recorded for Illness Coherence, Consequences and Emotional Representations subscales, with both Timeline: acute/chronic and cyclical; subscales having the lowest mean scores. Means and standard deviations for each of the seven subscales are presented in Table 3.

**Table 3. Means and Standard Deviations for IPQ-R Subscales**

Subscale	M (SD)
Personal Control	24.48 (3.51)
Treatment Control	20.22 (2.35)
Illness Coherence	18.43 (3.30)
Consequences	16.02 (4.75)
Emotional Representations	15.22 (4.52)
Timeline (Acute/Chronic)	12.24 (4.150)
Timeline (Cyclical)	10.98 (2.99)

In relation to the Causes subscale, four causes in particular were prominent in relation to the frequency of agreement from participants. The top cited cause was “Diet/Eating Habits” (80.4% agreement), followed by “Hereditary/It Runs in My Family” (63.0% agreement), “My Own Behaviour” (60.9% agreement) and “Ageing” (58.7% agreement). The remaining 14 causes listed were infrequently cited by participants in comparison to the above four causes. From the Exercise Belief Questionnaire the most noticeable behavioural beliefs to engaging in physical activity during pregnancy was; “Managing weight gain” (45.65%), while “Losing baby weight” (67.39%) was the most frequent behavioural beliefs for engaging in physical activity post pregnancy (Table 4).

**Table 4. Behavioural beliefs to engaging in physical activity during pregnancy and post pregnancy**

<b>Exercise Belief Questionnaire (N=46)</b>			<b>% (n)*</b>
<b>Physical activity during pregnancy</b>	Behavioural beliefs	Managing weight gain	45.64 (21)
		Manage diabetes/keep blood sugar down	36.96 (17)
		Keep fit	23.91 (11)
		Prepare body for labour	23.91 (11)
	Normative beliefs	Family	47.83 (22)
		Myself	28.26 (13)
		Baby	26.09 (12)
	Control beliefs	Pain/Discomfort	54.35 (25)
Tiredness/Fatigue/No energy		39.13 (18)	
<b>Physical activity post pregnancy</b>	Behavioural beliefs	Losing baby weight	67.39 (31)
		Getting back in shape/Improve fitness	32.61 (15)
		Relaxation/Reduce stress	32.61 (15)
	Normative beliefs	Family	50.00 (23)
		Baby/children	34.78 (16)
		Myself	23.91 (11)
	Control beliefs	Lack of time	39.13 (18)
		Tiredness/Fatigue/No energy	23.91 (11)
Pain/Discomfort		19.57 (9)	

\*Results are presented as % of N=46

## Discussion

This exploratory study describes physical activity scores among women with a diagnosis of GDM along with the psychological factors. Analysis of the data revealed that perceived benefits, barriers and exercise self-efficacy are significantly related to, and predict LTPA. Furthermore the Total MET-min/week score was not related to any of the psychological variables. Being physically active is important for any pregnant woman, however it may be more so, for women with GDM as they are at risk of type-2 diabetes later in life. Considering that approximately 40% of women affected by GDM may develop type-2 diabetes within four years<sup>7</sup>, addressing this public health issue is vital. Studies have shown that increasing physical activity can potentially reduce the onset of diabetes progression from GDM<sup>24</sup> with physical activity in the early stages of pregnancy reducing the risk of GDM by 48%<sup>25</sup>.

Results from the IPQ-R subscale indicate that diet was the most frequently cited cause, suggesting an awareness of control. If this awareness could be transferred into a positive change, through taking control of their dietary habits and being more physically active, women with GDM could vastly improve their outcomes. The specific barriers and benefits to exercise cited by participants from the Exercise Belief Questionnaire, are in line with those recorded in previous research. Lack of time, fatigue and being too tired were commonly reported barriers<sup>11,16</sup>. There is a need to address these barriers and discover ways for women to overcome them if they are to stand any chance of adopting and maintaining exercise behaviour.

The study has a few limitations. The IPAQ is a self-report measure and is subject to bias. The IPAQ requires participants to separate physical activity into domains and intensity which could be misleading. Secondly, having obtained only a 45.5% response rate of 46 participants results are not generalisable. Two importance factors were omitted from the survey; a record of pre-pregnancy physical activity, and gestational period. Throughout the course of a pregnancy, barriers, benefits and beliefs can shift. Having had a record of pre-pregnancy exercise levels would have added an extra dimension to the analysis. Future research should focus on motivators and facilitators to exercise, means of overcoming barriers and changing dysfunctional beliefs and perceptions. To achieve this successfully, a more holistic approach is necessary, where psychological constructs and cognitive strategies are discussed and explored. There is a huge potential for behavioural strategies such as goal-setting, along with the use of technology to influence these exercise determinants<sup>1</sup>. Further research is needed to identify which strategies might be successful in increasing physical activity behaviour among this population.

### Conflict of interest

The authors declare that they have no competing interests.



## Correspondence

Caragh Flannery, SPHeRE PhD Scholar, Health Behaviour Change Research Group/School of Psychology/Arts Millennium Building/NUI Galway/Ireland

Dept. Epidemiology & Public Health/University College Cork/Cork/Ireland

c.flannery13@nuigalway.ie

cflannery@ucc.ie

## References

1. Downs DS, Chasan-Taber L, Evenson KR, Leiferman J, Yeo S. Physical activity and pregnancy: past and present evidence and future recommendations. *Research quarterly for exercise and sport*. 2012;83(4):485-502.
2. Dennedy MC, Dunne F. The maternal and fetal impacts of obesity and gestational diabetes on pregnancy outcome. *Best Practice & Research Clinical Endocrinology & Metabolism*. 2010;24(4):573-89.
3. Coffey M. Gestational Diabetes Mellitus. . *World of Irish Nursing*,. 2006;14((11)):41-2.
4. Reece EA. The fetal and maternal consequences of gestational diabetes mellitus. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2010;23(3):199-203.
5. Hedderson M, Ferrara A. A call to increase physical activity among women of reproductive age: is it possible? *JAMA internal medicine*. 2014;174(7):1056-7.
6. Duncombe D, Wertheim EH, Skouteris H, Paxton SJ, Kelly L. Factors related to exercise over the course of pregnancy including women's beliefs about the safety of exercise during pregnancy. *Midwifery*. 2009;25(4):430-8.
7. Artal R, O'Toole M. Guidelines of the American College of Obstetricians and Gynecologists for exercise during pregnancy and the postpartum period. *Br J Sports Med*. 2003;37(1):6-12.
8. Fell DB, Joseph KS, Armson BA, Dodds L. The impact of pregnancy on physical activity level. *Matern Child Health J*. 2009;13(5):597-603. Epub 2008/08/23. doi: 10.1007/s10995-008-0404-7. PubMed PMID: 18719984.
9. Bungum TJ, Peaslee DL, Jackson AW, Perez MA. Exercise during pregnancy and type of delivery in nulliparae. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*. 2000;29(3):258-64.
10. Gynecologists ACoOa. Physical Activity and Exercise During Pregnancy and the Postpartum Period. <http://www.acog.org/Resources-And-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Physical-Activity-and-Exercise-During-Pregnancy-and-the-Postpartum-Period>. 2002.
11. Evenson KR, Moos M-K, Carrier K, Siega-Riz AM. Perceived barriers to physical activity among pregnant women. *Matern Child Health J*. 2009;13(3):364-75.
12. Walsh JM, McGowan C, Byrne J, McAuliffe FM. Prevalence of physical activity among healthy

pregnant women in Ireland. *Int J Gynaecol Obstet*. 2011;114(2):154-5.

13. Hegaard HK, Damm P, Hedegaard M, Henriksen TB, Ottesen B, Dykes A-K, Kjaergaard H. Sports and leisure time physical activity during pregnancy in nulliparous women. *Maternal and child health journal*. 2011;15(6):806-13.
14. Domingues MR, Barros AJ. Leisure-time physical activity during pregnancy in the 2004 Pelotas Birth Cohort Study. *Revista de Saúde Pública*. 2007;41(2):173-80.
15. Campbell F, Johnson M, Messina J, Guillaume L, Goyder E. Behavioural interventions for weight management in pregnancy: a systematic review of quantitative and qualitative data. *BMC Public Health*. 2011;11(1):1.
16. Cramp AG, Bray SR. A prospective examination of exercise and barrier self-efficacy to engage in leisure-time physical activity during pregnancy. *Annals of Behavioral Medicine*. 2009;37(3):325-34.
17. Leiferman J, Swibas T, Koiness K, Marshall JA, Dunn AL. My baby, my move: examination of perceived barriers and motivating factors related to antenatal physical activity. *Journal of Midwifery & Women's Health*. 2011;56(1):33-40.
18. Jette M, Sidney K, Blümchen G. Metabolic equivalents (METS) in exercise testing, exercise prescription, and evaluation of functional capacity. *Clinical cardiology*. 1990;13(8):555-65.
19. Weinman J, Petrie KJ, Moss-Morris R, Horne R. The illness perception questionnaire: a new method for assessing the cognitive representation of illness. *Psychology and health*. 1996;11(3):431-45.
20. Lau RR, Hartman KA, Ware JE. Health as a value: Methodological and theoretical considerations. *Health Psychology*. 1986;5(1):25.
21. Sechrist KR, Walker SN, Pender NJ. Development and psychometric evaluation of the exercise benefits/barriers scale. *Research in nursing & health*. 1987;10(6):357-65.
22. Downs DS, Ulbrecht JS. Understanding exercise beliefs and behaviors in women with gestational diabetes mellitus. *Diabetes care*. 2006;29(2):236-40.
23. Bandura A. *Social foundations of thought and action: A social cognitive theory*: Prentice-Hall, Inc; 1986.
24. Bao W, Tobias DK, Bowers K, Chavarro J, Vaag A, Grunnet LG, Strøm M, Mills J, Liu A, Kiely M. Physical activity and sedentary behaviors associated with risk of progression from gestational diabetes mellitus to type 2 diabetes mellitus: a prospective cohort study. *JAMA internal medicine*. 2014;174(7):1047-55.
25. Dempsey JC, Sorensen TK, Williams MA, Lee I-M, Miller RS, Dashow EE, Luthy DA. Prospective study of gestational diabetes mellitus risk in relation to maternal recreational physical activity before and during pregnancy. *American journal of epidemiology*. 2004;159(7):663-70.



[This Month](#) [Archive](#) [Subscribe](#) [For Authors](#) [About Us](#) [My CPD](#)  
[Hosted Content](#) [Home](#)

Theme by Tyler Moore