



Internet Gaming Disorder: The interplay between physical activity and user-avatar relationship

Journal:	<i>Behaviour & Information Technology</i>
Manuscript ID	TBIT-2017-0353.R1
Manuscript Type:	Original Research Study - Full Article
Keywords:	Internet Gaming Disorder, Proto-Self Presence, Physiological Activity, Emerging Adulthood, Massively Multiplayer Online games

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Internet Gaming Disorder:
The interplay between physical activity and user-avatar relationship

Word count: 9191 (including references)

ABSTRACT

Understanding both the risk and protective factors associated with Internet Gaming Disorder (IGD) has been viewed by many in the gaming studies field as an area of research priority. The present study focused on the potential risk and protective effects of user-avatar (game figure) relationship and physical activity (PA) respectively. To address these aims, a cross-sectional and a longitudinal mixed-methods design were combined (comprising both psychological and physiological assessments). A sample of 121 emerging adult gamers (18-29 years) residing in Australia, who played massively multiplayer online games, were assessed in relation to their IGD behaviors using the nine-item Internet Gaming Disorder Scale-Short Form. Additionally, the Proto-Self-Presence scale was used to evaluate the extent to which gamers identified with the body of their avatar. Finally, a PA monitor (*Fit-Bit Flex*) measured levels of energy consumed during real world daily activities (active minutes). A number of linear regressions and moderation analyses were conducted. Findings confirmed that Proto-Self-Presence functioned as an IGD risk factor and that PA acted protectively, weakening the association between Proto-Self-Presence and IGD behaviors. Implications of these findings are discussed in relation to IGD treatment and gaming development aspects.

Keywords: Internet Gaming Disorder; Proto-Self-Presence; Physical Activity; Emerging Adulthood; Massively Multiplayer Online games.

1. Introduction

Internet use has grown exponentially, constituting an inherent part of contemporary life (Anderson, Steen & Stavropoulos, 2016). In that context, the user-internet interplay (within the broader human computer interaction filed) has attracted the attention of researchers from a diverse range of scientific disciplines including psychology, information technology, computer science, and sociology (Lazar, Feng, & Hochheiser, 2017). One particularly popular and continuously expanding online application/activity that has gravitated research focus is Internet gaming (Andreassen et al., 2016). More specifically, the potentially positive (e.g., mental health improvements, cultural openness, a sense of meaning and accomplishment, collaboration and emotive stimulation; Smahel, Blinka, & Brown, 2012; Kaplan & Haenlein, 2010; Armitage, Claypool, & Branch, 2006; Billieux et al., 2015; Hsu, Wen, & Wu, 2009; Snodgrass, Lacy, Dengah, & Fagan, 2011) and negative effects (e.g., negative impact on identity formation, physical health and interpersonal relationships; Billieux et al., 2015; Hyun et al., 2015; Kuss, 2013; Hsu et al, 2009; Freeman, 2008) of Internet gaming, depending on the intensity of gaming involvement, have been demonstrated (Anderson et al., 2016). Additionally, the significance of the gamer-avatar (i.e., game character) association has been repeatedly highlighted, and associations with Internet Gaming Disorder (IGD) have been reported (Bessière, Seay & Kiesler, 2007; Caplan, Williams & Yee, 2009; Bainbridge, 2009; Williams, Yee, & Caplan, 2008. Corneliusen & Rettberg 2008; Constantiou, Legarth & Olsen, 2012). Despite these findings, further

1
2
3
4 research has been recommended in regard to the effects of the interplay of user-avatar
5 characteristics on the development of IGD behaviors (Stavropoulos et al., 2017a).

6
7
8 Driven by these recommendations, the present study explored the gamers' level
9 of physical-body identification with their avatars (Proto-Self-Presence [PSP]; Ratan &
10 Dawson, 2015) and their physiological activity (PA) as risk and protective factors of
11 IGD behaviors respectively. Addressing these questions appears important, given the
12 dearth of relevant studies and the significant potential implications for more effective
13 IGD prevention and intervention initiatives (e.g., informing more targeted and efficient
14 policies and programs).

23 1.1.1. Internet Gaming Disorder (IGD)

24
25 IGD was included in the latest (fifth) edition of the Diagnostic and Statistical
26 Manual of Mental Disorders (DSM-5) as an emerging disorder requiring further clinical
27 and research investigation by the American Psychiatric Association (APA, 2013). IGD
28 is described as a form of persistent and recurrent involvement with online gaming,
29 often leading to the decline of daily and work activities (i.e., neglecting daily routine
30 activities such as showering or eating; APA, 2013; Hawkins, Catalano, & Miller, 1992;
31 Kuss & Griffiths, 2012; Sanchis-Segura, & Spanagel, 2006; Pawlikowski & Brand,
32 2011). Accordingly, its prevalence and psychosocial impact have prompted much
33 research interest (e.g., Spekman, Knijn, Roelofsma, & Griffiths, 2013). However,
34 despite the increasing number of cross-sectional studies, there is a dearth of
35 longitudinal studies examining risk and protective factors (especially related to the
36 virtual-gaming context) of IGD behaviors (Kuss, 2013; Pontes & Griffiths, 2015). To
37 help prioritize practical knowledge concerning vulnerable IGD populations, the present
38 study focuses on emerging adulthood (as an age period; Spekman et al., 2013);
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 Stavropoulos et al., 2017a; Winkler et al, 2013) and Massively Multiplayer Online
5
6 (MMO) games (as the game genre; Billieux et al., 2015; Kuss, 2013; Stavropoulos et
7
8 al., 2017a; Young, 2009) because these features are believed to be risk factors in
9
10 developing IGD among the gamer population (Anderson et al., 2016).

11 12 **1.1.2. Emergent Adulthood as the Age Range of focus**

13
14 Emergent adulthood includes people aged between 18 and 29 years (Arnett,
15
16 Žukauskienė, & Sugimura, 2014) and has been identified as a risk period for addictions
17
18 more generally (including technological addictions; Arnett, 2000) and IGD in particular
19
20 (APA, 2013; Blinka, 2008; Freeman, 2008; Douglas et al, 2008; Substance Abuse and
21
22 Mental Health Services Administration, 2015). In Australia, 78% of online gamers are
23
24 emergent adults or adults, who predominantly play online videogames to positively
25
26 moderate their feelings and relieve boredom (Brand & Todhunter, 2015). However, to
27
28 date, there has been a dearth of longitudinal studies investigating IGD risk during this
29
30 particular developmental time (Spekman et al., 2013; Stavropoulos et al., 2017a;
31
32 Winkler et al, 2013).

33 34 **1.1.3. Game Genres**

35
36 Within the broader range of Internet gaming genres (Kuss, 2013; Thipse, 2016),
37
38 the present study focuses on Massively Multiplayer Online games (MMOs), as they
39
40 have consistently been associated with higher IGD risk (Billieux et al., 2015; Elliott et
41
42 al., 2012; Kuss, 2013; Lee et al., 2007; Stavropoulos et al., 2017a; Young, 2009).
43
44 MMOs are defined as games that are co-played online with others (on a massively
45
46 multi-player basis) and involve competitive and cooperative goals that are both
47
48 challenging and immersive (Steinkuehler & Williams, 2006; Lee, Cheung, & Chan,
49
50 2015). It should also be noted that MMO games are often used interchangeably with
51
52
53
54
55
56
57
58
59
60

1
2
3
4 that of Massively Multiplayer Online Role Playing Games (MMORPGs; Caplan,
5 Williams, Yee, 2009) although some MMO games may not involve role-playing.
6
7
8 Various factors have been suggested to explain the higher IGD risk in MMO games,
9
10 such as escapism motives, low achievement, and poor self-esteem in real life (Billieux
11 et al., 2015). Additionally, the higher IGD risk in MMO games has also been attributed
12
13 to the emotional attachment formed between the players and their avatars
14
15 (Badrinarayanan et al., 2014; Billieux et al., 2013; 2015; Hsu et al., 2009). More
16
17 specifically, players may find themselves becoming more immersed with their virtual
18
19 character, increasing their excessive gaming time (Badrinarayanan et al., 2014;
20
21 Stavropoulos et al., 2017a; Yee, 2006).
22
23
24

25 **1.2. Theoretical Background**

26 **1.2.1. Risk and Resilience Framework (RRF)**

27
28
29 To expand the knowledge considering the effect of the gamer-character
30
31 association on IGD behaviors, the present study adopts the risk and resilience
32
33 framework (RRF; Stanley, 2009). Resilience is defined as the combination of aspects
34
35 and experience (i.e., protective factors-resources) that allow a person to be unaffected
36
37 by something problematic (Masten, 2014). In contrast, risk is defined as involving
38
39 aspects and experiences of the individual that make him/her susceptible to problematic
40
41 conditions (Masten, 2014). In this context, the RRF suggests that the intensity of
42
43 psychopathological behaviors dynamically (and constantly) varies on a continuum
44
45 (minimum to maximum) based on the interplay between individual characteristics (e.g.,
46
47 age, gender, personality traits, etc.) and contextual factors (e.g., family, culture, etc.)
48
49 (Masten, 2014; Bronfenbrenner, 2006; Cicchetti & Toth, 2009; Hussain et al., 2015).
50
51 Given the importance of the gaming context in the development of IGD behaviors, the
52
53
54
55
56
57
58
59
60

1
2
3
4 RRF is enriched here by the inclusion of gaming factors (i.e., user-avatar relationship;
5
6 Andreassen et al., 2016). Within this broader framework, and to contribute to the field,
7
8 the present study emphasizes the effect of proto-self-presence as an IGD risk factor
9
10 (e.g., the level of physical-body identification between the gamers and their avatars;
11
12 Ratan & Dawson, 2015) and physical activity as an IGD resilience factor.
13

14
15 *-Figure 1. Risk and Resilience IGD Framework-*

16 17 **1.2.2. Proto-Self-Presence (PSP)**

18
19 The avatar is a complex multi-leveled tool enabling social and virtual interaction,
20
21 affecting the user emotionally, cognitively, and behaviorally (Ratan & Dawson, 2015).
22
23 The avatar may often merge with aspects of the user's identity (i.e., body, behavior, and
24
25 emotion) creating an overlap between reality and the virtual environment (Blinka,
26
27 2008). The concept of the 'Proteus effect' has been introduced to best describe how the
28
29 user's perception, knowledge, and experience merge with the avatar (Yee et al, 2009).
30
31 The 'Proteus effect' assumes that players are influenced by their avatars, changing how
32
33 they interact with others in real life (Yee, & Bailenson, 2007). One study found the
34
35 characteristics of the avatar to influence how the player perceived themselves (use of
36
37 tall avatars influenced taller self perception) and behaved (attractive avatars influenced
38
39 confident behavior during social interactions; Chandler et al., 2009). Subsequently,
40
41 virtual sensations (psychological or body-related) may often be mixed with those that
42
43 are physical and real (Billieux et al., 2015; Chandler et al., 2009). Additionally, it has
44
45 also been found that players who use avatars that are physically competent are also
46
47 inclined to perceive themselves as such, even if they lack such a competency in reality.
48
49 The latter has been described as a "de-individuation" (from someone's real self) aspect
50
51 involved with the 'Proteus effect' (Yee, & Bailenson, 2007).
52
53
54
55
56
57
58
59
60

1
2
3
4 In this context, the extent to which gamers identify with the body of their avatars
5
6 is conceptualized as PSP (Ratan & Dawson, 2015; Riva et al., 2004). Higher PSP levels
7
8 induce an ‘extension of the body’ experience in the world of the game (Luppicini,
9
10 2013; Ratan & Hasler, 2010; Riva et al., 2004). How the avatar’s influence poses a
11
12 potential IGD risk factor can be highlighted via the compensatory internet use model
13
14 (Kardefelt-Winther, 2014). This signifies behaviors of escaping and/or using the online
15
16 world to displace and/or address real life problems (Kardefelt-Winther, 2014) and
17
18 appears consistent with a significant body of literature (Bessièrè et al., 2007; Caplan et
19
20 al., 2009; Constantiou et al., 2012; Corneliussen & Rettberg 2008). Players may be
21
22 projecting their physical insecurities onto their in-game character to compensate for
23
24 what they may physically lack in the real world (Blinka, 2008; Hopp, Barker, & Weiss,
25
26 2015; Smahel et al., 2008). This can reach levels of deindividuation when the idealized
27
28 self of the player (projected onto the avatar) provides players with the perceptive
29
30 experiences of their ideal attributes (Yee, & Bailenson, 2007). Therefore, gamers are
31
32 likely to endorse their avatars with physical characteristics desired in real life,
33
34 becoming emotionally attached, as a means to attain an idealized self. This process may
35
36 reinforce excessive gaming involvement precipitating and perpetuating IGD symptoms.
37
38 Similarly, people who spend longer periods of time on-screen tend to experience a
39
40 more disadvantaged sense of self in reality (i.e., poorer outcomes in body composition,
41
42 fitness, self-esteem, self-worth, pro-social behavior, and academic achievement;
43
44 Tremblay et al., 2011). Thus, a cyclical interaction can be reinforced, understood
45
46 through the Selection Optimization and Compensation (SOC) model (Lerner, Freund,
47
48 Stefanis & Habermas, 2001). This portrays individuals directing their psychological
49
50 and physical efforts towards a goal or behavior that provides them with the greatest
51
52
53
54
55
56
57
58
59
60

benefit (in this case their avatar; Lerner, Freund, Stefanis & Habermas, 2001).
Consequently, the present study hypothesizes that higher PSP levels will accommodate
a fusion between the player's real (physical) self and virtual self (Yee et al., 2009)
potentially inducing IGD behaviors (Billieux et al., 2013; 2015; Hsu et al., 2009).

1.2.3. Moderation effect: Physical activity

Given that PSP, as an IGD risk factor, could be understood via compensating for physical disadvantages in the real world (Blinka, 2008; Lerner et al., 2001; Smahel et al., 2008), one could assume that improving physical activity (PA) could reduce the propensity for IGD behavior among players with higher PSP (Young, 2009). PA can be defined as any form of motion throughout the body that indicates significant movement, impacting or changing the individuals overall physical status or wellbeing (e.g., walking, bike riding, gardening, etc.; Fitbit, 2016; Kendzor et al., 2008; Kremer, Malkin, & Benshoff, 1995). Indeed, increasing PA in an individual's daily-life schedule by implementing an exercise routine has been found to reduce the severity of other addictive behaviors (Weinstock, Barry, & Petry, 2008), such as smoking (Kurti & Dallery, 2014), and alcohol consumption (Murphy et al., 1986; Read et al., 2001). Therefore, there are reasons to expect that PA may act as a moderator (buffer) between PSP and IGD (Bronfenbrenner, 2006; Cicchetti & Toth, 2009; Griffiths, 2005). This expectation is reinforced by consistent findings that have identified PA as a significant buffer in the associations between a range of maladaptive behaviors and physiological difficulties (McDonnall, 2011; Richardson et al., 2005; Schmidt, Beck, Rivkin, & Diestel, 2016; Sliter, Sinclari, Cheung, & McFadden, 2014; Sliter & Yuan, 2015; Torsheim et al., 2010).

1
2
3
4 The potential buffering effect of PA on IGD symptoms of players with higher
5
6 PSP levels can be supported on the basis of three theoretical hypotheses. First, *the*
7
8 *distraction hypothesis* suggests that diversions that draw attention from negative stimuli
9
10 could enable the maintenance, or the improvement of one's psychological status
11
12 (Festinger & Maccoby, 1964). Therefore, PA may facilitate competitive and
13
14 pleasurable emotions, redirecting interest from excessive Internet gaming involvement
15
16 (Read et al., 2001; Weinstock et al., 2008). Second, *the mastery hypothesis* emphasizes
17
18 the potential reduction of negative mood affects (that may precipitate and perpetuate
19
20 IGD), when an individual experiences a sense of accomplishment through PA-related
21
22 activities (Greist et al., 1979). A high level of PA performance, or reaching milestones,
23
24 has been associated with positive mood states (Feltz, 2007; Lerner, Freund, Stefanis &
25
26 Habermas, 2001), higher self-confidence, and personal judgements of worthiness
27
28 (Feltz, 2007; Taylor, 1987; Woodman & Hardy, 2003). These positive mood states
29
30 may later encourage further PA participation, thus acting as a buffer for IGD (Calzo et
31
32 al., 2014; Feltz, 2007). Finally, *the social interaction hypothesis* suggests that
33
34 socialization due to PA involvement can induce positive feelings (Bengston, Gans,
35
36 Putney & Silverstein, 2009). Studies contend that various forms of social exercise can
37
38 encourage positive group outcomes and maintenance of personal relationships, a
39
40 benefactor for the individual's overall mental wellbeing (Bengston et al., 2009; Duncan
41
42 & McAuley, 1993; McIntyre, Watson & Cunningham, 1990), that can influence further
43
44 participation and substitute gaming socialization. Therefore, it is hypothesized that
45
46 potentially beneficial outcomes of PA will compete with those attainable in Internet
47
48 Gaming. In this context, PA will anchor the player's presence to the physical world,
49
50 decreasing the effect of PSP in relation to IGD behaviors.
51
52
53
54
55
56
57
58
59
60

1.3. The Present Study

The present study¹ adopted a risk and resilience theoretical perspective that dimensionally conceptualizes IGD behaviors on a continuum from minimum to maximum, to explore the gamers' level of physical-body identification with their avatars (PSP) and PA, as risks and **resilience factors** of IGD behaviors respectively. To address these aims, the present study combined a cross-sectional and a longitudinal (three measurements over a period of three months, one month apart each) mixed methods design (involving both psychological and physiological assessments). More specifically, it examined a sample of Australian, emergent adults (18-29 years), MMO players, collected online and face-to-face. Based on the literature reviewed, the following hypotheses were formulated:

H1: **A higher level of physical-body identification between gamers and their avatars (PSP) will increase the risk IGD. Compensatory internet use behaviors may explain the association between higher PSP and IGD behaviors (Arnett, 2000; Blinka, 2008; Kardefelt-Winther, 2014; Smahel et al., 2008).** These behaviors can potentially generate a psychological attachment to the avatar as a means to experience an idealized self, increasing gaming exposure and IGD behaviors (Yee, & Bailenson, 2007).

H2: A higher PA will reduce (i.e., positively moderate) the IGD behaviors of gamers more physically-body identified with their avatars. PA has been repeatedly highlighted as an effective strategy of reducing addictive behaviors (Kurti & Dallery, 2014; Murphy et al., 1986), and a significant potential buffer of psychopathological symptoms in general (Richardson et al., 2005; Torsheim et al., 2010). PA combines a range of beneficial outcomes (i.e., sense of accomplishment, improved sense of body

1
2
3
4 image, etc.) that may compete with reasons for excessive MMO game engagement
5
6 (Bengston et al., 2009; Duncan & McAuley, 1993; Festinger & Maccoby, 1964).
7

8 **2. Materials and Methods**

9 **2.1. Sample**

10
11 The sample comprised 125 participants (64 online and 61 face-to-face) who were
12
13 emerging adults, aged between 18-29 years old, Australian residents, and played MMO
14
15 games (i.e., *World of Warcraft*, etc.; Arnett, Žukauskienė, & Sugimura, 2014)¹.
16
17 Participants in the online component of the study comprised 49 males and 15 females
18
19 aged from 18-29 years old ($M=23.34$ years, $SD=3.29$). Participants in the face-to-face
20
21 component of the study (three measurements over a period of three months, one month
22
23 apart each) comprised 45 males and 16 females aged 18-29 years old ($M=22.53$ years,
24
25 $SD=3.04$; see Figure 1). The combined use of online and offline data has been applied in
26
27 past studies (Orgad, 2005; Wilson, & Atkinson, 2005) and has been further advocated by
28
29 findings indicating equivalence between online and face to face collection methods
30
31 (Pettit, 2002; Weigold, Weigold, & Russel, 2013). In the present study, to ensure there
32
33 were no significant differences between the online and face-to-face samples in relation
34
35 to their demographic and internet use characteristics, independent sample t-tests and
36
37 chi-square analyses were additionally conducted. No significant differences were
38
39 detected concerning gender ($X^2=.21$, $DF=1$, $p=.89$), age, ($t_{(120)}=-.54$, $p=.59$), years
40
41 of internet use ($t_{(122)}=2.35$, $p=.06$), PSP ($t_{(119)}=1.09$, $p=.28$) and IGD scores ($t_{(111)}=-.14$,
42
43 $p=.89$). PA measurements were not collected for the online sample and therefore
44
45 relevant comparisons could not be calculated. Based on these, the present study
46
47 combined the face-to-face sample at Time Point 1 (TP1; $n=61$), with the online sample
48
49 ($n=64$) to address cross-sectional related questions (see Table 1 for sample details)².
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 The longitudinal (face-to-face/ offline) design was further assessed for attrition.
5
6 Time point 1 (TP1) comprised 61 participants, Time Point 2 (TP2) 56 participants
7
8 (8.20% attrition), and Time Point 3 (TP3) 43 participants (29.51% attrition from TP1).
9
10 The estimated maximum sampling errors with a size of 125 (cross-sectional sample) and
11
12 a size of 61 (longitudinal sample) are 8.77% and 12.55% ($Z = 1.96$, confidence level
13
14 95%) respectively.
15

16
17 -Table 1. *Sociodemographic Characteristics of the Participants in the Study-*

18
19
20 -Figure 2. Visual Representation of the Present Study-

22 23 **2.2. Instruments**

24
25 **2.2.1. Internet Gaming Disorder Scale – Short Form 9 (IGDS-SF9).** To
26
27 dimensionally examine IGD behaviors and reinforce assessment consistency among
28
29 researchers, the present study used the IGDS-SF9 (Pontes & Griffiths, 2015). The
30
31 IGDS-SF9 is based on the nine criteria in the DSM-5 for IGD (APA, 2013). It
32
33 comprises nine items that elicit information in relation to IGD behaviors (e.g., “Do you
34
35 feel more irritability, anxiety or even sadness when you try to either reduce or stop your
36
37 gaming activity?”) rated on a 5-point Likert scale (1 “Never” to 5 “Very Often”). To
38
39 calculate the final IGD score, each item score is added together resulting in a total
40
41 ranging from 9 to 45 with higher scores being indicative of IGD behaviors. Internal
42
43 reliability in the present study was high, with a Cronbach’s alpha of .88.
44
45

46
47 **2.2.2. Proto-Self-Presence Subscale (PSP) of the Self-Presence Questionnaire**
48
49 **(SPQ).** To assess the level of user-avatar physical-body identification, the PSP subscale
50
51 of the SPQ was used (Ratan & Dawson 2015). The SPQ is a 13-item self-report
52
53 instrument that assesses different aspects of the user-avatar relationship. Only the PSP
54
55 subscale was used in the present study. PSP describes the extent to which the mediated
56
57
58
59

1
2
3
4 self-representation in the game (i.e., avatar) is integrated into body schema of the user
5
6 and consists of five items (e.g., “When playing the game, how much do you feel like
7
8 your avatar is an extension of your body within the game?”). Participants respond to
9
10 each item on a 5-point Likert: 0 (“Not at all”), 1 (“Somewhat”), 2 (“Moderately”), 3
11
12 (“Very much”), and 4 (“Absolutely”). Scores of the five items are added to provide the
13
14 total PSP score ranging between 0 and 20, with higher scores indicating a higher
15
16 experienced level of PSP. Internal reliability of the PSP subscale in the present study
17
18 was high with a Cronbach’s alpha of .90.
19
20

21 **2.2.3. Physical activity: Active minutes.** To assess PA, the current study utilized
22
23 an activity monitor (i.e., a *Fitbit Flex*; FBF). Face-to-face participants were required to
24
25 consistently carry/wear the FBF for a period of three days for each of the three time
26
27 points assessed. The FBF constitutes an actuarial measurement that records the number
28
29 of steps, distance achieved, calories consumed, and accumulated ‘active minutes’ per
30
31 period of time chosen (Fitbit, 2016). More specifically, the number of ‘active minutes’
32
33 was selected as the PA indicator in the present study. Active minutes calculation relies
34
35 on recording the levels of energy-consumed-per-minute in offline physiological
36
37 activities. One active minute, as a unit of measure, is calculated using metabolic
38
39 equivalents that keep track of the level of energy expenditure (Fitbit, 2016). Therefore,
40
41 *Fitbit* active minute measurements rely on metabolic equivalents that record activities’
42
43 intensity. By measuring the intensity of the individual’s activities as opposed to keeping
44
45 track of distance covered or calories burned, active minutes become comparable against
46
47 users of different weights (Fitbit, 2016). The FBF validity and reliability were found to
48
49 be high with an error estimation of 17-18% (Alharbi, Bauman, Neubeck, & Gallagher,
50
51 2016; Bai et al., 2015; Diaz et al., 2015) and their use has been recommended in
52
53
54
55
56
57
58
59
60

1
2
3
4 medical practice to monitor treatment plans involving PA (Alharbi et al., 2016; Diaz et
5 al., 2015).

6
7
8 Studies have found that the FBF has an increased accuracy for longer periods of
9 time, compared to that of other activity tracking devices, partially due to its convenient
10 use (e.g., wrist-wear; Diaz et al., 2016). To increase the measurements' validity and
11 reliability, the present study adopted previous research recommendations to aggregate
12 measurements (i.e., produce mean scores) over longer periods of time (Diaz et al.,
13 2016). Therefore, the present study aggregated PA scores over the course of three days
14 per-time-point (to produce the mean of active minutes).

23 **2.3. Procedure**

24
25 This project was approved by the Human Research Ethics Committee of
26 Federation University Australia (HREC-A16-044, 2016). Participants were recruited in
27 the general community using traditional (i.e., information flyers) and electronic (i.e.,
28 email, social media) advertising methods³. The face-to-face, longitudinal component of
29 the study was conducted over three months, with the three time-point assessments
30 completed between June 2016 and September 2016. The process of data collection was
31 identical between the three time points and participants' measurements were matched
32 via the use of a re-identifiable code. A specially trained research team of five
33 undergraduate students and two postgraduate students collected the face-to-face data.

34
35
36 Considering the online collection (conducted between June 2016 and July 2016),
37 and based on the ethical approval received, eligible individuals interested in
38 participating and unable or unwilling to attend face-to-face testing sessions were invited
39 to register with the study via a *Survey-Monkey* link available on MMO gaming websites
40 and forums (i.e., <http://www.ausmmo.com.au>). The link directed individuals to the
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 plain language information statement. Individuals who chose to participate online had
5
6 to digitally provide informed consent (i.e., click a button). Physical measurements were
7
8 not included in the online collection.
9

10 **2.4. Analyses and statistical testing**

11
12 To assess both cross-sectionally and longitudinally the association between PSP
13 levels and IGD behaviors (H_1), offline and online questionnaire data were combined
14
15 (after controlling for significant differences) using the merge data command of SPSS 21
16
17 (see Figure 1). This method could only be applied to the questionnaire variables as PA
18
19 (i.e., active minutes) was only measured in the offline group. Two hierarchical linear
20
21 regression models were conducted on the cross-sectional and the longitudinal data
22
23 respectively. In relation to the cross-sectional analysis, and to control for the possible
24
25 confounding effects of gender and age on IGD behaviors (Andreassen et al., 2016), both
26
27 gender and age were inserted separately as IGD predictors during the first step. PSP was
28
29 then inserted as an IGD predictor at the second step of the model. In relation to the
30
31 longitudinal analysis, the same two-step hierarchical linear regression model was
32
33 calculated with the differences of (i) IGD behaviors assessed at TP3 being used as the
34
35 outcome variable, and (ii) PSP levels assessed at TP1 being used as the IGD predictor
36
37 variable at the second step. Potential confounding effects of age and gender were
38
39 (similarly to the cross-sectional analysis) controlled at the first step of the model (see
40
41
42
43
44
45 Table 2 for descriptive data of the sample).
46

47 To assess the potentially buffering (protective) effect of PA on the IGD behavior
48
49 of gamers, who are more physically-body identified with their avatars (present with
50
51 higher PSP levels) (H_2), a moderation analysis was conducted using only the
52
53 longitudinal data through the Process macro (Hayes, 2013) on SPSS 21. The use of the
54
55
56
57
58
59
60

1
2
3
4 longitudinal data only was based on recommendations supporting that causal
5
6 associations should be best addressed longitudinally (Winer et al., 2016). More
7
8 specifically, PSP at TP1 was used as the independent variable, PA (active minutes) at
9
10 TP2 were used as the moderator, and IGD behaviors at TP3 were used as the outcome
11
12 variable. Finally, the Johnson Neyman (J–N) technique was applied to derive regions of
13
14 significance (i.e., points of transition) of the moderation effect of PA on the association
15
16 between PSP and IGD behaviors (Preacher, Rucker, & Hayes, 2007).
17
18

19 3. Results

20
21
22 In the H_1 cross-sectional analysis, the slope of the two-step (step₁ gender and age,
23
24 step₂ PSP) hierarchical regression model calculated was significant ($F_{(3,118)} = 13.10, p <$
25
26 $.001$) and accounted for 25% of the IGD behaviors' variance ($R^2 = .25$). The addition of
27
28 PSP to the model accounted for 20% of the IGD behaviors' variance ($F_{change(1,118)} =$
29
30 $31.55, p < .001, R^2_{change} = .20$). More specifically, for each point of PSP increase, IGD
31
32 behaviors increased by .81 ($b = .81, p < .001$; see Table 2).
33
34

35 -Table 2. PSP predicting IGD (Cross-section)-

36
37 To complement and reinforce the direction of causality of the association between
38
39 PSP and IGD, the model was repeated using the longitudinal data (Winer et al., 2016).
40
41 In the H_1 longitudinal analysis, the slope of the two-step (step₁ gender and age, step₂
42
43 PSP) hierarchical regression model was (similarly to the cross-sectional model)
44
45 significant ($F_{(3,54)} = 10.36, p < .001$). The full model accounted for 37% of the variance
46
47 in the IGD ($R^2 = .37$). The addition of PSP to the model accounted for 36% of variance
48
49 ($F_{change(1,54)} = 30.55, p < .001, R^2_{change} = .36$). More specifically, for each point of
50
51 increase of PSP, IGD behaviors increased by .76 ($b = .76, p < .001$).
52
53

54 -Table 3. PSP predicting IGD (Longitudinal)-

1
2
3
4
5
6 The longitudinal data were chosen to address H_2 for two reasons. First, to
7
8 reinforce the inferences of causality assessed in the moderation analysis (Winer et al.,
9
10 2016) and second because PA measurements were not collected for the online cross-
11
12 sectional data. In the H_2 moderation analysis, the model aimed to estimate the risk
13
14 effect of PSP (IV) levels on IGD behaviors (DV) and how much, if at all, this effect
15
16 was buffered by higher levels of PA (Moderator) as measured by FBF active minutes.
17
18 Thus, PSP as the focal predictor (F), PA as the proposed moderator (M) and their
19
20 interaction (F x M) were included as predictors in an OLS regression predicting IGD
21
22 behaviors. Gender and age were also included as covariates in the model to control for
23
24 their effects. The equation estimated was the following:
25
26

$$IGD = a + b_1(PSP) + b_2(PA) + b_3(PSP \times PA) + b_4(Gender) + b_5(Age)$$

27
28 Findings indicated that the full model accounted for 41% of the variance of IGD
29
30 behaviors. The slope of the regression line for the overall model was significant ($F_{(5,52)}$
31
32 $= 7.35$, $p < .001$). An inspection of the moderation coefficient indicated that PSP and PA
33
34 significantly interacted (buffering effect) in predicting IGD behaviors, ($b_3 = -0.10$, $t_{(52)} = -$
35
36 2.05 , $p = .046$, LLCI = -0.01 , ULCI = -0.0005). This finding indicates that when gamers
37
38 present concurrently at a higher point (than the intercept) on PSP and at a higher point
39
40 on PA then their IGD scores reduce (i.e. the IGD behavior of those higher on PSP
41
42 decreases when they present with higher PA scores). The Johnson Neyman (J–N)
43
44 technique was then applied to examine how the effect of PSP on IGD differed
45
46 according to PA variations (Preacher, Rucker, & Hayes, 2007). Findings demonstrated
47
48 that the buffering effect of PA on the PSP-IGD association had a ceiling level at 144.92
49
50 active minutes. This means that while for PA values less than 144.92 active minutes,
51
52
53
54
55
56
57
58
59
60

1
2
3
4 the IGD behavior of gamers higher on PSP significantly decreased, the IGD behavior
5
6 related to higher levels of PSP did not further decrease after that point of active
7
8 minutes' (i.e., had reached a maximum level at 144.92 active minutes; see Figure 2).
9

10 -*Figure 2.* Moderation summary of PA (Active Minutes) on the relationship between
11
12 Proto-self and IGD-
13

14 **4. Discussion**

15
16 The interplay of user-avatar characteristics has been identified as an area of
17
18 research priority in relation to the development of IGD behaviors (Stavropoulos et al.,
19
20 2017a; Stavropoulos et al., 2017b). Consequently, the present study investigated
21
22 Massively-Multiplayer Online (MMO) gamers' level of physical-body identification
23
24 with their avatars (Proto-Self-Presence [PSP]; Ratan & Dawson, 2015) and their
25
26 physical activity (PA) levels as risk and protective factors of IGD behaviors
27
28 respectively. The study findings and their practical implications in planning IGD
29
30 prevention and intervention initiatives, as well as relevant future gaming developments
31
32 are discussed below.
33
34
35

36 **4.1. PSP and IGD**

37
38 In relation to H_1 , it was shown that higher PSP was associated with higher IGD
39
40 behaviors. Results additionally indicated that the association between PSP and IGD
41
42 behaviors remained significant over the period of time examined (i.e., three months).
43
44 These findings are in consensus with previous cross-sectional studies that have reported
45
46 associations between higher levels of user-avatar identification and excessive gaming
47
48 (Billieux et al., 2013; Blinka, 2008). As players establish a stronger presence in the
49
50 virtual world (via higher levels of body-identification with their avatars), they tend to
51
52 increase the amount of time spent gaming. Avatar control tasks and addressing gaming
53
54
55
56
57
58
59
60

1
2
3
4 challenges, that have been associated with higher PSP, could induce achievement and
5
6 gaming socialization experiences that may increase the players' gaming engagement
7
8 (Hsu et al., 2009; Luppici, 2013; Yee, 2006). This notion could be further reinforced
9
10 by the players' tendency to compensate for (perceived) real life body disadvantages
11
12 through their game character, as supported by the compensatory internet use model
13
14 (Kardefelt-Winther, 2014). Therefore – and based on the SOC model – it was expected
15
16 that players would concentrate their attention and efforts towards their virtual self, if
17
18 this was perceived as being more emotionally beneficial (Lerner et al., 2001). This
19
20 concurs with studies showing that an individuals' longer internet use associates with
21
22 poorer perception of their self in real life (Tremblay et al., 2011) and is encouraged
23
24 through more positive emotional gaming experiences (Banks & Bowman, 2016). Such
25
26 processes may exacerbate IGD behaviors by encouraging longer game play sessions that
27
28 could escalate the level that gamers emotionally rely on their in-game persona (Blinka,
29
30 2008; Hopp, Barker, & Weiss, 2015; Smahel et al., 2008).

31 32 33 34 **4.2. PA as a buffer of the risk effect of PSP on IGD**

35
36 In relation to H_2 , higher PA was shown to weaken the risk effect of PSP on IGD
37
38 behaviors. This finding concurs with past studies showing that PA significantly
39
40 moderates the associations between several psychopathological symptoms and behavior
41
42 (McDonnall, 2011; Richardson et al., 2005; Schmidt, Beck, Rivkin, & Diestel, 2016;
43
44 Sliter, Sinclari, Cheung, & McFadden, 2014; Sliter & Yuan, 2015; Torsheim et al.,
45
46 2010) and re-emphasizes the potential benefit of incorporating PA into medical and
47
48 psychological treatment programs (Kurti & Dallery, 2014). In line with this, the severity
49
50 of various forms of addictive behaviors has been found to decline via increased PA
51
52
53
54
55
56
57
58
59
60

(Kurti & Dallery, 2014; Murphy et al., 1986; Read et al., 2001; Weinstock, Barry, & Petry, 2008).

PA's buffering effect (on the higher IGD behavior of players experiencing higher PSP) could be explained on the basis of redirecting the player's attention via attaining positive PA outcomes and fulfilling goals by facilitating competitive and pleasurable physical activities (i.e., *the distraction hypothesis*; Festinger & Maccoby, 1964; Greist et al., 1979). These may increase players' sense of confidence in their real-world-self, thus competing with their gaming involvement (i.e., *the mastery hypothesis*; Greist et al., 1979). Furthermore, increased PA could involve group activities that initiate and strengthen personal relationships (i.e., *the social interaction hypothesis*) that reinforce players' real world identity, reducing their experienced PSP (Bengston et al., 2009; Duncan & McAuley, 1993; McIntyre, Watson & Cunningham, 1990). Subsequently, PA may put players (who are more physically identified with their avatars) in a positive real-world mood state that could later encourage further participation, thus acting as an IGD buffer (Calzo et al., 2014; Feltz, 2007). Furthermore, PA may reinforce real world physiological features and performance (Young, 2009), buffering the negative associations attributed to higher levels of PSP and reducing IGD vulnerability. Overall, it could be assumed that PA competes with attractive MMO features that involve improving personal, social, and game-related achievements and development (Andreassen et al., 2016; Bengston et al., 2009; Calzo et al., 2014; Feltz, 2007; Festinger & Maccoby, 1964). Consequently, PA appears to potentially counterbalance characteristics of MMO gaming in a way that could strengthen the players' physical identity and deter them from being more physically identified with their avatars (i.e., experience higher PSP).

1
2
3
4 Despite the significant moderation effect detected, findings demonstrated that
5
6 IGD behaviors of more physically/bodily identified with their avatars gamers did not
7
8 further decrease, after PA had reached significantly high levels. Therefore, PA appears
9
10 to have a maximum potential for reducing the PSP risk effect on IGD behaviors. This
11
12 could be interpreted as the result of user-avatar body identification (and the associated
13
14 IGD risk with it) not being exclusively associated to real world PA attractiveness
15
16 (Andreassen et al., 2016). Caplan's (2009) study was replicated indicating game-related
17
18 features (i.e., avatar graphs, gaming structure, etc.) contributed to the gamer's
19
20 immersion with their virtual character, increasing the amount of time within the virtual
21
22 environment (Andreassen et al., 2016). These gaming-related factors constitute
23
24 attractive features of the game context that may impose some level of IGD behavior,
25
26 independent of how PA may increase attraction to the real world.
27
28

29 30 **4.3. Strengths and limitations**

31
32 The present study contained several strengths: (a) a concurrent examination of
33
34 online and face-to-face participants; (b) a combination of cross-sectional and
35
36 longitudinal analyses; (c) an emphasis on an under-researched high risk population for
37
38 IGD (emergent adults, MMO gamers); (d) a focus on both individual and gaming
39
40 related factors, as well as their interplay in relation to IGD; and (e) the use of a mixed-
41
42 methods design that comprised both physical and psychological measurements.
43
44

45 However, there are several limitations that need to be considered. The most
46
47 notable limitation during data collection was the introduction of new expansions for
48
49 major online games (e.g., *Legion for World of Warcraft*) and the development of mobile
50
51 games (e.g., *Pokémon Go*). A previous study that surveyed *World of Warcraft* players,
52
53 found that the introduction of new gaming content increased player's exposure and
54
55
56
57
58
59
60

1
2
3
4 financial involvement (Constantiou, & Legarth, 2012). These motivations may have
5
6 affected data in a way that was outside the control of the current study. Another
7
8 limitation identified was that the assessment of IGD focused on a specific online
9
10 gaming genre (i.e., MMO games). There are many different online gaming genres, and
11
12 therefore characteristics of those who play them may vary accordingly. Furthermore, the
13
14 sample size being relatively small (N=125), suggesting that limitations may exist when
15
16 extrapolating and generalizing the results. The longitudinal design applied was over a
17
18 relatively short time frame (i.e., three months). In terms of the tools used, the FBF was
19
20 not a precise measure, like other physical activity trackers, the FBF simply provides
21
22 relatively accurate estimations (Alharbi et al., 2016; Bai et al, 2015; Diaz et al, 2015).
23
24 Additionally, the present study used self-report measures which may affect reliability as
25
26 participant responses can be influenced by mood and situational factors at the time
27
28 when the measurement was taken (Smith & Handler, 2007). Finally, different directions
29
30 of the causal relationships between the variables involved in the study could be
31
32 examined (i.e., PA moderated by PSP on IGD risk).
33
34
35

36 4.4. Implications

37
38 Considering IGD prevention, the findings suggest that individuals who experience
39
40 stronger PSP (i.e., are more bodily identified with their avatars) need to be prioritized,
41
42 especially when they exhibit lower levels of PA. Concerning treatment directions,
43
44 cognitive-behavior therapy may be useful to promote awareness in regard to the use of
45
46 IGD as a potential maladaptive emotional compensation strategy, emphasizing the
47
48 significance of the identification of gamers with their avatars. Interpersonal therapy may
49
50 prove beneficial when exploring the role of PSP on IGD behavior. Finally, it is
51
52
53
54
55
56
57
58
59
60

recommended that PA components should be included in the IGD treatment of players who present with higher PSP levels.

Conceptually, these implications resemble the foundations that comprise Tisseron's (2009) method of psychological IGD treatment. More specifically, Tisseron attempts to understand what facets of the player's characteristics are being compensated or are more appreciated within the virtual identity, and attempts to find ways to fulfill such gaps within the gamer's real identity (Tisseron, 2009).

Gaming implications can also be derived from the findings. Firstly, there is the notion of promoting healthy behaviors through games by combining virtual elements (e.g., goal orientation and avatar progression) with PA. This concept has already been highlighted through the mobile game *Pokémon Go* which encourages its player base to increase their PA to progress. In line with this development, gaming platforms like the virtual reality headsets (e.g., *Oculus Rift* and *Playstation VR*), *Nintendo Wii*, and *Microsoft Kinetic*, are some of the many examples of gaming organizations implementing PA components into their products. Additionally, the findings reiterate the reason for the increasing popularity for MMOs. Visual progression (i.e., experience bar, upgrading gear, expanding guild member count etc), character customization, and large communities are all attractive and rewarding experiences. Understanding the effects of these components may indicate why some games are more popular than others which can direct future gaming development and product choice.

4.5. Future research

Future research should further investigate the longitudinal interplay between individual (i.e., age, gender, societal, cultural background) and gaming-related factors (i.e., game genre, game structure, character development, etc.). Furthermore, increasing

1
2
3
4 the length of the longitudinal period studied could greatly benefit knowledge
5
6 considering time-related variations in the association between PSP and IGD, as well as
7
8 the PSP-PA interaction. Investigating other potential protective factors that might
9
10 reduce the association between PSP and IGD may also add to the extant knowledge.
11
12 Such findings can help inform the guidelines for more effective IGD treatment
13
14 programs. Given that IGD has been classified as a potentially significant public health
15
16 risk (APA, 2013), it is imperative to develop and expand our understanding of IGD to
17
18 better address it.
19
20
21
22

23 5. References

- 24
25 Adams, B. L., Stavropoulos, V., Burleigh, T. L., Liew, L. W., Beard, C. L., & Griffiths,
26
27 M. D. (2018). Internet Gaming Disorder Behaviors in Emergent Adulthood: a
28
29 Pilot Study Examining the Interplay Between Anxiety and Family Cohesion.
30
31 *International Journal of Mental Health and Addiction*, 1-17.
32
33
34 Alharbi, M., Bauman, A., Neubeck, L., & Gallagher, R. (2016). Validation of Fitbit-
35
36 Flex as a measure of free-living physical activity in a community-based phase III
37
38 cardiac rehabilitation population. *European Journal of Preventive Cardiology*,
39
40 33(124), 1-10. doi: 10.1177/2047/487316634883
41
42
43 American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental*
44
45 *disorders (5th ed.)*. Washington, DC
46
47 Anderson, E. L., Steen, E., & Stavropoulos. (2016). Internet use and problematic
48
49 internet use: A systematic review of longitudinal research trends in adolescence
50
51 and emergent adulthood. *International Journal of Adolescence and Youth*, 21(4).
52
53 doi: 10.1080/02673843.2016.1227716.
54
55
56
57
58
59
60

1
2
3
4 Armitage, G., Claypool, M., & Branch, P. (2006). *Networking and online games: Understanding and engineering multiplayer internet games*. Chichester: Wiley.
5
6
7
8 doi:10.1002/047003047X.
9

10
11 Andreassen, C. S., Griffiths, M. D., Kuss, D. J., Mazzoni, E., Billieux, J., Demetrovics,
12
13 Z., & Pallesen, S. (2016). The relationship between addictive use of social media
14
15 and video games and symptoms of psychiatric disorders: A large-scale cross-
16
17 sectional study. *Psychology of Addictive Behaviors*, 30 (2), 252-262. doi:
18
19 10.f037/abdb0000160
20

21
22 Arnett, J. J. (2000). Emerging adulthood: A theory of development from the late teens
23
24 through the twenties. *American Psychologist*, 55(1), 469-480. doi: 10.1037/0003-
25
26 066X.55.5.469
27

28
29 Arnett, J. J., Žukauskienė, R., & Sugimura, K. (2014). The new life stage of emerging
30
31 adulthood at ages 18–29 years: Implications for mental health. *The Lancet*
32
33 *Psychiatry*, 1(7), 569-576.
34

35
36 Badrinarayanan, V.A., Sierra, J.J., & Taute, H.A. (2014). Determinants and outcomes of
37
38 online brand tribalism: Exploring communities of Massive Multiplayer Online
39
40 Role Playing Games (MMORPGs). *Psychology and Marketing*, 31(10), 853-870.
41
42 doi: 10.1002/mar.20739.

43
44 Bai, Y., Welk, G.J., Nam, Y.H., Lee, J.A., Lee, J., Kim, Y., Meier, N.F. & Dixon, P.M.
45
46 (2015). Comparison of consumer and research monitors under semistructured
47
48 settings, *Medicine & Science in Sports & Exercise*, 48(1), 151-158. doi:
49
50 10.1249/MSS.00000000000000727
51

52 **Bainbridge, W. S. (Ed.). (2009). *Online worlds: Convergence of the real and the virtual*.**

53 **Springer Science & Business Media.**
54
55
56
57
58
59
60

1
2
3
4 Banks, J., & Bowman, N. D. (2016). Emotion, anthropomorphism, realism, control:
5
6 Validation of a merged metric for player-avatar interaction. *Computers in Human*
7
8 *Behavior*, 54(1), 215-223. doi: 10.1016/j.chb.2015.07.030.
9

10 Bessière, K., Seay, Fleming., & Kiesler, S. (2007). The ideal elf: Identity exploration in
11
12 world of warcraft. *CyberPsychology & Behavior*, 10(4), 530-535. doi:
13
14 10.1089/cpb.2007.9994
15

16
17 Bengston, V. L., Gans, D., Putney, N., & Silverstein, M. (2009). *Handbook of theories*
18
19 *of aging*, (2nd ed.). New York: Springer.
20

21 Billieux, J., Thornes, G., Khazall, Y., Zullino, D., Achab, S., & Van der Linden, M.
22
23 (2015). Problematic involvement in online games: A cluster analytic approach.
24
25 *Computers in Human Behavior*, 43(1), 242-250. doi: 10.1016/j.chb.2014.10.055.
26

27
28 Billieux, J., Van der Linden, M., Achab, S., Kazaal, Y., Paraskevopoulos, L., Zullino,
29
30 D., & Thorens, G. (2013). Why do you play World of Warcraft? An in-depth
31
32 exploration of self-reported motivations to play online and in-game behaviors in
33
34 the virtual world of Azeroth. *Computers in Human Behavior*, 29(1), 103-109. doi:
35
36 10.1016/j.chb.2012.07.021.
37

38 Blinka, L. (2008). The Relationship of Players to Their Avatars in MMORPGs:
39
40 Differences between Adolescents, Emerging Adults and Adults.
41
42 *Cyberpsychology: Journal of Psychosocial Research on Cyberspace*, 2(1), 1.
43
44 Retrieved January 24, 2016, from:
45
46 <http://cyberpsychology.eu/view.php?cisloclanku=2008060901&article=1>
47
48

49 Brand, J.E., & Todhunter, S. (2015). *Digital Australia 2016*. Eveleigh, NSW: IGEA.
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 Bronfenbrenner, U. & Morris, P. A. (2006). The bioecological model of human
5
6 development. In R. M. Lerner (Ed.), *Handbook of child psychology: Theoretical*
7
8 *models of human development (Sixth edition)* (pp. 793-828). Hoboken, NJ: Wiley.
9

10
11 Burleigh, T. L., Stavropoulos, V., Liew, L. W., Adams, B. L., & Griffiths, M. D. (2017).
12
13 Depression, Internet Gaming Disorder, and the Moderating Effect of the Gamer-
14
15 Avatar Relationship: an Exploratory Longitudinal Study. *International Journal of*
16
17 *Mental Health and Addiction*, 1-23. doi: 10.1007/s11469-017-9806-3
18

19
20 Caplan, S., Williams, D., & Yee, N. (2009). Problematic Internet use and psychosocial
21
22 well-being among MMO players. *Computers in Human Behavior*, 25(6), 1312-
23
24 1319. doi:10.1016/j.chb.2009.06.006.
25

26
27 Calzo, J. P., Roberts, A. L., Corliss, H. L., Blood, E. A., Kroshus, E., & Austin, S. B.
28
29 (2014). Physical activity disparities in heterosexual and sexual minority youth
30
31 ages 12-22 years old: Roles of childhood gender nonconformity and athletic self-
32
33 esteem. *Annals of Behavioural Medicine*, 47(1), 17-27. doi: 10.1007/s12160-013-
34
35 9570-y.
36

37
38 Caplan, S., Williams, D., & Yee, N. (2009). Problematic internet use and psychosocial
39
40 well-being among MMO players. *Computers in Human Behavior*, 25(6), 1312-
41
42 1319. doi: 101016/j.chb.2009.06.006
43

44
45 Chandler, J., Konrath, S., & Schwarz, N. (2009). Online and on my mind: Temporary
46
47 and chronic accessibility moderate the influence of media figures. *Media*
48
49 *Psychology*, 12(2), 210-226. doi: 10.1080/15213260902849935.
50

51
52 Cicchetti, D. & Toth, S. L. (2009). The past achievements and future promises of
53
54 developmental psychopathology: the coming of age of a discipline. *Journal of*
55
56

1
2
3
4 *Child Psychology and Psychiatry*, 50(1-2), 16-25. doi: 10.1111/j.1469-
5 7610.2008.01979.x.
6
7

8 Corneliussen, H., & Rettberg, J. W. (2008). *Digital culture, play, and identity: A World*
9 *of Warcraft reader*. Cambridge, MA: MIT Press.
10
11

12 Constantiou, I., & Legarth, M.F. (2012). What are the users' intentions towards real
13 money trading in massively multiplayer online games? *Electron Markets*, 22(1),
14 105-115. doi: 10.1007/s12525-011-0076-9
15
16
17
18

19 Diaz, K. M., Krupka, D. J., Chang, M. J., Peacock, J., Ma, Y., Goldsmith, J., Schwartz,
20 J.E. & Davidson, K.W. (2015). Fitbit: An accurate and reliable device for wireless
21 physical activity tracking. *International Journal of Cardiology*, 185(1), 138-140.
22 doi: 10.1016/j.ijcard.2015.03.038
23
24
25
26

27 Diaz, K. M., Krupka, D. J., Chang, M. J., Shaffer, J. A., Ma, Y., Goldsmith, J.,
28 Schwartz, J. E. & Davidson, K. W. (2016). Validation of the Fitbit One for
29 physical activity measurement at an upper torso attachment site. *BMC Research*
30 *Notes*, 9(213), 1-9. Doi: 10.1186/s13104-016-2020-8
31
32
33
34
35

36 Douglas, A. C., Mills, J. E., Niang, M., Stepchenkova, S., Byun, S., Ruffini, C., Lee, S.
37 K., Loutfi, J., Lee, J., Atallah, M., & Blanton, M. (2008). Internet addiction: Meta-
38 synthesis of qualitative research for the decade 1996-2006. *Computers in Human*
39 *Behavior*, 24(1), 3027-3044. doi: 10.1016/j.chb.2008.05.009.
40
41
42
43
44

45 Duncan, T. E. & McAuley, E. J. (1993). Social support and efficacy cognitions in
46 exercise adherence: A latent growth curve analysis. *Journal of Behavioral*
47 *Medicine*, 16(2), 199-218. doi: 10.1007/BF00844893
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Elliott, L., Golub, A., Ream, G., & Dunlap, E. (2012). Video game genre as a predictor
5 of problem use. *Cyberpsychology, Behavior, and Social Networking*, *15*(3), 155-
6 161. doi: 10.1089/cyber.2011.0387.
7
8
9
10 Feltz, D.L. (2007). Self-confidence and sports performance. *Exercise and Sports*
11 *Science Reviews*, *33*(41), 50-66. Retrieved January 24, 2016, from:
12 [http://t012.camel.ntupes.edu.tw/ezcatfiles/t012/img/img/171/Feltz1988_423-](http://t012.camel.ntupes.edu.tw/ezcatfiles/t012/img/img/171/Feltz1988_423-457_16_ESSR.pdf)
13 [457_16_ESSR.pdf](http://t012.camel.ntupes.edu.tw/ezcatfiles/t012/img/img/171/Feltz1988_423-457_16_ESSR.pdf)
14
15
16
17
18 Festinger, L., & Maccoby N. (1964). On resistance to persuasive communications.
19 *Journal of Abnormal and Social Psychology*, *68*(4), 359-366. doi:
20 10.1037/h0049073
21
22
23
24
25 Fitbit. (2016). *Make fitness a lifestyle with flex*. Retrieved May 4, 2016, from:
26 <https://www.fitbit.com/au/flex>
27
28
29
30 Freeman, C. B. (2008). Internet gaming addiction. *Journal for Nurse Practitioners*, *4*(1),
31 42-47. doi: 10.1016/j.nurpra.2007.10.006.
32
33
34 Greist, J. H., Klein, M. H., Eischens, R. R., Faris, J., Gurman, A. S., & Morgan, W. P.
35 (1979). Running as treatment for depression. *Comprehensive Psychiatry*, *20*(1),
36 41-54. doi: 10.1016/0010-440X(79)90058-0.
37
38
39
40 Griffiths, M. (2005). A 'components' model of addiction within a biopsychosocial
41 framework. *Journal of Substance Use*, *10*, 191-197. doi:
42 10.1080/14659890500114359
43
44
45
46
47 Hawkins, J. D., Catalano, R. F., & Miller, J. Y. (1992). Risk and protective factors for
48 alcohol and other drug problems in adolescence and early adulthood: Implications
49 for substance abuse prevention. *Psychological Bulletin*, *11* (1), 64-105. doi:
50 10.1037/0033-2909.112.1.64
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process*
5 *analysis: A regression-based approach*. New York: Guilford Press.
6
7
8 Hopp, T., Barker, V., & Weiss, A. S. (2015). Interdependent self-construal, self-
9 efficacy, and community involvement as predictors of perceived knowledge gain
10 among MMORPG players. *Cyberpsychology, Behavior, and Social Networking*,
11 *18*(8), 468-473. doi: 10.1089/cyber.2015.0073.
12
13
14
15
16 Hsu, S. H., Wen, M., & Wu, M. (2009). Exploring user experiences as predictors of
17 MMORPG addiction. *Computers & Education*, *53*(3), 990-999. doi:
18 10.1016/j.compedu.2009.05.016.
19
20
21
22
23 Hussain, Z., Williams, G. A., & Griffiths, M. D. (2015). An exploratory study of the
24 association between online gaming addiction and enjoyment motivations for
25 playing massively multiplayer online role-playing games. *Computers in Human*
26 *Behavior*, *50*(1), 221-230. doi: 10.1016/j.chb.2015.03.075.
27
28
29
30
31
32 Hyun, G. J., Han, D. H., Lee, Y. S., Kang, K. D., Yoo, S. K., Chung, U., & Renshaw, P.
33 F. (2015). Risk factors associated with online game addiction: A hierarchical
34 model. *Computers in Human Behavior*, *48*(1), 706-713. doi:
35 10.1016/j.chb.2015.02.008
36
37
38
39
40
41 Kaplan, A. M., & Haenlein, M. (2010). Users of the world, unite! The challenges and
42 opportunities of Social Media. *Business Horizons*, *53*(1), 59-68. doi:
43 10.1016/j.bushor.2009.09.003
44
45
46
47 Kardefelt-Winther, D. (2014). A conceptual and methodological critique of internet
48 addiction research: Towards a model of compensatory internet use. *Computers in*
49 *Human Behavior*, *31*(1), 351-354. doi: 10.1016/j.chb.2013.10.059.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Kendzor, D. E., Dubber, P. M., Olivier, J., Businelle, M. S., Grothe, K. B., & PATHS
5
6 Investigators. (2008). The influence of physical activity on alcohol consumption
7
8 among heavy drinkers participating in an alcohol treatment intervention. *Addictive*
9
10 *Behaviors*, 33(10), 1337-1343. doi:10.1016/j.addbeh.2008.06.013
11
12 Kremer, D., Malkin, M. J., & Benschoff, J. J. (1995). Physical activity programs offered
13
14 in substance abuse treatment facilities. *Journal of Substance Abuse Treatment*,
15
16 12(5), 327-333. doi: 10.1016/0740-5472(95)02008-3
17
18 Kurti, A. N., & Dallery, J. (2014). Effects of exercise on craving and cigarette smoking
19
20 in the human laboratory. *Addictive Behaviors*, 39(1), 1131-1137. doi:
21
22 10.1016/j.addbeh.2014.03.004
23
24 Kuss, D. J. (2013). Internet gaming addiction: Current perspective. *Psychology*
25
26 *Research and Behavior Management*, 6(1), 125-137. doi: 10.2147/PRBM.S39476.
27
28 Kuss, D. J., & Griffiths, M. D. (2012). Internet gaming addiction: A systematic review
29
30 of empirical research. *International Journal of Mental Health and Addiction*,
31
32 10(2), 278-296. doi: 10.1007/s11469-011-9318-5.
33
34
35
36 Lazar, J., Feng, J. H., & Hochheiser, H. (2017). *Research methods in human-computer*
37
38 *interaction*. New York: Morgan Kaufmann.
39
40 Lee, Z. W. Y., Cheung, C. M. K., & Chan, T. K. H. (2015). Massively multiplayer
41
42 online game addiction: Instrument development and validation. *Information &*
43
44 *Management*, 52(4), 413-430. doi: 10.1016/j.im.2015.01.006.
45
46 Lee, M., Ko, Y., Song, H., Kwon, K., Lee, H., Nam, M., & Jung, I. (2007).
47
48 Characteristics of internet use in relation to game genre in Korean adolescents.
49
50 *CyberPsychology & Behavior*, 10(2), 278-285. doi: 10.1089/cpb.2006.9958.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Lerner, R. M., Freund, A. M., Stefanis, I. D., & Habermas, T. (2001). Understanding
5
6 developmental regulation in adolescence: The use of the selection, optimization,
7
8 and compensation model. *Human Development*, 44(1), 29-50. Retrieved January
9
10 24, 2017, from www.karger.com/journals/hde
11
12
13 Luppicini, R. (2013). *Handbook of research on technoself: Identity in a technological*
14
15 *society*. Hershey, PA: Idea Group Publishing.
- 16
17 Masten, A. S. (2014) Invited commentary: Resilience and positive youth development
18
19 frameworks in developmental science. *Journal of Youth and Adolescence*, 43(6),
20
21 1018-1024. doi:10.1007/s10964-014-0118-7
22
- 23
24 McDonnall, M. C. (2011). Physical status as a moderator of depressive symptoms
25
26 among older adults with dual sensory loss. *Rehabilitation Psychology*, 56(1), 67-
27
28 76. doi: 10.1037/a0022696
29
- 30
31 Murphy, T. J., Pagano, R. R., & Marlatt, G. A. (1986). Lifestyle modification with
32
33 heavy alcohol drinkers: Effects of aerobic exercise and meditation. *Addictive*
34
35 *Behaviors*, 11(2), 175-186, doi: 10.1016/0306-4603(86)90043-2.
- 36
37 Orgad, S. (2005). Moving from online to offline relationships with research participants,
38
39 in C. Hine (ed.) *Virtual Methods: Issues in Social Research on the Internet* (pp.
40
41 51-65). Oxford: Berg
42
- 43
44 Pawlikowski, M. & Brand, M. (2011). Excessive internet gaming and decision making:
45
46 Do excessive world of warcraft players have problems in decision making under
47
48 risky conditions? *Psychiatry Research*, 188(3), 428-433. doi:
49
50 10.1016/j.psychres.2011.05.017.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Pontes, H. M. & Griffiths, M. D. (2015). Measuring DSM-5 internet gaming disorder:
5
6 Development and validation of a short psychometric scale. *Computers in Human*
7
8 *Behavior*, 45(1), 137-143. doi: 10.1016/j.chb.2014.12.006
9
- 10 Preacher, K. J., Rucker, D. D., & Hayes, A. F. (2007). Addressing moderated mediation
11
12 hypotheses: Theory, methods, and prescriptions. *Multivariate Behavioral*
13
14 *Research*, 42(1), 185-227.
15
- 16 Ratan, R. R., & Dawson, M. (2015). When Mii is me: A psychophysiological
17
18 examination of avatar self-relevance. *Communication Research*, 1(1), 1-30. doi:
19
20 10.1177/0093650215570652.
21
22
- 23 Ratan, R., & Hasler, B. S. (2010). Exploring self-presence in collaborative virtual
24
25 teams. *PsychNology Journal*, 8(1), 11-31.
26
- 27 Read, J. P., Brown, R. A., Marcus, B. H., Kahler, C. W., Ramsey, S. E., Dubreuil, M.
28
29 E., Jakicic, J. M., & Francione, C. (2001). Exercise attitudes and behaviors among
30
31 persons in treatment for alcohol use disorders. *Journal of Substance Abuse*
32
33 *Treatment*, 21(4), 199-206. doi: 10.1016/S0740-5472(01)00203-3
34
35
- 36 Richardson, R. C., Faulkner, G., McDevitt, J., Skrinar, G. S., Hutchinson, D. S. &
37
38 Piette, J. D. (2005). Integrating physical activity into mental health services for
39
40 persons with serious mental illness, *Psychiatric Services*, 56(3), 324-331. doi:
41
42 10.1176/appi.ps.56.3.324
43
44
- 45 Riva, G., Waterworth, J. A., & Waterworth, E. L. (2004). The layers of presence: A bio-
46
47 cultural approach to understanding presence in natural and mediated
48
49 environments. *CyberPsychology & Behavior*, 7(4), 402-416. doi:
50
51 10.1089/cpb.2004.7.402.
52
53
54
55
56
57
58
59

- 1
2
3
4 Substance Abuse and Mental health Services Administration (2015). *Substance use*
5 *disorders*. Retrieved March 3, 2016, from
6 <http://www.samhsa.gov/disorders/substance-use>
7
8
9
10 Sanchis-Segura, C., & Spanagel, R. (2006). Behavioural assessment of drug
11 reinforcement and addictive features in rodents: an overview. *Addiction Biology*,
12 *11*(1), 2-38. doi: 10.1111/j.1355-6215.2006.00012.x
13
14
15 Schmidt, K., Beck, R., Rivkin, W., & Diestel, S. (2016). Self-control demands at work
16 and psychological strain: The moderating role of physical fitness. *International*
17 *Journal of Stress Management*, *23*(3), 255-275. doi: 10.1037/str0000012
18
19
20 Sliter, K. A., Sinclari, R., Cheung, J., & McFadden, A. (2014) Initial evidence for the
21 buffering effect of physical activity on the relationship between workplace
22 stressors and individual outcomes. *International Journal of Stress Management*,
23 *21*(4), 348-360. doi: 10.1037/a0038110
24
25
26 Sliter, M., & Yuan, Z. (2015). Workout at work: Laboratory test of psychological and
27 performance outcomes of active workstations. *Journal of Occupational Health*
28 *Psychology*, *20*(2), 259-271. doi:10.1037/a0038175
29
30
31 Smahel, D., Blinka, L., & Brown, B. B. (2012). Associations between online friendship
32 and internet addiction among adolescents and emerging Adults. *Development*
33 *Psychology*, *48*(2), 381-388. doi: 10.1037/a0027025.
34
35
36 Smahel, D., Blinka, L., & Ledabyle, O. (2008). Playing MMORPGs: Connections
37 between addiction and identifying with a character. *CyberPsychology & Behavior*,
38 *11*(6). 715-718. doi: 10.1089/cpb.2007.0210.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Snodgrass, J. G., Lacy, M. G., Dengah, F. H. J., & Fagan, J. (2011). Enhancing one life
5 rather than living two: Playing MMOs with offline friends. *Computers in Human*
6 *Behavior*, 27(1), 1211-1222. doi: 10.1016/j.chb.2011.01.001
7
8
9
10 Spekman, M. L. C., Konijn, E.A., Roelofsma, P. H. M. P., & Griffiths, M. D. (2013).
11 Gaming addiction, definition and measurement: A large-scale empirical study.
12 *Computers in Human Behavior*, 29(6), 2150-2155. doi: 10.1016/j.chb.2013.05.015
13
14
15 Stanley, P. (2009). The risk and resilience framework and its implications for teachers
16 and schools. *Waikato Journal of Education*, 14(1), 139-153.
17
18
19 Stavropoulos, V., Kuss, D.J., Griffiths, M.D., Wilson, P. & Motti-Stefanidi, F. (2017a).
20 MMORPG gaming and hostility predict internet addiction symptoms in
21 adolescents: an empirical multilevel longitudinal study. *Addictive Behaviors*, 64,
22 294-300. doi: 10.1016/j.addbeh.2015.09.001
23
24
25
26
27
28
29
30 Stavropoulos, V., Gomez, R., Steen, E., Beard, C., Liew, L., & Griffiths, M. D. (2017b).
31 The longitudinal association between anxiety and Internet addiction in
32 adolescence: The moderating effect of classroom extraversion. *Journal of*
33 *Behavioral Addictions*, 6, 237-247. doi:10.1556/2006.6.2017.026
34
35
36
37
38
39 Steinkuehler, C., & Williams, D. (2006). Where everybody knows your (screen) name:
40 Online games as “third places”. *Journal of Computer-Mediated Communication*,
41 11(4), 885-909. doi: 10.1111/j.1083-6101.2006.00300.x
42
43
44
45 Taylor, J. (1987). Predicting athletic performance with self-confidence and somatic and
46 cognitive anxiety as a function of motor and physiological requirements in six
47 sports. *Journal of Personality*, 55(1), 139-153. doi: 10.1111/j.1467-
48 6494.1987.tb00432.x
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 Thipse, L. (2016). *5 points why video games are so popular across people of different*
5 *age groups*. Retrieved October 1, 2016, from
6 [https://www.linkedin.com/pulse/20141021074940-52500864-5-points-why-video-](https://www.linkedin.com/pulse/20141021074940-52500864-5-points-why-video-games-are-so-popular-across-people-of-different-age-groups)
7 [games-are-so-popular-across-people-of-different-age-groups](https://www.linkedin.com/pulse/20141021074940-52500864-5-points-why-video-games-are-so-popular-across-people-of-different-age-groups)
8
9
10
11
12
13 Tisseron, S. (2009). Jeux video: Entre nouvelle culture et séductions de la (dyade
14 numérique). *Psychiatre et Psychanalyste*, 15(1), 21-40. doi:
15 10.3917/psyt.151.0021
16
17
18
19
20
21 Torsheim, T., Eriksson, L., Schnohr, C. W., Hansen, F., Bjarnason, T., & Välimaa, R.
22 (2010). Screen-based activities and physical complaints among adolescents from
23 the Nordic countries. *BMC Public Health*, 10(324), 1-8. doi:10.1186/1471-2458-
24 10-324
25
26
27
28
29
30 Tremblay, M. S., LeBlanc, A. G., Kho, M. E., Saunders, T. J., Larouche, R., Colley, R.
31 C., Goldfield, G., & Gorber, S. C. (2011). Systematic review of sedentary
32 behaviour and health indicators in school-aged children and youth. *International*
33 *Journal of Behavioural Nutrition and Physical Activity*, 8(1), 98. doi:
34 10.1186/1479-5868-8-98
35
36
37
38
39
40
41 Weinstock, J., Barry, D., & Petry, N. M. (2008). Exercise-related activities are
42 associated with positive outcome in contingency management treatment for
43 substance use disorders. *Addictive Behaviors*, 33(8), 1072-1075. doi:
44 10.1016/j.addbeh.2008.03.011
45
46
47
48
49
50 Wilson, B., & Atkinson, M. (2005). Rave and straightedge, the virtual and the real:
51 Exploring online and offline experiences in canadian youth subcultures. *Youth and*
52 *Society*, 36(3), 276-311. doi: 10.1177/0044118X03260498
53
54
55
56
57
58
59
60

1
2
3
4 Williams, D., Yee, N., & Caplan, S. E. (2008). Who plays, how much, and why?

5
6 Debunking the stereotypical gamer profile. *Journal of Computer-Mediated*
7
8 *Communication*, 13(4), 993-1018. doi: 10.1111/j.1083-6101.2008.00428.x
9

10 Winer, E. S., Cervone, D., Bryant, J., McKinney, C., Liu, R. T., & Nadorff, M. R.

11
12 (2016). Distinguishing mediational models and analyses in clinical psychology:
13
14 Atemporal associations do not imply causation. *Journal of Clinical Psychology*,
15
16 72(9), 947-955. doi: 10.1002/jclp.22298
17
18

19 Winkler, A., Dorsing, B., Rief, W., Shen, Y., & Glombiewski, J. A. (2013). Treatment

20
21 of internet addiction: a meta-analysis. *Clinical Psychology Review*, 33(1), 317-
22
23 329. doi: 10.1016/j.cpr.2012.12.005.
24

25 Woodman, T. & Hardy, L. (2003). The relative impact of cognitive anxiety and self-

26
27 confidence upon sport performance: a meta-analysis. *Journal of Sports Sciences*,
28
29 21(6), 443-457. doi: 10.1080/0264041031000101809
30
31

32 Yee, N. (2006). Motivations for play in online games. *CyberPsychology & Behavior*,

33
34 9(6), 772-775.
35

36 Yee, N., & Bailenson, J. (2007). The Proteus Effect: The effect of transformed self-

37
38 representation on behavior. *Human Communication Research*, 33(1), 271-290.
39
40 doi: 10.1111/j.1468-2958.2007.00299.x.
41
42

43 Yee, N., Bailenson, J.N., & Ducheneaut, N. (2009). The Proteus Effect: Implications of

44
45 transformed digital self-representation on online and offline behavior,
46
47 *Communication Research*, 36(2), 285-312. doi: 10.1177/0093650208330254.
48


49 Young, K. (2009). Understanding online gaming addiction and treatment issues for

50
51 adolescents. *American Journal of Family Therapy*, 37(5), 355-372. doi:
52
53 10.1080/01926180902942191.
54
55
56
57
58
59
60

6. Footnotes

Note 1: The present study is part of a wider project of Federation University Australia that addresses the interplay between individual, Internet and proximal context factors in the development of Internet Gaming Disorder symptoms among emerging adults. Instruments used in the data include the: (1) Internet Gaming Disorder 9- Short Form (Pontes & Griffiths, 2015); (2) Beck Depression Inventory – 2nd edition (21 items) (Beck, Steer, & Brown, 1996); (3) Beck Anxiety Inventory (21 items) (Beck & Steer, 1990); (4) Hikikomori-Social Withdrawal Scale (5 items) (Teo et al., 2015); (5) Attention Deficit Hyperactivity Self-Report Scale (18 items) (Kessler et al., 2005); (6); Ten Item Personality Inventory (Gosling, Rentfrow, & Swann, 2003); (7) The Balanced Family Cohesion Scale (7 items) (Olson, 2000); (8) Presence Questionnaire (10 items) (Faiola, Newlon, Pfaff, & Smyslova, 2013); (9); Online Flow Questionnaire (5 items) (Chen, Wigand, & Nilan, 2000) (10) Self-Presence Questionnaire (Ratan & Hasler, 2010); (11) The Gaming-Contingent Self-Worth Scale (12 items) (Beard, & Wickham, 2016) and; (12) Demographic and Internet Use Questions. The battery of questionnaires was utilized for both online and face-to-face data collection. The use of the fitness tracker (Fitbit flex) was used only for face-to-face data collection. **It is noted that different parts of the present data has been used in two more published studies to address different research questions/hypotheses (Burleigh, Stavropoulos, Liew, Adams & Griffiths, 2017; Adams, Stavropoulos, Burleigh, Liew, Beard & Griffiths, 2018).** *Note 2:* Statistical power was computed separately for the cross-sectional and the longitudinal sample. A probability error of .05, power (1- β error probability) of .95 and effect size of .15. was used for the cross-sectional sample (N=125) resulting to an observed statistical power of $\text{Power}\alpha\text{-}\beta = .99$ ($F_{(122)}=3.92, \lambda=18.75$). A probability error of .05, power (1- β error probability) of .95 and effect size of .25 was used for the longitudinal sample (N = 61) resulting to an observed statistical power of $\text{Power}\alpha\text{-}\beta = .73$ ($\alpha_{\text{error probability}} = .27, \beta_{\text{error probability}} = .27, F_{(2,61)} = 1.34, \text{Pillai } V = .06, \lambda = 3.81$).

Note 3: In line with the approval received by the ethics committee of Federation University, the flyers: a) indicated that participants were required to participate on three separate measurement occasions approximately one month apart; b) included an email address to contact the investigators; and c) clearly described the process and stages of the data collection (face-to-face and online). MMO players, aged between 18-29 years old, interested in the study received the Plain Language Information Statement (PLIS). The PLIS clearly indicated that participation was voluntary and that participants could independently decide to withdraw from the study at any point. Individuals who choose to participate were required to provide informed consent.



27 *Figure 1. Risk and Resilience IGD Framework*

28
29 **Figure 1. Risk and Resilience IGD Framework**

30 190x128mm (120 x 120 DPI)

Table 1. Sociodemographic Characteristics of the Participants in the Study

Sociodemographic Variables		Face-to-Face (n = 61)	Online (n = 64)	Total (n = 125)
Gender	Male	45 (36.0%)	49 (39.2%)	94 (75.2%)
	Female	16 (12.8%)	15 (12.0%)	31 (24.8%)
Game Genre	MMOs	38 (30.4%)	43 (34.4%)	81 (64.8%)
	MMORPGs	23 (18.4%)	21 (16.8%)	44 (35.2%)
Highest level of Education	Year 7-10	4 (3.2%)	2 (1.6%)	6 (4.8%)
	Year 12	25 (20.0%)	21 (16.8%)	46 (36.8%)
	Tertiary Diploma	15 (12.3%)	17 (13.9%)	32 (26.2%)
	Undergraduate Degree	12 (9.8%)	15 (12.3%)	27 (22.1%)
Employment Status	Postgraduate Degree	5 (4.0%)	9 (7.2%)	14 (11.2%)
	Unemployed	1 (0.8%)	11 (8.8%)	12 (9.6%)
	Temporary Leave	2 (1.6%)	2 (1.6%)	4 (3.2%)
	Student	14 (11.2%)	10 (8.0%)	24 (19.2%)
	Casual Employment	15 (12.0%)	8 (6.4%)	23 (18.4%)
	Part-Time Employment	9 (7.2%)	8 (6.4%)	17 (13.6%)
Residing with	Full-Time Employment	20 (16.0%)	25 (20.0%)	45 (36.0%)
	Family of origin (two parents and siblings if any)	11 (9.1%)	23 (19.0%)	34 (28.1%)
	Mother and siblings if any (parents divorced/separated)	2 (1.7%)	5 (4.1%)	7 (5.8%)
	Mother and siblings if any (father passed away)	2 (1.6%)	1 (0.8%)	3 (2.4%)
	Father and siblings if any (parents divorced/separated)	0 (0.0%)	1 (0.8%)	1 (0.8%)
	With partner	18 (14.9%)	16 (13.2%)	34 (28.1%)
	With partner and siblings	1(0.8%)	1 (0.8%)	2 (1.7%)
	Alone	1 (0.8%)	3 (2.5%)	4 (3.3%)
With friends	13 (10.4%)	6 (4.8%)	19 (15.2%)	
Shared accommodation	13 (10.4%)	8 (6.4%)	21 (16.8%)	




Figure 2. Visual Representation of the Present Study

776x599mm (72 x 72 DPI)

Appendix A1. Descriptive statistics and reliability coefficients per item and questionnaire/ measurement used

Time point (TP)	Data collection type		Minimum	Maximum	Mean	SD	<i>Cronbach's α if Item removed</i>
TP1	Face-to-Face	IGD1	1.00	5.00	3.18	1.09	.88
		IGD2	1.00	5.00	2.20	1.13	.86
		IGD3	1.00	5.00	2.07	1.04	.86
		IGD4	1.00	5.00	1.83	1.07	.86
		IGD5	1.00	5.00	1.97	1.07	.88
		IGD6	1.00	5.00	1.95	1.19	.87
		IGD7	1.00	5.00	1.75	1.13	.85
		IGD8	1.00	5.00	3.02	1.23	.88
		IGD9	1.00	4.00	1.52	.98	.98
TP1 $N = 61$; TP1 Cronbach's $\alpha = .88$; TP1 Cronbach's α Based on Standardized Items = .88							
TP2	Face-to-Face	IGD1	1.00	5.00	2.92	1.11	.87
		IGD2	1.00	5.00	2.12	1.12	.86
		IGD3	1.00	5.00	2.02	.97	.85
		IGD4	1.00	4.00	1.84	.89	.87
		IGD5	1.00	5.00	1.95	1.14	.88
		IGD6	1.00	5.00	1.97	1.22	.86

IGD7	1.00	5.00	1.59	1.01	.86
IGD8	1.00	5.00	2.86	1.25	.87
IGD9	1.00	4.00	1.52	.88	.88

TP2 $N=56$; TP2 Cronbach's $\alpha=.88$; TP2 Cronbach's α Based on Standardized items= .88

TP3	Face-to-Face	IGD1	1.00	5.00	2.92	1.08	.83
		IGD2	1.00	5.00	2.03	1.11	.82
		IGD3	1.00	4.00	2.11	1.04	.81
		IGD4	1.00	5.00	1.74	.91	.82
		IGD5	1.00	5.00	2.01	.94	.83
		IGD6	1.00	5.00	1.87	1.05	.82
		IGD7	1.00	4.00	1.44	.74	.84
		IGD8	1.00	5.00	2.93	1.32	.81
		IGD9	1.00	4.00	1.39	.75	.84

TP3 $N=43$; TP3 Cronbach's $\alpha=.84$; TP3 Cronbach's α Based on Standardized items= .84

TP1	Face-to-Face	PSP1	.00	4.00	1.13	1.32	.88
		PSP2	.00	4.00	.72	.99	.87
		PSP3	.00	3.00	.92	1.09	.89
		PSP4	.00	4.00	.52	.99	.88

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

		PSP5	.00	4.00	.52	.89	.89
	TP1 <i>N</i> = 61; TP1 Cronbach's α = .90; TP1 Cronbach's α Based on Standardized items = .91						
TP2	Face-to-Face	PSP1	.00	4.00	.83	1.16	.91
		PSP2	.00	4.00	.72	1.11	.89
		PSP3	.00	3.00	.87	1.08	.92
		PSP4	.00	4.00	.50	1.01	.91
		PSP5	.00	4.00	.58	1.00	.91
	TP2 <i>N</i> = 56; TP2 Cronbach's α = .93; TP2 Cronbach's α Based on Standardized items = .93						
TP3	Face-to-Face	PSP1	.00	4.00	.86	1.16	.88
		PSP2	.00	4.00	.72	1.10	.89
		PSP3	.00	4.00	.81	1.06	.90
		PSP4	.00	4.00	.57	1.01	.88
		PSP5	.00	4.00	.74	1.03	.90
	TP3 <i>N</i> = 43; TP3 Cronbach's α = .91; TP3 Cronbach's α Based on Standardized items = .91						
TP1		Active minutes	.00	263.00	71.03	71.03	
TP2		Active minutes	.00	258.33	80.21	80.21	
TP3		Active minutes	.00	259.00	75.98	56.66	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

TP1	Online	IGD1	1.00	5.00	3.16	1.10	.94
		IGD2	1.00	5.00	2.15	1.19	.92
		IGD3	1.00	4.00	2.18	1.22	.92
		IGD4	1.00	5.00	2.00	1.18	.92
		IGD5	1.00	5.00	2.16	1.12	.93
		IGD6	1.00	5.00	2.17	1.42	.92
		IGD7	1.00	4.00	1.64	1.21	.92
		IGD8	1.00	5.00	2.84	1.49	.94
		IGD9	1.00	4.00	1.60	1.12	.92

20 TP1 Online $N = 64$; TP1 Online Cronbach's $\alpha = .93$; TP1 Online Cronbach's α Based on Standardized
21 items= .94
22

23
24
25
26
27
28
29
30
31
32

T1	Online	PSP1	.00	4.00	1.10	1.30	.91
		PSP2	.00	4.00	.56	1.01	.88
		PSP3	.00	4.00	.87	1.26	.89
		PSP4	.00	4.00	.37	.93	.88
		PSP5	.00	4.00	.32	.89	.91

33 T1 $N = 64$; T1 Cronbach's $\alpha = .91$; T1 Cronbach's α Based on Standardized items=.92
34
35
36

37
38
39
40
41
42
43
44
45
46
47

Table 2.PSP predicting IGD (Cross-section)

Variable	R^2	b	SE	p
Step 1	.05			
Age		-0.49*	0.23	-.190
Gender ^a		-2.18	1.72	-.110
Step 2	.20			
PSP		.81	.14	.001***
R^2	.25***			

Note, ^a0 = females, 1=males

* $p < .05$. ** $p < .01$. *** $p < .001$

Table 3.PSP predicting IGD (Longitudinal)

Variable	R^2	b	SE	p
Step 1	.006			
Age		-0.10	0.26	-0.050
Gender ^a		-0.78	1.83	-0.060
Step 2	.36			
PSP		0.76	0.14	.001***
R^2	.37**			

Note, ^a0 = females, 1=males

* $p < .05$. ** $p < .01$. *** $p < .001$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3. Moderation summary of PA (Active Minutes) on the relationship between Proto-self and IGD

141x81mm (96 x 96 DPI)

Appendix A2. Descriptive statistics of the aggregate study variables for offline and online data sets

Time point (TP)	Data collection type		Minimum	Maximum	Mean	SD
TP1	Face-to-Face	IGDT1	10.00	43.00	19.44	7.07
	TP1 <i>N</i> = 61					
TP2		IGDT2	9.00	38.00	18.80	6.91
	TP2 <i>N</i> = 56					
TP3		IGDT3	9.00	33.00	17.95	5.93
	TP3 <i>N</i> = 43					
TP1	Face-to-Face	PSPT1	.00	19.00	3.75	4.52
	TP1 <i>N</i> = 61					
TP2		PSPT2	.00	18.00	3.20	4.55
	TP2 <i>N</i> = 56					
TP3		PSPT3	.00	20.00	3.27	4.69
	TP3 <i>N</i> = 43					
TP1	Face-to-Face	ActiveMinT1	.00	263.00	71.03	66.77
	TP1 <i>N</i> = 61					
TP2		ActiveMinT2	.00	258.33	80.21	69.73
	TP2 <i>N</i> = 56					

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

TP3		ActiveMinT3	.00	259.00	75.98	56.66
	TP3 N = 43					
TP1	Online	IGDT1	9.00	45.00	19.70	8.91
	T1 N = 64					
TP1	Online	PSPT1	.00	20.00	3.26	4.72
	T1 N = 64					
