

1 **Effects of PAHs and dioxins on the earthworm *Eisenia andrei*: a**
2 **multivariate approach for biomarker interpretation**

3

4

5 Susanna Sforzini ^a, Michael N. Moore ^{a,b,c,d}, Marta Boeri ^a, Mauro Bencivenga^e, Aldo Viarengo ^a

6

7 ^a Department of Sciences and Technological Innovation (DiSIT), University of Piemonte Orientale

8 "A. Avogadro", V.le T. Michel 11, 15121 Alessandria, Italy

9 ^b European Centre for Environment & Human Health (ECEHH), University of Exeter Medical

10 School, Truro, TR1 3HD, UK

11 ^c Plymouth Marine Laboratory, Plymouth, PL1 3DH, UK

12 ^d School of Biological Science, University of Plymouth, Plymouth PL4 8AA, UK

13 ^e ISPRA, Via Vitaliano Brancati 60 Roma

14

15

16 Corresponding author:

17 Aldo Viarengo

18 Department of Sciences and Technological Innovation (DiSIT), University of Piemonte Orientale

19 "A. Avogadro", V.le T. Michel 11, 15121 Alessandria, Italy

20 Phone: + 390131360370

21 Fax: +390131360390

22 e-mail: viarengo@unipmn.it

23

24 **Abstract**

25 In this study, a battery of biomarkers was utilised to evaluate the stress syndrome induced in the
26 earthworm *Eisenia andrei* by exposure to environmentally realistic concentrations of
27 benzo[*a*]pyrene (B[*a*]P) and 2,3,7,8-tetrachlorodibenzo-*para*-dioxin (TCDD) in OECD soil. The set
28 of tests was then employed to assess the toxicity of field soils contaminated with organic xenobiotic
29 compounds (such as PAHs, dioxins and PCBs). The results highlighted an impairment of immune
30 and metabolic functions and genotoxic damage in worms exposed also to lower bioavailable
31 concentrations of toxic chemicals. Multivariate analysis of biomarker data showed that all different
32 contaminated soils had a detrimental effect on the earthworms. A separation between temporal and
33 concentration factors was also evident for B[*a*]P and TCDD treatments; and field contaminated
34 soils were further differentiated reflecting a diverse contamination. Multivariate analysis also
35 demonstrated that lysosomal membrane stability can be considered a prognostic indicator for worm
36 health status.

37

38 Capsule: Biomarkers were employed in *E. andrei* in laboratory and field studies. Multivariate
39 analysis ranked the stress syndrome in worms. Lysosomal stability is prognostic for health status.

40

41

42 Keywords: earthworms; lysosomal membrane stability; biomarkers; multivariate analysis

43

44 **1. Introduction**

45 Biomarkers are sensitive tools for detecting exposure and adverse effects of toxic chemicals both on
46 aquatic and terrestrial organisms (Moore et al., 2004; Peakall, 1994; Scott-Fordsmand and Weeks,
47 2000). The study of biological parameters at different level of functional complexity in diverse cells
48 and tissues of the organisms and with different meaning (i.e. biomarkers of stress, exposure and
49 genotoxicity) is useful to clarify the mechanisms of action of chemicals as well as to determine the
50 level of pollutant-induced stress syndrome in animals exposed to environmental matrices, where a
51 mixture of many different contaminants may be present (Asensio et al., 2013; Binelli et al., 2010;
52 Cajaraville et al., 2000; Sforzini et al., 2011; Turja et al., 2014; van der Oost et al., 2003; Viarengo
53 et al., 2007a, 2007b). However, this approach may result in an unclear correlation structure of the
54 data; an aspect that could be more evident analyzing results obtained in field studies than from
55 laboratory experiments, using known doses of a single substance. In addressing this problem,
56 previous studies increasingly suggest that multivariate analysis techniques are a useful tool for
57 interpreting multiple biomarker responses as they produce a two-dimensional pattern of the degree
58 of similarity between different groups of data (Allen and Moore, 2004; Astley et al., 1999; Bernet et
59 al., 2000; Brenner et al., 2014; Burgos et al., 2005; Galloway et al., 2004; Garmendia et al., 2011;
60 Sanchez et al., 2007). The use of multivariate analyses to identify prognostic biomarkers, useful to
61 provide better risk assessment at the early stages, is also of undoubted importance (Jenkins et al.,
62 2011; Moore et al., 2006; Ortiz et al., 2011).

63 In the last decades, earthworms acquired a position of growing importance in terrestrial
64 ecotoxicology. These animals have been shown to accumulate and respond to contaminants at
65 various levels of complexity ranging from the whole animal to the most sensitive molecular/cellular
66 changes (Caselli et al., 2006; Dimitrova et al., 2010; Hayashi et al., 2013; Santoyo et al., 2011;
67 Sforzini et al., 2012; Spurgeon and Hopkin, 1999; Ville et al., 1995; Yang et al., 2012).

68 In this study, a battery of biomarkers was utilised to evaluate the alterations in the health status
69 induced in the earthworm *Eisenia andrei* (Bouché, 1972) by exposure to environmentally relevant
70 concentrations of benzo[a]pyrene (B[a]P) and 2,3,7,8-tetrachlorodibenzo-*para*-dioxin (TCDD) in
71 OECD soil (OECD, 2004). These hazardous environmental chemicals, classified by the US EPA
72 (Environmental Protection Agency) as priority pollutants, are among the frequently occurring soils
73 contaminants (Martínez et al., 2006; USEPA, 2009; Wang et al., 2007). Stress responses at the
74 various levels of biological organisation (including lysosomal membrane stability, lipofuscin and
75 neutral lipid accumulation and tissue damage) were measured; the genotoxic effects caused by these
76 substances on worms, in terms of both DNA and chromosomal damage, have previously been
77 demonstrated (Sforzini et al., 2012). To verify the robustness of the selected multi-biomarkers in *E.*
78 *andrei* to be used for the assessment of polluted natural soils, the set of tests was then applied in a
79 field study conducted to evaluate the potential toxicity of soils contaminated by organic xenobiotic
80 compounds (such as PAHs, dioxins, PCBs), as a consequence of different anthropogenic activities.
81 Multiple biomarker responses in worms exposed to different chemical treatments and field
82 contaminated soils were analysed by multivariate statistics in order to identify any discernable
83 similarities or dissimilarities in multidimensional biomarker response patterns.
84 An overarching objective was the use of predictive models of lysosomal and other biomarker
85 reactions as both diagnostic and prognostic biomarkers for health status in the earthworms. For this
86 purpose, multivariate analysis has been previously used to develop statistical models to study the
87 role of lysosomal functions and responses to environmental variables, particularly chemical
88 pollutants (Allen & Moore, 2004; Moore et al., 2006).

89

90 **2. Materials and methods**

91 *2.1. Chemicals*

92 All chemicals were of analytical grade and purchased from Sigma-Aldrich Co. (St. Louis, MO,
93 USA), unless otherwise indicated.

94

95 *2.2. Animals*

96 Earthworms were cultured essentially as described in the OECD guideline (OECD, 2004).

97 Organisms were selected from a synchronised culture with an homogeneous age structure. Adult
98 worms with clitellum of similar size and weight (of 400 to 500 mg) were utilised in the
99 experiments.

100

101 *2.3. Artificial soil test*

102 The artificial soil tests were performed as described in the OECD guideline for the testing of
103 chemicals (OECD, 2004). The tests were performed under controlled conditions for a period of 10
104 and 28 d. The selected B[a]P and TCDD concentrations were i) beginning with the lowest, the
105 Italian law limit for residential areas, ii) for industrial areas, iii) five (B[a]P) and twenty (TCDD)
106 times higher than the latter (for details see Sforzini et al., 2012). Vitality and reproduction were
107 assessed at the end of the experiments.

108

109 *2.4. Field soils*

110 Five soils contaminated by organic xenobiotic compounds (i.e. PAHs, dioxins and PCBs) were
111 collected from different areas in the Campania region (Italy): Site 1 (4536773/446660) is located
112 close to the site of construction of an incinerator; Sites 2 and 3 (4538083/442802 and
113 4531906/424656, respectively) are industrial areas; Sites 4 and 5 (4540699/443848 and
114 4535670/440538, respectively) are areas subjected to uncontrolled waste fires. **The chemical
115 analysis of these samples were performed by ISPRA (...) and are a part of a larger study
116 published by Italian Ministry of the Environment (ISPRA, 2012).** As shown in Table 1, the

117 concentrations of the chemicals in the soils, in only a few cases, were higher than the limits for
118 residential areas set by Italian law, but always lower than the limits for industrial areas (ISPRA,
119 2012). A control field soil was collected from a site with no detectable soil contamination
120 (Reference site - 4569424/470553). Particle-size distribution, organic matter content and pH of soils
121 from the different Sites were homogeneous (ISPRA, 2012).
122 Soil samples were collected from the top 0-5 cm layer at each site after removal of surface
123 vegetation and litter. Each of these samples represented a composite of five subsamples from the
124 center and four corners of a square sampling grid. Soil subsamples at each location were
125 homogenized and transferred to clean containers. Soils were dried, sieved through a 2 mm mesh,
126 moistened with deionized water and then, for each soil replicate, ten worms were kept in 500 g of
127 soil placed in glass test containers. At least five replicates per soil were used. The test containers
128 were maintained in a climatized chamber with a temperature of 20 ± 1 °C. The test was performed
129 under controlled light-dark cycles (16 h light, 8 h dark) with illumination of 800 lx and for a period
130 of 10 days. Vitality was assessed at the end of the incubation.

131

132 *2.5. Biomarker tests*

133 A battery of biomarkers was used to evaluate the harmful effects induced in worms by increasing
134 concentrations of B[a]P, TCDD and field contaminated soils.

135 **Coelomocytes used for the determination of lysosomal membrane stability (LMS) and MNI**
136 **frequency were obtained by a non-invasive extrusion method i.e. ethanol extraction (Eyambe**
137 **et al., 1991; Fugère et al., 1996; Sforzini et al., 2012). Briefly, earthworms were rinsed in saline**
138 **(0.85 mg/ml NaCl at 4 °C) and the posterior portion was massaged to expel the gut contents of**
139 **the terminal part of the intestine. Then, animals were placed in 4 ml of cold extrusion medium**
140 **containing 5% ethanol, 2.5 mg/ml EDTA, 10 mg/ml of the mucolytic agent guaiacol glycerol**
141 **ether (pH 7.3). After 3 min, the earthworms were removed and 2 ml of Hanks' balanced salt**

142 **solution (HBSS) (Sigma product H6648) were added. The cell suspension was centrifuged at**
143 **150 × g at 4 °C for 2 min to remove mucus, and then for 10 min to recover the cells.**
144 **Chloragogenous tissue was used for the determination of LMS, lipofuscin and neutral lipid**
145 **lysosomal content and lysosomal/cytoplasm volume ratio. At the end of the incubation, worms**
146 **were washed and then kept on clean moist filter paper for 24 h to allow them to void their gut**
147 **contents Pieces of five earthworms (4-5 mm length, posterior to the clitellum), were placed on**
148 **an aluminium cryostat chuck and chilled in hexane at -70 °C as described previously (Sforzini**
149 **et al., 2014).**

150 LMS of amoeboid coelomocytes was evaluated by the neutral red retention time (NRRT) assay by
151 fluorescence microscopy (Sforzini et al., 2011).

152 The determination of LMS in the cells of chloragogenous tissue was performed on cryostat tissue
153 sections following essentially the method described by Moore (1988). This cytochemical assay is
154 based on acid labilization characteristics of latent hydrolase β -N-acetylhexosaminidase (NAH)
155 using naphthol AS-BI-N-acetyl- β -D glucosaminide as a substrate for NAH.

156 Lipofuscin and neutral lipid lysosomal content in the cells of the chloragogenous tissue as well as
157 the lysosomal/cytoplasmic volume ratio (L/C) of this tissue were evaluated as described by Sforzini
158 et al. (2011).

159 The micronucleus test in coelomocytes of worms exposed to field soils was performed following
160 the methods described in Sforzini et al. (2012).

161

162 *2.6. Univariate statistical analysis*

163 **At least five replicates per control and per treatment were analyzed. For the biomarker data**
164 **obtained in coelomocytes, each replicate consists of cells from two earthworms pooled**
165 **together; the two animals were collected from a separate replicate of soil, consisting of 10**
166 **earthworms incubated in 500 g of soil. For the biomarker data obtained in chloragogenous**

167 **tissue (cryostat sections), pieces of five earthworms, each one collected from a separate**
168 **replicate soil, were analysed.**

169 The non-parametric Mann-Whitney *U*-test was used to compare the data from treated earthworms
170 with those of the controls ones.

171

172 2.7. Multivariate analysis

173 Biomarker data for earthworms exposed to B[a]P, dioxin and field soils from Campania Region
174 were analysed using non-parametric multivariate analysis software, PRIMER v 6 (PRIMER-E Ltd.,
175 Plymouth, UK; Clarke, 1999). All data were log transformed [$\log_n(1+x)$] and standardised to the
176 same scale. Principal component analysis (PCA), hierarchical cluster analysis and non-metric multi-
177 dimensional scaling analysis (MDS), derived from Euclidean distance similarity matrices were used
178 to visualise dissimilarities between sample groups. The results were further tested for significance
179 using analysis of similarity (PRIMER v6 - ANOSIM), which is analogous to a univariate ANOVA
180 and reflects on differences between treatment groups in contrast to differences among replicates
181 within samples (the *R* statistic). Under the null hypothesis H_0 (“no difference between samples”), R
182 = 0 and this was tested by a non-parametric permutations approach; there should be little or no
183 effect on the average *R* value if the labels identifying which replicates belong to which samples are
184 randomly rearranged.

185 The PRIMER v6 - BIO-ENV routine (Spearman’s Rank Correlations, Rho) linking multivariate
186 biomarker response patterns was used to identify “influential biomarkers” - small subsets of
187 biomarkers capturing the full MDS biomarker response pattern.

188 Finally, in order to map integrated biomarker data onto “health status space” by using LMS; first
189 principal components (PC1) for the biomarker data were derived using PRIMER v6 and then
190 plotted against the LMS (as a measure of cellular well-being) values for each treatment/field sample
191 (Allen and Moore, 2004; Moore et al., 2006).

192

193 **3. Results**

194 *3.1. Biomarker responses*

195 The assessment of the different biomarker responses in worms exposed for 10 and 28 d to
196 increasing concentrations of B[a]P and TCDD in OECD soil demonstrated significant changes in
197 treated worms (Fig. 1); at the concentrations used no effect on vitality and reproduction rate was
198 found (data not shown).

199 LMS in coelomocytes of worms exposed for 10 d to both B[a]P and TCDD showed a significant
200 decrease with respect to controls at the two higher concentrations; in particular, very strong effects
201 were observed at 10 ppm B[a]P (-91% with respect to controls) and 0.1 and 2 ppb TCDD (-81%
202 and -99% with respect to controls respectively) (Fig. 1A, B). In animals exposed for 28 d,
203 significant changes were observed at all the concentrations of the two chemicals, with maximal
204 effect at the highest one, i.e. 50 ppm B[a]P and 2 ppb TCDD (-70% and -88% with respect to
205 controls respectively) (Fig. 1A, B).

206 The lysosomal responses in the cells of the chloragogenous tissue, in term of lipofuscin and neutral
207 lipid accumulations, highlighted relevant variations in worms exposed for 28 d to the higher
208 chemical concentrations (Fig. 1C, D and Fig. 1E, F respectively). In particular, a significant
209 increase in lipofuscin content was observed in worm exposed to 0.1 ppb TCDD (+78% with respect
210 to controls); the results of neutral lipid lysosomal content indicated a significant increase at 10 ppm
211 and 50 ppm B[a]P (+53% and +36% with respect to controls respectively), with maximal effect at
212 the highest dioxin concentration (+114% with respect to controls).

213 B[a]P and TCDD also caused an increase in the L/C (Fig. 1G, H). In particular, significant changes
214 were observed in worms exposed for 10 d to 10 ppm and 50 ppm B[a]P and to all the dioxin
215 concentrations; the alteration being particularly relevant at 0.1 and 2 ppb TCDD (+44% and +45%
216 with respect to controls). After 28 d, the values measured were significant at all the doses of both

217 chemicals, with greater effects in animals exposed to 10 and 50 ppm B[a]P (+52% and +64% with
218 respect to controls).

219 The results of biomarkers in worms exposed for 10 d to field soils collected in Campania,
220 highlighted significant alterations at the contaminated sites (Sites 1-5) with respect to reference site
221 (Fig. 2), without resulting in mortality (data not shown). It is important to point out that animals
222 exposed to the reference site soil did not show changes with respect to laboratory controls (data not
223 shown).

224 The results of LMS in the cells of the chloragogenous tissue indicated a significant reduction in
225 worms exposed to soils collected from all the contaminated sites (Site 1-5), with maximal effect at
226 Site 3 (Fig. 2A).

227 In the cells of the same tissue, significant changes were observed also in the other lysosomal
228 biomarkers. In particular, the results of lysosomal accumulation of lipofuscin highlighted a
229 significant increase in lipofuscin content in worms exposed to Sites 2, 4 and 5, with greater effect at
230 Site 2 (+173% with respect to the Reference Site); ~~In~~ In animals exposed to Site 1, the value was
231 significantly lower with respect to the Reference site (-78%) (Fig. 2B). Lysosomal accumulation of
232 neutral lipids showed significant changes only in worms exposed to Sites 2-3, with a slight increase
233 at Site 3 (+23%) and a marked reduction at Site 2 (-66%) with respect to the Reference site (Fig.
234 2C).

235 L/C in the chloragogenous tissue showed a significant increase in worms exposed to soils from all
236 the contaminated sites (Sites 1-5); the effect was greater at Sites 3 and 4 (+44% and +45% with
237 respect to Reference site respectively) (Fig. 2D).

238 The results of MN test in coelomocytes of worms exposed to soils from Sites 1-5 showed a
239 significant increase of chromosomal damage with respect to the Reference site; the MNi frequency
240 was higher at Sites 2 and 4 (7.5 %) (Fig. 2E).

241

242 3.2. Multivariate analysis of biomarker responses

243 Principal component analysis (PCA), multi-dimensional scaling (MDS) and hierarchical cluster
244 analysis of the biomarker responses in worms exposed to B[a]P and TCDD (in addition to the
245 results shown in this investigation, the data ~~of for~~ DNA damage and micronuclei induction obtained
246 in a previous study -Sforzini et al., 2012), showed that **both chemicals had a detrimental effect on**
247 **the earthworms. This effect is indicated by the (delete): there is a clear separation between the**
248 **clusters of for the control groups and the treatment groups. Due to the biological meaning of**
249 **these biomarkers of stress, the results could be interpreted as detrimental effects of both**
250 **chemicals on the earthworms** (Fig. 3). Analysis of similarity shows that these clusters are
251 significantly different (ANOSIM, R Statistic: B[a]P $R = 0.876$, $P < 0.001$; dioxin $R = 0.772$, $P <$
252 0.001). **The PCA analysis was also suitable to highlighted a separation between control worms**
253 **and animals those exposed to the lower chemical concentration. In particular, at 0.1 ppm**
254 **B[a]P after 28 d, the different distribution of the data seems to be due mainly to LMS and L/C**
255 **ratio. In dioxin exposed animals (lower dose after 28 d) the main contributions (separation**
256 **factors) seems to be MNi frequency and L/C ratio with a contribution of LMS.**

257 Multivariate analysis of the biomarker reactions from field soil samples collected in Campania also
258 showed clear differences between the reference and contaminated samples (ANOSIM, $R = 0.815$, P
259 < 0.001). Sites 1 and 5 (primarily PAHs) were clustered together and Site 2 (primarily dioxins and
260 furans) was clearly distinct from all the other samples (Fig. 3); **from this figure and the data**
261 **reported in Fig. 2, it is evident that the separation of Site 2 is mainly due to the huge-large**
262 **difference in LF (positive) and LN (negative) from all the other treatments.**

263 Multiple regression analysis of the biomarker data indicated that some of the biological parameters
264 are correlated (Fig. 4), although there is no consistent pattern of correlations across the various
265 treatments. However, the BIO-ENV routine indicated that the lysosomal parameters were influential
266 biomarkers in all three exposure studies. Combinations of lysosomal biomarkers showing

267 significant capture of the full MDS pattern are shown in Table 2. Experimental exposure of worms
268 to B[a]P and TCDD as well as to Campania soil samples resulted in LMS + lipofuscin (LF) + L/C
269 emerging as the strongest combinations of lysosomal biomarkers (Table 2).

270 By plotting LMS against the first principal component (PC1) of selected biomarker data (Fig. 5), we
271 effectively integrate the multi-biomarker results and the graph reflects the gradient of toxicity
272 between the samples. PC1 is a measure of the contaminant gradient with the left-hand side being the
273 most impacted and the right-hand side the least affected (Fig. 5). The correlation between LMS and
274 PC1 is highly significant for B[a]P and TCDD treatments. In the Campania data, the correlation is
275 also significant, but it is not as strong as in the experimental treatments in spiked soil (Fig.5).

276

277 4. Discussion

278 Hydrophobic organic compounds, such as polycyclic aromatic hydrocarbons (PAHs), dioxins and
279 furans (PCDD/Fs), and polychlorinated biphenyls (PCBs), are common contaminants in soils.
280 These toxic and genotoxic chemicals, which are generally strongly bound to soil particles, tend to
281 persist in the soil and can bioaccumulate, increasing the potential hazard for the ecosystem
282 (Baderna et al., 2013; Valentín et al., 2013).

283 In a previous study, we have demonstrated that the exposure of the earthworm *E. andrei* for 10 and
284 28 d to sublethal concentrations of B[a]P and TCDD has resulted in genotoxic effects (Sforzini et
285 al., 2012). In particular, even the lowest doses of the two chemicals utilised (representing the limit
286 for residential and industrial areas set by Italian law) induced in treated animals both DNA and
287 chromosomal damage. To further investigate the adverse effects caused by these chemicals on
288 worm health status, in this study the responses of well-established biomarkers of stress, namely
289 lysosomal membrane stability (LMS), the lysosomal lipofuscin content, the neutral lipid
290 accumulation in lysosomes (cell-level biomarkers); and the lysosomal/cytoplasmic volume ratio
291 (L/C, biomarker of tissue damage), were evaluated.

292 In B[a]P and TCDD-exposed worms, amoeboid coelomocytes (immunocompetent cells circulating
293 in the coelomic fluid - Adamowicz, 2005; Cooper and Roch, 2003), showed a significant reduction
294 of LMS; the effect was evident at all the different concentrations and at the shorter exposure time. **It**
295 **is interesting that at the medium concentration used for B[a]P and TCDD (10 ppm and 0.1**
296 **ppb, respectively), the strongest effect was observed after 10 d of exposure with a partial**
297 **recovery at 28 d. This effect is ~~of difficult to interpretation;~~ among ~~the~~ possible**
298 **explanations, ~~there are~~ include the biological responses of the animals, ~~also~~ related to the**
299 **turnover time of the coelomic cells in this ~~organisms~~ species of worm.** This type of subcellular
300 pathological reaction represents an extremely sensitive general index of cellular condition, able to
301 reveal the early adverse effects of pollutants in different animal models (Moore et al., 2012). The
302 neutral red retention time (NRRT) assay has been successfully applied for the *in vivo* evaluation of
303 LMS in fish hepatocytes, in the blood cells of a wide range of marine and freshwater invertebrates
304 as well in earthworms coelomocytes (Binelli et al., 2009; Camus et al., 2000; Canesi et al., 2006;
305 Guidi et al., 2010; Hauton et al., 1998; Lowe and Pipe, 1994; Moore et al., 2009; Sforzini et al.,
306 2011; Weeks and Svendsen, 1996; Winzer et al., 2002).

307 Chloragogenous tissue surrounding the intestine, where most of the key metabolic processes occur
308 (Prentø, 1987), also presented significant changes in treated animals. In particular, a significant
309 lysosomal accumulation of lipofuscin (end-products of membrane lipid peroxidation - Terman and
310 Brunk, 2004; Viarengo and Nott, 1993) and of neutral lipids (due to unbalanced fatty acid
311 metabolism - Lüllman-Rauch, 1979) was induced in the chloragocytes of worms exposed for 28 d
312 only to some concentrations. B[a]P and TCDD are known to provoke an increase of the cellular
313 levels of reactive oxygen species (ROS) (Gelboin, 1980; Lin et al., 2007); and there is evidence that
314 these chemicals can have an impact on lipid metabolism (Irigaray et al., 2006; Schiller et al., 1985).
315 However, we need to consider that in the cells of the chloragogenous tissue of *E. andrei* exposed to
316 different kinds of pollutants, accumulation of lipofuscin and neutral lipid ~~accumulation~~ in

317 lysosomes typically shows a bell-shaped trend (Gastaldi et al., 2007; Sforzini et al., 2011). This
318 ~~could~~can be partly explained by an enhanced rate of elimination of lipofuscin/neutral lipid-rich
319 residual bodies into the coelomic fluid as well as by an augmented rate of turnover of
320 chloragocytes, since several studies showed that chloragocyte depletion may occur in worms as a
321 way of eliminating toxic chemicals (Cancio et al., 1995; Fischer and Molnár, 1992). Furthermore,
322 when we also consider the results of the L/C, a biomarker reflecting the level of cellular autophagy
323 (Lowe et al., 1981), there is a clear indication that an impairment of the physiology of the whole
324 tissue was already occurring after 10 d at the lower doses of both chemicals.

325 Overall, the results obtained showed that environmentally realistic concentrations of B[a]P and
326 TCDD in OECD soils, without affecting vitality and reproduction of earthworms, provoked
327 significant alterations in the physiological status of the organisms. Relevant changes at cellular and
328 tissue levels were observed in animals exposed, for only 10 d, to lower contaminant chemical
329 concentrations.

330 To verify the robustness of the selected multi-biomarkers in *E. andrei* to be used for the assessment
331 of polluted natural soils, the set of tests was then applied in a field study conducted to evaluate the
332 potential toxicity of five soils contaminated by organic xenobiotic compounds (such as PAHs,
333 dioxins, PCBs), as a consequence of different anthropogenic activities. Chemical analysis revealed
334 that the concentrations of chemicals in soils were, in only a few cases, higher than the Italian law
335 limits for residential areas.

336 Worms incubated for 10 d in the different contaminated soils showed sublethal stress and genotoxic
337 effects. A significant decrease in LMS in the cells of the chloragogenous tissue (based on acid
338 labilization characteristics of the latent hydrolase β -N-acetylhexosaminidase - Moore, 1988;
339 Peeters-Joris, 2000) was observed at all sites; the alteration was particularly relevant in Site 3
340 (contaminated primarily with PCBs). Initial studies on the toxic effects of contaminants on
341 lysosomes were originally carried out to determine LMS by using histochemical procedures applied

342 on frozen tissue sections of fish liver or mussel hepatopancreas (Köhler, 1991; Moore, 1976, 1990).
343 This methodology is currently used by many researchers in the laboratory as well as in field studies
344 (Banni et al., 2014; Broeg et al., 1999; Domouhtsidou and Dimitriadis, 2001; Franzellitti et al.,
345 2012; Köhler et al., 2002; Roméo et al., 2000; Shaw et al., 2011; Viarengo et al., 1987). Once the
346 tissues are frozen, it is possible to preserve them until the analysis, an aspect that could be important
347 when large-scale biomonitoring studies are performed.

348 PCDD in field soils (Sites 2, 4) at concentrations similar to those spiked in OECD soil induced
349 (differently from that observed in dioxin treatments) a significant increase of lipofuscin content; a
350 minimal, although significant, change was observed also in worms exposed to Site 5 (primarily
351 contaminated with PAHs). The results of lysosomal accumulation of neutral lipids indicated a
352 significant increase in neutral lipid content in Site 3 (primarily with PCBs).

353 The bioavailable contaminants present in the polluted soils also induced in the chloragogenous
354 tissue a significant increase of L/C; the higher tissue alteration being detected Sites 3 and 4
355 (primarily contaminated with PCBs and PCDD-PCDF, respectively).

356 Finally, all contaminated soils resulted genotoxic, as indicated by the MN test showing a significant
357 increase in MNi frequency in coelomocytes of exposed worms. The MN test has emerged as an
358 alternative approach to the classical techniques for assessing cytogenetic damage (Schmid, 1975).
359 The test procedure is technically easier and faster than the analysis of chromosomal aberrations
360 during metaphase. The MN test has been widely applied in human and other mammalian cell types,
361 amphibians, fish and molluscs (Bolognesi and Hayashi, 2011; Fenech, 2000); and recently this
362 method has also been developed on *E. andrei* coelomocytes (Sforzini et al., 2012).

363 Taken together, the results obtained in the field study confirm that LMS and L/C as well as MNi
364 frequency are sensitive biomarkers able to highlight the stress syndrome induced in worms by
365 exposure to bioavailable chemicals present in soils. The evaluation of lipofuscin and neutral lipid
366 lysosomal accumulation (biomarkers widely used in monitoring studies, easy to perform and not

367 expensive) is important in order to highlight additional relevant adverse effects on animal health.
368 However, these parameters in worms have to be used together with LMS and L/C ratio (showing a
369 clear response profile over a stress gradient) to avoid a possible misinterpretation when negative
370 results are obtained.

371 Procedures capable of integrating and interpreting the biomarker responses within synthetic stress
372 indices have been developed (Beliaeff and Burgeot, 2002; Dagnino et al., 2007; Moore et al., 2004;
373 Narbonne et al., 1999; Sforzini et al., 2011; Viarengo et al., 2000). An addition useful tool for
374 interpreting multiple biomarker responses is multivariate statistics as they produce a two-
375 dimensional pattern of the degree of similarity between different groups of data (Allen and Moore,
376 2004; Astley et al., 1999; Bernet et al., 2000; Brenner et al., 2014; Burgos et al., 2005; Galloway et
377 al., 2004; Garmendia et al., 2011; Sanchez et al., 2007).

378 Multivariate analysis indicated that the battery of biomarkers deployed in this study (lysosomal
379 reactions and indicators of genotoxic damage) can effectively discriminate between experimental
380 treatments of earthworms exposed to B[a]P; and dioxin spiked ~~in~~ OECD soil and various types of
381 ~~polluted-contaminated~~ field soils (i.e. Campania samples). In particular, multivariate analysis of
382 biomarker data showed that all of the different contaminated soils had a detrimental effect on the
383 earthworms; control animals being clearly separated from the treated ones. Among these latter,
384 worms exposed for 28 d to the highest B[a]P and TCDD concentration as well to soil from Site 2
385 (primarily dioxins and furans) were the most distant from the controls, these animals showing
386 greater biological effects.

387 A separation between temporal and concentration factors were also evident for B[a]P and TCDD
388 treatments; while, as for the Campania samples, Sites 1 and 5 (primarily PAHs) were clustered
389 together and Site 2 (primarily dioxins and furans) was clearly distinct from all the other samples.

390 Many pollutants may exert toxicity and genotoxicity directly, as well as through oxidative stress.

391 The resulting (oxidative) damage to membranes, proteins and DNA contribute to decreased protein

392 synthesis, cell injury and physiological dysfunction (Kirchin et al., 1992; Lowe et al., 2006; Moore
393 et al., 2006; Viarengo, 1989). LMS in blue mussels is directly correlated with total oxyradical
394 scavenging capacity (TOSC), polyribosome formation (translational efficiency in initiation of
395 protein synthesis); and inversely proportional to DNA damage (micronuclei formation), lipofuscin
396 (age/stress pigment, ceroid lipofuscin) formation, lysosomal swelling and autophagic accumulation
397 of lipid (Dailianis et al., 2003; Kalpaxis et al., 2004; Krishnakumar et al., 1994; Moore et al., 2006;
398 Regoli, 2000).

399 In this study, multiple regression analysis of the biomarker data indicated that in all three exposure
400 studies LMS is significantly correlated to L/C, and both these lysosomal parameters to MNi
401 induction.

402 Although lipofuscin content showed no consistent pattern of correlations across the various
403 treatments, the BIO-ENV routine indicated that experimental exposure of worms to B[a]P and
404 TCDD as well as to Campania soil samples resulted in LMS + lipofuscin (LF) + L/C emerging as
405 the most effective combination of lysosomal biomarkers (Table 2).

406 The recent developments in many research fields are leading to the discovery of prognostic
407 biomarkers that could be suitable as risk indicator of pathologies (Berghella et al., 2014; Jenkins et
408 al., 2011; Moore et al., 2006; Ortiz et al., 2011). Many biomarkers probably only exhibit a response
409 in a part of the “health status space” (Allen and Moore, 2004; Depledge et al., 1993; Moore et al.,
410 2006); where they will indicate that a reaction has taken place and may even indicate health status
411 within a narrow range, or what has induced the response, but they do not generally indicate the
412 health status of the whole range from healthy to irreversible damage (Köhler et al., 2002). In terms
413 of environmental prognostics, the first stage is to relate biomarker responses to health status of
414 individual organisms by mapping the said responses against an integrated “health status” indicator
415 (Allen and Moore, 2004; Köhler et al., 2002; Moore et al., 2004, 2006).

416 Lysosomes have attracted a great interest in the field of ecotoxicology as they are the target of a
417 wide range of contaminants (Allison and Mallucci, 1964; Moore et al., 2009; Sforzini et al., 2014;
418 Viarengo et al., 1985, 2007b) and they are present in all nucleated cells. Lysosomes contain
419 numerous hydrolytic enzymes involved in diverse cellular processes including the degradation of
420 cellular and extracellular macromolecules (Moore, 1976; Pipe, 1993). The evidence is steadily
421 accumulating that LMS is a generic indicator of cellular health in eukaryotic cells, as is indicated by
422 studies with protozoans, coelenterates, annelids, crustaceans, molluscs, fish and mammals (Lin et
423 al., 2010; Moore et al., 2012; Sohaebuddin and Tang, 2013). This parameter is now considered a
424 highly sensitive biomarker that allows to follow the evolution of the stress syndrome from its early
425 phase to the development of pathological conditions (Moore, 1988; Moore et al., 2004). LMS has
426 been used in the liver cells of the flatfish flounder (*Platichthys flesus*) to predict the degree of liver
427 degeneration (from cell injury through to hepatocellular carcinoma) as a result of PAH and
428 organochlorine exposure (Köhler et al., 2002); and lysosomal integrity in hepatopancreatic digestive
429 cells of mussels is directly related to scope for growth; and also, in the digestive cells of oysters
430 (*Crassostrea virginica*) to larval viability (Allen and Moore, 2004; Ringwood et al., 2004).

431 A useful method of integrating biomarker data into a “health status space” involves the use of
432 Principal Components Analysis (PCA) to reduce the dimensionality of the problem to a simple two
433 dimensional representation (Allen and Moore, 2004; Chatfield and Collins, 1980). PCA is
434 commonly used as a cluster analysis tool and is designed to capture the variance in a dataset in
435 terms of principle components. Hence, by plotting the first principal component (PC1) of the
436 selected biomarker data against LMS, as an indicator of the gradient from health to pathology and
437 disease, we effectively integrated the multi-biomarker data and the graph reflects the gradient of
438 toxicity among the samples. **These results clearly showed-indicate that LMS is prognostic of for**
439 **the health status of the earthworms treated with benzo(a)pyrene and TCDD; and also in**
440 **worms exposed to contaminated field samples from Campania.; Ffurther research on the**

441 **second/third worm generation will clarify if this parameter is also prognostic of negative**
442 **effects at population level.**

443

444 **5. Conclusion**

445 Overall, the biomarkers utilised in this study were able to reveal in *E. andrei* the toxic and
446 genotoxic effects of even low levels (close to the Italian legal limits for contaminants in residential
447 areas) of bioavailable pollutants in OECD as well as in natural soils.

448 Multivariate statistical analysis proved to be a valuable additional tool for improving the
449 interpretation of multi-biomarker results in exposed worms. In particular, multivariate statistics
450 showed that the use of the selected parameters enabled us to distinguish between temporal and
451 concentration factors of chemicals' exposure as well as between different contaminated soils.

452 Among the different parameters evaluated, diagnostic of a stress syndrome in the organisms, we
453 have shown that LMS is a prognostic indicator for health of edaphic sentinel animals, such as
454 earthworms. The approach described here will facilitate the validation, and further the new
455 development of robust diagnostic and prognostic tools that can be used along with other chemical,
456 ecotoxicological and ecological tools as indices of sustainability.

457

458 **Acknowledgements**

459 This study was supported financially by C.I.P.E. (Comitato Interministeriale per la Programmazione
460 Economica) - Piemonte Region, Project Code C22 (R/02/16) and Irpa (ex Apat) research contract
461 "Valutazione del rischio ecologico in siti ad elevato inquinamento della Regione Campania".

462

463 **References**

464 Adamowicz, A., 2005. Morphology and ultrastructure of the earthworm *Dendrobaena veneta*
465 (Lumbricidae) coelomocytes. *Tissue Cell.* 37, 125-133.

466 Allen, J.I., Moore, M.N., 2004. Environmental prognostics: is the current use of biomarkers
467 appropriate for environmental risk evaluation. *Mar. Environ. Res.* 58, 227-232.

468 Allison, A.C., Mallucci, L., 1964. Uptake of hydrocarbon carcinogens by lysosomes. *Nature* 203,
469 1024-1027.

470 Asensio, V., Rodríguez-Ruiz, A., Garmendia, L., Andre, J., Kille, P., Morgan, A.J., Soto, M.,
471 Marigómez, I., 2013. Towards an integrative soil health assessment strategy: A three tier
472 (integrative biomarker response) approach with *Eisenia fetida* applied to soils subjected to
473 chronic metal pollution. *Sci. Total Environ.* 442, 344-365.

474 Astley, K.N., Meigh, H.C., Glegg, G.A., Braven, J., Depledge, M.H., 1999. Multi-variate Analysis
475 of Biomarker Responses in *Mytilus edulis* and *Carcinus maenas* from the Tees Estuary
476 (UK). *Mar. Pollut. Bull.* 39, 145-154.

477 Baderna, D., Colombo, A., Amodei, G., Cantù, S., Teoldi, F., Cambria, F., Rotella, G., Natolino, F.,
478 Lodi, M., Benfenati, E., 2013. Chemical-based risk assessment and in vitro models of
479 human health effects induced by organic pollutants in soils from the Olona Valley. *Sci.*
480 *Total Environ.* 463-464, 790-801.

481 Banni, M., Attig, H., Sforzini, S., Oliveri, C., Mignone, F., Boussetta, H., Viarengo, A., 2014.
482 Transcriptomic responses to heat stress and nickel in the mussel *Mytilus galloprovincialis*.
483 *Aquat. Toxicol.* 148, 104-112.

484 Beliaeff, B., Burgeot, T., 2002. Integrated biomarker response: a useful tool for ecological risk
485 assessment. *Environ. Toxicol. Chem.* 21, 1316-1322.

486 Berghella, A.M., Contasta, I., Marulli, G., D'Innocenzo, C., Garofalo, F., Gizzi, F., Bartolomucci,
487 M., Laglia, G., Valeri, M., Gizzi, M., Friscioni, M., Barone, M., Del Beato, T., Secinaro, E.,

488 Pellegrini, P., 2014. Ageing gender-specific "Biomarkers of Homeostasis", to protect
489 ourselves against the diseases of the old age. *Immun. Ageing* 11, 3.

490 Bernet, D., Schmidt-Posthaus, H., Wahli, T., Burkhardt-Holm, P., 2000. Effects of wastewater on
491 fish health: an integrated approach to biomarker responses in brown trout (*Salmo trutta* L.).
492 *J. Aquat. Ecosystem Stress Recovery* 8, 143-151.

493 Binelli, A., Cogni, D., Parolini, M., Riva, C., Provini, A., 2009. In vivo experiments for the
494 evaluation of genotoxic and cytotoxic effects of Triclosan in Zebra mussel hemocytes.
495 *Aquat. Toxicol.* 91, 238-244.

496 Binelli, A., Cogni, D., Parolini, M., Provini, A., 2010. Multi-biomarker approach to investigate the
497 state of contamination of the R. Lambro/R. Po confluence (Italy) by zebra mussel
498 (*Dreissena polymorpha*). *Chemosphere* 79, 518-528.

499 Bolognesi, C., Hayashi, M., 2011. Micronucleus assay in aquatic animals. *Mutagenesis* 26, 205-
500 213.

501 Bouché, M.B., 1972. Lombriciens de France, *Écologie et Systématique*. I.N.R.A. Publ. Ann. Zool.
502 *Ecol. Anim.* (no horse-serie) 72 (2), 671.

503 Brenner, M., Broeg, K., Frickenhaus, S., Buck, B.H., Koehler, A., 2014. Multi-biomarker approach
504 using the blue mussel (*Mytilus edulis* L.) to assess the quality of marine environments:
505 season and habitat-related impacts. *Mar. Environ. Res.* 95, 13-27.

506 Broeg, K., Zander, S., Diamant, A., Körting, W., Krüner, G., Paperna, I., Westernhagen, H., 1999.
507 The use of fish metabolic, pathological and parasitological indices in pollution monitoring I.
508 North Sea. *Helgol. Mar. Res.* 53, 171-194.

509 Burgos, M.G., Winters, C., Stürzenbaum, S.R., Randerson, P.F., Kille, P., Morgan, A.J., 2005. Cu
510 and Cd effects on the earthworm *Lumbricus rubellus* in the laboratory: multivariate
511 statistical analysis of relationships between exposure, biomarkers, and ecologically relevant
512 parameters. *Environ. Sci. Technol.* 39, 1757-1763.

513 Cajaraville, M.P., Bebianno, M.J., Blasco, J., Porte, C., Sarasquete, C., Viarengo, A., 2000. The use
514 of biomarkers to assess the impact of pollution in coastal environments of the Iberian
515 Peninsula: a practical approach. *Sci. Total Environ.* 247, 295-311.

516 Cancio, I., ap Gwynn, I., Ireland, M.P., Cajaraville, M.P., 1995. The effect of sublethal exposure on
517 the ultrastructure and on the distribution of acid phosphatase activity in chloragocytes of
518 earthworms (Annelida, Oligochaeta). *Histochem. J.* 27, 965-973.

519 Camus, L., Grøsvik, B.E., Børseth, J.F., Jones, M.B., Depledge, M.H., 2000. Stability of lysosomal
520 and cell membranes in haemocytes of the common mussel (*Mytilus edulis*): effect of low
521 temperatures. *Mar. Environ. Res.* 50, 325-329.

522 Canesi, L., Ciacci, C., Lorusso, L.C., Betti, M., Guarnieri, T., Tavolari, S., Gallo, G., 2006.
523 Immunomodulation by 17beta-estradiol in bivalve hemocytes. *Am. J. Physiol. Regul. Integr.*
524 *Comp. Physiol.* 291, R664-R673.

525 Caselli, F., Gastaldi, L., Gambi, N., Fabbri, E., 2006. In vitro characterization of cholinesterases in
526 the earthworm *Eisenia andrei*. *Comp. Biochem. Physiol. C* 143, 416-421.

527 Chatfield, C., Collins, A. J., 1980. Introduction to multivariate analysis. London: Chapman and
528 Hall.

529 Clarke, K.R., 1999. Nonmetric multivariate analysis in community-level ecotoxicology. *Environ.*
530 *Toxicol. Chem.* 18, 117-127.

531 Cooper, E.L., Roch, P., 2003. Earthworm immunity: a model of immune competence: The 7th
532 international symposium on earthworm ecology - Cardiff - Wales - 2002. *Pedobiologia* 47,
533 676-688.

534 Dagnino, A., Allen, J.I., Moore, M.N., Broeg, K., Canesi, L., Viarengo, A., 2007. Development of
535 an expert system for the integration of biomarker responses in mussels into an animal health
536 index. *Biomarkers* 12, 155-172.

537 Dailianis, S., Domouhtsidou, G.P., Raftopoulou, E., Kaloyianni, M., Dimitriadis, V.K., 2003.
538 Evaluation of neutral red retention assay, micronucleus test, acetylcholinesterase activity
539 and a signal transduction molecule (cAMP) in tissues of *Mytilus galloprovincialis* (L.), in
540 pollution monitoring. Mar. Environ. Res. 56, 443-470.

541 Depledge, M.H., Amaral-Mendes, J.J., Daniel, B., Halbrook, R.S., Kloepper-Sams, P., Moore,
542 M.N., Peakall, D.P., 1993. The conceptual basis of the biomarker approach, in Peakall,
543 D.G., Shugart, L.R. (Eds.), Biomarkers - research and application in the assessment of
544 environmental health. Springer, Berlin, Heidelberg, pp. 15-29.

545 Dimitrova, N., Dimitrov, S., Georgieva, D., Van Gestel, C.A.M., Hankard, P., Spurgeon, D., Li, H.,
546 Mekenyan, O., 2010. Elimination kinetic model for organic chemicals in earthworms. Sci.
547 Total Environ. 408, 3787-3793.

548 Domouhtsidou, G.P., Dimitriadis, V.K., 2001. Lysosomal and lipid alterations in the digestive gland
549 of mussels, *Mytilus galloprovincialis* (L.) as biomarkers of environmental stress. Environ.
550 Pollut. 115, 123-137.

551 Fenech, M., 2000. The in vitro micronucleus technique. Mutat. Res. 455, 81-95.

552 Fischer, E., Molnár, L., 1992. Environmental aspects of the chloragogenous tissue of earthworms.
553 Soil Biol. Biochem. 24, 1723-1727.

554 Franzellitti, S., Viarengo, A., Dinelli, E., Fabbri, E., 2012. Molecular and cellular effects induced by
555 hexavalent chromium in Mediterranean mussels. Aquat. Toxicol. 124-125, 125-132.

556 Galloway, T.S., Brown, R.J., Browne, M.A., Dissanayake, A., Lowe, D., Jones, M.B., Depledge,
557 M.H., 2004. A multibiomarker approach to environmental assessment. Environ. Sci.
558 Technol. 38, 1723-1731.

559 Garmendia, L., Soto, M., Ortiz-Zarragoitia, M., Orbea, A., Cajaraville, M.P., Marigómez, I., 2011.
560 Application of a battery of biomarkers in mussel digestive gland to assess long-term effects

561 of the Prestige oil spill in Galicia and Bay of Biscay: correlation and multivariate analysis. J.
562 Environ. Monit. 13, 933-942.

563 Gastaldi, L., Ranzato, E., Caprì, F., Hankard, P., Pérès, G., Canesi, L., Viarengo, A., Pons, G.,
564 2007. Application of a biomarker battery for the evaluation of the sublethal effects of
565 pollutants in the earthworm *Eisenia andrei*. Comp. Biochem. Physiol. C 146, 398-405.

566 Gelboin, H.V., 1980. Benzo(a)pyrene metabolism, activation and carcinogenesis: role and
567 regulation of mixed-function oxidases and related enzymes. Physiol. Rev. 60, 1107-1166.

568 Guidi, P., Frenzilli, G., Benedetti, M., Bernardeschi, M., Falleni, A., Fattorini, D., Regoli, F.,
569 Scarcelli, V., Nigro, M., 2010. Antioxidant, genotoxic and lysosomal biomarkers in the
570 freshwater bivalve (*Unio pictorum*) transplanted in a metal polluted river basin. Aquat.
571 Toxicol. 100, 75-83.

572 Hayashi, Y., Heckmann, L.-H., Simonsen, V., Scott-Fordsmand, J.J., 2013. Time-course profiling
573 of molecular stress responses to silver nanoparticles in the earthworm *Eisenia fetida*.
574 Ecotoxicol. Environ. Saf. 98, 219-226.

575 Hauton, C., Hawkins, L.E., Hutchinson, S., 1998. The use of the neutral red retention assay to
576 examine the effects of temperature and salinity on haemocytes of the European flat oyster
577 *Ostrea edulis* (L.). Comp. Biochem. Physiol. 119, 619-623.

578 Irigaray, P., Ogier, V., Jacquenet, S., Notet, V., Sibille, P., Méjean, L., Bihain, B.E., Yen, F.T.,
579 2006. Benzo[a]pyrene impairs β -adrenergic stimulation of adipose tissue lipolysis and
580 causes weight gain in mice. A novel molecular mechanism of toxicity for a common food
581 pollutant. FEBS J. 273, 1362-1372.

582 ISPRA, 2012. Diossine, furani e policlorobifenili. Indagine ambientale nella Regione Campania.
583 ISPRA, Quaderni - Laboratorio n. 1 / 2012, pp. 496.

584 Jenkins, M., Flynn, A., Smart, T., Harbron, C., Sabin, T., Ratnayake, J., Delmar, P., Herath, A.,
585 Jarvis, P., Matcham, J., 2011. A statistician's perspective on biomarkers in drug
586 development. *Pharm. Stat.* 10, 494-507.

587 Kalpaxis, D.L., Theos, C., Xaplanteri, M.A., Dinos, G.P., Catsiki, A.V., Leotsinidis, M., 2004.
588 Biomonitoring of Gulf of Patras N. Peloponnesus, Greece. Application of a biomarker suite
589 including evaluation of translation efficiency in *Mytilus galloprovincialis* cells. *Environ.*
590 *Res.* 94, 211-220.

591 Kirchin, M.A., Moore, M.N., Dean, R.T., Winston, G.W., 1992. The role of oxyradicals in
592 intracellular proteolysis and toxicity in mussels. *Mar. Environ. Res.* 34, 315-320.

593 Krishnakumar, P.K., Casillas, E., Varanasi, U., 1994. Effect of environmental contaminants on the
594 health of *Mytilus edulis* from Puget Sound, Washington, USA. I Cytochemical measures of
595 lysosomal responses in the digestive cells using automatic image analysis. *Mar. Ecol. Prog.*
596 *Ser.* 106, 249-261.

597 Köhler, A., 1991. Lysosomal perturbations in fish liver as indicator for toxic effects of
598 environmental pollution. *Comp. Biochem. Physiol.* 100C, 123-127.

599 Köhler, A., Wahl, E., Söffker, K., 2002. Functional and morphological changes of lysosomes as
600 prognostic biomarkers of toxic liver injury in a marine flatfish (*Platichthys flesus* (L.)).
601 *Environ. Toxicol. Chem.* 21, 2434-2444.

602 Lin, P.-H., Lin, C.-H., Huang, C.-C., Chuang, M.-C., Lin, P., 2007. 2,3,7,8-Tetrachlorodibenzo-p-
603 dioxin (TCDD) induces oxidative stress, DNA strand breaks, and poly(ADP-ribose)
604 polymerase-1 activation in human breast carcinoma cell lines. *Toxicol. Lett.* 172, 146-158.

605 Lin, Y., Epstein, D.L., Liton, P.B., 2010. Intralysosomal iron induces lysosomal membrane
606 permeabilization and cathepsin D-mediated cell death in trabecular meshwork cells exposed
607 to oxidative stress. *Invest. Ophthalmol. Vis. Sci.* 51, 6483-6495.

608 Lowe, D.M., Pipe, R.K., 1994. Contaminant induced lysosomal membrane damage in marine
609 mussel digestive cells: an in vitro study. *Aquat. Toxicol.* 30, 357-365.

610 Lowe, D.M., Moore, M.N., Clarke, R.K., 1981. Effects of oil on digestive cells in mussels:
611 quantitative alterations in cellular and lysosomal structure. *Aquat. Toxicol.* 1, 213-226.

612 Lowe, D.M., Moore, M.N., Readman, J.W., 2006. Pathological reactions and recovery of
613 hepatopancreatic digestive cells from the marine snail *Littorina littorea* following exposure
614 to a polycyclic aromatic hydrocarbon. *Mar. Environ. Res.* 61, 457-470.

615 Lüllman-Rauch, R., 1979. Drug-induced lysosomal storage disorders. *Lysosomes in Biology and*
616 *Pathology* 6, pp. 49-130.

617 Martínez, K., Abad, E., Rivera, J., 2006. Surveillance programme on dioxin levels in soils in the
618 Campo de Gibraltar (Southwest Spain). *Chemosphere* 65, 382-389.

619 Moore, M.N., 1976. Cytochemical demonstration of latency of lysosomal hydrolases in the
620 digestive cells of the common mussel, *Mytilus edulis*, and changes induced by thermal
621 stress. *Cell Tissue Res.* 175, 279-287.

622 Moore, M.N., 1988. Cytochemical responses of the lysosomal system and NADPH-
623 ferrihemoprotein reductase in molluscan digestive cells to environmental and experimental
624 exposure to xenobiotics. *Mar. Ecol. Prog. Ser.* 46, 81-89.

625 Moore, M.N., 1990. Lysosomal cytochemistry in marine environmental monitoring. *Histochem. J.*
626 22, 187-191.

627 Moore, M.N., Depledge, M.H., Readman, J.W., Leonard, P., 2004. An integrated biomarker-based
628 strategy for ecotoxicological evaluation of risk in environmental management. *Mutat. Res.*
629 552, 247-268.

630 Moore, M.N., Allen, J.I., McVeigh, A., 2006. Environmental prognostics: an integrated model
631 supporting lysosomal stress responses as predictive biomarkers of animal health status. *Mar.*
632 *Environ. Res.* 61, 278-304.

633 Moore, M.N., Koehler, A., Lowe, D., Viarengo, A., 2008. Lysosomes and autophagy in aquatic
634 animals, in: Klionsky, D. (Ed.), *Methods in Enzymology*. Academic Press/Elsevier,
635 Burlington, 451, pp. 582-620.

636 Moore, M.N., Readman, J.A., Readman, J.W., Lowe, D.M., Frickers, P.E., Beesley, A., 2009.
637 Lysosomal cytotoxicity of carbon nanoparticles in cells of the molluscan immune system: an
638 in vitro study. *Nanotoxicology* 3, 40-45.

639 Moore, M.N., Viarengo, A., Somerfield, P.J., Sforzini, S., 2012. Linking lysosomal biomarkers and
640 ecotoxicological effects at higher biological levels, in: Amiard-Triquet, C., Amiard, J.C.,
641 Rainbow, P.S. (Eds.), *Ecological Biomarkers: Indicators of Ecotoxicological Effects*. CRC
642 Press, Boca Raton, FL, pp. 107-130.

643 Narbonne, J.-F., Daubèze, M., Clérendeau, C., Garrigues, P., 1999. Scale of classification based on
644 biochemical markers in mussels: application to pollution monitoring in European coasts.
645 *Biomarkers* 4, 415-424.

646 OECD, 2004. Guideline for Testing of Chemicals. No. 222. Earthworm Reproduction Test (*Eisenia*
647 *fetida/andrei*). Organization for Economic Cooperation and Development, Paris.

648 Ortiz, A., Massy, Z.A., Fliser, D., Lindholm, B., Wiecek, A., Martínez-Castelao, A., Covic, A.,
649 Goldsmith, D., Süleymanlar, G., London, G.M., Zoccali, C., 2011. Clinical usefulness of
650 novel prognostic biomarkers in patients on hemodialysis. *Nat, Rev, Nephrol.* 8, 141-150.

651 Peakall, D.B., 1994. The role of biomarkers in environmental assessment (1). Introduction.
652 *Ecotoxicology* 3, 157-160.

653 Pipe, R.K., 1993. The generation of reactive oxygen metabolites by the haemocytes of the mussel
654 *Mytilus edulis*. *Dev. Comp. Immunol.* 16, 111-122.

655 Peeters-Joris, C., 2000. The lysosomes of earthworm chloragocytes: biochemical and morphological
656 characterization. *Comp. Biochem. Physiol. A* 126, 323-340.

- 657 Prentø, P., 1987. Distribution of 20 enzymes in the midgut region of the earthworm, *Lumbricus*
658 *terrestris* L., with particular emphasis on the physiological role of the chloragog tissue.
659 *Comp. Biochem. Physiol. A* 87, 135-142.
- 660 Regoli, F., 2000. Total oxyradical scavenging capacity (TOSC) in polluted and translocated
661 mussels: a predictive biomarker of oxidative stress. *Aquat. Toxicol.* 50, 351-361.
- 662 Ringwood, A.H., Hoguet, J., Keppler, C., Gielazyn, M., 2004. Linkages between cellular biomarker
663 responses and reproductive success in oysters, *Crassostrea virginica*. *Mar. Environ. Res.* 58,
664 151-155.
- 665 Roméo, M., Bennani, N., Gnassia-Barelli, M., Lafaurie, M., Girard, J.P., 2000. Cadmium and
666 copper display different responses towards oxidative stress in the kidney of the sea bass
667 *Dicentrarchus labrax*. *Aquat. Toxicol.* 48, 185-194.
- 668 Sanchez, W., Aït-Aïssa, S., Palluel, O., Ditche, J.M., Porcher, J.M., 2007. Preliminary investigation
669 of multi-biomarker responses in three-spined stickleback (*Gasterosteus aculeatus* L.)
670 sampled in contaminated streams. *Ecotoxicology* 16, 279-287.
- 671 Santoyo, M.M., Flores, C.R., Torres, A.L., Wrobel, K., Wrobel, K., 2011. Global DNA methylation
672 in earthworms: a candidate biomarker of epigenetic risks related to the presence of
673 metals/metalloids in terrestrial environments. *Environ. Pollut.* 159, 2387-2392.
- 674 Schiller, C.M., Adcock, C.M., Moore, R.A., Walden, R., 1985. Effect of 2,3,7,8-tetrachlorodibenzo-
675 p-dioxin (TCDD) and fasting on body weight and lipid parameters in rats. *Toxicol. Appl.*
676 *Pharmacol.* 81, 356-361.
- 677 Schmid, W., 1975. The micronucleus test. *Mutat. Res.* 31, 9-15.
- 678 Scott-Fordsmand, J.J., Weeks, J.M., 2000. Biomarkers in earthworms. *Rev. Environ. Contam.*
679 *Toxicol.* 165, 117-159.

680 Sforzini, S., Dagnino, A., Oliveri, L., Canesi, L., Viarengo, A., 2011. Effects of dioxin exposure in
681 *Eisenia andrei*: integration of biomarker data by an Expert System to rank the development
682 of pollutant-induced stress syndrome in earthworms. *Chemosphere* 85, 934-942.

683 Sforzini, S., Boeri, M., Dagnino, A., Oliveri, L., Bolognesi, C., Viarengo, A., 2012. Genotoxicity
684 assessment in *Eisenia andrei* coelomocytes: a study of the induction of DNA damage and
685 micronuclei in earthworms exposed to B[a]P and TCDD-spiked soils. *Mutat. Res.* 746, 35-
686 41.

687 Sforzini, S., Moore, M.N., Boeri, M., Benfenati, E., Colombo, A., Viarengo A., 2014.
688 Immunofluorescence detection and localization of B[a]P and TCDD in earthworm tissues.
689 *Chemosphere* 107, 282-289.

690 Shaw, J.P., Dondero, F., Moore, M.N., Negri, A., Dagnino, A., Readman, J.W., Lowe, D.R.,
691 Frickers, P.E., Beesley, A., Thain, J.E., Viarengo, A., 2011. Integration of biochemical,
692 histochemical and toxicogenomic indices for the assessment of health status of mussels from
693 the Tamar Estuary, U.K. *Marine Environ. Res.* 72, 13-24.

694 Sohaebuddin, S.K., Tang, L., 2013. A simple method to visualize and assess the integrity of
695 lysosomal membrane in mammalian cells using a fluorescent dye. *Methods Mol. Biol.* 991,
696 25-31.

697 Spurgeon, D.J., Hopkin, S.P., 1999. Comparisons of metal accumulation and excretion kinetics in
698 earthworms (*Eisenia fetida*) exposed to contaminated field and laboratory soils. *Appl. Soil*
699 *Ecol.* 11, 227-243.

700 Terman, A., Brunk, U.T., 2004. Lipofuscin. *Int. J. Biochem. Cell. Biol.* 36, 1400-1404.

701 Turja, R., Höher, N., Snoeijs, P., Baršienė, J., Butrimavičienė, L., Kuznetsova, T., Kholodkevich,
702 S.V., Devier, M.-H., Budzinski, H., Lehtonen, K.K., 2014. A multibiomarker approach to
703 the assessment of pollution impacts in two Baltic Sea coastal areas in Sweden using caged
704 mussels (*Mytilus trossulus*). *Sci. Total Environ.* 473-474, 398-409.

705 USEPA, 2009. Priority pollutants, Appendix A to 40 CFR, Part 423-126. US Environmental
706 Protection Agency, Washington, DC.

707 Valentín, L., Nousiainen, A., Mikkonen, A., 2013. Introduction to Organic Contaminants in Soil:
708 Concepts and Risks, in: Vicent, T., Caminal, G., Eljarrat, E., Barceló, D. (Eds.), Emerging
709 Organic Contaminants in Sludges: Analysis, Fate and Biological Treatment. Hdb Env.
710 Chem., 24, pp. 1-30.

711 van der Oost, R., Beyer, J., Vermeulen, N.P., 2003. Fish bioaccumulation and biomarkers in
712 environmental risk assessment: a review. Environ. Toxicol. Pharmacol. 13, 57-149.

713 Viarengo, A., 1989. Heavy metals in marine invertebrates: mechanisms of regulation and toxicity at
714 the cellular level. Rev. Aquat. Sci. 1, 295-317.

715 Viarengo, A., Nott, J.A., 1993. Mechanisms of heavy metal cation homeostasis in marine
716 invertebrates. Comp. Biochem. Physiol. C104, 355-372.

717 Viarengo, A., Moore, M.N., Pertica, M., Mancinelli, G., Zanicchi, G., Pipe, R.K., 1985.
718 Detoxification of copper in the cells of the digestive gland of mussel: the role of lysosomes
719 and thioneins. Sci. Tot. Environ. 44, 135-145.

720 Viarengo, A., Moore, M.N., Mancinelli, G., Mazzucotelli, A., Pipe, R.K., Farrar, S.V., 1987.
721 Metallothioneins and lysosomes in metal toxicity and accumulation in marine mussels: the
722 effect of cadmium in the presence and absence of phenanthrene. Mar. Biol. 94, 251-257.

723 Viarengo, A., Burlando, B., Giordana, A., Bolognesi, C., Gabrielides, G.P., 2000. Networking and
724 expert-system analysis: next frontier in biomonitoring. Mar. Environ. Res. 49, 483-486.

725 Viarengo, A., Dondero, F., Pampanin, D.M., Fabbri, R., Poggi, E., Malizia, M., Bolognesi, C.,
726 Perrone, E., Gollo, E., Cossa, G.P., 2007a. A biomonitoring study assessing the residual
727 biological effects of pollution caused by the HAVEN wreck on marine organisms in the
728 Ligurian Sea (Italy). Arch. Environ. Contam. Toxicol. 53, 607-616.

729 Viarengo, A., Lowe, D., Bolognesi, C., Fabbri, E., Koehler, A., 2007b. The use of biomarkers in
730 biomonitoring: A 2-tier approach assessing the level of pollutant-induced stress syndrome in
731 sentinel organisms. *Comp. Biochem. Physiol. C* 146, 281-300.

732 Ville, P., Roch, P., Cooper, E.L., Masson, P., Narbonne, J.F., 1995. PCBs increase molecular-
733 related activities (lysozyme, antibacterial, hemolysis, proteases) but inhibit macrophage-
734 related functions (phagocytosis, wound healing) in earthworms. *J. Invertebr. Pathol.* 65,
735 217-24.

736 Wang, Z., Chen, J., Yang, P., Qiao, X., Tian, F., 2007. Polycyclic aromatic hydrocarbons in Dalian
737 soils: distribution and toxicity assessment. *J. Environ. Monit.* 9, 199-204.

738 Weeks, J.M., Svendsen, C., 1996. Neutral red retention by lysosomes from earthworm (*Lumbricus*
739 *rubellus*) coelomocytes: a simple biomarker of exposure to soil copper. *Environ. Toxicol.*
740 *Chem.* 15, 1801-1805.

741 Winzer, K., Van Noorden, C.J.F, Köhler, A., 2002. Sex-specific biotransformation and
742 detoxification after xenobiotic exposure of primary cultured hepatocytes of European
743 flounder (*Platichthys flesus* L.). *Aquat. Toxicol.* 59, 17-33.

744 Yang, X., Song, Y., Kai, J., Cao, X., 2012. Enzymatic biomarkers of earthworms *Eisenia fetida* in
745 response to individual and combined cadmium and pyrene. *Ecotoxicol. Environ. Saf.* 86,
746 162-167.
747

748 Fig. 1. Biomarker responses in *E. andrei* after exposure of worms for 10 d (grey columns) and 28 d
749 (black columns) to different concentrations of B[a]P and TCDD spiked in OECD soil. A, B)
750 Lysosomal membrane stability; C, D) lipofuscin content; E, F) neutral lipid accumulation; G, H)
751 lysosomal/cytoplasmic volume ratio. Data, expressed as percent change with respect to control
752 values, represent the mean \pm SD of at least five replicates. * indicates statistically significant
753 differences ($p < 0.05$ Mann-Whitney *U*-test).

754

755 Fig. 2. Biomarker responses in *E. andrei* after exposure of worms for 10 d to soils collected from
756 different areas in Campania region (Italy). A) Lysosomal membrane stability in the cells of
757 chloragogenous tissue (based on latency of β -*N*-acetylhexosaminidase); B) lipofuscin content; C)
758 neutral lipid accumulation; D) lysosomal/cytoplasmic volume ratio; E) micronuclei frequency.
759 Data, expressed as percent change with respect to control values, represent the mean \pm SD of at
760 least five replicates. * indicates statistically significant differences ($p < 0.05$ Mann-Whitney *U*-test).

761


762 Fig. 3. Principal Component Analysis (PCA) with superimposed Cluster Analysis showing
763 Resemblance Levels (distance), and vectors for the individual biomarkers.

764






765 Fig. 4. Regressions for B[a]P, TCDD and Campania data with correlation coefficients below each
766 set of plots. *Critical values for the Correlation coefficients are $r = 0.381$ for $P = 0.05$ (Campania),
767 and $r = 0.349$ for $P = 0.05$ (B[a]P and dioxin).

768




769 Fig. 5. Lysosomal membrane stability as an integrated indicator of health plotted against the first
770 principal component eigenvectors (PC 1) for all the remaining cellular biomarkers. The data has
771 been log transformed and normalised; and open triangles shown on the B[a]P plot are outliers not
772 included in the analysis.



773 Fig. 1.
774






775 Fig. 2.
776



777

778 Fig. 3.

779




780


781 Fig. 4.

782


BENZO(a)PYRENE



DIOXIN



Campania Soil Samples



784 Fig. 5.
785