



# Phytotoxin produced by the netted scab pathogen, *Streptomyces turgidiscabies* strain 65, isolated in Sweden

Masahiro Natsume<sup>1</sup> · Asaho Nagagata<sup>2</sup> · Marja Aittamaa<sup>3</sup> · Naoko Okaniwa<sup>2</sup> · Panu Somervuo<sup>3</sup> · Hans-Peter Fiedler<sup>4</sup> · Jan F. Kreuze<sup>5,6</sup> · Veli-Matti Rokka<sup>7</sup> · Hans Bång<sup>8</sup> · Hiroshi Kawaide<sup>1</sup> · Jari P. T. Valkonen<sup>3,5</sup>

Received: 25 June 2017 / Accepted: 12 November 2017 / Published online: 20 January 2018  
© The Author(s) 2018. This article is an open access publication

## Abstract

*Streptomyces* spp. are a highly diverse group of bacteria most of which are soil-inhabiting saprophytes. A few are plant pathogens that produce a family of phytotoxins called thaxtomins and cause significant economic losses, e.g., by reducing the marketability of potato tubers (*Solanum tuberosum*). In northern Europe, *S. scabies*, *S. turgidiscabies* and *S. europaeiscabiei* are the most common plant pathogenic species. In this study, a *Streptomyces* strain isolated from a netted scab lesion on a tuber of potato cv. Bintje in northern Sweden was identified as *S. turgidiscabies* but was found to differ in the genomic region carrying genes required for thaxtomin biosynthesis. Our results showed that the strain did not produce thaxtomin but rather phytotoxin fridamycin E, which is an anthraquinone novel to plant pathogenic *Streptomyces* spp. Fridamycin E was shown to reduce or inhibit sprouting of potato microtubers in vitro. While fridamycin E is known to have antibiotic activity against Gram-positive bacteria, the inhibitory activity of fridamycin E on plant growth is a novel finding.

**Keywords** Netted scab · Potato · *Streptomyces turgidiscabies* · *Streptomyces reticuliscabiei* · Fridamycin E · Phytotoxin

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s10327-018-0765-8>) contains supplementary material, which is available to authorized users.

✉ Masahiro Natsume  
natsume@cc.tuat.ac.jp

✉ Jari P. T. Valkonen  
jari.valkonen@helsinki.fi

<sup>1</sup> Division of Bioregulation and Biointeraction, Institute of Agriculture, Tokyo University of Agriculture and Technology, Saiwai-cho, Fuchu, Tokyo 183-8509, Japan

<sup>2</sup> Graduate School of Agriculture, Tokyo University of Agriculture and Technology, Saiwai-cho, Fuchu, Tokyo 183-8509, Japan

<sup>3</sup> Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland

<sup>4</sup> Department of Microbiology/Biotechnology, University of Tübingen, Tübingen, Germany

<sup>5</sup> Department of Plant Biology and Forest Genetics, Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden

<sup>6</sup> International Potato Center (CIP), Lima, Peru

<sup>7</sup> AgriFood Research Centre Finland, Jokioinen, Finland

<sup>8</sup> Svenskt Potatisutsäde AB, Umeå, Sweden

## Introduction

Bacteria in the genus *Streptomyces* are Gram-positive and have been classified into 767 species and subspecies (DSMZ 2017). Most species are soil-inhabiting saprophytes, and only a few are plant pathogenic. The streptomycetes that infect potato (*Solanum tuberosum* L.) cause economic losses through reduction of the marketable yield. Diseased potato tubers are characterized by diverse lesions with superficial, raised, erumpent or pitted shapes, that are collectively called common scab to distinguish them from powdery scab, which is caused by the protist *Spongospora subterranea* (Loria et al. 1997; Loria 2001; Wanner and Kirk 2015). Loria et al. (1997) proposed classifying the lesions into two types: one typified by raised or pitted lesions (so-called “common scab”) and the other characterized by superficial lesions.

The first *Streptomyces* species described as the causal agent of common scab was *S. scabies* (syn. *scabiei*), which has a worldwide distribution in potato production areas (Lambert and Loria 1989). Another, more recently described and widely distributed species, *S. turgidiscabies*, causes all the aforementioned symptoms on potato tubers (Aittamaa et al. 2010; Hiltunen et al. 2005; Kreuze et al. 1999; Miyajima et al. 1998; Takeuchi et al. 1996). The types of lesions

induced by these species depend on the phytotoxins and virulence factors carried by the species and, also, on the timing of infection, potato cultivar and environmental conditions (Bignell et al. 2014; Loria et al. 1997; Natsume et al. 2017). Therefore, it is difficult to distinguish scab pathogens based on disease symptoms.

The common scab symptoms caused by *S. scabies* and *S. turgidiscabies* are attributable to thaxtomin A (Fig. 1) (Goyer et al. 1998; Lawrence et al. 1990; Toth et al. 1998), a cyclic dipeptidic phytotoxin produced by these bacteria (King et al. 1989; King and Calhoun 2009). The genes required for thaxtomin synthesis, virulence and colonization of tubers are located in two or three genomic regions in *S. scabies* and *S. turgidiscabies* and are collectively referred to as a pathogenicity island (PAI) (Aittamaa et al. 2010; Kers et al. 2005). One region of the PAI contains genes for thaxtomin synthesis and is called the toxicogenic region (Lerat et al. 2009). The *txtD* gene in the region encodes a nitric oxide (NO) synthase that generates NO from L-arginine (Kers et al. 2004), while the *txtE* gene product catalyzes nitration of L-tryptophan by NO and O<sub>2</sub> to produce 4-nitro-L-tryptophan (Barry et al. 2012). The *txtA* and *txtB* genes encode non-ribosomal peptide synthetases that combine 4-nitro-L-tryptophan and phenylalanine to form the cyclic dipeptide skeleton of thaxtomin (Healy et al. 2000). The *txtC* gene encodes a cytochrome P450-type monooxygenase that is required for postcyclization hydroxylation of cyclic dipeptide (Healy et al. 2002). Expression of these genes is regulated by an AraC/XylS-type regulator, *txtR*, which is also located in the PAI region (Joshi et al. 2007a). The toxicogenic region of PAI shows wide genetic variability among the strains of *S. turgidiscabies* (Aittamaa et al. 2010), possibly because the PAI or parts of it may be exchanged among *Streptomyces* strains and species (Kers et al. 2005).

A variety of superficial deformations are observed in potatoes in various areas of the world. Superficial scab symptoms called netted scab occur in Europe and russet scab occurs in Japan and North America (Loria et al. 1997). Classification and identification of the causal agents of netted and russet scab have gained less attention, probably because these symptoms are less severe than common scab. In Japan, a *Streptomyces* species that induces tortoiseshell-like cracks

in the periderm of the potato was isolated and named “*S. cheloniumii*” (Oniki et al. 1986; Suzui et al. 1988). “*S. cheloniumii*” was found to be closely related to *S. diastatochromogenes* by a homology search of 16S rDNA sequences (Suzui et al. 2004). This species is distinguished from other phytopathogenic *Streptomyces* spp. by its reddish aerial mycelia, spiny spores forming a spiral spore chain, and no production of melanin or other diffusible pigments (Oniki et al. 1986; Suzui et al. 2004), but an authoritative description of the species has not been published. The Canadian russet scab agent is discerned from the Japanese strain by gray aerial mycelia and smooth surface-spores forming flexuous spore chains (Faucher et al. 1993). The bacterium shows highest similarity to cluster group 14 of Williams et al. (1983), *S. aureofaciens*, by numerical classification. Phenotypic studies on *Streptomyces* sp. isolated in France showed that strains isolated from netted scab tissue were different from the pathogens of common scab or agents isolated from russet scab in Canada (Bouček-Mechiche et al. 1998). The *Streptomyces* strains causing netted scab in France were described as a new species, *S. reticuliscabiei* (type strain DSM41804; CFBP 4531) (Bouček-Mechiche et al. 2000a, b), but were later found to belong to *S. turgidiscabies* (Bouček-Mechiche et al. 2006). This strain is characterized by the production of light grey spores that make flexuous chains, and the lack of melanin production (Bouček-Mechiche et al. 2000a). The strain is pathogenic to only few potato cultivars such as Bintje, Désirée and Carmine and does not infect carrot or radish (Bouček-Mechiche et al. 2000b). Other *Streptomyces* spp. causing common scab have a wide host range among root crops, including carrot and radish (Wanner and Kirk 2015).

Among the *Streptomyces* spp. that produce superficial symptoms, “*S. cheloniumii*” was found to produce phytotoxin FD-891 (Fig. 1), which induces necrosis in potato tuber slices (Natsume et al. 2005). The phytotoxin FD-891 was detected in potatoes infected by the pathogen and was also produced by other pathogenic strains of “*S. cheloniumii*” collected from different geographical areas in Japan.

In northern Sweden and Finland, *S. turgidiscabies* is a typical causal agent of common scab (Aittamaa et al. 2010; Kreuze et al. 1999; Lehtonen et al. 2004). Surveys of potato




Fig. 1 Structure of phytotoxins produced by phytopathogenic *Streptomyces* spp

scab pathogens in these regions revealed a few tubers with netted scab symptoms (Bång 1979; Lindholm et al. 1997). *Streptomyces* sp. strain 65 isolated from a netted scab-affected tuber of potato cv. Bintje in northern Sweden caused netted scab following inoculation of healthy Bintje plants under controlled conditions (Fig. 2).

We hypothesized that the different types of scab lesions on potato are caused by different phytotoxins and have been searching for them in *Streptomyces* strains (Natsume et al. 1998, 2005, 2017). The aim of this study was to make a molecular level comparison of *Streptomyces* sp. strain 65 with common scab pathogens previously detected in northern Sweden and Finland and to analyze the phytotoxins produced by strain 65.

## Materials and methods

### Taxonomical identification of *streptomyces* sp. Strain 65

Phytopathogenic *Streptomyces* sp. strain 65 was isolated from a netted scab-affected tuber of potato cv. Bintje in northern Sweden (Lehtonen et al. 2004). Isolation of DNA and amplification by polymerase chain reaction (PCR) with species-specific primers for *S. turgidiscabies* and *S. scabies* were carried out as described previously (Kreuze et al. 1999; Lehtonen et al. 2004).

### Microarray analysis

Microarray analysis of *Streptomyces* sp. strain 65 was carried out with 131 probes designated for the predicted genes of *S. scabies* to target the genes involved in the thaxtomin biosynthesis and genes previously shown to be related to pathogenicity, including *necl1*, *tomA* and *fas* (Aittamaa et al. 2010). We used genomic DNA for the

analysis because the aim was to analyze genomic differences rather than differences in gene expression.

### Phytotoxicity tests and HPLC analysis of phytotoxins

Phytotoxicity of microbial extracts and fractions during purification was examined by application to rape (*Brassica rapa* L.) seedlings and monitoring their growth, as reported previously (Natsume et al. 2005). Possible effects of isolated phytotoxin on potato were studied by applying phytotoxin in methanol at different concentrations (3.6 or 1.2  $\mu\text{g}/\mu\text{L}$ ) on microtubers of potato cv. Bintje grown *in vitro*. The microtubers (10–15 mm in diameter) were placed on water agar in Petri dishes, and 50  $\mu\text{L}$  of phytotoxin solution was applied to each microtuber. The lids of Petri dishes were closed and tubers were grown in a growth chamber at 22 °C for 3 weeks.

Thaxtomin A production by *Streptomyces* sp. strain 65 and *S. turgidiscabies* NBRC 16080<sup>T</sup> was analyzed using HPLC system LC-10 (Shimadzu, Kyoto, Japan) as described previously (Natsume et al. 1998, 2005).


### Isolation and identification of the phytotoxin produced by *streptomyces* sp. Strain 65

*Streptomyces* sp. strain 65 was cultured with shaking in potato peel broth (Beauséjour et al. 1999) at 28 °C for 14 days. The cultured material was filtered to separate the mycelial mass and filtrate. Mycelia were soaked in acetone for 2 days, the residue was filtered out, and the acetone extract was evaporated. The residual aqueous mycelial extract and culture filtrate were treated with chloroform at pH 3 to yield a chloroform-soluble crude extract. Phytotoxin was purified from the extract by column chromatography and preparative TLC as shown in Fig. 3.

**Fig. 2** Netted scab symptoms caused by *Streptomyces turgidiscabies* strain 65 in potato cv. Bintje



**Fig. 3** Purification of the phytotoxin produced by *S. turgidiscabies* strain 65. Isolation steps (a), chromatogram and fractionation in preparative TLC (b) and growth inhibitory activity of fractionated samples (c). Each fraction from the preparative TLC equivalent to 20 mL of cultured material was used for the growth inhibitory assay



## Results

### Identification of strain 65 as *S. turgidiscabies*


The 16S rRNA gene from strain 65 was amplified by PCR with specific primers for *S. turgidiscabies* but not with the primers specific to *S. scabiei* (Fig. 4). On the other hand, genes *necl* and *tmp* located in the “colonization region” of the PAI were not amplified from strain 65 with the primers designed for *S. turgidiscabies*. The 16S rRNA gene sequence of strain 65 was determined (NCBI accession no. AJ278759) and found to be identical or nearly identical to many strains of *S. turgidiscabies* that cause common scab

on potato (Aittamaa et al. 2010). Furthermore, the 16S rRNA gene sequence of strain 65 was 98–99% identical to that of *S. reticuliscabiei* strain CFBP4531 and DSM41804 that cause netted scab on potatoes in France. Phylogenetic comparison indicated that strain 65 and *S. reticuliscabiei* are closely related and belong to *S. turgidiscabies* (Fig. 5).

### Microarray analysis of thaxtomin biosynthetic genes of *S. turgidiscabies* strain 65

Genes involved in thaxtomin biosynthesis and pathogenicity of *S. turgidiscabies* strain 65 were studied using microarray analysis and compared with (1) *S. turgidiscabies* strains






**Fig. 4** PCR analysis of selected genes located in the pathogenicity island using **a** universal *Streptomyces* primers for the 16S gene (16S); **b** species-specific primers for *S. scabies* (Scab); **c** specific primers for *S. turgidiscabies* (Turg); and primers for **d** the *necl* gene (Nec) and **e** *tnp*, a transposase pseudogene located upstream and separated by an intergenic region from *necl* (Tnp). The numbers above the lanes identify the tested *Streptomyces* strains described previously (Lehtonen et al. 2004)

32 and 300 which produce thaxtomin A but differ in having high and low virulence on potato tubers, respectively (Hiltunen et al. 2005); (2) *S. reticuliscabiei* DSM 41,804 isolated from a netted scab lesion on potato cv. Bintje in France (Bouchek-Mechiche et al. 2000a), and (3) the type strain of *S. scabies* (ATCC49173). In *S. turgidiscabies* strain 32, all probes generated strong signals, e.g., for the genes *txtA*, *txtB*, *txtC*, *txtD* and *txtR* involved in thaxtomin biosynthesis and gene *necl* involved in virulence (Fig. 6) (Bukhalid et al. 1998; Joshi et al. 2007b). Presence of the aforementioned genes was detected also in *S. scabies* ATCC49173. In contrast, *S. turgidiscabies* strain 65 and *S. reticuliscabiei* DSM41804 showed no signal for these genes.

### Phytotoxin production of *S. turgidiscabies* strains 65 and NBRC 16080<sup>T</sup>

The phytotoxicity of crude chloroform extracts of strain 65 and the type strain of *S. turgidiscabies*, NBRC 16080, cultured on oatmeal agar (Babcock et al. 1993) was examined on rape seedlings. Extracts of both strains showed strong growth inhibitory activity at the dose corresponding to the amount of extract obtained from 10 mL of bacterial culture (Fig. S1).



**Fig. 5** Clustering of *Streptomyces* strains by neighbor-joining analysis based on sequences of the 16S RNA cistrons. The tree was constructed using the Kimura two-parameter model (Kimura 1980). Bootstrap values were determined by 1000 replicates; values greater than 70% are shown. Bar indicates the Kimura nucleotide differences (Knucl). All strains isolated from scabby potato tubers, except strain 65, have been described previously (Aittamaa et al. 2010). The upper-

most cluster contains strains of *S. turgidiscabies*, including DSM1804 originally described as *S. reticuliscabiei* (Bouchek-Mechiche et al. 2000a, 2006). Strain 266 belongs to an unknown species related to *S. turgidiscabies*. The cluster in the middle contains strains of *S. scabies*, including the type strain ATCC49173. Unknown nonpathogenic *Streptomyces* strains form the clade at the bottom of the tree

**Fig. 6** Microarray analysis: Heat map indicating mean signal densities (red color) given by probes that target different genes in the pathogenicity island (PAI) region of *Streptomyces* strains. The microarray included 1–5 probes for 30 open reading frames (ORFs) in the three characterized PAI regions of *S. turgidiscabies* strain Car8 (separated by dotted lines) (Kers et al. 2005; Aittamaa et al. 2010). PAI regions I and II contain nine genes typical to the “colonization region” (CR), such as *nec1* contributing to virulence, whereas PAI III includes the genes needed for thaxtomin synthesis and is called a “toxicogenic region” (TR) (Lerat et al. 2009). (Color figure online)

The chloroform extract was purified by silica-gel column chromatography and analyzed by HPLC. Thaxtomin A was detected in the extract of strain NBRC 16080<sup>T</sup> but not in that of strain 65 (Fig. S2). The result is consistent with previous results in microarray analysis showing that the toxicogenic region genes for thaxtomin production differ between strain 65 with thaxtomin-producing *Streptomyces* strains.


### Isolation of phytotoxin produced by *S. turgidiscabies* strain 65

Comparison of phytotoxin production in agar and liquid cultures with oatmeal, oat bran and potato peel media (Babcock et al. 1993; Beauséjour et al. 1999; Goyer et al. 1998) showed that shake culture in potato peel broth provided high phytotoxin productivity and facilitated mass culture (Fig. S3), and this culture method was selected for further experiments.

Phytotoxin was purified as outlined in Fig. 3a, b according to the bioassay. In the first column chromatography on silica gel, 10% MeOH in CHCl<sub>3</sub> fraction showed phytotoxic activity. Under these conditions, thaxtomin A was recovered in 20% MeOH in CHCl<sub>3</sub> fraction and, thus, the presence of new phytotoxin was determined. In this purification, 37.1 mg of orange-yellow compound was isolated from 8 L of culture. Only one fraction showed phytotoxicity at each chromatographic step. This newly isolated phytotoxin had an *R<sub>f</sub>* value of 0.30 that was indistinguishable from thaxtomin A on silica gel TLC (solvent: 10% MeOH in CHCl<sub>3</sub>). Both phytotoxins had a very similar yellow color.

### Structural determination of the phytotoxin

HPLC-PDA analysis of the isolated phytotoxin showed a single peak (Fig. S4), and the UV spectrum of the peak showed  $\lambda_{\max}$  at 433, 289, 254 and 215 nm (Fig. S5). This UV spectrum is characteristic of the presence of a 1,5-dihydroxyanthraquinone chromophore such as  $\alpha$ -citromycinone (Johdo et al. 1991). LC-ESI-MS analysis of the isolated phytotoxin shows peaks at *m/z* 735, 730, 379, 357 and 339, which can be assigned to ions [2M+Na]<sup>+</sup>, [2M+H<sub>2</sub>O]<sup>+</sup>, [M+Na]<sup>+</sup>, [M+H]<sup>+</sup> and [M+H-H<sub>2</sub>O]<sup>+</sup>. From these data, the molecular weight of the phytotoxin was determined



to be 356. The molecular formula  $C_{19}H_{16}O_7$  was deduced from high-resolution MS data,  $m/z$  357.09855 (calculated for  $C_{19}H_{17}O_7$ : 357.09743).

The presence of one carboxyl group was demonstrated by disappearance of the peak in the mass chromatogram at  $m/z$  357 and appearance of a peak at  $m/z$  371 by LC–ESI–MS analysis of phytotoxin treated with trimethylsilyldiazomethane in methanol–benzene (Hashimoto et al. 1981).

The  $^1H$ -NMR spectrum (600 MHz,  $CD_3OD$ , Fig. S6) revealed three adjacent aromatic protons ( $\delta$  7.72 [d,  $J=8.1$  Hz], 7.62 [dd,  $J=8.1, 8.1$  Hz] and 7.21 [d,  $J=8.1$  Hz]); 2 aromatic protons with *ortho*-coupling ( $\delta$  7.66 [d,  $J=8.1$  Hz] and 7.65 [d,  $J=8.1$  Hz]); 2 pairs of isolated methylene protons ( $\delta$  2.93 [d,  $J=13.2$  Hz] 2.90 [d,  $J=13.2$  Hz] and  $\delta$  2.28 [d,  $J=15.4$  Hz], 2.22 [d,  $J=15.4$  Hz]); and a quarternary methyl signal ( $\delta$  1.13, 3H, s).

Searches in the SciFinder database based on the aforementioned structural characteristics identified fridamycin E as the structural candidate. Comparison of isolated phytotoxin with the authentic fridamycin E in the HPLC–UV–Vis database system (Fiedler 1993) showed an identical HPLC chromatogram, UV spectrum and ESI–MS spectra (positive and negative ion modes) (Fig. S7). We thus identified the phytotoxin produced by *S. turgidiscabies* strain 65 as fridamycin E (Fig. 1). The absolute stereochemistry of isolated fridamycin E was the same as previously reported (Matsumoto et al. 1991) by the positive optical rotation of the isolated sample ( $[\alpha]_D^{28} +4.3^\circ$  [ $c=0.49$ , dioxane]).

### Phytotoxicity of fridamycin E

The purified phytotoxin caused a 17% reduction in root growth and a 14% reduction in shoot growth at 14  $\mu\text{g}/\text{mL}$  in rape seedlings and a 96% reduction in root growth and a 75% reduction in shoot growth at 100  $\mu\text{g}/\text{mL}$  (Fig. 3c). Thaxtomin A caused a 24% reduction in root growth and a 25% reduction in shoot growth in rape seedlings at 0.16  $\mu\text{g}/\text{mL}$ , which indicates that the phytotoxicity of fridamycin E is approximately 1% of that of thaxtomin A.

Application of fridamycin E to potato microtubers at a concentration of 3.6  $\mu\text{g}/\mu\text{L}$  reduced the growth of sprouts or inhibited sprouting of potato microtubers grown *in vitro*. Typical netted scab symptoms were not observed on the microtubers, but a few small scab-like lesions were observed under the present experimental conditions (Fig. 7). Application of fridamycin E at a lower concentration (1.2  $\mu\text{g}/\mu\text{L}$ ) caused only minor or no reduction in growth of sprouts and no symptoms on the microtubers (data not shown).




**Fig. 7** Inhibited sprouting and sprout elongation on microtubers of potato cv. Bintje following application of fridamycin E in methanol at 3.6  $\mu\text{g}/\mu\text{L}$  (50  $\mu\text{L}/\text{tuber}$ ) (to the left) compared with microtubers treated with methanol only (right)

### Effect of plant extracts on phytotoxin production

Requirements of plant extracts such as oatmeal for thaxtomin production by *S. scabies* have been reported (Beauséjour et al. 1999), and effective ingredients were shown to be suberin and cellobiose (Beauséjour et al. 1999; Johnson et al. 2007; Lerat et al. 2010). We examined the requirements for phytotoxin production in *S. turgidiscabies* strain 65 and NBRC16080<sup>T</sup> by culturing on oatmeal agar and starch–polypeptone agar (Sakai et al. 1984). Crude chloroform extracts of strain NBRC16080 prepared from oatmeal agar strongly inhibited growth of rape seedlings but those from starch–polypeptone medium did not (Fig. 8). Thaxtomin A content in oatmeal agar culture was determined to be 0.39  $\mu\text{g}/\text{mL}$  of cultured material by HPLC analysis of the partially purified extracts. Both extracts of strain 65 showed comparable growth inhibitory activity (Fig. 8), and fridamycin E production was determined to be 0.83  $\mu\text{g}/\text{mL}$  in both media. Thus, fridamycin E production by strain 65 does not require any plant extract.

**Fig. 8** Effect of plant extract on production of phytotoxin. *Ctrl* control, *OA* oatmeal agar, *SPA* starch-polypeptone agar. Growth inhibitory activity of the crude  $\text{CHCl}_3$  extract was examined at the dose equivalent to 10 mL of cultured material



## Discussion

*Streptomyces* on potato tubers cause a severe symptom with raised, erumpent, or pitted lesions and milder symptoms characterized by superficial lesions (Loria et al. 1997). Various symptoms caused by *Streptomyces* spp. are considered to be related to the level of virulence of the pathogen, the amount of phytotoxin (thaxtomin) produced, infection timing in relation to developmental stage of tubers, genetic differences among potato varieties and environmental conditions. The finding that potato cv. Green Mountain is very susceptible to common scab but highly resistant to netted scab (Loria et al. 1997) suggests that the pathogenicity determinants of *Streptomyces* strains causing netted scab differ from those causing common scab.

Extensive investigations of potato diseases caused by *Streptomyces* in northern Sweden revealed a strain causing only superficial lesions on tubers of potato cv. Bintje (Lehtonen et al. 2004). Genetic analysis of the bacterium in this study showed that the strain belongs to species *S. turgidiscabies* but that its toxicogenic region in PAI, including the thaxtomin biosynthetic gene cluster, differs from that of *S. turgidiscabies* which causes deep pitting and other severe scab symptoms. Consistent with these findings, *S. turgidiscabies* strain 65 did not produce thaxtomin but did produce a newly identified phytotoxin called fridamycin E. Fridamycin E is the aglycone of fridamycin A (= vineomycinone B<sub>2</sub>) (Matsumoto et al. 1991), but there is no report on its biosynthetic pathway or biosynthetic genes.

Application of fridamycin E to potato microtubers inhibited sprouting or reduced the growth of sprouts in a concentration-dependent manner. Antibiotic activity of fridamycin E against Gram-positive bacteria has been reported (Matsumoto et al. 1991), but this is the first report of phytotoxicity associated with fridamycin E.

Application of fridamycin E to potato microtubers did not cause any obvious netted scab symptom. One reason may be that we applied the phytotoxin once rather than repeatedly over time. For the formation of netted scab symptoms,

continuous production of phytotoxin on the epidermis by the pathogen may be required. Further experimentation will be needed to confirm whether fridamycin E is essential for development of netted scab and whether repeated applications of fridamycin E are required.

We previously found that bacteria isolated from russet scab lesions in Japan produces another phytotoxin FD-891 (Natsume et al. 2005). It has been suggested that the pathogenicity determinants of netted scab differ from those of common scab because resistance of potato cultivars to the netted scab pathogens is not correlated with resistance to the common scab pathogen, *S. scabies* (Loria et al. 1997). Identification of a novel phytotoxic macrolide, FD-891, in “*S. cheloniumii*” (Natsume et al. 2005) and an anthraquinone phytotoxin, fridamycin E, in *S. turgidiscabies* strain 65 supports the hypothesis. The results of the present study, along with our previous findings on FD-891, are important for demonstrating that the toxins produced by the pathogen determine the lesion type. Recently, we have shown that phytotoxin concanamycin is involved in determination of lesion type of common scab along with thaxtomin (Natsume et al. 2017).

*S. reticuliscabiei* was, at first, reported as the agent of netted scab in France (Bouček-Mechiche et al. 2000a, b) but was later identified as strain *S. turgidiscabies* by DNA–DNA hybridization and phylogenetic comparisons of 16S rRNA gene sequences (Bouček-Mechiche et al. 2006). However, because the symptoms caused by *S. reticuliscabiei* are distinct from the symptoms typically caused by *S. turgidiscabies*, Bouček-Mechiche et al. (2006) proposed maintaining the species name. Strain 65 of *S. turgidiscabies* causes symptoms similar to *S. reticuliscabiei* in potato cv. Bintje, but both *S. reticuliscabiei* and strain 65 are also pathogenic on other potato varieties (Kreuze et al. 1999; Bouček-Mechiche et al. 2000b; Lehtonen et al. 2004). Hence, it is worthwhile to consider whether both *S. reticuliscabiei* and strain 65 could be considered as strains of *S. turgidiscabies*. Whether *S. reticuliscabiei* produces fridamycin E is an interesting topic for further study.



To prove that fridamycin E is the causal agent of netted scab in northern Europe, we are now studying (1) pathogenicity of the fridamycin E-nonproducing mutants and (2) detection of fridamycin E in the lesions of netted scab caused by *S. turgidiscabies* strain 65.

**Acknowledgements** This collaborative research began on the occasion of International Potato Scab Symposium held in Sapporo in 2004. The authors express their deep appreciation to Prof. Emeritus Shigeo Naito of Hokkaido University, chair of the organizing committee.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Aittamaa M, Somervuo P, Laakso I, Auvinen P, Valkonen JPT (2010) Microarray-based comparison of genetic differences between strains of *Streptomyces turgidiscabies* with focus on the pathogenicity island. *Mol Plant Pathol* 11:733–746
- Babcock MJ, Eckwall EC, Schottel JL (1993) Production and regulation of potato-scab-inducing phytotoxins by *Streptomyces scabies*. *J Gen Microbiol* 139:1579–1586
- Bång H (1979) Studies on potato russet scab. I. A characterization of different isolates from northern Sweden. *Acta Agric Scand* 29:145–150
- Barry SM, Kers JA, Johnson EG, Song L, Aston PR, Patel B, Krasnoff SB, Crane BR, Gibson DM, Loria R, Challis GL (2012) Cytochrome P450-catalyzed l-tryptophan nitration in thaxtomin phytotoxin biosynthesis. *Nat Chem Biol* 8:814–816
- Beauséjour J, Goyer C, Vachon J, Beaulieu C (1999) Production of thaxtomin A by *Streptomyces scabies* strains in plant extract containing medium. *Can J Microbiol* 45:764–768
- Bignell DRD, Fyans JK, Cheng Z (2014) Phytotoxins produced by plant pathogenic *Streptomyces* species. *Appl Microbiol* 116:223–235
- Bouček-Mechiche K, Guérin C, Jouan B, Gardan L (1998) *Streptomyces* species isolated from potato scabs in France: numerical analysis of “Biotype-100” carbon source assimilation data. *Res Microbiol* 149:653–663
- Bouček-Mechiche K, Gardan L, Normand P, Jouan B (2000a) DNA relatedness among strains of *Streptomyces* pathogenic to potato in France: description of three new species, *S. europaeiscabiei* sp. nov. and *S. stelliscabiei* sp. nov. associated with common scab, and *S. reticuliscabiei* sp. nov. associated with netted scab. *Int J Syst Evol Microbiol* 50:91–99
- Bouček-Mechiche K, Pasco C, Andrivon D, Jouan B (2000b) Differences in host range, pathogenicity to potato cultivars and response to soil temperature among *Streptomyces* species causing common and netted scab in France. *Plant Pathol* 49:3–10
- Bouček-Mechiche K, Gardan L, Andrivon D, Normand P (2006) *Streptomyces turgidiscabies* and *Streptomyces reticuliscabiei*: one genomic species, two pathogenic groups. *Int J Syst Evol Microbiol* 56:2771–2776
- Bukhalid RA, Chung SY, Loria R (1998) *necl*, a gene conferring a necrogenic phenotype, is conserved in plant-pathogenic *Streptomyces* spp. and linked to a transposase pseudogene. *Mol Plant-Microbe Interact* 11:960–967
- DSMZ (2017) Catalogue of microorganisms. <https://www.dsmz.de/catalogues/catalogue-microorganisms.html>. Accessed 24 Jun 2017
- Faucher E, Otrysko B, Paradis É, Hodge NC, Stall RE, Beaulieu C (1993) Characterization of streptomycetes causing russet scab in Québec. *Plant Dis* 77:1217–1220
- Fiedler H-P (1993) Biosynthetic capacities of actinomycetes. 1. Screening for secondary metabolites by HPLC and UV-visible absorbance spectral libraries. *Nat Prod Lett* 2:119–128
- Goyer C, Vachon J, Beaulieu C (1998) Pathogenicity of *Streptomyces scabies* mutants altered in thaxtomin A production. *Phytopathology* 88:442–445
- Hashimoto N, Aoyama T, Shioiri T (1981) New methods and reagents in organic synthesis. 14. A simple efficient preparation of methyl esters with trimethylsilyldiazomethane (TMSCHN<sub>2</sub>) and its application to gas chromatographic analysis of fatty acids. *Chem Pharm Bull* 29:1475–1478
- Healy FG, Wach M, Krasnoff SB, Gibson DM, Loria R (2000) The *txtAB* genes of the plant pathogen *Streptomyces acidiscabies* encode a peptide synthetase required for phytotoxin thaxtomin A production and pathogenicity. *Mol Microbiol* 38:794–804
- Healy FG, Krasnoff SB, Wach M, Gibson DM, Loria R (2002) Involvement of a cytochrome P450 monooxygenase in thaxtomin A biosynthesis by *Streptomyces acidiscabies*. *J Bacteriol* 184:2019–2029
- Hiltunen LH, Weckman A, Ylhäinen A, Rita H, Richter E, Valkonen JPT (2005) Responses of potato cultivars to the common scab pathogens, *Streptomyces scabies* and *S. turgidiscabies*. *Ann Appl Biol* 146:395–403
- Johdo O, Ishikura T, Yoshimoto A, Takeuchi T (1991) Anthracycline metabolites from *Streptomyces violaceus* A262 I. Isolation of antibiotic-blocked mutants from *Streptomyces violaceus* A262. *J Antibiot* 44:1110–1120
- Johnson EG, Joshi MV, Gibson DM, Loria R (2007) Celooligosaccharides released from host plants induce pathogenicity in scab-causing *Streptomyces* species. *Physiol Mol Plant Pathol* 71:18–25
- Joshi MV, Bingell DRD, Johnson EG, Sparks JP, Gibson DM, Loria R (2007a) The AraC/XylS regulator TxtR modulates thaxtomin biosynthesis and virulence in *Streptomyces scabies*. *Mol Microbiol* 66:633–642
- Joshi M, Rong X, Moll S, Kers J, Franco C, Loria R (2007b) *Streptomyces turgidiscabies* secretes a novel virulence protein, Nec1, which facilitates infection. *Mol Plant-Microbe Interact* 20:599–608
- Kers JA, Wach MJ, Krasnoff SB, Widom J, Cameron KD, Bukhalid RA, Gibson DM, Crane BR, Loria R (2004) Nitration of a peptide phytotoxin by bacterial nitric oxide synthase. *Nature* 429:79–82
- Kers JA, Cameron KD, Joshi MV, Bukhalid RA, Morello JE, Waach MJ, Gibson DM, Loria R (2005) A large, mobile pathogenicity island confers plant pathogenicity on *Streptomyces* species. *Mol Microbiol* 55:1025–1033
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16:111–120
- King RR, Calhoun LA (2009) The thaxtomin phytotoxins: sources, synthesis, biosynthesis, biotransformation and biological activity. *Phytochemistry* 70:833–841
- King RR, Lawrence CH, Clark MC, Calhoun LA (1989) Isolation and characterization of pytoxins associated with *Streptomyces scabies*. *J Chem Soc Chem Commun* 1989:849–850
- Kreuze JF, Suomalainen S, Paulin L, Valkonen JPT (1999) Phylogenetic analysis of 16S rRNA genes and PCR analysis of the *necl* gene from *Streptomyces* spp. causing common scab, pitted scab, and netted scab in Finland. *Phytopathology* 89:462–469
- Lambert DH, Loria R (1989) *Streptomyces scabies* sp. nov., nom. rev. *Int J Syst Bacteriol* 39:387–392

- Lawrence CH, Clark MC, King RR (1990) Induction of common scab symptoms in aseptically cultured potato tubers by the vivotoxin, thaxtomin. *Phytopathology* 80:606–608
- Lehtonen MJ, Rantala H, Kreuze JF, Bång H, Kuisma L, Koski P, Virtanen E, Vihlman K, Valkonen JPT (2004) Occurrence and survival of potato scab pathogens (*Streptomyces* species) on tuber lesions: quick diagnosis based on a PCR-based assay. *Plant Pathol* 53:280–287
- Lerat S, Simao-Beaunoir A, Beaulieu C (2009) Genetic and physiological determinants of *Streptomyces scabies* pathogenicity. *Mol Plant Pathol* 10:579–585
- Lerat S, Simao-Beaunoir A-M, Wu R, Beaudoin N, Beaulieu C (2010) Involvement of the plant polymer suberin and the disaccharide cellobiose in triggering thaxtomin A biosynthesis, a phytotoxin produced by the pathogenic agent *Streptomyces scabies*. *Phytopathology* 100:91–96
- Lindholm P, Korttemaa H, Kokkola M, Haahtela K, Salkinoja-Salonen M, Valkonen JPT (1997) *Streptomyces* spp. isolated from potato scab lesions under Nordic conditions. *Plant Dis* 81:1317–1322
- Loria R (2001) Common scab. In: Stevenson WR, Loria R, Franc GD, Weingartner DP (eds) *Compendium of potato diseases*, 2nd edn. APS Press, St Paul, pp 14–15
- Loria R, Bukhalid RA, Fry BA, King RR (1997) Plant pathogenicity in the genus *Streptomyces*. *Plant Dis* 81:836–845
- Matsumoto T, Jona H, Katsuki M, Suzuki K (1991) Efficient method for introducing vineomycin-fridamycin-type side chain. Total synthesis of fridamycin E. *Tetrahedron Lett* 32:5103–5106
- Miyajima K, Tanaka F, Takeuchi T, Kuninaga S (1998) *Streptomyces turgidiscabies* sp. nov. *Int J Syst Bacteriol* 48:492–502
- Natsume M, Yamada A, Tashiro N, Abe H (1998) Differential production of the phytotoxins thaxtomin A and concanamycins A and B by potato common scab-causing *Streptomyces* spp. *Ann Phytopathol Soc Jpn* 64:202–204
- Natsume M, Komiya M, Koyanagi F, Tashiro N, Kawaide H, Abe H (2005) Phytotoxin produced by *Streptomyces* sp. causing potato russet scab in Japan. *J Gen Plant Pathol* 71:364–369
- Natsume M, Tashiro N, Doi A, Nishi Y, Kawaide H (2017) Effects of concanamycins produced by *Streptomyces scabies* on lesion type of common scab of potato. *J Gen Plant Pathol* 83:78–82
- Oniki M, Suzui T, Araki T, Sonoda R, Chiba T, Takeda T (1986) Causal agent of russet scab of potato (Japanese with English abstract). *Bull Natl Inst Agroenviron Sci* 2:45–59
- Sakai R, Kawamura H, Mino Y, Emami-Saravi R, Tani A (1984) Toxin production by *Streptomyces* spp. associated with scab of potato tuber and sugar beet I. Effect of carbon and nitrogen sources. *Ann Phytopathol Soc Jpn* 50:646–468
- Suzui T, Miyashita K, Tashiro N (1988) *Streptomyces cheloniumii* sp. nov., a new species causing russet scab of potato. Abstracts of Papers, 5th International Congress of Plant Pathology, Kyoto, p 177
- Suzui T, Tashiro N, Miyashita K, Manabe K, Matsuyama N (2004) Studies on the causal organism, ecology and control of potato russet scab in Japan. *Proc Int Potato Scab Sym*, Sapporo, pp 339–343
- Takeuchi T, Sawada H, Tanaka F, Matsuda I (1996) Phylogenetic analysis of *Streptomyces* spp. causing potato scab based on 16S rRNA sequences. *Int J Syst Bacteriol* 46:476–479
- Toth L, Akino S, Kobayashi K, Doi A, Tanaka F, Ogoshi A (1998) Production of thaxtomin A by *Streptomyces turgidiscabies*. *Soil Microorg* 51:29–34
- Wanner LA, Kirk WW (2015) *Streptomyces*—from basic microbiology to role as a plant pathogen. *Am J Potato Res* 92:236–242
- Williams ST, Goodfellow M, Alderson G, Wellington EMH, Sneath PHA, Sackin MJ (1983) Numerical classification of *Streptomyces* and related genera. *J Gen Microbiol* 129:1743–1813