

ORIGINAL REPORT

EFFECTIVENESS OF MULTIDISCIPLINARY EARLY REHABILITATION IN REDUCING BEHAVIOUR-RELATED RISK FACTORS

Mikhail Saltychev, MD¹, Katri Laimi, PhD^{2,3}, Ashraf El-Metwally, PhD^{4,5}, Tuula Oksanen, PhD^{6,7}, Jaana Pentti, BSc⁶, Marianna Virtanen, PhD⁶, Anne Kouvonen, PhD⁸, Mika Kivimäki, PhD^{6,9} and Jussi Vahtera, PhD^{3,6}

From the ¹Department of Rehabilitation, ²Department of Physical and Rehabilitation Medicine, Turku University Hospital, ³Department of Public Health, University of Turku and Turku University Hospital, Turku, Finland, ⁴King Abdullah International Medical Research Center, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Kingdom of Saudi Arabia, ⁵Epidemiology Group, University of Aberdeen, Aberdeen, UK, ⁶Finnish Institute of Occupational Health, Helsinki, Finland, ⁷Department of Society, Human Development and Health, Harvard School of Public Health, Boston, MA, USA, ⁸Warsaw School of Social Sciences and Humanities, Wroclaw Faculty, Wroclaw, Poland and ⁹Department of Behavioral Sciences, University of Helsinki, Finland

Objective: To evaluate the effect of a 4-week primary prevention programme on health-risk behaviours amongst employees at increased risk of work incapacity.

Methods: Based on survey data and health records from 53,416 public sector employees in Finland, we identified 872 employees who participated in early rehabilitation after the baseline survey. We selected 2,440 propensity-score-matched controls for these rehabilitants. Changes in the prevalence of physical inactivity, obesity, heavy drinking, and smoking, as well as in the intensity of leisure-time physical activity, weight, and alcohol consumption after the intervention were examined between the baseline and two subsequent surveys representing short-term (mean follow-up 1.7 years) and long-term (mean 5.8 years) follow-ups.

Results: There were no statistically significant differences between the rehabilitants and controls in terms of changes in weight, alcohol consumption, intensity of leisure-time physical activity, or prevalence of obesity, heavy drinking and physical inactivity during short-term or long-term follow-ups. During short-term follow-up, a higher rate of smoking cessation was observed for rehabilitants than controls (31.7% vs. 20.2%, $p=0.037$).

Conclusion: Vocationally oriented multidisciplinary early rehabilitation had little effect on health risk behaviours.

Key words: health behaviour change; health risk behaviour; propensity score; modifiable risks.

J Rehabil Med 2012; 44: 370–377

Guarantor: Jussi Vahtera, Department of Public Health, University of Turku, Turku, Finland. E-mail: jussi.vahtera@ttl.fi

Submitted May 8, 2011; accepted December 13, 2011

INTRODUCTION

Tobacco smoking, alcohol consumption, obesity, poor diet and physical inactivity are the most important potentially modifiable behaviour-related risk factors responsible for over half of the disease burden in developed countries (1). Furthermore, a

particular health risk behaviour rarely occurs alone. Instead, they tend to co-occur, causing multiplied risk of disease and premature death (2, 3). For example, only 3% of all adults in the USA have succeeded to avoid all 4 of the major health risk behaviours: smoking, overweight, physical inactivity and poor diet (4).

In addition to the co-occurrence of health risk behaviours, a co-variation of multiple health behavioural change has been observed. This implies that successful improvement in one health behaviour can lead to improvement in other health behaviours (5). Thus, public health professionals are trying to develop effective and efficient measures to reduce not just one, but multiple health risk behaviours simultaneously over the course of the same intervention programme. These measures should also take into account the contextual environment in which the person lives and works (6, 7). Multiple health behavioural changes to prevent cardiovascular diseases have been studied widely during the last few decades (8), but the evidence on successful measures to affect health risk behaviours remains inconclusive (2, 5, 9, 10).

Multidisciplinary prevention or rehabilitation programmes may constitute a means of helping participants make positive changes in their health behaviours, as these changes are associated with better health, assuring the work ability of the participants (11–16). In the present study, we examined the most common multidisciplinary rehabilitation programme in Finland, which aims at preventing long-term work disability. The programme includes physical and psychological training targeted at achieving multiple positive health behavioural changes. Although this preventive programme has been used since the early 1980s, only a few studies have evaluated its effectiveness in terms of health behavioural changes. These studies, mostly conducted without a control group and based on a relatively small number of participants (17–21), found no improvement in physical activity levels among participants (18, 19). However, we are not aware of previous studies that have simultaneously examined the effectiveness of this programme

on multiple behaviour-related risk factors, such as physical inactivity, smoking, overweight, and alcohol abuse. We have reported previously that the programme is of limited value in reducing the risk of a long-term work disability (22); the hazard ratio for incident all-cause work disability was 0.98 (95% confidence interval (CI) 0.76–1.25) in cases compared with their propensity score matched controls. The corresponding hazard ratios for diagnosis-specific work disability was 0.86 (95% CI 0.57–1.30) for musculoskeletal diseases and 1.08 (95% CI 0.67–1.74) for mental disorders.

The objective of this study was to evaluate the short- and long-term effects of a vocationally oriented, in-patient, multi-disciplinary preventive programme on smoking cessation, weight change, increase in leisure-time physical activity, and decrease in alcohol consumption.

METHODS

Study population

Finnish Public Sector Study. The present study is part of the Finnish Public Sector Study (FPSS), which is an ongoing prospective study amongst employees working in 10 towns and 21 hospitals. The study was approved by the ethics committee of the Finnish Institute of Occupational Health. The study covers all 151,618 employees with a ≥6 month job contract in any year from 1991/1996 to 2005 in 10 towns and 6 hospital districts in Finland. Questionnaire surveys targeted 94,494 employees who were working in the years 1997–1998 (phase 1, sub-cohort), 2000–2002 (phase 2), or 2004–2005 (phase 3). All of the respondents have been followed with repeated surveys (2006, phase 4; 2008–2009, phase 5; 2010, phase 6). In phases 1–3 and phase 5, identifiable questionnaire data on psychosocial factors at work, individual factors, health, and health behaviours were gathered. Through the use of the unique personal identification codes that are assigned to all citizens in Finland, all of the participants have been linked to employers’ records and national health registers.

Case-control selection procedure. We used data from phases 1–3 and 5 and included those who responded to 3 consequent identifiable surveys. The first survey response was considered as the baseline (i.e. 1997–1998 or 2000–02) and the following surveys as the first (2000–2002 or 2004–2005) and second follow-ups (2004–2005 or 2008–2009). The earliest survey response was considered for those who participated in both baseline surveys. This approach yielded a sample of 53,416 employees (response rate 70%, 81% of whom were women). We excluded participants who had been granted any rehabilitation by the Social Insurance Institution of Finland (SII) before the baseline survey ($n=4,176$) or with missing data on any of the matching variables ($n=3,699$). Eligible cases were those who had entered vocationally oriented medical rehabilitation (VOMR) between the first (baseline) and second (first follow-up) survey. By using propensity score matching (see the next section), we identified a study population of 872 cases and 2440 non-cases of VOMR (Fig. 1).

Propensity score matching. We used the propensity score approach to approximate the exchangeability of the comparison groups, which, in theory, would differ only in the receipt of VOMR (23, 24). Propensity score is the conditional probability of being assigned “treatment”, here VOMR, given the observed covariates. In order to calculate the propensity score, we used binary logistic regression models for VOMR (dichotomous outcome) including 24 pre-treatment variables known to be associated with rehabilitation and the health-risk behaviours available in the data, and their interactions with gender, socio-economic status, and age group. Once we had estimated the propensity score, each case was matched with 1–3 controls (non-VOMR recipients) according to a pre-defined calliper width of +0.01, and the unmatched cases were dis-




Fig. 1. Case-control selection procedure. SII: Social Insurance Institution of Finland; VOMR: vocationally oriented medical rehabilitation.

carded, resulting in a total of 3,312 subjects (872 participants and 2,440 propensity-score-matched controls) for the statistical analysis (Table 1). The balance achieved by matching was studied using the χ^2 test.

Intervention: vocationally oriented multidisciplinary rehabilitation

In Finland, VOMR is the most common rehabilitation programme sponsored by SII. It accounts for 12.3% of SII’s annual rehabilitation budget (25). In 2009, the median age of the VOMR participants was 50 years (25).

VOMR is a group-based multidisciplinary rehabilitation programme targeted towards employees at workplaces and occupations in which workers are subjected to considerable physical, mental or social strain that may lead to health problems and a deterioration in work capacity. Briefly, one of its selection criteria is the absence of recent long-term sick-leaves or severe long-standing illnesses.

The VOMR programme contains 3–4 periods of extensive multimodal and multiprofessional inpatient rehabilitation (a total of 15–21 days) implemented as group-based (8–10 persons) and individual sessions of supervised activity 4–6 h per day. Between the inpatient periods, the participants are expected to follow an individual exercise plan at home, which usually consists of self-reliant physical activities and psychological exercises. The participants are guided by professionals, the aim being better aerobic capacity, muscle strength and endurance through the adoption of more regular and comprehensive leisure-time physical activity and healthy dietary habits, cessation of smoking, and reduction in alcohol consumption (26).

Outcome: Change in behaviour-related risk factors

Using 3 repeated questionnaire surveys, we measured two types of outcome. Firstly, we measured the changes in the prevalences of obesity, leisure-time physical inactivity, smoking, and heavy drinking. Secondly, we measured the changes in weight, intensity of leisure-time physical activity, and alcohol consumption.

The body mass index (BMI; kg/m²) was derived from self-reported weight and height and dichotomized to indicate obesity (BMI ≥30). The change in self-reported weight (kg) was measured for those who were obese or non-obese at baseline.

The participants’ reports of the quantity of their physical activity equivalent to walking, brisk walking, jogging, or running was used to estimate the metabolic equivalent of task (MET) hours per day. Physical inactivity was defined as ≤2 MET h (no/yes) (27). The change in MET hours/day was assessed for those who were physically active and for those who were physically inactive at baseline.

Table I. Definitions of the variables used in the propensity score matching

Variable	Definition
Demographic characteristics	
Age group ^a	≤40 years, 41–50 years, or ≥51 years
Gender ^a	Male/female
Occupational grade ^a	International Standard Classification of Occupations (ISCO) grades: 1–2 – managers and professionals, 3 – technicians and associate professionals, 4 – clerks, 5 – service workers, or 6–9 – manual workers
Educational level ^b	High school: yes/no
Marital status ^b	Married or cohabiting vs single, divorced or widowed
Type of employer ^a	Municipality or hospital district
Residential region ^a	Southern, Northern, or Central Finland
Work characteristics	
Type of job contract ^a	Permanent/temporary
Length of the contract ^a	< 1 year, 1–9 years, or ≥ 10 years
Shift work ^b	Yes/no
Intention to leave work ^b	Continue the job, switch jobs, or give up the job
Job insecurity ^b	Two questions, which estimate the threat of long-term unemployment or dismissal (from 1 = very little to 5 = very much). A mean score was computed and divided into tertiles.
Job control ^b	9 items derived from the Job Content Questionnaire
Health behaviour	
Current smoking ^b	Yes/no
Heavy drinking ^b	Yes (>210 g pure alcohol per week)/no
Obesity ^b	Yes (body mass index (BMI) ≥30 kg/m ²)/no
Physical inactivity ^b	Yes (≤2 metabolic equivalent task (MET) h/day)/no
Health indicators	
Sickness absence ^c	Yes (>60 absence days during the last 3 years)/no (0 or 1–60 absence days)
Psychological distress ^b	Yes (psychological distress for at least 4 items of GHQ ^f)/no
Anxiety ^b	6-item Trait Anxiety Inventory: low, intermediate, or high anxiety
Self-rated health ^b	Sub-optimal (average or worse)/optimal (good or very good health)
Baseline physical health ^d	Yes (presence of a chronic disease ^g)/no
Use of prescribed painkillers ^e	Yes (>30 defined daily dosages during the survey year (ATC ^h code N02 and M01A))/no
Use of prescribed antidepressants ^e	Yes (>30 defined daily dosages during the survey year (ATC ^h code N06A))/no

^aObtained from employers' records; ^bObtained from survey; ^cObtained from the Sickness Absence Register of the Finnish Social Insurance Institution; ^dObtained from the Drug Reimbursement Register of the Finnish Social Insurance Institution; ^eObtained from the Drug Prescription Register of the Finnish Social Insurance Institution; ^f12-item General Health Questionnaire (GHQ); ^gIndicated by special reimbursement for the medical treatment of hypertension, cardiac failure, ischaemic heart disease, diabetes, asthma or other chronic obstructive lung disease, and rheumatoid arthritis; ^hAnatomical therapeutic chemical.

Smoking status (never, ex-smoker and current smoker) was operationalized as current smoking (yes/no). The change in smoking status amongst the baseline smokers was used to identify smoking cessation.

Participants reported their mean weekly consumption of beer, wine and spirits in units. The units were converted into grams of pure alcohol, and > 210 g pure alcohol per week was considered the cut-off for heavy drinking (no/yes) (28, 29). Binge drinking was determined by enquiring whether the participant had passed out due to heavy alcohol consumption once or more often during the past 12 months (30). The Finnish term used for "pass-out" refers to alcohol-related loss of consciousness, but without reference to loss of memory (blackout) (30). The change in alcohol consumption (g/week) was measured by the heavy drinking status at baseline.

Statistical analysis

We applied repeated-measures log-binomial regression analysis with the generalized estimating equations (GEE) method for studying the changes in the prevalence of the behaviour-related risk factors of the participants and controls (31, 32). The results were reported as percentages of the estimated prevalence, prevalence ratios and their 95% confidence limits and *p*-values.

We performed a repeated-measures analysis of variance to study the changes in the continuous variables (weight, physical activity and alcohol consumption) and reported the results as the mean values, their 95% confidence limits and *p*-values. Data on the baseline characteristics were gathered before the beginning of the intervention

(mean 1.8 years (SD 1.07)). The short-term follow-up ended with the first and the long-term follow-up with the second follow-up survey. The follow-up began immediately after the beginning of the rehabilitation. The first follow-up survey (short-term follow-up) took place on average 1.7 years (SD 1.01, range 0.003–4.5 years) after the start of the rehabilitation; for 19% of the rehabilitants, the rehabilitation programme was not yet completed. The second follow-up survey (long-term follow-up) was performed on average 5.8 years (SD 1.13, range 3.12–9.16) after the rehabilitation.

All of the statistical analyses were performed using SAS[®] 9.2 software (SAS Institute, Cary, NC, USA).

RESULTS

The prevalences of the health risk behaviours were relatively low amongst the participants before the intervention: only 13% were smokers, 22% were physically inactive, 10% were obese and 7% were heavy drinking. Three-quarters of the participants reported good self-rated health and a similar proportion did not use prescribed painkillers. Only 5% of the participants used prescribed antidepressants (Table II). Supporting the success of the propensity score matching, no differences were observed between the participants and their controls with respect to the

individual variables used to calculate the propensity score, except for employer type (Table II).

Changes in the prevalences of behaviour-related risk factors

Fig. 2 shows the changes in the prevalence of behaviour-related risk factors shortly after the intervention (mean 1.7 years (SD 1.01, range 0.003–4.55)) and in the long-term follow-up (mean 5.8 years (SD 1.13, range 3.12–9.16)). The prevalence of obesity increased amongst the participants and controls (by 6.4% and 4.1%, respectively) during an entire follow-up without a significant difference between the two groups. The prevalence of smoking decreased in both groups during an entire follow-up, amongst the participants by 4.5% and amongst the controls by 3.8%. Although the participants quit smoking ($p=0.037$) more often than the controls did during a short-term follow-up, this difference disappeared during a long-term follow-up. The prevalence of physical inactivity increased slightly amongst the participants and controls in the long-term follow-up (by 2.5% and 3.1%, respectively) without a significant difference between the groups ($p=0.68$). The prevalence of heavy and binge drinking remained almost unchanged in both groups.

Changes in weight, intensity of physical activity, and alcohol consumption

Table III shows the changes in weight (kg), intensity of physical activity (MET h/day), and alcohol consumption (g/week). There was a minor increase in body weight amongst the participants who were not obese at baseline when they were

Table II. Differences between the cases and controls as regards the covariates after the propensity score matching

Covariates used in the propensity score matching	Participants (n=872) n (%)	Controls (n=2440) n (%)	χ^2 -test p-value
Propensity score, mean (range)	0.111 (0.001–0.548)	0.093 (0.001–0.549)	
Gender			0.162
Women	789 (90)	2,166 (89)	
Men	83 (10)	274 (11)	
Age group, years			0.055
≤40	138 (16)	410 (17)	
41–50	532 (61)	1,558 (64)	
≥51	202 (23)	472 (19)	
ISCO grade ^a			0.946
1–2 (managers)	310 (36)	894 (37)	
3	223 (26)	594 (24)	
4	41 (5)	109 (4)	
5	205 (24)	585 (24)	
6–9 (manual)	93 (11)	258 (11)	
Type of employer			0.002
Municipality	590 (68)	1,785 (73)	
Hospital district	282 (32)	655 (27)	
Residential region			0.353
Southern Finland	500 (57)	1,396 (57)	
Central Finland	259 (30)	684 (28)	
Northern Finland	113 (13)	360 (15)	

Marital status			0.869
Married/cohabiting	672 (77)	1,887 (77)	
Single	200 (23)	553 (23)	
High educational level			0.440
No	406 (47)	1,099 (45)	
Yes	466 (53)	1,341 (55)	
Job contract			0.665
Permanent	810 (93)	2,277 (93)	
Fixed-term	62 (7)	163 (7)	
Length of job contract, years			0.286
<1	156 (18)	480 (20)	
1–9	599 (69)	1,673 (69)	
≥10	117 (13)	287 (12)	
Shift work			0.679
No	560 (64)	1,586 (65)	
Yes	312 (36)	854 (35)	
Intention to leave work			0.450
Continue the job	462 (53)	1,260 (52)	
Switch jobs	221 (25)	672 (28)	
Give up the job	189 (22)	508 (21)	
Job insecurity			0.916
Low	396 (45)	1,090 (45)	
Intermediate	352 (40)	1,004 (41)	
High	124 (14)	346 (14)	
Job control			0.594
Low	266 (31)	753 (31)	
Intermediate	305 (35)	889 (36)	
High	301 (35)	798 (33)	
Obesity			0.963
No	781 (90)	2,184 (90)	
Yes	91 (10)	256 (10)	
Smoking			0.353
No	758 (87)	2,090 (86)	
Yes	114 (13)	350 (14)	
Alcohol consumption, g/week			0.814
0–210	807 (93)	2,264 (93)	
>210	65 (7)	176 (7)	
Physical inactivity			0.897
No	683 (78)	1,906 (78)	
Yes	189 (22)	534 (22)	
Sickness absence, days/3 years			0.505
No	567 (65)	1,627 (67)	
1–59	247 (28)	674 (28)	
≥60	58 (7)	139 (6)	
Chronic medical conditions			0.473
No	799 (92)	2,216 (91)	
Yes	73 (8)	224 (9)	
Use of antidepressants			0.565
No	790 (95)	2,160 (95)	
Yes	38 (5)	116 (5)	
Use of painkillers			0.753
No	623 (75)	1,725 (76)	
Yes	205 (25)	551 (24)	
Anxiety			0.905
Low	242 (28)	662 (27)	
Intermediate	342 (39)	954 (39)	
High	288 (33)	824 (34)	
Sub-optimal self-rated health			0.906
No	675 (77)	1,884 (77)	
Yes	197 (23)	556 (23)	
Psychological distress			0.183
No	640 (73)	1,733 (71)	
Yes	232 (27)	707 (29)	

^aISCO: International Standard Classification of Occupations.




Fig. 2. Prevalence of participants' behaviour-related risk factors at baseline and at short-term (mean 20 months) and long-term (mean 70 months) follow-ups in comparison with controls.

compared with their controls (mean increase by 3.5 kg and 2.8 kg, respectively) in the long-term follow-up ($p=0.005$). Amongst the participants who were obese at baseline, no significant difference in weight change was observed. The physically inactive cases and controls at baseline increased their physical activity in the short-term follow-up (mean 1.66 and 1.69 MET h/day, respectively), and the improvement also lasted through the entire follow-up (mean 1.46 and 1.71 MET h/day, respectively), with no difference between the two groups. At the same time, the cases and controls physically active at baseline slightly reduced their activity throughout the entire follow-up. Also in relation to changes in alcohol consumption, no significant difference was observed between the two groups during follow-up.

As it is possible that the longest duration of the short-term follow-up in some cases exceeded the shortest duration of the long-term follow-up, we ran a sensitivity analysis including the 750 cases with short-term follow-up of 3 years or less, and the long-term follow-up was at least 4 years. The results did not differ from those of the main analyses: we observed no benefits other than a higher likelihood of quitting smoking in the short-term among the cases (data not shown).

DISCUSSION

In this prospective cohort study of 872 public sector employees who underwent an in-patient primary prevention programme to reduce the risk of early retirement on health grounds, no

Table III. Changes in the behaviour-related risk factors of the participants and controls

Type of change and risks before rehabilitation	Short-term follow-up			Long-term follow-up		
	Mean	95% CI	<i>p</i> -value	Mean	95% CI	<i>p</i> -value
Increase in weight, kg						
No obesity at baseline						
Participants (781)	1.80	1.51 to 2.10	0.010	3.45	3.08 to 3.83	0.005
Controls (2184)	1.36	1.20 to 1.52		2.83	2.61 to 3.05	
Obesity at baseline						
Participants (91)	-0.26	-1.98 to 1.46	0.651	0.03	-2.23 to 2.30	0.053
Controls (256)	0.18	-0.63 to 1.00		2.47	1.51 to 3.42	
Increase in physical activity, MET h/day						
No physical inactivity at baseline						
Participants (683)	-0.59	-0.89 to -0.29	0.835	-0.76	-1.10 to -0.41	0.275
Controls (1906)	-0.63	-0.81 to -0.44		-0.98	-1.16 to -0.79	
Physical inactivity at baseline						
Participants (189)	1.66	1.30 to 2.02	0.901	1.46	1.16 to 1.77	0.206
Controls (534)	1.69	1.45 to 1.92		1.71	1.48 to 1.95	
Increase in alcohol consumption, g/week						
No heavy drinking at baseline						
Participants (807)	9.46	5.31 to 13.6	0.402	8.08	3.44 to 12.7	0.145
Controls (2264)	11.68	8.55 to 15.8		12.11	9.35 to 14.87	
Heavy drinking at baseline						
Participants (65)	-37.34	-75.7 to 1.07	0.054	-86.35	-116 to -56.7	0.679
Controls (176)	-84.16	-112 to -56.1		-95.22	-125 to -65.4	

MET: metabolic equivalent of task; CI: confidence interval.

evidence of a favourable change in behaviour-related health risks was found when they were compared with their 2,440 propensity-score-matched controls, except for a temporary decrease in smoking. The aim of the programme was to prevent long-term work disability by helping the participants to adopt a healthier lifestyle composed of more regular and more comprehensive leisure-time physical activities, healthy dietary habits, a reduction in or cessation of smoking, and a reduction in alcohol consumption.

Our finding that obesity, leisure-time physical inactivity, and heavy drinking had not changed in comparison with the controls' respective behaviours after the rehabilitation is in line with the results of most previous studies on modifiable health risk behaviours, which also reported only minor or null effects of individual-based primary prevention in a low-risk population (2, 33–35). Improving participants' motivations and skills to endure more-intense leisure-time physical activity is the main tool used by VOMR teams to achieve changes in other health behaviours as well. However, as in previous studies of VOMR and other preventive rehabilitation programmes, VOMR seemed to be ineffective in improving the leisure-time physical activity of the participants (18, 19, 36). We are not aware of previous studies on the effectiveness of VOMR with respect to smoking, weight control, or alcohol consumption with which to compare these results. In relation to smoking, our results are in accordance with the results of a previous study reporting that multidisciplinary rehabilitation may be effective in helping participants to quit smoking (37). However, the improvement found in our study was not sustained in the long-term.

A failure to select participants at increased risk of early retirement is a potential explanation for the inability of VOMR to

contribute to health behavioural changes in our study. Indeed, VOMR is more likely to be granted to employees with few known risk factors (38). Failed selection may partially explain the inability of VOMR to affect the health behaviours of the participants. Because all of the 4 health risk behaviours studied have been shown to have an impact on work disability (11–16), the selection of low-risk participants to VOMR is potentially weakening the possibilities of the intervention to decrease the risk of work disability, the ultimate goal of VOMR.

The strengths of this prospective study are its large study population and its use of propensity score matching based on an explicit counterfactual framework. Although the distribution of the covariates used to derive the propensity score is the same for cases and controls, propensity-based matching might not remove bias due to unmeasured confounding when a strong selection bias exists (39). Information on patient prognosis and physician decision-making behaviours is typically not available in observational data-sets, such as ours, although the likelihood of being treated depends on clinical judgment and referral selection. Inability to take these factors into account is likely to result in an overestimation of the benefit in case lower-risk patients were selected for treatment and an underestimation if higher-risk participants entered rehabilitation. However, propensity score matching is likely to reduce such bias, given that the distribution of unmeasured prognostic factors are more likely to be similar when considering therapies with similar clinical indications and risk. Under such conditions, randomized clinical trials and observational studies have been shown to present the greatest similarities (40, 41). Because we focused on a low-risk population (22, 38) in terms of work disability, strong selection bias and major confounding from unmeasured factors seem unlikely. The study population consisted only of

full-time public sector employees from a developed European country, and therefore the generalizability of the findings may be reduced. Our study assessed the effectiveness of VOMR in achieving health behavioural changes only, and hence it is possible that this type of a preventive intervention programme affects other risk factors of early retirement on health grounds, such as levels of experienced work stress, anxiety, psychological distress, or self-rated health.

In conclusion, our results suggest that long-term change in behaviour-related risk factors, such as smoking, obesity, physical inactivity, and heavy drinking, may not be achieved by the vocationally oriented multidisciplinary rehabilitation programme widely used in Finland. Further research is needed to identify the reasons for the apparent ineffectiveness of this costly early intervention programme and to develop more effective preventive measures to reduce modifiable risks in working populations.

ACKNOWLEDGEMENTS

The Finnish Public Sector Study is supported by the Academy of Finland (projects 124271, 124322, and 132944) and the participating organizations.

REFERENCES

- World Health Organization (WHO). World health report 2002 – reducing risks, promoting healthy life. Geneva: WHO; 2002.
- Prochaska JJ, Spring B, Nigg CR. Multiple health behavior change research: an introduction and overview. *Prev Med* 2008; 46: 181–188.
- Kouvonen A, Kivimäki M, Vaananen A, Heponiemi T, Elovainio M, Ala-Mursula L, et al. Job strain and adverse health behaviors: the Finnish Public Sector Study. *J Occup Environ Med* 2007; 49: 68–74.
- Reeves MJ, Rafferty AP. Healthy lifestyle characteristics among adults in the United States, 2000. *Arch Intern Med* 2005; 165: 854–857.
- Noar SM, Chabot M, Zimmerman RS. Applying health behavior theory to multiple behavior change: considerations and approaches. *Prev Med* 2008; 46: 275–280.
- Nieuwenhuijsen ER, Zemper E, Miner KR, Epstein M. Health behavior change models and theories: contributions to rehabilitation. *Disabil Rehabil* 2006; 28: 245–256.
- Virtanen M, Kivimäki M, Kouvonen A, Elovainio M, Linna A, Oksanen T, et al. Average household income, crime, and smoking behaviour in a local area: the Finnish 10-Town study. *Soc Sci Med* 2007; 64: 1904–1913.
- Labarthe DR. Prevention of cardiovascular risk factors in the first place. *Prev Med* 1999; 29 (6 Pt 2): S72–S78.
- Goldstein MG, Whitlock EP, DePue J. Multiple behavioral risk factor interventions in primary care. Summary of research evidence. *Am J Prev Med* 2004; 27 Suppl 2: 61–79.
- Prochaska JO. Multiple health behavior research represents the future of preventive medicine. *Prev Med* 2008; 46: 281–285.
- Albertsen K, Lund T, Christensen KB, Kristensen TS, Villadsen E. Predictors of disability pension over a 10-year period for men and women. *Scand J Public Health* 2007; 35: 78–85.
- Kouvonen A, Kivimäki M, Virtanen M, Pentti J, Vahtera J. Work stress, smoking status, and smoking intensity: an observational study of 46,190 employees. *J Epidemiol Community Health* 2005; 59: 63–69.
- Skillgate E, Vingard E, Josephson M, Holm LW, Alfredsson L. Is smoking and alcohol consumption associated with long-term sick leave due to nonspecific back or neck pain among employees in the public sector? Results of a three-year follow-up cohort study. *J Rehabil Med* 2009; 41: 550–556.
- Robroek SJ, van den Berg TI, Plat JF, Burdorf A. The role of obesity and lifestyle behaviours in a productive workforce. *Occup Environ Med* 2010.
- Ferrie JE, Head J, Shipley MJ, Vahtera J, Marmot MG, Kivimäki M. BMI, obesity, and sickness absence in the Whitehall II study. *Obesity (Silver Spring)* 2007; 15: 1554–1564.
- Vahtera J, Poikolainen K, Kivimäki M, Ala-Mursula L, Pentti J. Alcohol intake and sickness absence: a curvilinear relation. *Am J Epidemiol* 2002; 156: 969–976.
- Arokoski JP, Juntunen M, Luikku J. Use of health-care services, work absenteeism, leisure-time physical activity, musculoskeletal symptoms and physical performance after vocationally oriented medical rehabilitation – description of the courses and a one-and-a-half-year follow-up study with farmers, loggers, police officers and hairdressers. *Int J Rehabil Res* 2002; 25: 119–131.
- Holopainen K, Nevala N, Kuronen P, Arokoski JP. Effects of vocationally oriented medical rehabilitation for aircraft maintenance personnel – a preliminary study of long-term effects with 5-year follow-up. *J Occup Rehabil* 2004; 14: 233–242.
- Leino P, Kivekäs J, Hänninen K. Effects of work-oriented fitness courses in lumberjacks with low back pain. *J Occup Rehabil* 1994; 4: 67–76.
- Nevala-Puranen N. Effects of occupationally-oriented rehabilitation on farmers' work techniques, musculoskeletal symptoms, and work ability. *J Occup Rehabil* 1996; 6: 191–200.
- Väyrynen S, Könönen U. Short and long-term effects of a training programme on work postures in rehabilitees: a pilot study of loggers suffering from back troubles. *International J Ind Ergon* 1991; 7: 103–109.
- Saltychev M, Laimi K, El-Metwally A, Oksanen T, Pentti J, Virtanen M, et al. Effectiveness of multidisciplinary primary prevention in decreasing the risk of work disability in low-risk population. *Scand J Work, Environ Health*. 2011 May 12 pii 3169 [Epub ahead of print].
- Pearl J. Understanding propensity scores. Causality: models, reasoning, and inference. Cambridge: Cambridge University Press; 2009.
- Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983; 70: 41–55.
- Social Insurance Institution of Finland. Statistical yearbook of the Social Insurance Institution, Finland, 2009. Helsinki: The Social Insurance Institution of Finland; 2010.
- Suoyrjö H, Oksanen T, Hinkka K, Kivimäki M, Klaukka T, Pentti J, et al. The effectiveness of vocationally oriented multidisciplinary intervention on sickness absence and early retirement among employees at risk: an observational study. *Occup Environ Med* 2009; 66: 235–242.
- Kujala UM, Kaprio J, Koskenvuo M. Modifiable risk factors as predictors of all-cause mortality: the roles of genetics and childhood environment. *Am J Epidemiol* 2002; 156: 985–993.
- Rehn N, Room R, Edwards G. Alcohol in the European region – consumption, harm and policies. Copenhagen: World Health Organization, Regional Office for Europe; 2001.
- Kouvonen A, Kivimäki M, Elovainio M, Väänänen A, De Vogli R, Heponiemi T, et al. Low organisational justice and heavy drinking: a prospective cohort study. *Occup Environ Med* 2008; 65: 44–45.
- Paljärvi T, Koskenvuo M, Poikolainen K, Kauhanen J, Sillanmäki L, Mäkelä P. Binge drinking and depressive symptoms: a 5-year population-based cohort study. *Addiction* 2009; 104: 1168–1178.
- Lipsitz SR, Kim K, Zhao L. Analysis of repeated categorical data using generalized estimating equations. *Stat Med* 1994; 13:

- 1149–1163.
32. Spiegelman D, Hertzmark E. Easy SAS calculations for risk or prevalence ratios and differences. *Am J Epidemiol* 2005; 162: 199–200.
 33. Ketola E, Sipila R, Makela M. Effectiveness of individual lifestyle interventions in reducing cardiovascular disease and risk factors. *Ann Med* 2000; 32: 239–251.
 34. Ebrahim S, Beswick A, Burke M, Davey Smith G. Multiple risk factor interventions for primary prevention of coronary heart disease. *Cochrane Database Syst Rev* 2006 (4): CD001561.
 35. Summerbell CD, Waters E, Edmunds LD, Kelly S, Brown T, Campbell KJ. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2005 (3): CD001871.
 36. Bernaards CM, Ariens GA, Knol DL, Hildebrandt VH. The effectiveness of a work style intervention and a lifestyle physical activity intervention on the recovery from neck and upper limb symptoms in computer workers. *Pain* 2007; 132: 142–153.
 37. Paone G, Serpilli M, Girardi E, Conti V, Principe R, Puglisi G, et al. The combination of a smoking cessation programme with rehabilitation increases stop-smoking rate. *J Rehabil Med* 2008; 40: 672–677.
 38. Saltychev M, Laimi K, Oksanen T, Pentti J, Virtanen M, Kivimäki M, et al. Predictive factors of future participation in rehabilitation in working population: the Finnish public sector study. *J Rehabil Med* 2011; 43: 404–410.
 39. Stukel TA, Fisher ES, Wennberg DE, Alter DA, Gottlieb DJ, Vermeulen MJ. Analysis of observational studies in the presence of treatment selection bias: effects of invasive cardiac management on AMI survival using propensity score and instrumental variable methods. *JAMA* 2007; 297: 278–285.
 40. Benson K, Hartz AJ. A comparison of observational studies and randomized, controlled trials. *N Engl J Med* 2000; 342: 1878–1886.
 41. Concato J, Shah N, Horwitz RI. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000; 342: 1887–1892.