



Cyclotomic shuffles

Oleg Ogievetsky, Varvara Petrova

► **To cite this version:**

Oleg Ogievetsky, Varvara Petrova. Cyclotomic shuffles. 12th International Workshop on Supersymmetries and Quantum Symmetries, Jul 2017, Dubna, Russia. pp.867-872, 10.1134/S1063779618050325 . hal-01777546

HAL Id: hal-01777546

<https://hal.archives-ouvertes.fr/hal-01777546>

Submitted on 24 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cyclotomic shuffles

Oleg Ogievetsky^{*1,2} and Varvara Petrova¹

¹Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France

²Kazan Federal University, Kremlevskaya 17, Kazan 420008, Russia

Abstract

Analogues of 1-shuffle elements for complex reflection groups of type $G(m, 1, n)$ are introduced. A geometric interpretation for $G(m, 1, n)$ in terms of rotational permutations of polygonal cards is given. We compute the eigenvalues, and their multiplicities, of the 1-shuffle element in the algebra of the group $G(m, 1, n)$. Considering shuffling as a random walk on the group $G(m, 1, n)$, we estimate the rate of convergence to randomness of the corresponding Markov chain. We report on the spectrum of the 1-shuffle analogue in the cyclotomic Hecke algebra $H(m, 1, n)$ for $m = 2$ and small n .

PACS number: 02.10.Hh

Introduction

In 1988, N. Wallach considered an element of the group algebra of the symmetric group S_n which is the sum of cycles $(12 \dots i)$ where i ranges from 1 to n . He discovered in [20] that the operator of the left multiplication by this element in the group algebra $\mathbb{Z}S_n$ is diagonalizable with eigenvalues

$$0, 1, 2, \dots, n - 2, n. \quad (1)$$

The sum of cycles $(12 \dots i)$ denoted $\mathbb{III}_{1, n-1}$ appears in different circumstances and is called 1-shuffle element. In particular, it describes all possible ways of removing the top card from the deck of n cards and inserting it back in the deck at a random position.

Investigating the repeated top-to-random shuffling as a random walk on S_n , P. Diaconis et al. [5] (see also R. Phatarfod [18]) found that the multiplicity of the eigenvalue i in (1) is equal to the number of permutations in S_n with i fixed points, explaining the absence of $n - 1$ in (1).

The q -deformation of the result of N. Wallach for the q -analogue ${}^q\mathbb{III}_{1, n-1}$ in the Hecke algebra $H_n(q)$ was proposed by G. Lusztig in [11]. He established that the spectrum of the operator $L_{\mathbb{III}_{1, n-1}}$ of the left multiplication by $\mathbb{III}_{1, n-1}$ consists of the q -numbers

$$q^{j-1}[j]_q := 1 + q^2 + q^4 + \dots + q^{2j-2}, \quad j = 0, 1, \dots, n - 2, n. \quad (2)$$

Later, A. Isaev and O. Ogievetsky considered shuffle elements $\mathbb{III}_{p, q}$ in the braid group ring $\mathbb{Z}B_n$. With the help of baxterized elements [9], they constructed additive and multiplicative analogues of $\mathbb{III}_{p, q}$ in Hecke and Birman-Murakami-Wenzl algebras. The multiplicities of the eigenvalues in (2) have been established therein by taking the trace of $L_{\mathbb{III}_{1, n-1}} : H_n(q) \rightarrow$

^{*}Also at Lebedev Institute, Moscow, Russia.

$H_n(q)$, using the fact that the q -numbers (2) are linearly independent over \mathbb{Z} as polynomials in q . For generic q , the multiplicities turn out to be the same as for the symmetric group.

In the present paper we propose polygonal analogues of cards that we call m -cards. We introduce elements ${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}$, which realise the analogues of top to random shuffling on m -cards. The elements ${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}$ belong to the group algebra of complex reflection groups of type $G(m, 1, n)$. We adopt the approach of [5] (for details see [6]) and of [9] to compute the spectrum and the multiplicities of the eigenvalues of $L_{{}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}}$. The obtained result for the multiplicities is expressed in terms of the so-called m -derangements numbers. Asymptotic convergence to randomness in the shuffling the m -cards is briefly analysed. We give a preliminary result on the spectrum of $L_{{}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}}$ in the cyclotomic Hecke algebra $H(m, 1, n)$, which is a deformation of the group algebra of the complex reflection group.

1 Complex reflection groups $G(m, 1, n)$

A finite complex reflection group is a finite subgroup of $\mathrm{GL}_n(\mathbb{C})$ generated by complex reflections, that is, elements $\tau \in \mathrm{GL}_n(\mathbb{C})$ of finite order such that $\mathrm{Ker}(\tau - id)$ is a hyperplane. The finite complex reflection groups have been classified by Shephard and Todd (1954) into an infinite family of groups $G(m, p, n)$ where m, p, n are positive integers such that p divides m , and 34 exceptional groups. The group $G(m, 1, n) \subset \mathrm{GL}_n(\mathbb{C})$ is formed by all matrices with exactly one nonzero entry in each row and column; nonzero entries are m -th roots of unity; it is generated by the elements t, s_1, \dots, s_{n-1} with the defining relations:

$$\begin{cases} s_{i+1}s_i s_{i+1} = s_i s_{i+1} s_i & \text{for } i = 1, \dots, n-2, \\ s_i s_j = s_j s_i & \text{for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 1, \\ s_i^2 = 1 & \text{for } i = 1, \dots, n-1 \end{cases} \quad (3)$$

and

$$\begin{cases} s_1 t s_1 t = t s_1 t s_1, \\ s_i t = t s_i & \text{for } i > 1, \\ t^m = 1, \end{cases} \quad (4)$$

or equivalently by $t_1, \dots, t_n, s_1, \dots, s_{n-1}$ with the defining relations (3) and

$$\begin{cases} t_i^m = 1 & \text{for } i = 1, \dots, n, \\ t_i t_j = t_j t_i & \text{for } i, j = 1, \dots, n, \\ s_i t_j = t_{\pi_i(j)} s_i & \text{for } i = 1, \dots, n-1, j = 1, \dots, n. \end{cases} \quad (5)$$

Here $t_1 := t$, $t_{i+1} := s_i t_i s_i$ for $i = 1, \dots, n-1$ and π_i is the transposition $(i, i+1)$.

In particular, the group $G(1, 1, n)$ is isomorphic to the symmetric group S_n and the group $G(2, 1, n)$ to the hyperoctahedral group B_n , the Coxeter group of type B .

The group $G(m, 1, n)$ also admits the following description, see e.g. [16]. Let C_m be the cyclic group of order m with a generator γ . Let E be the set $\{(x, \rho) \mid x \in \mathbf{n}, \rho \in C_m\}$ where $\mathbf{n} = \{1, 2, \dots, n\}$. Define the action of C_m on E by:

$$\rho \cdot (x, \rho') = (x, \rho \rho'), \quad \forall \rho, \rho' \in C_m \text{ and } \forall x \in \mathbf{n}.$$

Let $\mathrm{Perm}(E)$ be the group of all permutations of the set E . Denote by $\mathrm{Perm}^0(E)$ the subgroup of $\mathrm{Perm}(E)$ consisting of elements $\pi \in \mathrm{Perm}(E)$ such that:

$$\pi(x, \rho) = \rho \cdot \pi(x, 1), \quad \forall \rho \in C_m \text{ and } \forall x \in \mathbf{n}.$$

The group $G(m, 1, n)$ is isomorphic to $\text{Perm}^0(E)$ via the map ϕ such that:

$$\phi(t)(x, 1) = \begin{cases} (x, \gamma) & \text{if } x = 1, \\ (x, 1) & \text{otherwise;} \end{cases}$$

$$\phi(s_i)(x, 1) = (\pi_i(x), 1) \quad \text{for } x \in \mathbf{n} \text{ and } i = 1, \dots, n-1.$$

We propose a definition of *polygonal* cards (or *m-cards*) and interpret $\text{Perm}^0(E)$, and thus $G(m, 1, n)$, as the group of “rotational” permutations of polygonal cards.

Definition. An *n-deck* \mathfrak{X} of *m-cards* is an *n-tuple* $\mathfrak{X} = ((x_1, \gamma^{k_1}), (x_2, \gamma^{k_2}), \dots, (x_n, \gamma^{k_n}))$ where (x_1, \dots, x_n) is a permutation of $(1, \dots, n)$ and $\gamma^{k_j} \in C_m$, $j \in \mathbf{n}$.

Let M be an *m-gon* with a distinguished vertex. We interpret $(x, 1)$ as an *m-gon* M , on which the number x is printed, with the distinguished vertex pointing to the north. The *m-card* (x, γ^k) is the card $(x, 1)$ rotated clockwise by an angle $\frac{2\pi}{m}k$ in the plane of M .

Abstractly, a 1-card (a usual card) is a point and a 2-card is a segment with two vertices. In the context of card shuffling, we shall slightly transgress this picture and conceive of 1 and 2-cards as the rectangular cards, as illustrated in figure 1.




FIG. 1: *m-cards*

Let $((1, 1), \dots, (n, 1))$ be an initial deck. We identify a deck \mathfrak{X}

with a “rotational” permutation of the form: $g = \begin{pmatrix} (1, 1) & \dots & (n, 1) \\ (x_1, \gamma^{k_1}) & \dots & (x_n, \gamma^{k_n}) \end{pmatrix}$, meaning that the *j*-th position in the deck gets occupied by the x_j -th card rotated clockwise by an angle $\frac{2\pi k_j}{m}$. The group $\text{Perm}^0(E)$ describes $G(m, 1, n)$ as the group of all possible rotational permutations of the *n-deck* of *m-cards*. In particular, for $i = 1, \dots, n-1$, the generator s_i permutes the *i*-th and $(i+1)$ -st *m-cards* without any turn. For $i = 1, \dots, n$, the element t_i rotates the *i*-th *m-card* clockwise by an angle $\frac{2\pi}{m}$.

Schreier coset graph for the chain of groups $G(m, 1, n)$

Given a group G with a finite generating set \mathcal{G} , and a subgroup $W \subset G$ of finite index, the (right) Schreier coset graph of $(G, W; \mathcal{G})$ is an edge-labeled graph whose vertices are the right W -cosets and edges are of the form (Wg, Wgx) where $x \in \mathcal{G}$ and $g \in G$; the edge (Wg, Wgx) has label x . The graph depicts the action of G on cosets of W by right multiplication and is equivalent to the coset table obtained by the Todd-Coxeter algorithm [4].

Figure 2 is the Schreier coset graph of $(G(m, 1, n), W; \mathcal{G})$ where $\mathcal{G} = \{t, s_1, \dots, s_{n-1}\}$ and W is generated by the elements t, s_1, \dots, s_{n-2} . An unoriented edge represents a pair of edges with opposite directions. For simplicity, only non-trivial actions are drawn (if $Wgx = Wg$ then (Wg, Wgx) is a loop and we do not draw it). At each vertex of the oriented *m-gon* in the middle of the figure starts a tail with $n-1$ edges. All tails are identically (edges-)labelled. The graph provides a “normal form with respect to W ” [16].

Proposition 1. Any element $x \in G(m, 1, n)$ can be uniquely written in the form: $x = ws_{n-1} \dots s_1 t^\alpha s_1 \dots s_j$ with $j \in \{0, \dots, n-1\}$, $\alpha \in \{0, \dots, m-1\}$ and $w \in W \simeq G(m, 1, n-1)$ (by convention, the empty product is equal to 1).




FIG. 2: Schreier coset graph for $G(m, 1, n)$ with respect to W [16].

Starting with a normal form (powers of t) for $G(m, 1, 1) \simeq C_m$, we build recursively [16] the global normal form for elements of $G(m, 1, n)$ for any n . We have an ascending chain:

$$G(m, 1, 0) \subset G(m, 1, 1) \subset G(m, 1, 2) \subset \dots$$

In the computational context, in order to build an algorithm that takes as input a word in generators of $G(m, 1, n+1)$ and their inverses, and returns its normal form one only needs to know the normal form of the words $\psi_{kj}^{(n)} x = s_n \cdots s_1 t^k s_1 \cdots s_j x$ for all generators x of $G(m, 1, n+1)$, $k = 0, \dots, m-1$ and $j = 0, \dots, n$. We established the following recursive list of rewriting rules (the equalities are to be understood as “replace the left hand side by the right hand side”).

Proposition 2. *The rewriting rules for $G(m, 1, n+1)$ with $n > 1, m \geq 1$ include (additionally to the defining relations (3)-(4)) the rewriting rules for $G(m, 1, n)$ and the following:*

$$\begin{cases} s_n s_{n-1} \cdots s_j s_n = s_{n-1} s_n s_{n-1} \cdots s_j, & j = 1, \dots, n-2, \\ \psi_{kj}^{(n)} s_n = s_{n-1} \psi_{kj}^{(n)}, & k = 0, \dots, m-1, j = 0, 1, \dots, n-2. \end{cases}$$

For $G(m, 1, 2)$ the rewriting rules (in addition to the defining relations) are:

$$s_1 t^k s_1 t = t s_1 t^k s_1, \quad \forall k = 2, \dots, m-1.$$

2 Shuffle elements in $G(m, 1, n)$ group algebra

We start by recalling that a (p, q) -shuffle, where p and q are non-negative integers, is a permutation $\sigma \in S_{p+q}$ of the set $\{1, \dots, p+q\}$ such that $\sigma(1) < \sigma(2) < \cdots < \sigma(p)$ and $\sigma(p+1) < \cdots < \sigma(p+q)$. When $p = 1$ or $q = 1$, the (p, q) -shuffle is called 1-shuffle. The sum over all (p, q) -shuffles with p, q fixed is called shuffle element and is denoted by $\mathbb{I}\mathbb{I}\mathbb{I}_{p,q}$.

Let $w^{\uparrow \ell}$ denote the image of $w \in S_n$ under the homomorphism $S_n \rightarrow S_{n+\ell}$ sending $\pi_i = (i, i+1)$ to $\pi_{i+\ell}$, $i = 1, \dots, n-1$. Shuffles $\mathbb{I}\mathbb{I}\mathbb{I}_{p,q}$ can be defined inductively by any of the recurrence relations (analogues of the Pascal rule):

$$\begin{aligned} \mathbb{I}\mathbb{I}\mathbb{I}_{p,q} &= \mathbb{I}\mathbb{I}\mathbb{I}_{p-1,q} + \mathbb{I}\mathbb{I}\mathbb{I}_{p,q-1} \pi_{p+q-1} \cdots \pi_q \\ &= \mathbb{I}\mathbb{I}\mathbb{I}_{p,q-1}^{\uparrow 1} + \mathbb{I}\mathbb{I}\mathbb{I}_{p-1,q}^{\uparrow 1} \pi_1 \cdots \pi_q, \end{aligned}$$

with $\mathbb{I}\mathbb{I}\mathbb{I}_{0,q} = \mathbb{I}\mathbb{I}\mathbb{I}_{q,0} = 1$ for any $q \geq 0$. Let $\tilde{\Sigma}_n := \sum_{\sigma \in S_n} \sigma$ be the symmetrizer, $\sigma \tilde{\Sigma}_n = \tilde{\Sigma}_n, \forall \sigma \in S_n$. Then $\tilde{\Sigma}_{q+p} = \mathbb{I}\mathbb{I}\mathbb{I}_{p,q} \tilde{\Sigma}_q \tilde{\Sigma}_p^{\uparrow q}$, in particular,

$$\tilde{\Sigma}_n = \mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} \tilde{\Sigma}_{n-1}. \quad (6)$$

By their combinatorial structure and relationship with the structure of products, shuffle elements appear in many constructions in homotopy theory and higher category theory. In particular, they are involved in the product of simplices [12], in the study of multiple polylogarithms [2,3], in the bialgebras of type 1 [13], in the construction of the Hopf algebra structure on the tensor algebra $T(V)$ of a vector space V [7], in the construction of the standard complex for quantum Lie algebras [8], etc.

In the present paper we consider the analogues of the 1-shuffle elements $\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}$ in $\mathbb{Z}G(m, 1, n)$ of the form

$${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} := \mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} \cdot {}^{(m)}\mathcal{T}_n,$$

where $\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} = 1 + s_{n-1} + s_{n-2}s_{n-1} + \dots + s_1 \cdots s_{n-1}$ and ${}^{(m)}\mathcal{T}_n := 1 + t_n + t_n^2 + \dots + t_n^{m-1}$. The element ${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}$ describes all possible ways of rotating the n -th m -card and inserting it at random position in the deck; we found that ${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}$ satisfies:

$${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}^2 = {}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} \left(m + {}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-2} \right). \quad (7)$$

Let ${}^{(m)}\tilde{\Sigma}_n$ denote the sum of all elements of the group $G(m, 1, n)$. The recurrent relation similar to (6) holds:

$${}^{(m)}\tilde{\Sigma}_n = {}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} \cdot {}^{(m)}\tilde{\Sigma}_{n-1}.$$

Making use of the relation (7) we prove by induction on n that

$$\prod_{i=0}^{n-1} \left({}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} - im \right) = {}^{(m)}\tilde{\Sigma}_n. \quad (8)$$

Since ${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} \cdot {}^{(m)}\tilde{\Sigma}_n = mn \cdot {}^{(m)}\tilde{\Sigma}_n$, we obtain, multiplying (8) by $\left({}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} - nm \right)$,

$$\prod_{i=0}^n \left({}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} - im \right) = 0. \quad (9)$$

For a given $x \in \mathbb{Z}G(m, 1, n)$, let L_x be the map $\mathbb{Z}G(m, 1, n) \rightarrow \mathbb{Z}G(m, 1, n)$, $y \mapsto xy$. We calculate the multiplicities of the eigenvalues of $L_{{}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1}}$ following the approach Garsia and Wallach developed for the symmetric group in [6]. Let α and β be two words in some alphabet. We denote by $\alpha \uplus \beta$ the formal sum of all words that can be obtained by shuffling α and β as it is done with two decks of cards. We define for $0 \leq a \leq n$:

$$B_a = \sum_{\alpha \in G(m, 1, a)} \alpha \uplus \beta_{a,n},$$

where α is viewed as an a -deck of m -cards and $\beta_{a,n}$ is the deck $((a+1, 1), (a+2, 1), \dots, (n, 1))$. In particular, $B_0 = 1$. Since $B_1 = {}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{n-1,1} = w \cdot {}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{1,n-1} \cdot w^{-1}$ where w is the longest element of the symmetric group $S_n = G(1, 1, n)$, it has the same spectrum as ${}^{(m)}\mathbb{I}\mathbb{I}\mathbb{I}_{n-1,1}$. The element B_a is composed of $\frac{n!}{(n-a)!} m^a$ decks that may be grouped according to the number and the angle of the top card. We get $am + 1$ groups labelled by $(1, \rho), (2, \rho), \dots, (a, \rho)$ and $(a+1, 1)$, with $\rho \in C_m$. Upon multiplication by B_1 , each group (k, ρ) , $1 \leq k \leq a$, $\rho \in C_m$, yields a term B_a , and the group $(a+1, 1)$ gives the term B_{a+1} . Thus, $B_1 B_a = am B_a + B_{a+1}$ or $B_{a+1} = (B_1 - am) B_a$, $0 \leq a \leq n-1$. Iterating, we obtain

$$B_a = B_1 (B_1 - m) (B_1 - 2m) \cdots (B_1 - (a-1)m), \quad 0 \leq a \leq n. \quad (10)$$

Applying the Stirling inversion to (10), we find, for $0 \leq k \leq n$:

$$B_1^k = m^k \sum_{i=0}^n i^k \mathbf{E}_i, \quad \text{where } \mathbf{E}_i = \frac{1}{i!} \sum_{a=i}^n \frac{(-1)^{a-i}}{(a-i)! m^a} B_a. \quad (11)$$

Substituting into the identity $B_1 B_1^k = B_1^{k+1}$ the expressions (11) for B_1^k and B_1^{k+1} we obtain:

$$\sum_{i=0}^n i^k (B_1 \mathbf{E}_i - im \mathbf{E}_i) = 0, \quad k = 0, \dots, n.$$

Since the matrix $M_{ik} := \{i^k\}_{i,k=0}^n$ is invertible, we get $B_1 \mathbf{E}_i = im \mathbf{E}_i$, $i = 0, \dots, n$.

Let \mathcal{B} be the \mathbb{Q} -subalgebra of $\mathbb{Q}G(m, 1, n)$, generated by B_1 . By (10), \mathcal{B} is a linear span of B_a , $a = 0, \dots, n$. By (11), \mathcal{B} is a linear span of \mathbf{E}_a , $a = 0, \dots, n$. Therefore \mathbf{E}_i and \mathbf{E}_j commute. We have

$$B_1 \mathbf{E}_i \mathbf{E}_j = m i \mathbf{E}_i \mathbf{E}_j = m j \mathbf{E}_i \mathbf{E}_j, \quad 0 \leq i, j \leq n,$$

whence $\mathbf{E}_i \mathbf{E}_j = 0$, $0 \leq i \neq j \leq n$. On the other hand, the equality $\sum_{i=0}^n \mathbf{E}_i = 1$ obtained from (11) for $k = 0$ implies that $\mathbf{E}_i^2 = \mathbf{E}_i$, $i = 0, \dots, n$. Operators \mathbf{E}_i , $i = 0, \dots, n$, are therefore the orthogonal projections onto the eigenspaces of B_1 . The multiplicity of the eigenvalue im in (9) is thus given by the trace of the matrix $L_{\mathbf{E}_i}$. Since the trace of any element except the identity vanishes in the regular representation, we get:

$$\text{tr}(L_{\mathbf{E}_i}) = \frac{1}{i!} \sum_{a=i}^n \frac{(-1)^{a-i}}{(a-i)! m^a} \text{tr}(L_{B_a}) = \binom{n}{i} m^{n-i} (n-i)! \sum_{a=0}^{n-i} \frac{(-1)^a}{m^a a!}.$$

The obtained multiplicities have the combinatorial meaning. We shall say that an element $g \in G(m, 1, n)$ has i fixed points if it fixes i m -cards. For a given n the m -derangement number is the number $d_{m,n}$ of elements in $G(m, 1, n)$ with no fixed points. We have

$$d_{m,n} = m^n n! \sum_{k=0}^n \frac{(-1)^k}{k! m^k}.$$

We summarize the obtained results.

Theorem 3. *The operator $L_{(m)\text{III}_{1,n-1}} : \mathbb{Z}G(m, 1, n) \rightarrow \mathbb{Z}G(m, 1, n)$ is diagonalizable with eigenvalues im , $i = 0, 1, \dots, n-1, n$. The multiplicity of the eigenvalue im equals $\binom{n}{i} d_{m,n-i}$, the number of elements of $G(m, 1, n)$ with i fixed points.*

Remark. The spectrum of $L_{(1)\text{III}_{1,n-1}}$ is $\{0, 1, \dots, n-2, n\}$, because $d_{1,1} = 0$; the spectrum has a gap at $n-1$. For $m > 1$, the gap disappears. This result proves important in estimating how close k repeated shuffles, randomly picked among the set of elements of $G(m, 1, n)$ whose sum is $(m)\text{III}_{1,n-1}$, get the n -deck of m -cards to being randomized. Indeed, under a suitable normalization, the matrix of $L_{(m)\text{III}_{1,n-1}}$ is a matrix P of transition probabilities. As it belongs to the convex envelope of the set of permutation matrices, the matrix P is doubly stochastic. The underlying Markov chain is ergodic and its unique stationary probability distribution is uniform. Using the spectral representation formula for the k -step transition matrix P^k [10,19], one can show that the convergence to the stationarity occurs exponentially fast with the rate of convergence governed by the second largest eigenvalue: there exists a constant $\alpha > 0$ such that:

$$\forall k \geq 1, \quad \left| P_{ij}^k - \frac{1}{m^n n!} \right| \leq \begin{cases} \alpha \left(\frac{n-2}{n} \right)^k & \text{if } m = 1, \\ \alpha \left(\frac{n-1}{n} \right)^k & \text{if } m > 1. \end{cases}$$

Cyclotomic algebras. The cyclotomic Hecke algebra $H(m, 1, n)$ is the associative algebra over $\mathbb{C} [q^\pm, v_1^\pm, \dots, v_m^\pm]$ generated by $\tau, \sigma_1, \dots, \sigma_{n-1}$ with the defining relations:

$$\left\{ \begin{array}{ll} \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} & \text{for all } i = 1, \dots, n-2, \\ \sigma_i \sigma_j = \sigma_j \sigma_i & \text{for all } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 1, \\ \tau \sigma_1 \tau \sigma_1 = \sigma_1 \tau \sigma_1 \tau & \\ \tau \sigma_i = \sigma_i \tau & \text{for } i > 1, \\ \sigma_i^2 = (q - q^{-1}) \sigma_i + 1 & \text{for all } i = 1, \dots, n-1, \\ (\tau - v_1) \cdots (\tau - v_m) = 0. & \end{array} \right.$$

The algebra $H(m, 1, n)$ is a deformation of the complex reflection group algebra $\mathbb{C}G(m, 1, n)$. The representation theory of $H(m, 1, n)$ has been constructed in [1]; the representation theory of $G(m, 1, n)$ and $H(m, 1, n)$ was reobtained in [14, 15] from the analysis of the spectrum of the so called Jucys-Murphy elements, following the approach of [17].

Consider the algebra $H(2, 1, n)$ with parameters $v_1 = p, v_2 = -p^{-1}$. Let elements τ_i be recursively defined by, cf (5):

$$\tau_i = \sigma_{i-1}^{-1} \tau_{i-1} \sigma_{i-1}, \quad i = 2, \dots, n,$$

with $\tau_1 = \tau$. We propose the following analogue of ${}^{(2)}\mathbb{III}_{1, n-1}$ for the algebra $H(2, 1, n)$:

$${}^{(2)}\mathbb{III}_{1, n-1; q, p} := \left(1 + q\sigma_{n-1} + q^2\sigma_{n-2}\sigma_{n-1} + \cdots + q^{n-1}\sigma_1 \cdots \sigma_{n-1} \right) (1 + p\tau_n). \quad (12)$$

Our calculations of the spectrum of (12) in various representations of $H(2, 1, n)$ for small n lead to the following conjecture.

Conjecture. *The spectrum of the operator of the left multiplication by ${}^{(2)}\mathbb{III}_{1, n-1; q, p}$ in the algebra $H(2, 1, n)$ consists of the values:*

$$\left(p^2 + q^{2(j-\ell)} \right) q^{\ell-1} [\ell]_q,$$

where $q^{\ell-1} [\ell]_q$ are the q -numbers defined in (2), $\ell = 0, 1, \dots, n$ and $j = \ell, \ell + 1, \dots, n - 2, n$.

Remark. We numerically observed (for $n \leq 5$) that the eigenvalues $0, (p^2 + q^2)q^{n-2} [n-1]_q$ and $(p^2 + 1)q^{n-1} [n]_q$ have the same multiplicities as their respective counterparts $0, 2(n-1), 2n$ in the classical situation, for the group $G(m, 1, n)$. This is not the case for the other eigenvalues.

Acknowledgements. The work of O. O. was supported by the Program of Competitive Growth of Kazan Federal University and by the grant RFBR 17-01-00585. The work of V. P. has been carried out thanks to the support of the A*MIDEX grant (N^o ANR-11-IDEX-0001-02) funded by the French Government “Investissements d’Avenir” program.

References

- [1] Ariki S. and Koike K., A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr S_n$ and construction of its irreducible representations // Adv. in Math. 1994. V. 106 (2). P. 216.
- [2] Borwein J. M., Bradley D. M., Broadhurst D. J., and Lisoněk P., Combinatorial aspects of multiple zeta values // electronic j. of combinatorics 1998. V. 5 (1). P. R38.

- [3] *Bowman D. and Bradley D.*, Multiple Polylogarithms: A Brief Survey // Contemporary Mathematics 2001. V. 291. P. 71.
- [4] *Coxeter H. and Todd J.*, A practical method for enumerating cosets of a finite abstract group // Proc. Edinburgh Math. Soc. 1936. V. 5. P. 26.
- [5] *Diaconis P., Fill J. A., and Pitman J.*, Analysis of top to random shuffles // Combinatorics, Probability and Computing 1992. V. 1. P. 135.
- [6] *Garsia A. and Wallach N.*, Qsym over Sym is free // Journal of Combinatorial Theory, Series A 2003. V. 104. P. 217.
- [7] *Grapperon T. and Ogievetsky O. V.*, Braidings of tensor spaces // Letters in Mathematical Physics 2012. V. 100. P.17.
- [8] *Isaev A. P. and Ogievetsky O. V.*, BRST operator for quantum Lie algebras: explicit formula // International Journal of Modern Physics A 2004. V. 19. P. 240.
- [9] *Isaev A. P. and Ogievetsky O. V.*, Braids, shuffles and symmetrizers // J. Phys. A: Math. Theor. 2009. V. 42. P. 1.
- [10] *Lorek P.*, Speed of convergence to stationarity for stochastically monotone Markov chains // PhD thesis, Mathematical Institute University of Wroclaw, 2007.
- [11] *Lusztig G.*, A q-analogue of an identity of N. Wallach // Studies in Lie Theory 2006. P. 405.
- [12] *Mac Lane S.*, Homology, Classics in Mathematics 114, Springer, 1963.
- [13] *Nichols W. D.*, Bialgebras of type one // Communications in Algebra 1978. V. 6. P. 1521.
- [14] *Ogievetsky O. V. and Poulain d'Andecy L.*, On representations of cyclotomic Hecke algebras // Modern Physics Letters A 2011. V. 26 (11). P. 795.
- [15] *Ogievetsky O. V. and Poulain d'Andecy L.*, An inductive approach to representations of complex reflection groups $G(m, 1, n)$ // Theoretical and Mathematical Physics 2013. V. 174 (1). P. 95.
- [16] *Ogievetsky O. V. and Poulain d'Andecy L.*, Induced representations and traces for chains of affine and cyclotomic Hecke algebras // Journal of Geometry and Physics 2015. V. 87. P. 354.
- [17] *Okounkov A. and Vershik A.*, A new approach to representation theory of symmetric groups II // Selecta Math (New series) 1996. V. 2 (4). P. 581.
- [18] *Phatarfod R. M.*, On the matrix occurring in a linear search problem // Journal of applied probability 1991. V. 28. P. 336.
- [19] *Seneta E.*, Non-negative Matrices and Markov Chains, Springer, 2006.
- [20] *Wallach N. R.*, Lie algebra cohomology and holomorphic continuation of generalized Jacquet integrals // Representations of Lie Groups, Kyoto, Hiroshima 1986. V. 14. P. 123.